

## The determination of gibberellic acid effects on seed germination of *Echinacea purpurea* (L.) Moench under salt stress

Ruvejde Tuncturk<sup>1,\*</sup> 

Serhat Kipcak<sup>1</sup> 

Mahdi Ghiyasi<sup>2</sup> 

Murat Tuncturk<sup>1</sup> 

<sup>1</sup>Van Yuzuncu Yil University, Faculty of Agriculture, Department of Field Crops, Van-Turkey

<sup>2</sup>Urmia University, Faculty of Agriculture, Department of Field Crops, Urmia-Iran

\*Corresponding Author: ruveydetuncturk@yyu.edu.tr

### Abstract

The research was conducted to determine the effect of gibberellic acid pre-treatments on germination of *Echinacea purpurea* seeds under salt stress. In the study, three different gibberellic acid concentrations (0, 100, 200 and 300 ppm) were pre-treated. Seeds planted in petri dishes were left to germinate after salt applications at 0, 50 and 100 mM (NaCl) concentrations. Responses of *echinacea* to priming and salt stress treatments were observed on the bases of some growth and viability (radicle and hypocotyl length, radicle fresh and dry weight, hypocotyl fresh and dry weight, germination power, germination speed, mean germination time, germination and sensitivity index) parameters.

According to the result of the research; when the salt concentrations increase, the germination and growth parameters of *echinacea* seeds were inhibited. It was conducted determined that increasing doses of gibberellic acid pre-treatments had significant and positive effects on the germination and growth parameters of *echinacea* seeds under salt stress conditions. In physiological enhancement of *echinacea* seeds, the best results were obtained from the 300 ppm GA<sub>3</sub>+0 mM (control) salt combination.

**Keywords:** Gibberellic acid, Salt, *Echinacea* (*Echinacea purpurea* L.), Germination

Received: 20 March 2018



Accepted: 27 May 2019



Published: 28 June 2019

### Introduction

*Echinacea* which is a medicinal and aromatic plant, has been used for traditional medicine because of its pharmacological properties in many countries. The first reports of the use of the plants belonging to *Echinacea* genus date to the beginning of the twentieth century, in the Ethnobotanical studies made by WHO, documented uses as a primitive antibiotic use against snakebites to heal wounds (Anonymus, 1999). Although medicinal plants have been used for thousands of years in every culture of the world, serious scientific researches supporting their therapeutic value have begun around 1960 (Lopez and Shepard, 2007). *Echinacea* is an indigenous herbal plant genus of North America and occupies an important place among medicinal plants due to its immunostimulatory properties against respiratory ailments. The genus *Echinacea* is one of the prominent genera of the plants that are utilized in medicinal preparations and drugs (Ahmad et al., 2017). *Echinacea* plant which is

perennials has 9 species. Being commonly used in traditional medicine, *E. purpurea*, *E. pallida* and *E. angustifolia* have been the most investigated of these species (Ivanova et al., 2014). *Echinacea* which belongs to the family *Asteraceae* (*Compositae*) is a plant commonly found in the world. *Echinacea purpurea* (L.) Moench has been traditionally also used for the treatment of toothache, snake bite, bowel pain, skin disorders, chronic arthritis, seizure and cancer (Grimm and Muller, 1999). Salinity is considered to be one of the environmental stresses that reduce the growth and efficiency of most glycophytic plants worldwide. Induces both osmotic and ionic stresses leading to deterioration of many physiological and biochemical processes including salinity, water relations, ionic homeostasis, gas exchange and mineral nutrition (Parida and Das, 2005; Munns and Tester, 2008). Gibberellins are involved in the stimulation of enzymes involved in seed germination.

### Cite this article as:


Tuncturk, R., Kipcak, S., Ghiyasi, M., Tuncturk, M. (2019). The determination of gibberellic acid effects on seed germination of *Echinacea purpurea* (L.) Moench under salt stress. Int. J. Agric. Environ. Food Sci., 3(2), 101-105. DOI: <https://dx.doi.org/10.31015/jaefs.2019.2.8>

Year: 2019 Volume: 3 Issue: 2 (June) Pages: 101-105

Available online at : <http://www.jaefs.com> - <http://dergipark.gov.tr/jaefs>

Copyright © 2019 International Journal of Agriculture, Environment and Food Sciences (Int. J. Agric. Environ. Food Sci.)

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC-by 4.0) License



Gibberellic acid (GA<sub>3</sub>) is one of the hormones proposed to control primary dormancy by inducing germination. Cavusoglu et al. (2007) reported that GA<sub>3</sub> was the most effective salt stress reducing agent among plant growth regulators used in the research.

This study was carried out to detect the effects of salt doses with pre-applications in various levels of Gibberellic acid (GA<sub>3</sub>) on germination of *Echinacea purpurea* L. seeds.

### Materials and Methods

In order to evaluate the effect of seed priming on the germination and seedlings' growth traits of *Echinacea purpurea* L. under different salinity stress levels, a factorial experiment was conducted in completely randomized design (CRD) with four replications, at the laboratory of Seed Science and Technology in the Department of Field Crops, Faculty of Agriculture, Van Yuzuncu Yil University, Turkey in 2018. Three salt levels (0, 50, 100 NaCl mM) and four gibberellic acid (GA<sub>3</sub>) levels (0, 100, 200 and 300 ppm) were used in the experiments. Seeds were rinsed with water and distilled water, then their surface sterilized in 20% sodium hypochlorite (vol/vol) for 5 minutes, followed by 70% alcohol (vol/vol) treatment for 1 minutes, after then, were thoroughly rinsed with sterile deionized water before priming.

The seed lots were imbibed within different doses gibberellic acid (GA<sub>3</sub>) in priming treatments. Treated seed lots with GA<sub>3</sub> were kept in darkness in an incubator at 25 ± 0.5 °C for 24 h. Seeds, removed from the solutions (GA<sub>3</sub>) were air dried for 24 hours at 25 °C to reach the original moisture content (12-13 %) and then proceed for germination test as suggested by ISTA. Unprimed seeds were used as control. Treated seeds were placed in 9 cm diameter sterile petri dishes with filter paper and 20 seeds were placed in each petri dish and 5ml of salt solution was added dish to all petri except for control treatments, to the control treatments was given distilled water (5 ml), then petri dishes were placed in a totally dark incubator which stable at temperature (25±0.5°C). As a final step, after 14 days all parameters were observed, measured and recorded. The root and shoot vigors were calculated as the sum of total root length (cm) and shoot length (cm) of all the seedlings of a replicate divided by the number of seedlings. Fresh root and shoot were then placed in a hot air oven (70 °C for 24 hours) to dry (Anonymus, 2012). Root and shoot dry masses were weighed germination rate, germination power, germination index, mean germination time and sensitivity index were calculated with the following formulas. It was considered that number of germinated seeds on 7<sup>th</sup> day as "germination rate" and number of germinated seeds on 14<sup>th</sup> day as "germination power". Germination rate (GR) was calculated using the Equation 1 (Akinci and Caliskan, 2010).

GR = Total seeds germinated after day 14/Total number of planted seeds (Equation 1)

Germination index (GI) was calculated using the Equation 2 (Wang et al., 2004).

$$GI = \sum(G_i/T_t) \quad (\text{Equation 2})$$

GI: Germination index; G<sub>i</sub>: i. Days germinated seed rate; T<sub>t</sub>: count day

Mean germination time (MGT) was calculated using the Equation 3 (Ellis and Roberts, 1980).

$$MGT = \sum(fx) / \sum f \quad (\text{Equation 3})$$

f: Number of seeds germinated x: germination day

Sensitivity index (SI) was calculated using the Equation 4

(Foolad and Lin, 1997).

SI = MGT in the salt application / MGT in the control application (Equation 4)

Data were analyzed statistically by using analysis of variance with COSTAT (version 6.3) software. The variance analysis of data was performed (ANOVA) and the multiple comparison of the means was made according to the Duncan test (Duzgunes et al., 1987).

### Results and Discussion

According to the research results, salinity and seed priming showed a significant effect on all parameters. Only, the effect of gibberellic acid on the radicle length wasn't found insignificant. Furthermore, salinity stress x seed priming interaction that it was found important with regard to parameters such as dry radicle weight (mg) and sensitivity index (%). Salinity stress x seed priming Interaction in terms of dry radicle weight seem seeds primed with GA<sub>3</sub> unlike untreated seeds had the highest germination value at all salinity levels. While the highest sensitivity index was obtained as 1.54 % from seeds primed with 100 ppm GA<sub>3</sub>, in 100 mM salt stress conditions, the highest dry radicle weight was reported as 5.70 mg from seeds primed with 300 ppm GA<sub>3</sub> in non-stress (0 mM NaCl) conditions.

Highest germination power, germination rate and germination index were determined as 92.50, 66.25 and 14.64 % from 0 mM (control) salt treatments, also the lowest values were as obtained as 72.08, 13.75 and 8.56 % from 100 mM NaCl applications. In addition, the effect of gibberellic acid on the germination rate was found significant and the best mean (47.77 %) was determined from 300 ppm GA<sub>3</sub>, lowest germination rate (33.88 %) is due to control of GA<sub>3</sub>. But, among the first three doses of gibberellic acid there isn't any a different as statistically. Based on statistical analysis, salinity significantly decreased mean germination time of seeds primed with 200 and 300 ppm GA<sub>3</sub> applications. In salt doses of 50 mM and 100 mM, the applications of gibberellic acid in concentrations of 100, 200 and 300 ppm caused to the increase in germination power or rate, respectively. In the result of this research, mean germination time (6.37 day) has shortened with control (0 mM) salt application. The longest average germination time recorded as 9.57 day from 100 mM NaCl treatments. As for the sensitivity index, increasing salt applications have also increased these values. Sensitivity index value is 1.46 % in 100 mM salt dose and 0.46 % in 0 mM salt dose. In this study, we observed that germination percentage delayed and decreased with increasing NaCl concentration and drastically reduced at 100 mM NaCl. The results showed that echinacea is highly sensitive to salinity in the germination stage. In carried previous work by Yuonesi and Moradi (2015), reported that is consistent with the hypothesis that under salinity stress, priming can prepare a suitable metabolic reaction in seeds and can improve seed germination performance and seedling establishment. These results agree with Singh and Jakar (2018), in *Vigna mungo* and Fardus et al. (2018) in wheat, they suggested that salinity may influence germination by decreasing the water uptake, and germination percentage declined with increase of salinity level. However, osmotic stresses reduce germination percentage (Cornelia and Bandici 2008; Maghsoudi and Arvin 2010) although, the seeds pre-treated with GA<sub>3</sub> solutions exhibited higher germination percentage. Sabra et al. (2012) reported that

photosynthetic activity in echinacea was affected by salinity and Zollinger et al. (2007) also found a similar reduction in photosynthetic rate and stomatal conductance in *E. purpurea* irrigated with increasing concentrations of salinity. In addition, Sanam et al. (2014), different levels of salinity significantly reduced germination rate, germination vigor and index. The results also showed that by increasing salinity levels, the percentage of germination and normal seedlings significantly decreased and the mean time to germination increased, compared to the control treatment. But the seeds treated with GA<sub>3</sub> showed higher viability and better performance under salinity stress condition. Ali et al. (2012) and Zadeh et al. (2015) reported that application of GA<sub>3</sub> enhanced growth parameters (shoot length, shoot fresh weight, shoot dry weight, root length, root fresh weight, root dry weight, leaf area etc.) under saline condition. These results are consistent with those of Jamil and Rha (2007) in

sugar beet, Erdemli and Kaya (2015) in sunflower, Yıldız et al. (2017) in sweet william, Cavusoglu et al. (2007) in radish seed, Singh and Jakar (2018), in *Vigna mungo* and Zadeh et al. (2015) in echinacea. Promotion in seed germination with GA<sub>3</sub> may be due to increased uptake of oxygen and α-amylase activity.

Yalcın (2010) and Yildiz et al. (2017) recorded that the positive and significant effects of GA<sub>3</sub> applications on germination power were found. Jamil and Rha (2007) reported that water uptake of primed seeds also increased significantly with increasing concentration of GA<sub>3</sub> as compared to control. Erdemli and Kaya (2015), recorded that it was concluded that seed treatment with GA<sub>3</sub> can be beneficial for decreasing the effect of abiotic stress conditions on germination. It was seen that as gibberellic acid dose increases germination parameters were improve when compared to control.

Table 1. Effect of gibberellic acid treatments on germination of echinacea seeds under salt stress

Applications		Germination power (%)	Germination rate (%)	Mean germination time (day)	Germination index (%)	Sensitivity index (%)
Salt Doses	GA3					
Control (T0)	GA0 (control)	90.00	68.33	6.50	15.51	-
	GA100	91.66	66.66	6.62	13.39	1.01 de
	GA200	95.33	60.00	6.54	14.27	1.00 de
	GA300	95.00	70.00	5.81	15.41	0.89 e
T0 Means		92.5 a	66.25 a	6.37 c	14.64 a	0.72 c
50 mM (T5)	GA0 (control)	88.33	26.66	8.29	11.78	1.41 ab
	GA100	83.33	40.00	8.05	12.48	1.23 c
	GA200	86.66	41.66	7.28	12.10	1.15 cd
	GA300	88.33	50.00	7.51	13.55	1.15 cd
T5 Means		85.41 a	39.58 b	7.78 b	12.48 b	1.23 b
100 mM (T10)	GA0 (control)	70.00	6.66	9.99	7.60	1.53 a
	GA100	76.66	10.00	10.03	9.93	1.54 a
	GA200	76.66	15.00	9.84	7.81	1.50 a
	GA300	70.00	23.33	8.43	8.91	1.29 bc
T10 Means		72.08 b	13.75 c	9.57 a	8.56 c	1.46 a
GA <sub>3</sub> Doses Means	GA0 (control)	82.77	33.88 b	8.26 a	11.63 ab	0.98 c
	GA100	83.88	38.88 b	8.23 a	11.94 ab	1.26 a
	GA200	82.22	38.88 b	7.89 ab	11.39 b	1.20 ab
	GA300	84.44	47.77 a	7.25 b	12.62 a	1.11 b
CV (%)		10.28	19.39	8.37	9.76	11.20

\*There is no significant difference between the means indicated with same letter (5%).

Table 2. Effect of gibberellic acid treatments on some characteristics of echinacea seedlings

Applications		Radicula length (cm)	Plumula length (cm)	Fresh radicle weight (mg)	Fresh plumula weight (mg)	Dry radicle weight (mg)	Dry plumula weight (mg)
Salt Doses	GA3						
Control (T0)	GA0 (control)	1.23	2.65	57.66	374.7	5.20 ab	29.26
	GA100	1.71	3.01	59.43	388.66	5.03 a-d	32.0
	GA200	1.67	3.13	64.66	417.6	5.13a-c	33.66
	GA300	1.63	3.43	63.90	441.23	5.70 a	34.76
T0 Means		1.56 a	3.05 a	61.41 a	405.5 a	5.26 a	32.42 a
50 mM (T5)	GA0 (control)	1.03	1.66	54.53	354.63	4.33 c-f	24.20
	GA100	0.78	2.36	57.26	357.06	4.23 d-f	26.33
	GA200	1.10	2.46	63.86	397.23	4.63 b-e	24.96
	GA300	1.28	2.53	60.23	432.96	4.03 ef	30.30
T5 Means		1.05 b	2.25 b	58.97 a	385.47a	4.30 b	26.45 b
100 mM (T10)	GA0 (control)	0.50	0.91	33.33	165.5	1.63 g	21.76
	GA100	0.46	1.15	39.20	188.80	3.80 f	21.66
	GA200	0.70	1.10	40.66	195.7	3.66 f	23.10
	GA300	0.72	1.48	40.93	224.76	3.73 f	23.70
T10 Means		0.59 c	1.16 c	38.53 b	193.69 b	3.20 c	22.55 c
GA <sub>3</sub> Doses Means	GA0 (control)	0.92 b	1.74 c	48.51 c	298.27 b	3.72 b	25.07 b
	GA100	0.98 ab	2.17 b	51.96 bc	311.51 ab	4.35 a	26.66 ab
	GA200	1.15 ab	2.23 b	56.40 a	336.84 ab	4.47 a	27.24 ab
	GA300	1.21 a	2.48 a	55.02 ab	366.32 a	4.48 a	29.58 a
CV (%)		9.12	11.51	7.98	7.98	8.01	8.20

\*There is no significant difference between the means indicated with same letter (5%).

The effect germination inhibiting in seeds of salt has been revealed in many studies. It has been reported that it was caused to the inhibition of water intake of high salt concentration on the decrease of germination rate, toxicity of salt and the can not be activated due to salt stress of required enzymes during germination (Mansour, 1994; Essa, 2002; Sadeghian and Yavari 2004). However, Seyedi et al. (2012) reported that, with increasing salinity stress germination characteristics such as germination percentage, germination rate and seedling fresh weight decreased. All of the early growth parameters have decreased as parallel with salt application.

Priming of seeds with different materials particularly  $GA_3$  was useful for alleviating salt stress effects and improving germination and seedling establishment under salt stress. Under salinity condition, primed seeds possessed more germination and emergence than control. According to the results of the study; the highest values for investigated characters such as radicle and seedling length, fresh radicle and seedling weight and dry radicle and seedling weight were determined as 1.56 cm, 3.05 cm, 61.41 mg, 405.5 mg, 5.26 mg and 32.42 mg from control applications (0 mM NaCl), respectively. Also, the most negative results were determined as 0.59 cm, 1.16 cm, 38.53 mg, 193.69 mg, 3.20 mg and 22.55 mg from 100 mM NaCl applications, respectively. In terms of gibberellic acid applications the most positive results for the characters such as radicle and seedling length, fresh radicle and seedling weight and dry radicle and seedling weight were obtained from 300 ppm  $GA_3$  applications as 1.21 cm, 2.48 cm, 56.40 mg, 366.32 mg, 4.48 mg and 29.58 mg, respectively. In this experiment, we observed that, growth parameters reduced with increasing salinity (Table 2) which was similar with the report of Sabra et al. (2012) in echinacea species, Nazarian (2016) in rapeseed, Salahuddin et al. (2017), in mungbean, Fardus et al. (2018) in wheat and Singh and Jakhar (2018), in *Vigna mungo*.

## Conclusions

In the trial conducted to evaluate some germination and growth characteristics (radicle length, shoot length, radicle age and dry weight, shoot age and dry weight, germination strength, germination rate, average germination time, germination and sensitivity index) of echinacea was concluded that germination and growth parameters of echinacea seeds were inhibited as the salt concentration increases. Echinacea seeds exposed to the salt stress and gibberellic acid concentrations were found to have a significant positive effect on germination and growth properties of echinacea. The best results were obtained from the combination of 300 ppm  $GA_3$  and 0 mM (control) salt in the improving as physiological of echinacea seeds. As a result, it was determined that  $GA_3$  treatments eliminated the germination-inhibiting effects caused by salt stress and shortened the average germination time.

## References

Ahmad, K. M., Zafar, Z., Mirza, J. I. (2017). Effect of plant growth regulators on shoot organogenesis and somatic embryogenesis of *Echinacea purpurea* L. Pak. j. life soc. Sci. 15(2): 107-113. [[Google Scholar](#)]

Akinci, I.E., Caliskan, U. (2010). Effect of lead on seed germination and tolerance levels in some summer vegetables. *Ekoloji Journal*, 19, 164-172. [[Google Scholar](#)] [[CrossRef](#)]

Ali, H. M., Siddiqui, M. H., Basalah, M. O., Al-Wahaibi, M. H., Sakran, A. M., Al-Amri, A. (2012). Effects of gibberellic acid on growth and photosynthetic pigments of *Hibiscus sabdariffa* L. under salt stress. *African Journal of Biotechnology* Vol. 11(4), pp. 800-804. [[Google Scholar](#)] [[CrossRef](#)]

Anonymus (1999). World Health Organization. WHO Monographs on Selected Medicinal Plants, vol. 1. Malta: WHO. [[Google Scholar](#)]

Anonymus (2012). ISTA (International Rules for Seed Testing). Edition 2012. International Seed Testing Association, Bassersdorf, Switzerland. [[Google Scholar](#)]

Cavusoglu, K., Kılıç, S., Kabar, K. (2007). Some morphological and anatomical observations in alleviating salt stress with gibberellic acid, kinetin and ethylene during the germination of barley seeds. *SDU, Faculty of Science and Letters Science Journal (E-Journal)*, 2 (1): 27-40. [[Google Scholar](#)] [[CrossRef](#)]

Cornelia, P., Bandici, Gh. E. (2008). The effect of the salicylic acid on the peroxidase activity and on the growth of the barley (*Hordeum vulgare*) seedling under salt stress. *Bulletin UASVM, Agriculture* 65: 208-211. [[Google Scholar](#)] [[CrossRef](#)]

Duzgunes, O., Kesici, T., Kavuncu, O., Gurbuz, F. (1987). Research and experimental methods. *Statistical Methods-II*. Ankara University, Agr Fac Press, 1021, 295. [[Google Scholar](#)]

Ellis, R.H., Roberts, E.H. (1980). Towards a rational basis for testing seed quality. In *Seed Production* (Ed: P.D. Hebblethwaite), 605-635, Butterworths, London. [[Google Scholar](#)]

Erdemli, H., Kaya, M. D. (2015). The effects of gibberellic acid doses on yield and germination under abiotic stress conditions in sunflower (*Helianthus annuus* L.). *Field Crops Central Research Institute Journal*, 24 (1): 38-46. [[Google Scholar](#)] [[CrossRef](#)]

Essa, T.A. (2002). Effect of salinity stress on growth and nutrient composition of three soybean (*Glycine max* L. Merrill) cultivars. *Journal of Agronomy and Crop Sci.* 188, 86-93. [[Google Scholar](#)] [[CrossRef](#)]

Fardus, J., Matin, M. A., Hasanuzzaman, M., Hossain, M. A., Hasanuzzaman, M. (2018). Salicylic acid-induced improvement in germination and growth parameters of wheat under salinity stress. *The Journal of Animal & Plant Sciences*, 28 (1): 197-207. [[Google Scholar](#)]

Foolad, M.R., Lin, G.Y. (1997). Genetic potential for salt tolerance during germination in *Lycopersicon* species. *HortScience*, 32, 296-300. [[Google Scholar](#)] [[CrossRef](#)]

Grimm, W., Muller, H. H. A. (1999). Randomized controlled trial of the effect of fluid extract of echinacea purpurea on the incidence and severity of colds and respiratory infections. *Am. J. Med.* 106:138-43. [[Google Scholar](#)] [[CrossRef](#)]

Ivanova, R. V., Kraptchev, B., Stancheva, I., Geneva, M., Iliev, I., Georgiev, G. (2014). Utilization of related wild species (*Echinacea purpurea*) for genetic enhancement of cultivated sunflower (*Helianthus annuus* L.). *Turk. J. Agric. For.* 38: 15-22. [[Google Scholar](#)] [[CrossRef](#)]

Jamil, M., Rha, E. S. (2007). Gibberellic acid ( $GA_3$ ) enhance seed water uptake, germination and early seedling growth in sugar beet under salt stress. *Pakistan Journal of Biological Sciences* 10 (4): 654-658. [[Google Scholar](#)] [[CrossRef](#)]

Lopez, R., Shepard, B. M., 2007. Arthropods Associated with Medicinal Plants In Coastal South Carolina. *Insect Science*. 14, 519-524. [[Google Scholar](#)] [[CrossRef](#)]

Maghsoudi, K., M.J. Arvin (2010). Salicylic acid and osmotic stress effects on seed germination and seedling growth of wheat (*Triticum aestivum* L.) cultivars. *Plant Ecophysiol.* 2:7-11. [[Google Scholar](#)]

Mansour, M.M.F. (1994). Changes in growth, osmotic potential and cell permeability of wheat cultivars under salt stress. *Biological Plant*, 36: 429-434. [[Google Scholar](#)] [[CrossRef](#)]

Munns, R., Tester, M. (2008). Mechanisms of salinity tolerance. *Plant Biol.* 59, 651-681. [[Google Scholar](#)] [[CrossRef](#)]

Nazarian, G. R. (2016). The Effects of Priming by Salicylic Acid

- Acid Under Salt Stress on Some Morphological and Physiological Characteristics of Canola Plant. PhD Thesis, Department of Field Crops, Ege University Graduate School of Science.
- Parida, A.K., Das, A.B. (2005). Salt tolerance and salinity effects on plants: a review. *Ecotoxicol. Environ. Saf.* 60, 324-349. [[PubMed](#)] [[CrossRef](#)]
- Sabra, A., Daayf, F., Renault, S. (2012). Differential physiological and biochemical responses of three Echinacea species to salinity stress. *Scientia Horticulturae* 135: 23-31. [[ScienceDirect](#)] [[CrossRef](#)]
- Sadeghian, S.Y., Yavari, N. (2004). Effect of water-deficit stress on germination and early seedling growth in sugar beet. *Journal of Agronomy and Crop Science* 190,138-144. [[Wiley Online Library](#)] [[CrossRef](#)]
- Salahuddin, M., Nawaz, F., Shahbaz, M., Naeem, M., Zulfiqar, B., Shabbir, R. N., Hussain, R. A. (2017). Effect of exogenous nitric oxide (NO) supply on germination and seedling growth of mungbean (cv. Nm-54) under salinity stress. *Legume Research*, 40 (5): 846-852. [[Google Scholar](#)] [[CrossRef](#)]
- Sanam, S. A., Pirdashti, M. H., Mirjalili, M. H., Hashempour, A. (2014). Effect of exogenous nitric oxide on germination and some of biochemical characteristics of purple coneflower (*Echinacea purpurea* L.) in saline condition. *Zīst/shināsī-i Giyāhī-i Irān*, Vol 6, Iss 20, Pp 55-74. [[Google Scholar](#)]
- Seyedi, M., Hamzei, J., Fathi, H., Bourbour, A., Dadrasi, V. (2012). Effect of seed priming with zinc sulfate on germination characteristics and seedling growth of chickpea (*Cicer arietinum* L.) under salinity stress. *International Journal of Agriculture: Research and Review* 2 (3) Electronic Center for International Scientific Information, 108-114. [[Google Scholar](#)]
- Singh, S., Jakhar, S. (2018). 24-Epibrassinolide mediated changes on germination and early seedling parameters of *Vigna mungo* (L.) Hepper Var. Shekhar-2 under Salinity Stress. *Pertanika J. Trop. Agric. Sci.* 41 (1): 485-494. [[Google Scholar](#)]
- Wang, Y.R., Yu, L., Nan, Z.B., Liu, Y.L. (2004). Vigor Tests used to rank seed lot quality and predict field emergence in four forage species. *Crop Sci.*, 44 (2), 535-541. [[Google Scholar](#)]
- Yalcın, O. F. (2010). The Investigation Of Effects Of Gibberellic Acid And IAA (Indol-3-Acetic Acid) On Germination Of Seed Of *Silene Salsuginea* Hub.-Mor. Under Salt Stress. Selçuk University, Institute of Science, Ms. Thesis.
- Younesi, O., Moradi, A. (2015). Effect of different priming methods on germination and seedling establishment of two medicinal plants under salt stress conditions. 48 (3) *Iasi: Editura "Ion Ionescu de la Brad"*, 43-51. [[Google Scholar](#)] [[CrossRef](#)]
- Yıldız, S., Karagoz, F. P., Dursun, A. (2017). Germination in salt stress of sweet william (*dianthus barbatus* l.) seeds applied pretreatment of gibberellic acid. *Atatürk Univ., J. Of The Agricultural Faculty*, 48 (1): 1-7. [[CABDirect](#)]
- Zadeh, S. Y., Ramin, A. A., Baninasab, B. (2015). Effect of gibberellic acid, stratification and salinity on seed germination of *Echinacea purpurea*cv. *Magnus*. *Herba Polonica*; 61(3): 13-22. [[Google Scholar](#)] [[CrossRef](#)]
- Zollinger, N., Koenig, R., Cerny-Koenig, T., Kjelgren, R. (2007). Relative salinity tolerance of intermountain western United States native herbaceous perennials. *HortScience* 42, 529-534. [[Google Scholar](#)] [[CrossRef](#)]