

THE BOVIDAE FAUNA OF MIDDLE SINAP OF TURKEY

Türkiye Orta Sinap Bovidae Faunası

Ibrahim Tekkaya
M.T.A. Enstitüsü, Ankara

ABSTRACT. — *The Bovidae fauna of the Middle Sinap are very interesting because, the representatives of this fauna are helpful in establishing further the association of species and genera between Europe and Asia. For this reason, the Bovidae fauna of the Middle Sinap are very important in correlating the faunal scale between east and west (Asia and Europe) are established by Palaeoreas, Pseudotragus, Tragoreas and Helicotragus. In addition to these, there is the association of species between Europe and Asia established by Piotoryx carolinae Major, Protoryx longiceps Pilgrim and Hopwood, Gazella deperdita Gervais and Gazella gaudryi Schlosser. On the other hand, these representatives generally migrated from the middle Anatolia (Middle Sinap) to Aegean district and Asia.*

ÖZ. — *Orta Sinap Bovidae faunası çok ilginçtir. Çünkü, bu fauna topluluğu gerek Avrupa ve gerekse Asya genus ve espes birliginin tesisinde büyük rol oynamaktadır. Bu sebepten Orta Sinap Bovidae faunası doğu ile batı arasındaki gerçek bir faunik skalanın kurulmasında ilgi çekicidir. Bu temsilcilerden Palaeoreas, Pseudotragus, Tragoreas ve Helicotragus ile doğu ve batı (Asya ve Avrupa) arasında genus birligi tesis edilmiştir. Piotoryx carolinae Major, Protoryx longiceps Pilgrim ve Hopwood, Gazella deperdita Gervais ve Gazella gaudryi Schlosser ile de Avrupa ve Asya arasında espes birligi kurulmuştur. Bu nedenle Orta Sinap Bovidae fawnası Türkiye için olduğu kadar Asya-Avrupa faunik birliginin kuruluşunda da büyük önem taşır.*

INTRODUCTION

The Bovidae family discovered in the Middle Sinap series¹ was a very interesting development for Anatolian paleontology. This family is equally important for Asia and Europe because, these representatives of the Middle Sinap are Asiatic like Maragha fauna and European like Samos, Pikermi and Salonica fauna. There are 81 extinct genera of this family, but we dis-

(1) The detailed study on Middle Sinap faunas is in preparation.

covered 8 genera in the Middle Sinap series; one of them is a new genus, 15 species which belong to these genera and are distributed systematically in the following subfamilies:

Subfam. Bovinae

Palaeoreas brachyceras Ozansoy

Palaeoreas elegans Ozansoy

Subfam. Hippotraginae

Protoryx carolinae Major n. var.

Protoryx longiceps Pilgrim and Hopwood n. var.

Protoryx capricornis n. sp.

Pseudotragus parallelocornis n. sp.

Tragoreas sinapensis n. sp.

Subfam. Antilopinae

Helicotragus incarinatus Ozansoy

Sinapocerus ozansoyi n.gen. n.sp.

Gazella deperdita Gervais n. var.

Gazella ancyrensis Tekkaya

Gazella cf. ancyrensis Tekkaya

Gazella gaudryi Schlosser

Gazella sp.

Incertia sedis.

Qurliqnoria şenyüreki Ozansoy

Palaeoreas

The correlation was carried out morphologically, biometrically and graphically on the specimens of the Middle Sinap and the other representatives of the other areas of Turkey, Europe and Asia. We especially considered the stratigraphical periods and systematical points. In the light of present knowledge we investigated the association of species and genera, the faunical migration of the Middle Sinap and the other areas of Turkey, including Asia and Europe. In this way, the evolutional continuity of the genus was determined.

According to published records, *Palaeoreas* is generally found in Thrace (Turkey) and in the other parts of Anatolia, in Aegean district and in France. *Palaeoreas lindermayeri* Gaudry was found in Mont Léberon (France), In Salonica, in Pikermi and in Samos (Greece), in Thrace-Küçük-çekmece and Dardanelles in Turkey (Gaudry, 1862-67; Major, 1891; Pilgrim and Hopwood, 1928; Arambourg and Piveteau, 1929; Malik and Nafiz, 1933; Melentis, 1970; Gentry, 1971; Tekkaya, 1973b). There are new species in the Middle Sinap series in Anatolia. They are *Palaeoreas brachyceras* Ozansoy and *Palaeoreas elegans* Ozansoy. Later, *Palaeoreas* sp. was found in Manisa, in Uşak, and at different places in and round Ankara (Ozansoy, 1957, 1965; Yalçınlar, 1946, 1947). *Palaeoreas* is not found in the other parts of Europe and Asia. For this reason, it has a very important place in the evolution of mammals. In our opinion, this genus lived during the Pliocene only in the Aegean area, in western and middle parts of Anatolia. It never occurred in the Miocene or in the upper Pliocene.

Protoryx

It lived in Pikermi, Samos and Salonica, in Greece, at Maragha in Iran, in Turkey and in China. *Protoryx carolinae* Major in Maragha and Salonica, *Protoryx carolinae* Major, *Protoryx hentscheli* Schlosser, *Protoryx carolinae* Major n. var. *laticeps*, *Protoryx* cf. *carolinae* Major and *Protoryx longiceps* Pilgrim and Hopwood was discovered in Samos. *Protoryx carolinae* Major was also found in China (Mecquenem, 1924, 1925; Andree, 1926; Pilgrim and Hopwood, 1928; Arambourg and Piveteau, 1929; Major, 1891; Bohlin, 1936; Schlosser, 1904; Gentry, 1971; Tekkaya, 1972). *Protoryx carolinae* Major was first discovered at Çobanpinar (Ankara), Turkey, by Ozansoy (1957, 1965, 1969). *Protoryx carolinae* Major n. var., *Protoryx longiceps* Pilgrim and Hopwood n. var., *Protoryx capricornis* Tekkaya were found by the author in the Middle Sinap series. Bohlin (1935), Teilhard and Trassaert (1938) uncertainly recorded this genus in China. According to Gentry (1971) some *Protoryx* and *Pseudotragus* species belonging to *Pachytragus laticeps* Andree can be noted as follows:

Protoryx carolinae Major (Mecquenem, 1924; Major, 1891)

Protoryx carolinae var. *laticeps* Andree (Andree, 1926)

Protoryx cf. *carolinae* Major (Schlosser, 1904)

Protoryx longiceps Major (Major, 1891)

Protoryx gaudryi Major (Major, 1891)

- Protoryx hippolyte* Major (Major, 1891)
- Protoryx hentscheli* Schlosser (Schlosser, 1904; Andree, 1926;
Pilgrim and Hopwood, 1928)
- Protoryx hentscheli* var. *tenuicornis* Andree (Andree, 1926;
Pilgrim and Hopwood, 1928)
- Protoryx crassicornis* Andree (Andree, 1926)
- Protoryx carolinae* var. *crassicornis* Andree (Pilgrim and Hopwood,
1928)
- Protoryx longicemp* Pilgrim and Hopwood (Pilgrim and Hopwood,
1928)
- Protoryx laticeps* Andree (Pilgrim and Hopwood, 1928)
- Pseudotragus capricornis* var. *hippolyte* (Pilgrim and Hopwood, 1928)

Protoryx has not been found in Europe and it was discovered only in the Aegean region, in Asia minor (Anatolia) and in Iran. However, this genus may have migrated from middle Anatolia or the area of comprising middle Anatolia to the Aegean district and Asia. *Protoryx* lived during the Lower Pliocene and it has never been found in the biozones of Miocene or the Upper Pliocene.

Pseudotragus

This genus is known in Samos, Muğla and China. *Pseudotragus longiceps* Andree, *Pseudotragus capricornis* Schlosser, *Pseudotragus capricornis* var. *Mppolyte* Schlosser were found in Samos (Schlosser, ibid; Andree, ibid; Pilgrim and Hopwood, ibid). *Pseudotragus* cf. *longicornis* Pilgrim and Hopwood was first discovered in Muğla (Ozan soy, 1951, 1967).

In the Middle Sinap, we also found a new species which was called *Pseudotragus parallelcornis* n. sp. In our opinion, this species is very important for the Anatolia region because there exists only one representative, *Pseudotragus capricornis* Schlosser from China. And according to Gentry (1971), *Pseudotragus capricornis* Schlosser (Andree, 1926) and *Pseudotragus longicornis* Andree (Andree, 1926) belong to *Pachytragus crassicornis* Schlosser. It is unknown whether *Pseudotragus* lived in Europe or whether it lived only during the Lower Pliocene in the Aegean region and in Anatolia. For this reason, it is very interesting to show with this species the faunal association of these areas. This genus may have migrated from middle

Anatolia or the area of surrounding of the middle Anatolia to the southern part of Anatolia or Samos island and then to China.

Tragoreas

It was found in the Middle Sinap series, in Samos, in Sebastopol and in China. *Tragoreas sinapensis* n. sp. was discovered the first time in the Middle Sinap, in Turkey. There are *Tragoreas oryxodes* Schlosser and *Tragoreas* sp. in Samos (Schlosser, 1904). Borissiak discovered a new species called *Tragoreas leskewitschi* Borissiak in Sebastopol, in 1914. ?*Tragoreas lagreliae* Bohlin and ?*Tragoreas palaeosinensis* (Schlosser) Bohlin lived in China. *Tragoreas* is known in Europe, more than this genus may be distributed to the eastern area of Sebastopol or the upper region of Black Sea, and later the first group may have migrated from here to middle Anatolia (Orta Sinap), and then from the middle Anatolia to Samos island. The second group migrated from the Black Sea region to China.

Helicotragus

Helicotragus rotundicornis Weithofer (Syn. *Helicophora rotundicornis* Weithofer) was found in Pikermi, in Salonica and in Maragha (Mecquenem, 1925; Arambourg and Piveteau, 1929; Major, 1891; Rodler and Weithofer, 1890; Melentis, 1970). At the same time *Helicotragus fraasi* Andree (Syn. *Helicoceras fraasi* Andree) was found in Samos by Andree, in 1926. *Helicotragus rotundicornis* Weithofer has a very extensive distribution in Turkey. This species was discovered in Thrace-Küçükçekmece (Major, 1891; Malik and Nafiz, 1933). Şenyürek found the same species in Elmadağı (Şenyürek, 1952). Ozansoy discovered a new species called *Helicotragus incarinatus* Ozansoy at the Middle Sinap (Ozansoy, 1957, 1965), Gentry (1971) mentioned all the *Helicoreas rotundicorne* Weithofer, *Helicophora rotundicornis* Weithofer, *Helicotragus rotundicornis* Weithofer and *Helicoceras fraasi* Andree belong to *Prostrepisceros rotundicornis* (Weithofer). This genus is unknown in Europe and Asia. *Prostrepisceros* (*Helicotragus*) lived only in Aegean and in Anatolia regions. This genus may have migrated from the middle Anatolia to the eastern part of Anatolia and then Iran and to the western part of Anatolia and then to Samos.

Sinapocerus

This is a new genus and new species. Bohlin found some horn-cores in Tsaidam in China (Bohlin, 1937). But the horn-core of Tsaidam is unlike

the horn-cores of Sinap. Prof. Ozansoy discovered the same horn-cores in the Middle Sinap series but he called these materials only Antilope gen. and sp. indet. These horn-cores are never comparable or similar to the other genus of Bovidae in Pliocene. For this reason, we called these materials *Sinapocerus ozansoyi* n. gen and n. sp.

Gazella

The genus *Gazella* is still living, but some species of *Gazella* are extinct. *Gazella stehlini* Thenius has reportedly occurred in the upper Miocene. It lived in the Rhône and in Saône regions in France, in la Chaux-de-Fondes district in Switzerland and in Vienna, in Austria (Stehlin, 1937; Thenius, 1951; Gentry, 1964). At the same time, Thenius stated that this species had lived in Nikolsburg, in Czechoslovakia and in Chios, in Greece (Thenius, 1952; Papp and Thenius, 1959). *Gazella anglica* Newton (Newton, 1884) and *Gazella daviesi* Hinton (Hinton, 1908) were found in the Pliocene deposits in England. *Gazella deperdita* Gervais and *Gazella burbonica* Deperet were discovered in Teruel and in Alcoy district in Spain (Crusafont and Truyols, 1954). *Gazella deperdita* Gervais lived in Croix-Rousse in Lyons and in Mont Léberon in France (Gaudry, 1873; Pilgrim and Hopwood, 1928; Major, 1891; Heintz, 1969, 1971). According to some writers *Gazella baltavarensis* Benda, *Gazella capricornis* Rodler and Weithofer, *Gazella gaudryi* Schlosser, *Gazella deperdita* Gervais lived in the Baltavar region in Hungary (Major 1891; Pilgrim and Hopwood, 1928). *Gazella capricornis* Rodler and Weithofer, *Gazella deperdita* Gervais, *Gazella mytilinii* Pilgrim *Gazella schlosseri* Pavlow have been discovered in Romania and Ukraine (Pilgrim and Hopwood, 1928). Kretzoi mentioned that *Gazella* lived near the Karpad region. *Gazella gaudryi* Schlosser, *Gazella longicornis* Andree, *Gazella schlosseri* Andree, *Gazella mytilinii* Pilgrim, *Gazella deperdita* Gervais were found in Samos, *Gazella deperdita* Gervais, *Gazella capricornis* Rodler and Weithofer were discovered in Pikermi and *Gazella gaudryi* Schlosser, *Gazella deperdita* Gervais were found in Salonica (Andree, 1926; Melentis, 1970; Gaudry 1862-67; Gervais, 1859; Arambourg and Piveteau, 1929; Pilgrim and Hopwood, 1928; Schlosser, 1904; Major, 1891).

There are many species of *Gazella* in Turkey. *Gazella pererdita* Gervais and *Gazella gaudryi* Schlosser were found in Thrace. *Gazella burbonica* Deperet was discovered in Eskişehir. *Gazella otkuni* Ozansoy, *Gazella gaudryi* Schlosser were found in Muğla, in Akçaköy of Uşak and in Eşme of Manisa. *Gazella gaudryi* Schlosser, *Gazella pererdita* Gervais, *Gazella cf. pererdita* Gervais, *Gazella eleanorae* Şenyürek, *Gazella proatlantica* Ozansoy, *Gazella aencyrensis* Tekkaya in the Middle Sinap, in Kavaklıdere, in Elmadağı and in Ayaş in the Ankara region. After the later *Gazella* sp. was discovered in Akdere in the Kayseri region and in Istanbul (Ozansoy, 1951, 1956, 1957, 1969; Yalçınlar, 1946, 1947, 1952; Malik and Nafiz, 1933; Kansu, 1936; Leuchs, 1949; Thenius, 1949, 1951; Şenyürek, 1952, 1953, 1954; Becker-Platen and Sickenberg, 1968; İzbırak and Yalçınlar, 1951; Tekkaya, 1969, 1970, 1973a, 1973b).

Gazella gaudryi Schlosser and *Gazella pererdita* Gervais were discovered in Kerkük (Piveteau, 1935). *Gazella gaudryi* Schlosser was found in Sahandağ and *Gazella pererdita* Gervais, *Gazella capricornis* Rodler and Weithofer in Maragha in Iran (Mecquenem, 1924, 1925; Pilgrim and Hopwood, 1928; Rodler and Weithofer, 1890). Some writers reported that *Gazella pererdita* Gervais lived in Kazachstan and *Gazella gaudryi* Schlosser, *Gazella paeotehensis* Teilhard and Young were found in the western part of Mongolia (Thenius, 1949; Jaworowska, 1970). *Gazella dorcadoides* Schlosser, *Gazella gaudryi* Schlosser, *Gazella paeotehensis* Teilhard and Young, *Gazella* (protetracerus) *gaudryi* Schlosser, *Gazella blacki* Teilhard and Young, *Gazella cf. blacki* Teilhard and Young, *Gazella sinensis* Teilhard and Piveteau, *Gazella cf. sinensis* Teilhard and Piveteau, *Gazella cf. subgutturosa* Guldenst, *Gazella kueitensis* Bohlin, *Gazella paragutturosa* Bohlin lived in Pliocene and Pleistocene periods in China (Teilhard and Young, 1931; Teilhard and Trassaert, 1938; Teilhard and Piveteau, 1930; Bohlin, 1938, 1941). *Gazella lydekkeri* Pilgrim and *Gazella cf. lydekkeri* Pilgrim were found in Siwalik series in India (Pilgrim, 1937, 1939).

According to Arambourg (1959) some species of *Gazella* lived in North Africa, one of them, *Gazella praegaudryi* Arambourg occurred in the upper Miocene and *Gazella sitifense* Pomel and *Gazella thomasi* Pomel were found in the Pleistocene series.

Gazella, widely distributed during the Pliocene and Pleistocene ages in Europe and Asia (including Turkey) has conclusively established the genus association between Europe and Asia. At the same time, *Gazella gaudryi* Schlosser and *Gazella deperdita* Gervais have established the species west-east association between Europe and Asia. For this reason, these species have been characterized during the Lower Pliocene, from Europe to Asia. These two species have been found in the Pliocene age in Turkey. The other *Gazella* species have almost local characteristics, therefore, they have not migrated to these continents. Teilhard and Trassaert (1938) discussed evolutionary points of *Gazella* in China. According to these writers, the first group of *Gazella gaudryi* Schlosser is characterized only the Lower Pliocene in Shensi, Shianshiang, Kansu, Tsaidam and Shansi (zone I) localities. The second group of *Gazella blacki* Teilhard and Young lived in the Middle Pliocene in Shiashaung, Shouyangyütze, Kansu, Taiku, Mienchich-hsinam and Shansi (zone II) localities. The other groups of *Gazella sinensis* Teilhard and Piveteau, *Gazella subgutturosa* Guldenst occurred in the Villefranchian Age in Taiku, Yangton-yeh, Taoping, Wuhsiang, Fushan, Szechuan, Honan, Fukien, Yonan, Mienchich-hsinan, Nihowan, Yang-shaotsun, Kueite-tal, Hsi-kou and Shansi (zone III) localities in China.


Gazella ancyrensis Tekkaya belongs to the group of *Gazella gaudryi* Schlosser. For this reason, *Gazella ancyrensis* Tekkaya characterizes only the Lower Pliocene.

Qurliqnoria chensi Bohlin lived merely during the Pliocene age in China (Bohlin, 1935). *Qurliqnoria şenyüreki* Ozansoy was first discovered in the Middle Sinap series by Ozansoy (1957, 1965). *Qurliqnoria* has made the faunal connection between Anatolia and China.


CONCLUSION

It appears that, considering the extension of Bovidae fauna of the Middle Sinap, Anatolia was placed between Europe and Asia during Pliocene. Probably due to the widely-varied ecological zones of Anatolia at that time, representatives of the Eurasian fauna seem to have come together in Middle Sinap times.

Distribution of the Fossil Bovidae Faunas of Eurasia.



The Stratigraphic position of the Bovidae faunas of Middle Sinap



The distribution list of the Fossil Bovidae faunas of Eurasia

- 1 — *Palaeoreas brachyceras* Ozansoy
- 2 — *Palaeoreas elegans* Ozansoy
- 3 — *Palaeoreas lindermayeri* Gaudry
- 4 — *Protoryx carolinae* Major
- 5 — *Protoryx carolinae* Major n. var. *laticeps*
- 6 — *Protoryx hentscheli* Schlosser
- 7 — *Protoryx cf. carolinae* Major
- 8 — *Protoryx longiceps* Pilgrim and Hopwood
- 9 — *Protoryx carolinae* Major n. var.
- 10 — *Protoryx longiceps* Pilgrim and Hopwood n. var.
- 11 — *Protoryx capricornis* m sp.
- 12 — *Pseudotragus longiceps* Andree
- 13 — *Pseudotragus capricornis* Schlosser
- 14 — *Pseudotragus capricornis* var. *hippolyte* Schlosser
- 15 — *Pseudotragus cf. longicornis* Pilgrim and Hopwood
- 16 — *Pseudotragus parallelocornis* n. sp.
- 17 — *Tragoreas sinapensis* n. sp.
- 18 — *Tragoreas oryxoides* Schlosser
- 19 — *Tragoreas leskewitschi* Borissiak
- 20 — ? *Tragoreas lagrelii* Bohlin
- 21 — ?*Tragoreas palaeosinensis* (Schlosser)
- 22 — *Helicotragus rotundicornis* Weithofer

- 23 — *Helicotragus fraasi* Andree
- 24 — *Helicotragus incarinatus* Ozansoy
- 25 — *Sinapocerus ozansoyi* n.gen. n.sp.
- 26 — *Gazella stehlini* Thenius
- 27 — *Gazella anglica* Newton
- 28 — *Gazella daviesi* Hinton
- 29 — *Gazella deperdita* Gervais
- 30 — *Gazella burbonica* Deperet
- 31 — *Gazella baltavarensis* Benda Bohlin
- 32 — *Gazella capricornis* Rodler and Weithofer
- 33 — *Gazella gaudryi* Schlosser
- 34 — *Gazella mytilinii* Pilgrim
- 35 — *Gazella schlosseri* Pavlow
- 36 — *Gazella longicornis* Andree
- 37 — *Gazella otkuni* Ozansoy
- 38 — *Gazella cf. deperdita* Gervais
- 39 — *Gazella eleanorea* Şenyürek
- 40 — *Gazella proatlantica* Ozansoy
- 41 — *Gazella ancyrensis* Tekkaya
- 42 — *Gazella* sp.
- 43 — *Gazella paeotehensis* Teilhard and Young
- 44 — *Gazella dorcadoides* Schlosser
- 45 — *Gazella (prototracerus) gaudryi* Schlosser
- 46 — *Gazella blacki* Teilhard and Young
- 47 — *Gazella cf. blacki* Teilhard and Young
- 48 — *Gazella sinensis* Teilhard and Piveteau
- 49 — *Gazella cf. sinensis* Teilhard and Piveteau
- 50 — *Gazella cf. subgutturosa* Guldenst
- 51 — *Gazella kueitensis* Bohlin
- 52 — *Gazella paragutturosa* Bohlin
- 53 — *Qurliqnoria chensi* Bohlin
- 54 — *Qurliqnoria şenyüreki* Ozansoy

Yayına verildiği tarih: 15.1.1974

REFERENCES

- Andree, J., 1926, Neue Cavicornier aus dem Pliocen von Samos, *Palaeontographica*, Stuttgart, Bd. 67, 135-175 p.
- Arambourg, C., 1959, Vertébrés continentaux du Miocène supérieur de l'Afrique du Nord, *Mém. carte géol. Algérie*, 1-159 p.
- Arambourg, C., Piveteau, J., 1929, Les Vertébrés du Pontien de Salonique, Paris, Ann. Pal., t. XVIII.
- Becker, Platen, J. D. - Sickenberg, O., 1968, Die unterpleistozaenen kiese von Es-kişehir (Anatolien) und ihre Säugetierfauna, Hannover, Mitt. Geol. Inst Techn. Univ., 7-20 p.
- Bohlin, B., 1935, Cavicornier der Hipparion-fauna Nord-chinas, Peking, *Pal. sinica*, ser. C, vol. IX, Fase. 4,
- 1936, Bemerkungen über einige Pontischen Antilopen-Gattungen, Stockholm, *Ark. Zool.*, No. 28,
- 1937, Eine Tertiaere Säugetier-fauna aus Tsaidam, Peking, *Pal. Sinica*, Ser. C, vol. XIV, Fase. 1,
- 1938, Einige Jungteriaere und pleistozaene Cavicornier aus Nord-China, Uppsala, Nov. Act. Reg. Soc. Sci.,
- 1941, *Gazella* (*Protetracerus*) *gaudryi* (Schlosser) and *Gazella dorcadoides* Schlosser, Upsala, *Bull. Geol. Inst. Univ.*
- Borissiak, A., 1914, Mammifères fossiles de Sébastopol, Paris, *Mem. du comt. Geol. Nouvelle ser.*
- Crusafont, P. - Truyols, J., 1954, *Sinopsis estratigrafico, Paleontologica de Vallés-Penedés*, Sabadell.
- Gaudry, A., 1962-67, *Animaux fossiles et Géologie de L'Attique*, Paris, 1873, *Animaux fossiles du Mont Léberon*, Librairie de la Soc. Géol. de France, Paris.
- Gentry, A. W., 1964, Skull characters of African Gazelles, *Ann. Mag. Nat. Hist.*, 1971, The earliest Goats and other Antelopes from the Samos Hipparion fauna, *Bull. Br. Mus. Nat. Hist. (Geol.) London*, vol. 20, No. 6.
- Gervais, P., 1859, *Zoologie et Paléontologie Françaises*, Paris, 544 p.
- Heintz, E., 1969, Le dimorphisme sexuel des appendices frontaux chez *Gazella perperdita* Gervais (Bovidae, Artiodactyla, Mammalia) et signification phylogénique mammalia, Paris, T. 33, No. 4,
- 1971, *Gazella perperdita* (Gervais) 1847 (Bovidae, Artiodactyla, Mammalia) du Pontien du Mont Léberon Vaucluse, France, *Ann. Pal. Vertèbres*, Tom. LVH, Fasc. 2, 209-229 p.
- Hinton, M. A., 1908, Note on *Gazella daviesi* Hinton, an Antelope from the Nördlich crag. *Geol. Mag.*, London, 445 p.
- İzbırak, R. - Yalçınlar, İ., 1951, Kayseri kuzeyinde üst Miosen'e ait omurgalılar, T.J.K., Cilt III, Sayı 1.

- Jawarowska, Z. K., 1970, Result of the Polish-Mongolian, Palaeontological expedition, Ak. Sic. Inst. Paleo. Warszawa, Part II, 95-104 p.
- Kansu, A. Ş.; ,1936, Mitteilung über Palaontologische und Praehistorische Funde in der Umgebung von Ankara, Ülkü Dergisi.
- Leuchs, K., 1939, Beitrage zur Tertiargliederung in Innerana tolian, Berlin z. Deutsch. Geol. Ges., 785 p.
- Major, F. J., 1891, Le gesement ossifere de Mitylini, Etude Geol. Jaleon. et Botanique, Laussanne, 94 p.
- Malik, A. - Nafiz, H., 1933, Küçükçekmece fosil fıkralı hayvanlar mecması, Darülfünun Geol. Enst. No. 8, İstanbul.
- Mecquenem, de R., 1924, Contribution a l'étude des fossiles de Maragha, Ann. Pal. Paris, T. XIII, 28 p.,
1925, Fossiles de Maraha, Ann. de Pal. Paris, T. XIV, Fasc. 1, 1-36 p.
- Melentis, J. K., 1970, Die Pikermifauna von Halmyropotamus (Euböa-Griechenland) Ann. Geol. des Pays Heleniques, premier série., Tom dix neuviene, Athènes, 285-404 p.
- Newton, E. T., 1884, On the occurrence of Antelope remains in newer, Pliocene Beds in Britain, with description of a new species, *Gazella anglica*, Quart. Geol. Soc. London, 280-293 p.
- Ozan soy, F., 1951, Muğla Pontien Memeli faunası, T.J.K., vol. III, Sıra No. 1, 1956, Ankara çevresi neojeninin Jeolojik katları ve Memeli fosil faunaları hakkında, M.T.A. derleme, rop. No. 2357, Ankara (neşredilmemiş), 1957, Türkiye Tersier Memeli faunaları ve Stratigrafik revizyonu, M.T.A. Dergisi, No. 49, Ankara.
- 1965, Etude des gisement continentaux et des Mammifères de Cenozoïque de Turquie, Mem. Soc. Geol. France, Paris, 92 p.,
- 1969, Biozonların ışığında Geyve - Gölpazarı - Göynük - Mudurnu - Nallıhan kuzeyi çevresinde Kretase-Senozoik sınırı ve Senozoik tortulları hakkında, M.T.A. derleme, Rop. No. 4081, Ankara (neşredilmemiş).
- Pilgrim, G. E., 1937, Siwalik Antelopes and Oxen in the American Museum, Bull. Ame. Mus. Nat. Hist. vol. 72, 731-873 p.,
- 1939, The fossil Bovidae of India, Pal. Indica, 356 p.
- Pilgrim, G. E. - Hopwood, A. T., 1928, Catalogue of the Pontien. Bovidae of Europe, Br. Mus. Nat. Hist. 30 p.
- Piveteau, J., 1935, Les Mammifères du Pontien de l'Irak, Bull. Soc. Geol. France, Paris, T. 5.
- Rodler, A. - Weithofer, K. A., 1890, Die Viederkauer der Fauna von Maragha, Denkschr. k. Ak. Wiss. Mat. Nat. Kl. Bd. LVII, Abt. II, Vienne, 753-772 p.
- Schlosser, M., 1904, Die fossilen Cavicornis von Samos, Beitrage zur Palaen. und Geol. Bd. XVII
- Stehlin, H. G., 1937, Bemerkungen über die Miocaenen Hirschgenera Stephanoceras

- mas und Lagomeryx, verhandl. Naturforsch. Ges. Basel.
- Şenyürek, M., 1952, A study of the Pontien fauna of Gökdere (Elmadağı) South-east of Ankara, Belleten, Ankara, Cilt XVI, Sayı 64,
- 1953 a, A note a new species of Gazella from the Pontien of Küçükkyozgat, A.Ü.D.T.C.F. Dergisi, ai t XI, Sayı 1,
- 1953 b, A horn-core of *Gazella copricornis* Rodler and Weithofer found at Küçükkyozgat, Bull. Geol. Soc. Turkey, Ankara, vol. IV, Sayı 2, 141-144 p.,
- 1954, Pontien çağda Ankara civarında yaşamış olan bir ceylân türü, Coğ. Mes. Haft. İstanbul, 67-72 p.
- Teilhard de Chardin, P. - Piveteau, J., 1930, Les Mammifères fossiles de Nihowan (China), Ann. Pal. Paris, T. XIX, 134 p.
- Teilhard de Chardin, P. - Trassaert, M., 1938, Cavicornia of South eastern Shansi, Pal. Sinica, Peking, New Series C, No. 6, 98 p.
- Teilhard de Chardin, P. - Young, C. C., 1931, Fossil Mammals from the late Cenozoik of Nortem China, Pal. Sinica, Peking, Serie C, vol. IX, Fasc. 1, 66 p.
- Tekkaya, İ., 1969, Konya Kayadibi faunası hakkında rapor, M.T.A. Dergisi, Ankara, No. 73, 155-160 p.; Preliminary report on the Bovidae fauna from Kayadibi, Konya, Bull. M.T.A., No. 73, bir boynuz, M.T.A. Dergisi, Ankara, No. 74; Horn-core of *Gazella perdeperita* Gervais (n. var.) from Middle Sinap, Bull. M.T.A., No. 74.
- Tekkaya, İ. - Şen, Ş. - Atalay, Z., 1972, Kırık Memeli Faunası hakkında bir inceleme, M.T.A. Dergisi, Ankara, No. 78.
- Tekkaya, İ. 1973 a, Orta Sinap'ta yeni bir *Gazella* türü, M.T.A. Dergisi, Ankara, No. 80; Une Nouvelle espèce de *Gazella* de Sinap Moyen, Bull, M.T.A., No. 80,
- 1973 b, Gülpınar'daki Fosil Bovidae kalıntıları hakkında bir not, T.J.K., Ankara, No. 16/2; A note on the remains of Fossil Bovidae of Gülpınar, Bull. Geol. Soc. Turkey, No. 16/2.
- Thenius, E., 1949, Über die Säugetierfauna aus dem Unterplizaen von İlhan bei Ankara (Türkei), Öst. Ak. Wiss. Mat. Nat. Klass. Abt. 1, Bd. 158,
- 1951, *Gazella* cf. *deperdita* aus dem Mitteleuropaeschen Vidobonien und das Auftreten der Hipparrionfauna, Eclog. Geol. Helv., 381-394 p.,
- 1952, Teratologische erscheinungen an Hipparrionzähnen, Eclog. Geol. Helv., 381-394 p., 1959, Tertiär, Stuttgart, 328 p.
- Yalçınlar, İ., 1946, Eşme civarında Miosen'e ait Omurgalı faunası, İ.Ü.F.F. Mec, Seri B, Cilt XI, Sayı 2, 1947, Yukarı Gediz vadisinin Miosen'e ait Vertebra fosilleri, T.J.K., Ankara, Cilt 1, Sayı 1, 1952, Les vertèbres fossiles Neogenes de la Turquie occidentale, Bull. Mus. Nat. Hist. Nat. Ser. 2, Cilt XXIV, Sayı 4.