

## Comparative Study of Lipid and Fatty Acid Profile in Liver Tissues of Male and Female *Silurus triostegus* During the Catching Seasons

Semra Kaçar<sup>1</sup> , Mehmet Başhan<sup>2</sup> 

**Cite this article as:** Kacar, S., & Bashan, M. (2021). comparative study of lipid and fatty acid profile in liver tissues of male and female *Silurus triostegus* during the catching seasons. *Aquatic Sciences and Engineering*, 36(4), 166-174.

### ABSTRACT

In this study, seasonal variations of total fatty acid (FA), phospholipid (PL) and triacylglycerol (TAG) compositions in liver tissues of catfish (*Silurus triostegus*) were investigated. Samples of *S. triostegus* were obtained from Atatürk Dam Lake, Turkey, in two month periods during one year as from May. The major components were palmitic acid (16:0), stearic acid (18:0), oleic acid (18:1n-9), palmitoleic acid (16:1n-7), docosahexaenoic acid (DHA, 22:6n-3) and arachidonic acid (AA, 20:4n-6) in total lipid, 16:0, 18:0, 18:1n-9, eicosapentaenoic acid (EPA, 20:5n-3), DHA and AA in PL, 16:0, 16:1n-7, 18:1n-9, linoleic acid (LA, 18:2n-6), AA, EPA, DHA and docosapentaenoic acid (22:5n-3) in TAG extracted from the liver of *S. triostegus* in all seasons. N-3/n-6 ratio was found 2.00-2.61 in females, 1.15-2.75 in males in total lipid. The highest lipid content was found in May (2.39%) in the females. In the males, the highest level was found in September (2.91%). In TAG fraction, the component with the lowest ratio in both sexes' TAG is PUFA. In PL fraction, SFA, MUFA, and PUFA percentages were found at similar rates in both sexes in all months.

**Keywords:** *Silurus triostegus*, fatty acid, phospholipid, triacylglycerol, comparison, male, female, season, total lipid

ORCID IDs of the author:  
S.K. 0000-0002-9869-9045;  
M.B. 0000-0002-1228-9548

<sup>1</sup>Department of Nutrition and Dietetics,  
School of Health, Mardin Artuklu  
University, Mardin, Turkey

<sup>2</sup>Dicle University, Faculty of Science,  
Department of Biology, Diyarbakir, Turkey

Submitted:  
17.01.2021

Revision Requested:  
17.03.2021

Last Revision Received:  
19.03.2021

Accepted:  
20.03.2021

Online Published:  
03.05.2021

Correspondence:  
Semra Kaçar  
E-mail:  
[semrakacar21@gmail.com](mailto:semrakacar21@gmail.com)

### INTRODUCTION

Mesopotamia catfish (*S. triostegus*) is a catfish species from the Siluridae family which is found in the Mesopotamia region namely Syria, Iraq, Iran, and Turkey. In *S. triostegus* living in the Atatürk Dam Lake, it was determined that egg laying started in May and continued until the end of June by utilizing parameters such as gonadosomatic index values and direct observation of gonads. Consequently, it can be said that the reproduction period is May, June, and July (Oymak *et al.*, 2001).

Catfish meat is delicious, and has high protein content. It is a carnivore and aggressive species. Their nutrients usually are water insects and their larvae, worms, frogs, and tadpoles (Geldiay & Balık 1996).

Today, there is a great interest in fish and fish oil due to the polyunsaturated fatty acids (PUFA) they contain. Analyzes usually constitute the fish muscle that forms the food. On the other hand, the fish liver, i.e., the main organ for long-chain PUFA (LC-PUFA), has not been analyzed much (Ackman *et al.*, 2002). Fish liver is the source of the essential oils for the prevention of problems related to vision and growth (Njinkoué *et al.*, 2002). Moreover, livers of some fish species are used for muscle pain and rheumatism in Pakistan (Saify *et al.*, 2003).

The liver is an important organ in terms of lipid metabolism. This organ also has an important role in the uptake, oxidation, and transformation of fatty acids (FA) and the supply of long-chain highly unsaturated fatty acids to other tissues (Rincon-Sanchez *et al.*, 1992).



Phospholipids (PL) and triacylglycerides (TAG) have different roles in fish metabolism. Phospholipids are the main components of cell membrane and structure, and serve as precursors of eicosanoids with the C20 PUFA they contain. Meanwhile, TAG function as energy reserves (Sargent et al., 1995; Kiessling et al., 2001).

Fish oil and fatty acid composition are the most changed biochemical compounds according to the ecological factors and the physiological state of the fish. The total fat content and fatty acid composition in fish vary depending on species, sex, season, nutritional environment, nutrient difference, water temperature, water pollution, and whether the species are in the culture or natural form. Fat and fatty acids differ structurally in different fish species. If the same species of fish live in different geographical regions, they may differ in terms of fatty acid diversity (Ackman et al, 2002; Akpınar et al., 2009; Kayhan et al., 2014). This difference is also seen in different organs of the fish (Crowford et al., 1986; Suzuki et al., 1986).

Total, TAG, and PL fatty acids of *S. triostegus* liver have not studied before. Therefore, in this study, it was aimed to compare the seasonal changes of the total, TAG, and PL fatty acid contents of liver tissues of female and male individuals of *S. triostegus* living in Atatürk Dam Lake.

## MATERIALS AND METHODS

Samples of *S. triostegus* fish species were gathered from the Atatürk Dam Lake in one year period by using a fishing net (between May, 2008 and March, 2009). On the same day, the caught fish samples were placed in heat-insulated protective containers filled with ice, and brought to the laboratory. The measurements of height and weight of the samples were made. Weight measurements were noted in grams, and height measurements were taken in cm using the fork length of the fish (Kaçar et al., 2016). In this study, three male and three female fish were used. The sexes of fish samples were detected. After determining the wet weights of the liver samples taken out, they were put in tubes and stored in the chloroform-methanol mixture at -80 °C until analysis.

### Lipid extraction and conversion of fatty acids to methyl esters

Liver samples homogenized in the chloroform-methanol mixture (Folch et al., 1957). The thin layer chromatography technique was used to fractionate the total lipids in the samples. For this purpose, the mixture of 30 g of silica gel and 50 ml of pure water was applied as a thin layer to the plates of 20X20 size, and they were dried in an oven at 100 °C for one hour. Total lipid extracts of the samples were spotted onto the plates in a single row. The total lipids were run in a mixture of petroleum ether-diethyl ether-acetic acid (80:20:1). After drying the plates in the air, 27' dichlorofluorescein was sprayed to make the lipid fractions visible under the UV lamp. The bands of phospholipids and triacylglycerol fractions that were determined by means of standards were scraped and transferred to the reaction tubes. To each fraction, 3 ml of methanol and 3-5 drops of sulfuric acid were added dropwise, and they were all heated at 85 °C under reflux for 2 hours (Stanley-Samuelson and Dadd, 1983). The solution was extracted with methyl esters using hexane. Gas chromatography instrument with a FID detector was used for the analysis of fatty acid methyl esters.

### Gas chromatography conditions

Fatty acid analyses of the oil samples converted to methyl esters were performed by using flame ionization detector and DB-23 capillary column in HP6890 model Gas Chromatography device. The features used were as follows. Detector temperature: 280 °C, injector temperature: 270 °C, and injection: split-model 1/20. Gas flow rates were; carrier gas: 2.8 ml/min (constant flow model), hydrogen: 30 ml/min, and air: 300 ml/min. Column (oven) temperature: at 130 °C, stand-by time 1 minute; to 170 °C with 6.5 °C/min, to 215 °C with 2.75 °C/min, standby time 12 minutes; to 230 °C with 40 °C/min, standby time 3 minutes; total analysis time: 38.8 minutes. For example, 1 microliter was injected into the device. A mixture of methyl esters of fatty acids was used as the standard in the detection of fatty acids. Chromatograms of fatty acids methyl esters and total fatty acids were obtained from the computer by the software HP 3365 Chem Station.

### Evaluation of data

SPSS 16 computer program was used to compare fatty acid percent rates. All data obtained in our study were obtained from the average of three replicates. In the gas chromatographic analysis of fatty acid methyl esters, three samples of each period were injected separately, and the three values of the same fatty acid were averaged. Comparison of fatty acid percentages was made by one-way analysis of variance. Differences were determined by Tukey HSD test. As a result of the statistics, it was accepted that the differences were significant when the data were  $p < 0.05$ .

## RESULT AND DISCUSSION

### Lipid content

The quantity of total lipids ranging from 1.29 to 2.39% (breeding period) in *S. triostegus* females decreased in July, September, and January. In male fish, this ratio was found between 0.50-2.91% (post-breeding period). In males, the amount of lipid decreased significantly after September, in November and January. It increased in March (pre-breeding period) and in May (breeding period) (Table 1).

In freshwater fish, the lipid content of the liver varies depending on the season, feeding cycle, and reproductive status (Ackman et al., 2002).

**Table 1.** Total lipid of liver tissue of female and male *S. triostegus*.

	Total lipid (%)	
	Female	Male
May	2.39±0.65a	2.14±0.45a
July	1.78±0.38b	1.04±0.12b
September	1.61±0.40b	2.91±0.13a
November	2.03±0.35a	0.50±0.07c
January	1.29±0.21b	0.60±0.14c
March	1.74±0.41b	2.09±0.52a

Superscript letters (a,b,c) denote significant differences ( $p < 0.05$ ) in lipid content among months.

In the current study, it was found that liver lipid content decreased in the reproductive period of July. The decrease in total lipid and total fatty acid levels in the liver and muscles of the fish in the reproductive period shows that they obtain the energy they need from these lipids. During the reproductive period, the lipids in the liver and muscle are mobilized to the gonad for gonad development (Castell et al., 1972).

In previous studies, the liver lipid content of *Cyprinus carpio* increased in the spring (Kminikova et al., 2001; Akpınar, 1986a). Kandemir & Polat (2007) found the maximum total lipid amount in the liver of *Oncorhynchus mykiss* in the autumn season. In the present study, the highest lipid content of female fish was found in May in accordance with the previous studies.

The amount of lipid in the *S. triostegus* males decreased significantly in November and January, and increased in March and May. Jangaard et al., (1967) reported that seasonal changes in lipid level in liver and other organs are caused by irregular seasonal changes in fish feeding and water temperature.

In our study, the amount of lipid increased in both sexes in the autumn months, which was the post-reproductive period. Stored lipids vary during the breeding and feeding period. Especially when fish find enough food, they can control their reproduction and lipid storage period. The lipid storage cycle is directly related to the abundance of food. Lipid variation is high throughout the year if nutrients are abundant, whereas lipid variation is low if nutrients are scarce (Kluytmans & Zandee, 1973, Ackman & Eaton, 1976, Kinsella et al., 1977, Mute et al., 1989).

### The FA composition of total lipid

The total fatty acid content of female and male *S. triostegus* liver tissue is given in Table 2. The highest saturated fatty acids (SFA) in females were found in the summer whereas in males, they were found in spring which is the reproductive period. The lowest SFA was found in January in both sexes. The SFA was found to have 16:0 and 18:0 most commonly. The 16:0 ratio with saturated fatty acids decreased in males and females in January and in the pre-breeding period in March. In males, both components were found to be the highest in May, which is in the breeding period. The 18:0 were highest in females during November, yet no statistically significant difference was found in other seasons. This fatty acid in males showed fluctuations throughout the year. In males and females, monounsaturated fatty acids (MUFA) in liver tissue increased in January.

The MUFA and 18:1n-9 ratio was high in January and March, and decreased in September in both males and females. In September, the amount of 16:1n-7 in females reached the lowest value. This fatty acid did not show a significant change except June and September. The highest value for males was found in January. Also, 18:1n-9 did not differ in females in all seasons. In males, it is low in May and September and close to each other. The PUFA in both sexes was found to be high in September. It fell in females in May and in July in males. The 20:5n-3, 22:5n-3, and 22:6n-3 are major n-3PUFA while 18:2n-6 and 20:4n-6 are dominant n-6PUFA. Moreover, the 20:5n-3 was found to decrease in September after the breeding period, and has the highest value in

winter in female *S. triostegus* liver tissue. It was also observed that it decreased in July and increased in May in males. In May, when the arachidonic acid was the lowest in females, it was the highest in males. Docosahexaenoic acid (DHA) increased in both sexes in July and September. In females, in May, July, and November the highest SFA, in January the highest PUFA and the lowest MUFA was found while in males, in September the highest PUFA and in July, November, January, and March the highest MUFA was determined. The dominant fatty acids throughout the year are 16:0 from SFA, 18:1n-9 from MUFA, and DHA from PUFA. N-3/n-6 ratio was found 2.00-2.61 in females and 1.15-2.75 in males. In both sexes, the highest value was determined in the same month, and it was close to each other.

Fish are not only important protein sources but also contain nutritionally valuable lipids. Analyses usually focus on the fish muscle that forms the food. But the fish liver is the main organ of long-chain PUFA and has not been analyzed much (Ackman et al., 2002).

In fish liver, lipids major fatty acids are similar to the ones in fish muscle. Most of the fish that have been studied have 16:0 most among the saturated fatty acids, 18:1n-9 among the monounsaturated, and 22:6n-3 among the polyunsaturated (Kminikova et al., 2001; Uysal et al., 2006; Akpınar et al., 2009; Njinkoué et al., 2002; Aras et al., 2003b). Similar results were found in *S. triostegus* liver. However, quantitative fatty acid content in the liver varies. In accordance with the present study, in the previous studies, 14:0 and 18:0 found least among the saturated fatty acids, 16:1n-7 among the monounsaturated fatty acids, 18:2n-6 and 18:3n-3 among the highly unsaturated fatty acids, 20:3n-6 and 20:4n-6 acids which are the precursors of eicosanoids (Tufan et al., 2013; Misir et al., 2016; Kaçar & Başhan, 2017).

*Sander lucioperca's* n-3 fatty acids increased most in November when the temperature fell (Uysal et al., 2006). In both sexes of *S. triostegus*, PUFA increased in the autumn and winter when the temperature fell (Kaçar et al., 2016).

In the previous studies, in line with the current research, it was observed that long-chain unsaturated fatty acids have changed more than saturated fatty acids. It was concluded that gonad development and breeding periods had directly an effect on these changes (Akpınar, 1986b).

The data indicate that the ratio of n-3/n-6 in total lipids of the liver may be different in both sexes. This depends on the ratios of 18:3n-3, 20:5n-3, 22:5n-3, and 22:6n-3, forming n-3PUFA and 18:2n-6, 20:3n-6, and 20:4n-6, forming n-6PUFA.

In the current investigation, it was determined that the fatty acid content of fish liver lipids is affected by the sex and the season, thus, the reproductive period.

### The FA composition of TAG fraction

Table 3 shows the TAG fatty acid content of male and female *S. triostegus* liver. Palmitic acid and therefore the SFA ratio in males increased in September, which is the post-breeding period, and decreased in March, just before the breeding season. There was no significant difference in females for 16:0 throughout the year.

**Table 2.** Seasonal variations of fatty acid composition of total lipid of liver tissue from female and male *S. triostegus* (% of total FA)\*.

Fatty Acids	May		July		September		November		January		March	
	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male
10:00	-	-	-	-	0.03±0.01	-	-	-	-	-	-	-
12:00	0.05±0.02a	0.06±0.02a	-	-	0.01±0.01b	-	-	0.01±0.01b	0.12±0.05c	0.12±0.04c	0.07±0.04d	0.11±0.01c
13:00	0.32±0.03a	0.13±0.05b	0.21±0.02c	0.15±0.03b	-	-	-	-	1.14±0.16d	1.26±0.16d	1.98±0.67ab	1.96±0.17ab
14:00	1.59±0.16a	2.47±0.21b	1.75±0.11c	1.19±0.11d	1.47±0.17a	2.07±0.25ab	1.10±0.10d	1.45±0.17a	0.35±0.02c	0.25±0.02c	0.47±0.05e	0.60±0.05b
15:00	0.72±0.07a	0.54±0.04b	0.26±0.02c	0.17±0.03d	0.47±0.03e	0.54±0.04b	0.01±0.01f	0.11±0.03d	19.49±1.09b	17.28±1.17c	19.91±1.14b	18.43±1.08b
16:00	25.53±1.26a	23.46±1.26a	26.85±1.20a	20.48±1.28b	24.60±1.22a	20.89±1.09b	22.49±1.25b	21.00±1.30b	0.36±0.02d	0.27±0.03b	0.56±0.05e	0.64±0.05e
17:00	0.42±0.03a	0.42±0.04a	0.18±0.01b	0.23±0.01b	0.28±0.02b	0.51±0.05c	0.31±0.03d	0.26±0.01b	8.95±0.88a	4.84±0.45d	9.21±0.91a	5.97±0.57d
18:00	9.95±0.99a	9.94±0.99a	10.47±1.10a	11.55±1.01a	10.66±1.01a	8.58±0.87a	14.15±1.14b	7.78±0.77c	30.41±1.30b	24.02±1.28c	32.20±1.44b	27.71±1.29c
ΣSFA***	38.58±1.37a	36.96±1.33a	39.72±1.40a	33.77±1.44b	37.49±1.33a	32.69±1.29b	38.06±1.36a	30.61±1.38b	7.40±0.59a	8.60±0.83a	7.72±0.69a	6.64±0.63b
16:1n-7	7.97±0.77a	5.77±0.55b	4.11±0.41b	4.25±0.43b	3.26±0.30c	5.68±0.45b	7.41±0.66a	5.55±0.56b	21.36±1.17a	30.79±1.36b	23.52±1.22a	29.10±1.28b
18:1n-9	22.24±1.28a	21.80±1.26a	20.93±1.20a	30.91±1.30b	20.57±1.19a	21.59±1.27 <sup>a</sup>	20.13±1.10a	30.01±1.38b	3.21±0.33b	4.78±0.43d	1.93±0.17c	2.70±0.295c
20:1n-9	0.71±0.07a	0.86±0.05a	2.17±0.29b	3.16±0.32b	1.64±0.11c	1.31±0.10c	2.07±0.24b	0.99±0.08a	31.97±1.38a	44.17±1.54d	33.17±1.42e	38.44±1.45b
ΣMUFA	30.92±1.33a	28.43±1.27a	27.21±1.28a	38.32±1.43b	25.47±1.26c	28.58±1.26a	29.61±1.30a	36.55±1.33b	1.94±0.18c	1.55±0.22c	2.14±0.25a	3.80±0.36b
18:2n-6	2.52±0.24a	3.51±0.32b	1.40±0.11c	0.88±0.07d	1.69±0.16c	2.96±0.27a	2.55±0.22a	2.79±0.22a	1.34±0.16a	0.79±0.07ab	1.14±0.18a	2.32±0.25c
18:3n-3	1.22±0.17a	1.22±0.13a	0.57±0.05b	0.49±0.05b	0.93±0.08a	1.93±0.16c	1.20±0.12a	3.29±0.37c	0.54±0.05a	0.70±0.05b	0.72±0.07b	1.26±0.17d
20:2n-6	0.45±0.04a	0.49±0.03a	0.61±0.06b	0.34±0.02c	0.76±0.07b	0.80±0.07b	0.90±0.09b	0.38±0.04c	0.57±0.05a	0.68±0.06b	0.46±0.04a	0.72±0.07b
20:3n-6	0.48±0.04a	0.54±0.04a	0.46±0.03a	0.51±0.03a	0.58±0.05a	0.70±0.07b	1.25±0.13c	0.33±0.02d	7.09±0.78c	5.28±0.58a	7.05±0.71c	4.96±0.44a
20:4n-6	4.37±0.39a	11.42±1.01b	7.78±0.76c	5.75±0.56a	9.29±0.91c	8.93±0.88c	4.64±0.44a	6.26±0.65ac	5.10±0.56b	3.86±0.36a	4.68±0.41a	5.42±0.55b
20:5n-3	3.97±0.36a	5.38±0.59b	2.08±0.23c	1.97±0.17c	1.74±0.14c	4.68±0.42a	4.56±0.48a	2.84±0.21c	5.80±0.59b	3.30±0.31c	4.62±0.48b	3.51±0.33b
22:5n-3	1.05±0.10a	3.84±0.37b	2.28±0.22c	2.21±0.20c	3.03±0.38c	4.47±0.45b	3.13±0.33c	2.05±0.36c	14.27±1.14c	14.69±1.04c	13.80±1.13c	10.86±0.98b
22:6n-3	15.42±1.05a	8.05±0.86b	17.80±1.17a	15.67±1.15a	18.96±1.15a	14.21±1.07c	14.02±1.04c	13.91±1.19c	36.65±1.34c	30.85±1.27a	34.61±1.42aa	32.85±1.43a
ΣPUFA	30.48±1.30a	34.45±1.39a	32.98±1.22a	27.82±1.22b	36.98±1.33c	38.68±1.44c	32.25±1.38a	31.85±1.30c	26.51±1.29c	22.64±1.20a	24.24±1.09a	22.11±1.38a
n-3	21.66±1.27a	18.49±1.08b	22.73±1.20a	20.34±1.29a	24.66±1.30a	25.29±1.33a	22.91±1.25a	22.09±1.32a	10.14±1.10c	8.21±0.87a	10.37±1.00c	10.74±1.10c
n-6	8.82±0.89a	15.96±1.05b	10.25±0.99c	7.48±0.77a	12.32±1.02c	13.39±1.04c	9.34±0.87a	9.76±0.98a	2.61	2.75	2.33	2.05
n-3/n-6	2.45	1.15	2.21	2.71	2.00	1.88	2.45	2.26	2.61	2.75	2.33	2.05

\* Means are the averages of 3 replicates \*\* Values reported are means ±standard deviation; means followed by different letters in same line are significantly different (p<0.05) by Tukey's test \*\*\* SFA: saturated fatty acids, MUFA: monounsaturated fatty acids, PUFA: polyunsaturated fatty acids

Table 3. Seasonal variations of fatty acid composition in triacylglycerol fraction of liver tissue from female and male *S. triostegus* (% of total FA)\*.

Fatty Acids	May		July		September		November		January		March	
	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male
10:00	0.18±0.01a	0.30±0.03b	0.48±0.03b	0.11±0.01c	-	-	-	-	-	-	-	-
12:00	0.49±0.03a	0.14±0.02b	0.19±0.01c	0.16±0.05b	-	-	-	-	-	-	-	-
13:00	0.94±0.08a	0.34±0.05b	0.50±0.04c	0.33±0.06b	0.66±0.05c	0.77±0.06c	0.92±0.08a	1.95±0.19c	2.01±0.20c	2.35±0.26b	3.73±0.33a	2.96±0.30a
14:00	4.00±0.40a	2.61±0.22b	2.51±0.25b	2.69±0.26b	2.84±0.24b	4.56±0.44a	2.70±0.22b	0.31±0.03b	0.74±0.07a	0.40±0.02b	1.09±0.10a	0.70±0.06a
15:00	0.92±0.07a	0.52±0.04b	0.47±0.04b	0.35±0.03b	0.88±0.07a	1.16±0.16c	0.40±0.03b	27.42±1.30a	27.16±1.28a	26.48±1.27a	29.07±1.28a	20.22±1.27b
16:00	30.30±1.33a	21.88±1.20b	30.78±1.30a	21.30±1.19b	28.02±1.27a	28.52±1.27a	30.27±1.29a	0.16±0.01e	0.55±0.04b	0.25±0.01c	0.59±0.05b	0.61±0.05b
17:00	0.94±0.09a	0.40±0.04b	0.44±0.04b	0.27±0.01c	0.30±0.03c	0.21±0.08d	1.03±0.10a	6.55±0.56b	4.82±0.40ab	3.75±0.31c	5.19±0.51ab	4.06±0.40a
18:00	4.09±0.43a	6.69±0.65b	3.62±0.34c	5.23±0.52ab	5.41±0.54ab	7.24±0.77b	4.72±0.44a	36.39±1.44d	35.28±1.33d	33.23±1.36b	39.67±1.42a	28.55±1.29b
ΣSFA***	41.86±1.40a	32.88±1.33b	38.99±1.33a	30.44±1.30b	38.11±1.38a	42.46±1.40a	40.04±1.44a	14.00±1.14d	14.05±1.14d	15.99±1.05c	29.44±1.30a	11.09±0.99a
16:1n-7	9.01±0.07a	7.32±0.67b	10.54±1.08a	16.74±1.06c	11.62±1.10a	13.99±1.03d	15.17±1.05c	32.80±1.29b	29.60±1.34a	32.76±1.39b	29.44±1.30a	30.78±1.30ab
18:1n-9	27.56±1.28a	28.74±1.27a	26.91±1.22a	33.69±1.33b	26.81±1.25a	26.63±1.25a	26.58±1.39a	0.42±0.04d	1.74±0.20b	2.09±0.02b	1.07±0.17a	1.69±0.11b
20:1n-9	0.92±0.08a	1.23±0.10a	1.88±0.16b	1.07±0.09a	1.47±0.11c	0.83±0.07a	1.35±0.19a	47.22±1.52d	45.39±1.52d	50.84±1.56b	43.10±1.55c	43.56±1.43c
ΣMUFA	37.49±1.38a	37.29±1.33a	39.33±1.33a	51.50±1.59b	39.90±1.40a	41.45±1.44c	43.10±1.41c	2.50±0.22a	2.40±0.28a	1.93±0.19b	2.86±0.27a	5.02±0.56b
18:2n-6	3.05±0.34a	4.31±0.46b	2.83±0.22a	3.89±0.32a	2.54±0.35a	3.88±0.33a	2.57±0.29a	2.27±0.28b	1.39±0.16a	3.95±0.30d	1.27±0.18ac	2.62±0.20b
18:3n-3	1.77±0.11a	2.14±0.21b	1.07±0.16c	2.50±0.28b	1.45±0.19a	3.38±0.33d	1.07±0.20c	0.15±0.01d	0.67±0.05b	0.33±0.02a	0.83±0.07bc	1.81±0.11e
20:2n-6	0.25±0.02a	0.33±0.03a	0.50±0.04b	0.36±0.02a	1.07±0.19c	0.40±0.03b	0.42±0.04b	0.18±0.01d	0.27±0.02a	0.50±0.04b	0.61±0.05b	0.86±0.07e
20:3n-6	0.30±0.02a	0.63±0.05b	0.46±0.03a	0.43±0.04a	0.29±0.05a	0.14±0.01c	0.24±0.03a	2.31±0.30b	3.71±0.32c	2.80±0.31b	3.03±0.33b	3.92±0.38c
20:4n-6	4.53±0.44a	5.97±0.55a	5.38±0.50a	2.45±0.22b	3.68±0.36c	2.47±0.28b	2.50±0.20b	1.31±0.17c	2.35±0.22a	1.48±0.15c	1.74±0.19e	2.59±0.27a
20:5n-3	2.54±0.29a	5.73±0.55b	2.64±0.23a	1.28±0.11c	3.49±0.44d	1.75±0.10e	1.86±0.16e	0.26±0.01d	2.38±0.27a	0.94±0.07b	2.08±0.31a	4.76±0.04e
22:5n-3	2.23±0.20a	2.81±0.29a	2.32±0.22a	1.12±0.10b	1.85±0.16c	1.01±0.13b	1.30±0.15c	6.50±0.68a	6.07±0.60a	3.90±0.33d	4.76±0.45a	6.21±0.56a
22:6n-3	5.89±0.55a	7.83±0.71b	6.38±0.34a	5.93±0.56a	7.52±0.77b	2.98±0.23c	6.83±0.66a	15.48±1.05c	19.24±1.09a	15.83±1.03c	17.18±1.17c	27.79±1.22d
ΣPUFA	20.56±1.22a	29.75±1.28b	21.58±1.39a	17.96±1.07c	21.89±1.29a	16.01±1.16c	16.79±1.06c	10.34±1.00a	12.19±1.03a	10.27±1.07a	9.85±0.99a	16.18±1.10b
n-3	12.43±1.02a	18.51±1.08b	12.41±1.12a	10.83±1.10a	14.31±1.04c	9.12±0.99a	11.06±1.10a	5.73±0.51c	7.05±0.69a	5.56±0.45c	7.33±0.77a	11.61±1.03b
n-6	8.13±0.78a	11.24±1.12b	9.17±0.88a	7.13±0.88a	7.58±0.70a	6.89±0.65c	5.73±0.51c	2.01	1.72	1.84	1.34	1.39
n-3/n-6	1.52	1.64	1.35	1.51	1.88	1.32	1.93	2.01	1.72	1.84	1.34	1.39

\* Means are the averages of 3 replicates \*\* Values reported are means ±standard deviation; means followed by different letters in same line are significantly different (p<0.05) by Tukey's test \*\*\* SFA: saturated fatty acids, MUFA: monounsaturated fatty acids, PUFA: polyunsaturated fatty acids

**Table 4.** Seasonal variations of fatty acid composition in phospholipid fraction of liver tissue from female and male *S. triostegus* (% of total FA)\*.

Fatty Acids	May		July		September		November		January		March	
	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male
14:00	0.45±0.03a	1.79±0.10b	1.44±0.12b	1.15±0.11c	1.61±0.16b	1.89±0.13b	1.00±0.10c	1.47±0.16b	1.01±0.34c	1.18±0.18c	1.25±0.33c	1.41±0.19b
15:00	0.40±0.03a	0.49±0.04a	0.24±0.02b	0.26±0.02b	0.58±0.05a	0.57±0.03a	0.25±0.02b	0.11±0.01c	0.44±0.03a	0.36±0.03ab	0.43±0.02a	0.74±0.06d
16:00	22.75±1.20a	20.98±1.20a	20.34±1.19a	23.47±1.19a	24.79±1.22a	23.99±1.29a	23.79±1.28a	22.66±1.22a	20.63±1.18a	26.56±1.32b	20.31±1.10a	23.51±1.22a
17:00	0.18±0.01a	0.48±0.04b	0.15±0.02a	0.17±0.01a	0.28±0.03c	0.34±0.03b	0.40±0.04b	0.27±0.02c	0.62±0.05b	0.40±0.03b	0.85±0.07d	0.99±0.01d
18:00	11.71±1.09a	12.56±1.02a	14.74±1.14b	11.38±1.01a	11.86±1.01a	12.78±1.13a	14.43±1.09b	10.05±0.98a	13.81±1.12b	9.00±1.01a	14.68±1.17b	9.13±0.97a
ΣSFA***	35.49±1.36a	36.30±1.33a	36.91±1.39a	36.43±1.35a	39.12±1.40a	39.57±1.40a	39.87±1.33a	34.56±1.29a	36.50±1.45a	37.50±1.30a	37.52±1.37a	35.78±1.34a
16:1n-7	2.40±0.29a	3.36±0.34a	3.21±0.31a	3.07±0.32a	3.01±0.33a	3.76±0.31a	3.26±0.27a	5.58±0.54b	2.02±0.20a	4.04±0.40a	3.88±0.41a	3.32±0.32a
18:1n-9	16.01±1.17a	15.74±1.05a	17.24±1.15a	19.83±1.09b	17.05±1.08a	15.66±1.15a	14.48±1.17a	16.72±1.06a	19.16±1.09b	17.00±1.07a	15.66±1.63a	15.09±1.15a
20:1n-9	0.62±0.05a	0.90±0.09b	2.56±0.21c	1.08±0.10b	0.98±0.07b	1.31±0.11b	1.40±0.10b	0.52±0.04a	2.05±0.23c	1.03±0.15b	1.13±0.18b	1.68±0.13d
ΣMUFA	19.03±1.09a	20.00±1.20a	23.01±1.22a	23.98±1.26a	21.04±1.20a	20.73±1.19a	19.14±1.09a	22.82±1.29a	23.23±1.29a	22.07±1.21a	20.67±1.30a	20.09±1.25a
18:2n-6	1.58±0.16a	1.24±0.11a	0.79±0.07b	1.72±0.16a	1.42±0.11a	2.18±0.20c	2.58±0.28c	1.24±0.13a	1.66±0.27a	1.22±0.16a	1.39±0.16a	2.05±0.20c
18:3n-3	0.47±0.05a	0.53±0.04a	0.44±0.03a	0.47±0.03a	0.75±0.06b	1.34±0.12c	0.68±0.05b	0.22±0.02d	0.72±0.07b	0.59±0.04b	0.51±0.03b	0.78±0.06b
20:2n-6	0.42±0.04a	0.69±0.05b	0.88±0.07c	0.25±0.02d	0.73±0.05c	0.84±0.07c	0.49±0.04a	0.26±0.02d	0.53±0.04a	0.50±0.03a	0.47±0.03a	0.73±0.07c
20:3n-6	0.76±0.07a	0.83±0.05a	0.59±0.03b	0.55±0.04b	0.57±0.04b	0.71±0.06a	1.25±0.12c	0.31±0.02d	0.46±0.03b	0.75±0.06a	0.28±0.01d	0.50±0.04b
20:4n-6	12.82±1.02a	12.54±1.02a	9.03±0.99b	9.33±0.98b	9.34±0.97b	10.02±1.00b	10.53±1.13b	12.46±1.03a	7.46±0.77c	8.25±0.83bc	10.80±1.01b	8.83±0.78c
20:5n-3	5.23±0.55a	5.35±0.54a	2.08±0.22b	4.69±0.43a	4.02±0.40a	4.46±0.34a	4.55±0.34a	4.86±0.44a	5.11±0.51a	5.31±0.54a	5.28±0.50a	8.89±0.87c
22:5n-3	4.23±0.33a	4.84±0.47a	4.70±0.45a	3.00±0.31a	3.35±0.31a	4.54±0.42a	3.04±0.39a	3.39±0.33a	3.59±0.54a	2.00±0.21b	3.54±0.33a	3.64±0.35a
22:6n-3	19.89±1.09a	17.60±1.07b	21.40±1.20a	19.49±1.09a	19.60±1.20a	15.53±1.15b	17.79±1.07b	19.79±1.20a	20.69±1.27a	21.78±1.21c	19.45±1.08a	18.62±1.08a
ΣPUFA	45.40±1.46a	43.62±1.43a	39.91±1.38b	39.50±1.40b	39.78±1.35b	39.62±1.38b	40.91±1.44b	42.53±1.44a	40.22±1.39b	40.40±1.40b	41.72±1.50b	44.04±1.41a
n-3	29.82±1.22a	28.32±1.22a	28.62±1.34a	27.65±1.26a	27.72±1.28a	25.87±1.34a	26.06±1.27a	28.26±1.21a	30.11±1.31a	29.68±1.30a	28.78±1.32a	31.93±1.34b
n-6	15.58±1.05a	15.30±1.54a	11.29±1.01b	11.85±1.01b	12.06±1.12b	13.75±1.03a	14.85±1.04a	14.27±1.14a	10.11±1.10b	10.72±1.01b	12.94±1.02b	12.11±1.12b
n-3/n-6	1.91	1.85	2.53	2.33	2.29	1.88	1.75	1.98	2.97	2.76	2.22	2.63

\* Means are the averages of 3 replicates \*\* Values reported are means ± standard deviation; means followed by different letters in same line are significantly different (p<0.05) by Tukey's test \*\*\* SFA: saturated fatty acids, MUFA: monounsaturated fatty acids, PUFA: polyunsaturated fatty acids

In males, it was low in May, July, and March, and it was close to each other, while in September, November, and January it was high and close to each other. Palmitoleic acid and MUFA ratio were found to be highest in females in November and January and in males in July and January. These components decreased in both sexes in May, which is the reproductive period. The 18:1n-9 and 16:1n-7 are the most frequently detected MUFA in both sexes. In addition, 20:1n-9 were detected in both sexes, albeit in small amounts.

In the female TAG fraction, SFA was found most in May, and MUFA was found in November, January, and March. The ratio of these components in the other two months is similar. In males, MUFA was found in all months most commonly in males except for September. The component with the lowest ratio in both sexes' TAG is PUFA. It is noteworthy that MUFA ratios are higher in this fraction. In females, the PUFA ranged 16.79 to 21.89% and in males ranged from 15.48 to 29.75% through the year. The 20:4n-6 and DHA were found as major PUFA. The ratio of n-3/n-6 in the liver TAG fraction of *S. triostegus* was 1.34-1.93 in females and 1.32-2.01 in males. The highest n-3/n-6 ratio was determined in November in both sexes.

Fat-free fish store 50-80% of their fat in their livers in the form of TAG, which are good sources for fat-soluble vitamins, especially A and D vitamins (Jacquot, 1961).

In fish caught in nature, in TAG mostly monoenes, then saturated followed by PUFA are found (Henderson & Tocher, 1987). Based on this, it has been found that fish species accumulate mainly SFA and MUFA as stored lipids (Pinela et al., 2009). The reason for the high rate of monoenes in triacylglycerol is that among the components in this group, the ratio of 18:1n-9 is higher. In addition, n-3HUFAs were found to be low in this fraction since the amount of n-3 components such as 20:5n-3 and 22:6n-3 was low (Cejas et al., 2003).

Fatty acids such as palmitic acid, 16:1n-7, and 18:1n-9 are found to be excessively present in storage lipids (Ackman 1967). In the present research, MUFA were found in the liver TAG of both sexes most, then SFA followed by PUFA were found. Similar results were found by Kozlova & Khotimchenko (2000) as well. In May (reproduction period) the percentage of MUFA decreased. Their amounts may be reduced because these components are mobilized to the gonad for reproduction.

In *Comephorus baicalensis*, MUFA were found to be the most, SFA next, and then PUFA the least. Among the monounsaturated fatty acids 18:1n-9 were found most. The 16:0 was determined most within the SFA group. Other major fatty acids are 16:1n-7, 18:1n-7, and 20:5n-3 (Kozlova & Khotimchenko, 2000). Ackman et al., (2002) identified the 16:0, 18:1n-9, 16:1n-7, 18:2n-2, 18:3n-3, 20:4n-6, 20:5n-3, and 22:6n-3 as the major fatty acids. This finding is consistent with the findings of the current study as well.

#### The FA composition of PL fraction

In the liver PL fraction of *S. triostegus*, the seasonal rates of SFA was 35.49-39.87% in females and 34.56-39.57% in males, MUFA was 19.03-23.23% in females and 20.0-23.98% in males, and PUFA was 39.78-45.40% in females and 39.50-44.04% in males. As can

be seen from the data, SFA, MUFA, and PUFA percentages were found at similar rates in both sexes in all months. Similarly, the percentages of major components such as 16:0, 18:1n-9, 20:4n-6, 20:5n-3, and 22:6n-3 did not differ within months. In both sexes of *S. triostegus*, PUFA found most, SFA next, and MUFA least, in all seasons. In this fish, the ratio of n-3/n-6 was 1.75-2.97 in females and 1.85-2.76 in males. In both sexes, the n-3/n-6 rate increased in January (Table 4).

The SFA rate is reduced by exposure to low temperature, yet the proportion of unsaturated fatty acids increases (Jobling and Bendiksen 2003). The increased unsaturated fatty acids are either monoene or polyene (Wallaert and Babin, 1994, Fodor et al., 1995, Logue et al., 2000).

The increase of the ambient temperature reduces the accumulation of n-3 in PL (Delgado et al., 1994). The change in the salinity and temperature of water affects the length and the degree of unsaturation of the fatty acids in the membrane PL in poikilotherms. (Cordier et al., 2002).

In this study, it was observed that PUFA decreased in July and September when the temperature was high.

The 16:0 is dominant in SFA and its ratio in fish tissues is not affected by nutrition (Ackman et al., 1975) Ackman et al., (2002) detected that 16:0, 18:0, 18:1n-9, 16:1n-7, 20:4n-6, 20:5n-3, and 22:6n-3 were dominant in the PL fraction. The results are consistent with the results of the current study. Buzzi et al., (1997) stated that in the liver of *Exos lucius*, linolenic acid was converted into EPA and DHA. This transformation is very important in fish physiology (Arts et al., 2001).

AA was found as major n-6 PUFA in both sexes in all seasons. AA is the precursor molecule for the synthesis of thromboxane and prostaglandins (Bell et al., 1994).

In the case of migration and gonad development, the amount of EPA and DHA is mostly maintained compared to MUFA. The fact that the percentages of these two components do not change much is necessary to maintain the cell membrane structure and function (Sargent et al., 1995, Cejas et al., 2004).

#### CONCLUSION

In the present study, it was found that the amount of DHA and EPA in both sexes did not differ much by season either. PUFA was detected most in the PL fraction in both sexes. According to these results, it can be said that the lipid metabolism of the fish changes depending on the spawning and season.

**Conflict of interests:** The authors declare that they have no conflict of interest.

**Ethics committee approval:-**

**Funding:** This study was financed by the Dicle University Scientific Research Foundation (DUAPK-08-FF-07).

**Acknowledgements:-**

**Disclosure: -**

## REFERENCES

- Ackman, R.G. (1967). Characteristics of the fatty acid composition and biochemistry of some fresh-water fish oils and lipids in comparison with marine oils and lipids. *Comparative Biochemistry and Physiology*, 22,907-922. [CrossRef]
- Ackman, R.G., Eaton, C.A., & Linne, B.A. (1975). Differentiation of freshwater characteristics of fatty acids in marine specimens of the Atlantic sturgeon (*Acipenser oxyrinchus*). *Fishery Bulletin*, 73,838-845.
- Ackman, R.G., & Eaton, C.A. (1976). Fatty acid composition of the decapod shrimp, *Pandalus borealis*, in relation to that of the euphasiid, *Meganyctiphanes noruegica*. *Journal of the Fisheries Research Board of Canada*, 33,1634-1638. [CrossRef]
- Ackman, R.G., Mcleod, C., Rakshit, S., & Mısra, K.K. (2002). Lipids and fatty acids of five freshwater food fishes of India. *Journal of Food Lipids*, 9 (2),127-145. [CrossRef]
- Akpınar, M.A. (1986a). *Cyprinus carpio* L. (Osteichthyes:Cyprinidae)'nın karaciğer ve kasındaki total lipit ve total yağ asidinin mevsimsel değişimi. C.Ü. Fen Edebiyat Fakültesi Fen Bilimleri Dergisi, 4,33-42.
- Akpınar, M.A. (1986b). *Cyprinus carpio* L. (Osteichthyes:Cyprinidae)'nın karaciğer yağ asitlerinin mevsimsel değişimi. *Doğa TU Biyoloji*, 10(3),232-239.
- Akpınar, M.A., Görgün, S., & Akpınar, A.E. 2009. A comparative analysis of the fatty acid profiles in the liver and muscles of male and female *Salmo trutta macrostigma*. *Food Chemistry*, 112,6-8. [CrossRef]
- Aras, N.M., Haliloğlu, H.I., Ayık, Ö., & Yetim, H. (2003). Comparison of fatty acid profiles of different tissues of mature trout (*Salmo trutta labrax*, Pallas, 1811) caught from Kazandere Creek in the Çoruh Region, Erzurum, Turkey. *Turkish Journal of Veterinary and Animal Science*, 27,311-316.
- Arts, M.T., Ackman, R.G., & Holub, B.J. (2001). Essential fatty acids in aquatic ecosystems: A crucial link between diet and human health and evolution. *Canadian Journal of Fisheries and Aquatic Science*, 58,122-137. [CrossRef]
- Bell, J.G., Tocher, D.R., McDonald, F.M., & Sargent, J.R. (1994). Effect of supplementation with (20:3n-6), (20:4n-6) and (20:5n-3) on the production of prostaglandin-e and prostaglandin-f on the 1-series, 2-series and 3-series in turbot (*Scophthalmus maximus*) brain astroglial cells in primary culture. *Biochimica et Biophysica Acta*, 1211,335-342. [CrossRef]
- Buzzi, M., Henderson, R.J., & Sargent, J.R. (1997). The biosynthesis of docosahexaenoic acid [22:6(n-3)] from linolenic acid in primary hepatocytes isolated from wild Northern pike. *Journal of Fish Biology*, 51(1),1197-1208. [CrossRef]
- Castell, J.D., Lee, D.J., & Sinnhuber, R.O. (1972). Essential fatty acids in the diet of rainbow trout (*Salmo gairdneri*): Lipid metabolism and fatty acid composition. *The Journal of Nutrition*, 102,77-86. [CrossRef]
- Cejas, J.R., Almansa, E., Villamandos, J.E., Badia, P., Bolanos, A., & Lorenzo, A. (2003). Lipid and fatty acid composition of ovaries from wild fish and ovaries and eggs from captive fish of white sea bream (*Diplodus sargus*). *Aquaculture*, 216(1-4), 299-313. [CrossRef]
- Cejas, J.R., Almansa, E., Jerez, S., Bolanos, A., Samper, M., & Lorenzo, A. (2004). Lipid and fatty acid composition of muscle and liver from wild and captive mature female broodstocks of white seabream, *Diplodus sargus*. *Comparative Biochemistry and Physiology*, 138B,91-102. [CrossRef]
- Cordier, M., Brichon, G., Weber, J.M., & Zwingelstein, G. (2002). Changes in the fatty acid composition of phospholipids in tissues of farmed sea bass (*Dicentrarchus labrax*) during annual cycle. Roles of environmental temperature and salinity. *Comparative Biochemistry and Physiology*, 133B,281-288. [CrossRef]
- Crowford, R.H., Cusack, R.R., & Parlee, T.R. (1986). Lipid content and energy expenditure in the spawning migration of alewife (*Alosa pseudoharengus*) and blueback herring (*Alosa aestivalis*). *Canadian Journal of Zoology*, 64,1902-1907. [CrossRef]
- Delgado, A., Estevez, A., Hortelano, P., & Alejandre, M.J. (1994). Analyses of fatty acids from different lipids in liver and muscle of sea bass (*Dicentrarchus labrax* L.). Influence of temperature and fasting. *Comparative Biochemistry and Physiology*, 108A,673-680. [CrossRef]
- Fodor, E., Jones, R.H., Buda, C., Kitajka, K., Dey, I., & Farkas, T. (1995). Molecular architecture and biophysical properties of phospholipids during thermal adaptation in fish: An experimental and model study. *Lipids*, 30(12),1119-1126. [CrossRef]
- Folch, J., Lees, M., & Sladane-Stanley, G.H.A. (1957). Simple method for the isolation and purification of total lipids from animal tissues. *The Journal of Biological Chemistry*, 226, 497-509. [CrossRef]
- Geldiay, R., & Balık, S. (1996). Türkiye tatlısu balıkları. Ege Üni. Su Ürünleri Fak. Yay. No: 46 (II. Baskı), İzmir.
- Henderson, R.J., & Tocher, D.R. (1987). The lipid composition and biochemistry of freshwater fish. *Progress in Lipid Research*, 26, 281-347. [CrossRef]
- Jacquot, R. 1961. Organic constituents of fish and other aquatic animals. Fish as food. Borgstrom. G. (Ed.). Academic Press, p. 145-209. New York and London. [CrossRef]
- Jangaard, R.G., Ackman, R., & Sipos, J.C. (1967). Seasonal changes in fatty acid composition of cod liver, flesh, roe and milt lipids. *Journal of the Fisheries Research Board of Canada*, 24,613-627. [CrossRef]
- Jobling, M., & Bendiksen, E.A. (2003). Dietary lipids and temperature interact to influence tissue fatty acid compositions of Atlantic salmon, *Salmo salar* L. parr. *Aquaculture Research*, 34,1423-1441. [CrossRef]
- Kaçar, S., Başhan, M. & Oymak, S.A. (2016). Effect of seasonal variation on lipid and fatty acid profile in muscle tissue of male and female *Silurus triostegus*. *Journal of Food Science and Technology*, 53, 2913-2922. [CrossRef]
- Kaçar, S., & Başhan, M. (2017). Variations in the fatty acid compositions of the liver and gonad tissue of spiny eel (*Mastacembelus mastacembelus*) from Atatürk Dam Lake. *Turkish Journal of Biochemistry*, 42(6),617-623. [CrossRef]
- Kandemir, Ş., & Polat, N. (2007). Seasonal variation of total lipid and total fatty acid in muscle and liver of rainbow trout (*Oncorhynchus mykiss* W. 1792) reared in Derbent Dam Lake. *Turkish Journal of Fisheries and Aquatic Science*, 7,27-33.
- Kayhan, H., Başhan, M., & Kaçar, S. (2014). Seasonal variations in the fatty acid composition of phospholipids and triacylglycerols of brown trout. *European Journal of Lipid Science and Technology*, 117(5), 738-744. [CrossRef]
- Kiessling, A., Pickova, J., Johansson, L., Asgard, T., Storebakken, T., & Kiessling, K.H. (2001). Changes in fatty acid composition in muscle and adipose tissue of farmed rainbow trout (*Oncorhynchus mykiss*) in relation to ration and age. *Food Chemistry*, 73, 271-284. [CrossRef]
- Kinsella, J.E., Shimp, J.L., & Mai, J. (1977). Fatty acid content and composition of fresh water finfish. *Journal of the American Oil Chemists' Society*, 54,424-429. [CrossRef]
- Kluytmans, J.H.F.M., & Zandee, D.I. (1973). Lipid metabolism in the Northern pike (*Exos lucius* L.) I. The fatty compositions of the Northern pike. *Comparative Biochemistry and Physiology*, 44B,451-458. [CrossRef]
- Kminkova, M., Winterova, R., & Kucera, J. (2001). Fatty acids in lipids of carp (*Cyprinus carpio*) tissues. *Czech Journal of Food Science*, 19,177-181. [CrossRef]
- Kozlova, T.A., & Khotimchenko, S.V. (2000). Lipids and fatty acids of two pelagic cottoid fishes (*Comephorus spp.*) endemic to Lake Baikal. *Comparative Biochemistry and Physiology*, 126B,477-485. [CrossRef]
- Logue, J.A., DeVries, A.L., Fodor, E., & Cossins, A.R. (2000). Lipid compositional correlates of temperature-adaptive interspecific differences in membrane physical structure. *Journal of Experimental Biology*, 203,2105-2115.
- Misir, G.B., Tufan, B., & Köse, S. (2016). Variations in Total Lipid and Fatty Acid Contents of Edible Muscle, Liver, and Roes of Spotless Shad,



- Alosa immaculata*, During Catching Season in Black Sea, Journal of Aquatic Food Product Technology, 25(1),2-14. [CrossRef]
- Mute, P., Agren, J.J., Lindovist, O.V, & Hanninen, O. (1989). Fatty acid composition of vendace (*Coregonus albula* L.) muscle and its plankton feed. Comparative Biochemistry and Physiology, 92B,75-79. [CrossRef]
- Njinkoué, J-M., Barnathan, G., Miralles, J., Gaydou, E.M, & Samb, A. (2002). Lipids and fatty acids in muscle, liver and skin of three edible fish from the Senegalese coast: *Sardinella maderensis*, *Sardinella aurita* and *Cephalopholis taeniops*. Comparative Biochemistry and Physiology, 131B,395-402. [CrossRef]
- Oymak, S.A., Solak, K, & Ünlü, E. (2001). Some biological characteristics of *Silurus triostegus* Heckel, 1843 from Atatürk Dam Lake (Turkey). Turkish Journal of Zoology, 25,139-148.
- Pinela, S., Quintella, B.R., de Almeida, P.R, & Lança, M.J. (2009). Comparison of the fatty acid profile of muscle neutral lipids and phospholipids of up-river anadromous sea lamprey (*Petromyzon marinus* L.) from three Portuguese River Basins. Scientia Marina, 73,785-795. [CrossRef]
- Rincon-Sanchez, A.R., Hernandez, A., Lopez, M.L, & Mendoza-Figueroa, T. (1992). Synthesis and secretion of lipids by long-term cultures of female rat hepatocytes. Biology of the Cell, 76,131-138. [CrossRef]
- Saify, Z.S., Akhtar, S., Khan, K.M., Perveen, S., Ayattollahi, S.A.M., Hassan, S., Arif, M., Haider, S.M., Ahmad, F., Siddiqui, S, & Khan, M.Z. (2003). A study on the fatty acid composition of fish liver oil from two marine fish, *Eusphyra blochii* and *Carcharhinus bleekeri*. Turkish Journal of Chemistry, 27,251-258.
- Sargent, J.R., Bell, J.G., Bell, M.V., Henderson, R.J, & Tocher, D.R. (1995). Requirements criteria for essential fatty acids. Journal of Applied Ichthyology, 11,183-198. [CrossRef]
- Stanley-Samuels, D.W, & Dadd, R.H. (1983). Long-chain polyunsaturated fatty acids: patterns of occurrence in insects. Journal of Food Science and Technology, 13, 549-558. [CrossRef]
- Suzuki, H., Okazaki, K., Hayakawa, S., Wada, S, & Tamura, S. (1986). Influence of commercial dietary fatty acids on PUFA of cultured freshwater fish and comparison with those of wild fish of the same species. Journal of Agricultural and Food Chemistry, 34,58-60. [CrossRef]
- Tufan, B., Koral, S, & Köse, S. (2013). The Variations in Proximate Chemical Composition and Fatty Acid Profile in Different Parts of the Thornback Ray (*Raja clavata*) Caught from Black Sea, Turkey, Journal of Aquatic Food Product Technology, 22(1),83-95. [CrossRef]
- Uysal, K., Yerlikaya, A., Aksoylar, M.Y., Yöntem, M, & Ulupinar, M. (2006). Variations in fatty acids composition of pikeperch (*Sander lucioperca*) liver with respect to gonad maturation. Ecology of Freshwater Fish, 15,441-445. [CrossRef]
- Wallaert, C, & Babin, P.J. (1994). Thermal adaptation affects the fatty acid composition of plasma phospholipids in trout. Lipids, 29,373-376. [CrossRef]