

Mindfulness in the Relationship Between Perceived Stress and Quality of Life in Pediatric Asthma

Pediyatrik Astımda Algılanan Stres ve Yaşam Kalitesi Arasındaki İlişkide Bilinçli Farkındalık

Ayşe Sena Ayhan¹ , Zekeriya Deniz Aktan¹ , Zeynep Ülker Tamay² 

¹Işık University, Department of Clinical Psychology, Social Sciences Institute, İstanbul, Türkiye

²Istanbul University, İstanbul Faculty of Medicine, Department of Pediatrics, Division of Pediatric Allergy and Immunology, İstanbul, Türkiye

ORCID ID: A.S.A. 0000-0002-1192-2575; Z.D.A. 0000-0003-1757-2024; Z.Ü.T. 0000-0002-3200-5493

Citation/Atf: Ayhan AS, Deniz Aktan Z, Ülker Tamay Z. Mindfulness in the relationship between perceived stress and quality of life in pediatric asthma. Çocuk Dergisi - Journal of Child 2023;23(1):27-34. <https://doi.org/10.26650/jchild.2023.875601>

ABSTRACT

Objective: The purpose of this study is to analyze the possible mediator effect of mindfulness in the relationship between perceived stress and quality of life in pediatric asthma.

Material and Method: The sample of this study consisted of 100 asthmatic children aged between 9-12 years who applied to the outpatient clinic of Istanbul University, Istanbul Faculty of Medicine, Division of Pediatric Allergy. Sociodemographic information forms, Perceived Stress Scale in Children (8-11 years), Child and Adolescent Mindfulness Measure (CAMM), and Pediatric Asthma Quality of Life Questionnaire (PAQLQ) were used as data collection tools. Linear hierarchical regression analysis was used during the process of analyzing data.

Results: It has been identified that mindfulness has a partial mediator effect on the relationship between perceived stress and quality of life ($p = 0.000$). The presence of a partial mediator effect of mindfulness has been determined in the relationship between perceived stress and symptoms which is the subscale of quality of life ($p = 0.000$). Finally, it has been demonstrated that mindfulness has a partial mediator effect on the relationship between perceived stress and emotional function which is another subscale of quality of life ($p = 0.000$). The mediating role of mindfulness in the relationship between perceived stress and activity limitations could not be analyzed due to the lack of a significant correlation between activity limitations which is the subscale of quality of life and mindfulness ($p=0.178$).

Conclusion: It can be helpful to add psychotherapy interventions involving mindfulness practices to asthma treatment for better control of the disease in children.

Keywords: Asthma, Child, Mental Health, Mindfulness, Quality of Life

ÖZ

Amaç: Bu çalışmanın amacı pediyatrik astımda algılanan stres düzeyi ile yaşam kalitesi arasındaki ilişkide bilinçli farkındalığın olası aracı etkisini analiz etmektir.

Gereç ve Yöntem: Bu çalışmanın örneklemini İstanbul Üniversitesi İstanbul Tıp Fakültesi Hastanesi "Çocuk Alerji Polikliniği"ne başvuran 9-12 yaş arası 100 astımlı çocuk oluşturdu. Veri toplama araçları olarak sosyodemografik bilgi formları, Çocuklarda Algılanan Stres Ölçeği (8-11 yaş) (ÇASÖ), Çocuk ve Ergen Bilinçli Farkındalık Ölçeği (ÇEBFÖ) ve Pediyatrik Astım Yaşam Kalitesi Ölçeği (PAYKÖ) kullanılmıştır. Verilerin analiz edilmesi sürecinde doğrusal hiyerarşik regresyon analizi kullanılmıştır.

Bulgular: Bilinçli farkındalığın, algılanan stres ile yaşam kalitesi arasındaki ilişkide kısmi medyatör etkisinin olduğu belirlenmiştir ($p=0.000$). Algılanan stres ile yaşam kalitesinin alt boyutu olan semptomlar arasındaki ilişkide bilinçli farkındalığın kısmi medyatör etkisinin varlığı tespit edilmiştir ($p=0.000$). Son olarak bilinçli farkındalığın, algılanan stres ile yaşam kalitesinin bir diğer alt boyutu olan duygusal işlev arasındaki ilişkide kısmi medyatör etkisinin olduğu gösterilmiştir ($p=0.000$). Bilinçli farkındalık ve yaşam kalitesinin alt boyutlarından biri olan faaliyet kısıtlanması arasında korelatif ilişki bulunmadığı için bilinçli farkındalığın algılanan stres ve yaşam kalitesi arasındaki ilişkinin üzerindeki aracı etkisi analiz edilememiştir ($p=0.178$).

Sonuç: Bilinçli farkındalık uygulamaları içeren psikoterapi uygulamalarının çocuklarda astım tedavisine eklenmesi hastalığın daha iyi kontrol altına alınmasına yardımcı olabilir.

Anahtar Kelimeler: Astım, Çocuk, Ruh Sağlığı, Bilinçli Farkındalık, Yaşam Kalitesi

Corresponding Author/Sorumlu Yazar: Mahmut Caner Us E-mail: mcanerus@gmail.com

Submitted/Başvuru: 20.01.2023 • **Accepted/Kabul:** 13.02.2023 • **Published Online/Online Yayın:** 27.03.2023



This work is licensed under Creative Commons Attribution-NonCommercial 4.0 International License

1. INTRODUCTION

Asthma, characterized as a chronic airway inflammation, is the most common chronic disease in childhood.¹ Asthma often starts in childhood although it could be observed in all stages of life. The disease often presents with typical symptoms such as wheezing, chest tightness, shortness of breath, and cough.² Although there are considerable differences in asthma prevalence between countries and regions, the prevalence of childhood asthma is approximately 9.4% worldwide³ and 13.36% in Turkey.⁴

Childhood asthma attacks can be triggered by a variety of environmental factors. Recent studies have shown that psychological factors such as somatization, behavior problems, attention deficit and hyperactivity disorder (ADHD), depression, anxiety, and stress also are crucial in the development of asthma attacks.⁵⁻¹⁰ Psychological triggers such as stress have a negative impact on children's quality of life.¹¹ For instance, facing difficulties in conducting physical and social activities can make children feel different and incompetent from their peers.¹²

Children's quality of life is also affected by asthma control and poor asthma control leads to more symptoms. Asthma control means minimizing symptoms that interfere with daily living such as coughing, shortness of breath after exercise or during the night, and preventing future asthma attacks.¹³ Components such as mindfulness can be helpful in asthma control by reducing stress levels and thus can increase the quality of life of asthmatic children.

Mindfulness is the awareness arising from paying attention, intentionally, in the present moment without judgment. It has been found that there is a positive correlation between mindfulness and perceived health with physical well-being.¹⁴ A study performed in adults also showed that as mindfulness rises, perceived stress on psychological well-being decreases.¹⁴

Even though there are several studies focusing on mindfulness and asthma in adults, there are limited studies performed on children with asthma. There are some studies that separately analyze the mindfulness, perceived stress, and quality of life of people with asthma, but as far as we know, there is no study investigating the effects of all these variables in children with asthma. The purpose of this study is to evaluate if mindfulness has a mediator effect on the relationship between perceived stress and quality of life in children with asthma.

2. MATERIAL AND METHOD

This research was created in a cross-sectional design. The data collection process was carried out in January, February, and March 2020. Children and parents/guardians read and signed the informed consent form to indicate that they voluntarily participated in the study.

2.1. Population and Sample

The participants who were followed-up patients of the outpatient clinic of İstanbul University, İstanbul Faculty of

Medicine, Division of Pediatric Allergy were enrolled in the study consecutively. Asthmatic children aged between 9-12 years were included. Patients who had concomitant psychiatric diagnoses according to DSM 5 and/or using psychiatric medicine and/or receiving psychological therapy were excluded from the study.

2.3. Measures

2.3.1. Demographic Information Forms

Two demographic information forms were used in the study, one including participants' characteristics such as age, sex, and educational level, and the other including the parents'/ caregivers' characteristics.

2.3.2. Determining Asthma Control Level of The Participants

The Level of asthma control of participants was evaluated by physicians according to the "Global Initiative for Asthma" (GINA) criteria. Asthma symptom control including daily symptoms, activity limitation, night symptoms, need for rescue therapy (bronchodilator) over the last 4 weeks and pulmonary function tests (PEF and FEV1) were assessed. In addition, the patient's past asthma attacks were asked about and evaluated.

2.3.3. Perceived Stress Scale in Children (8-11 Years)

The scale was developed to assess children's perceived stress levels.¹⁵ The scale's internal consistency reliability is 0.76 and the test-retest correlation is 0.79. The Turkish version of the scale's internal consistency reliability is 0.76 and the test-retest correlation is 0.71.¹⁶ It consists of 9 questions whose answers are given on a 4-point Likert scale. The highest point means the highest perceived stress. The minimum and maximum points are 9 and 36, respectively. Although the "Perceived Stress Scale in Children" was developed for children aged between 8-11 years, it was applied to 12-year-old children in this study. The Cronbach alpha (α) internal consistency reliability of the scale in this study was found as 0.71.

2.3.4. Child and Adolescent Mindfulness Measure (CAMM)

CAMM was developed to measure the acceptance and focus of the current moment without trying to change the emotions and thoughts in children aged above 9 years old.¹⁷ The scale's Cronbach alpha (α) internal consistency coefficient is 0.80. The Turkish version of the scale was assessed in 211 adolescents aged between 14-18 years, and its internal consistency coefficient was found as 0.80.¹⁸ The English and Turkish version of the scale consists of 10 questions whose answers are scored as 0 (never true), 1 (rarely true), 2 (sometimes true), 3 (often true) and 4 (always true). High scores indicate a high level of mindfulness. The minimum and maximum points are 0 and 40, respectively. The Cronbach alpha (α) coefficient was 0.68 in this study.

2.3.5. Pediatric Asthma Quality of Life Questionnaire (PAQLQ)

The scale was developed to determine the level of educational, physical, social, and emotional difficulties experienced by

Table 1: Sample characteristics according to asthma control

Characteristics	Total	Uncontrolled	Partially-controlled	Controlled
	(n = 100) n %	(n=51) n %	(n=36) n %	(n=13) n %
Sex				
Female	31 (31)	53.8 (7)	33.3 (12)	23.5 (12)
Male	69 (69)	46.2 (6)	66.7 (24)	76.5 (39)
Age				
9	42 (42)	53.8 (7)	38.9 (14)	41.2 (21)
10	28 (28)	23.1 (3)	30.6 (11)	27.5 (14)
11	9 (9)	7.7 (1)	11.1 (4)	7.8 (4)
12	21 (21)	15.4 (2)	19.4 (7)	23.5 (12)
Total number of siblings				
1	7 (7)	0	11.1 (4)	5.9 (1)
2	56 (56)	38.5 (5)	47.2 (17)	66.7 (34)
3	22 (22)	46.2 (6)	25 (9)	13.7 (7)
4	12 (12)	15.4 (2)	11.1 (4)	11.8 (6)
5	2 (2)	0	5.6 (2)	0
6	1 (1)	0	0	2 (1)
Order of birth				
1	39 (39)	38.5 (5)	33.3 (12)	43.1 (22)
2	43 (43)	38.5 (5)	44.4 (16)	43.1 (22)
3	14 (14)	23.1 (3)	13.9 (5)	11.8 (6)
4	2 (2)	0	2.8 (1)	2 (1)
5	2 (2)	0	5.6 (2)	0
Education level of the child				
Primary school	46 (46)	53.8 (7)	44.4 (16)	45.1 (23)
Middle school	54 (54)	46.2 (6)	55.6 (20)	54.9 (28)
Grade level				
3	20 (20)	15.4 (2)	19.4 (7)	21.6 (11)
4	25 (25)	38.5 (5)	25 (9)	21.6 (11)
5	15 (15)	7.7 (1)	13.9 (5)	17.6 (9)
6	19 (19)	23.1 (3)	19.4 (7)	17.6 (9)
7	21 (21)	15.4 (2)	22.2 (8)	21.6 (11)
School type				
State	98 (98)	100 (13)	97.2 (35)	98 (50)
Private	2 (2)	0	2.8 (1)	2 (1)
Place of residence				
Istanbul	98 (98)	100 (13)	100 (36)	96.1 (49)
Yalova	2 (2)	0	0	3.9 (2)
Academic success				
Very Low	1 (1)	2.8 (1)	0	0
Low	6 (6)	2.8 (1)	7.7 (1)	7.8 (4)
Middle	25 (25)	25 (9)	23.1 (3)	25.5 (13)
High	53 (53)	55.6 (20)	53.8 (7)	51 (26)
Very High	15 (15)	13.9 (5)	15.4 (2)	15.7 (8)
Educational level (a parent/guardian)				
Illiterate	1 (1)	0	0	2 (1)
Literate	4 (4)	0	11.1 (4)	0
Primary school	40 (40)	69.2 (9)	30.6 (11)	39.2 (20)
Middle school	17 (17)	0	19.4 (7)	19.6 (10)
High School	33 (33)	23.1 (3)	38.9 (14)	31.4 (16)
Junior College	1 (1)	0	0	2 (1)
University	4 (4)	7.7 (1)	0	5.9 (3)
Master	0	0	0	0
Educational level (other parent/guardian)				
Illiterate	1 (1)	0	0	2 (1)
Literate	4 (4)	0	5.6 (2)	3.9 (2)
Primary school	47 (47)	61.5 (8)	50 (18)	41.2 (21)
Middle school	14 (14)	23.1 (3)	13.9 (5)	11.8 (6)

High School	26 (26)	15.4 (2)	22.2 (8)	31.4 (16)
Junior College	1 (1)	0	2.8 (1)	7.8 (4)
University	6 (6)	0	5.6 (2)	0
Master	1 (1)	0	0	2 (1)
Monthly Income				
0 -2500 TL	39 (39)	46.2 (6)	38.9 (14)	37.3 (19)
2501-5000 TL	54 (54)	38.5 (5)	58.3 (21)	54.9 (28)
5001-7500 TL	2 (2)	0	2.8 (1)	3.9 (2)
7501-10000 TL	5 (5)	15.4 (2)	0	3.9 (2)
Perceived Income Level				
Very Low	9 (9)	15.4 (2)	13.9 (5)	3.9 (2)
Low	24 (24)	46.2 (6)	22.2 (8)	19.6 (10)
Middle	64 (64)	30.8 (4)	63.9 (23)	72.5 (37)
High	3 (3)	7.7 (1)	0	3.9 (2)
Very High	0	0	0	0

Table 2: Relationships between CAMM, perceived stress, PAQLQ, activity limitation, symptoms, emotional function according to Pearson correlation analysis (n = 100)

Scale	CAMM	Perceived Stress	PAQLQ	Activity Limitation	Symptoms	Emotional Function
CAMM	1					
Perceived Stress	-0.501**	1				
PAQLQ	0.307**	-0.474**	1			
Activity Limitation	0.136	-0.321**	0.806**	1		
Symptoms	0.256*	-0.399**	0.94**	0.679**	1	
Emotional Function	0.381**	-0.514**	0.844**	0.598**	0.65**	1

Note: CAMM: Child and Adolescent Mindfulness Measure, PAQLQ: Pediatric Asthma Quality of Life Questionnaire, *p<.05: statistically significant, **p<.01: statistically significant

children aged 7-17 years with asthma in their daily life.¹⁹ It consists of 23 items and some of them are answered on a 7-point Likert scale. The scale has 3 subscales consisting of activity limitation, symptoms, and emotional function. The scores range from 23 to 161. A high score shows a high level of life quality, whereas a low score indicates a low level of life quality. The Cronbach alpha (α) value of the quality of life scale was 0.89. Additionally, the Cronbach alpha (α) coefficient of activity limitation was 0.75, the Cronbach alpha (α) value of symptoms was 0.76 and finally, the Cronbach alpha (α) coefficient of emotional function subscale was 0.87. The Turkish validity and reliability study was conducted on 122 children between the ages of 7-16 years and the scale was found as valid and reliable.²⁰ The Turkish version of the scale's Cronbach alpha (α) internal consistency coefficient of the activity limitation subscale, symptoms subscale, and emotional function subscale was 0.80, 0.90, and 0.86, respectively.

2.4. Ethics

Ethics Committee Approval was obtained from Işık University on 02.09.2019. Işık University Clinical Psychology Graduate Coordination Office has given permission to conduct the research at the outpatient clinic of İstanbul University, İstanbul Faculty of Medicine, Division of Pediatric Allergy.

2.5. Statistical Analysis

Data were analyzed with SPSS version 23. Firstly, correlational relationships between variables were examined by Pearson correlation analysis. In the next step, regression analysis was conducted to control if there is a significant mediator effect in correlative relationships. The mediator variable analysis included four considerations.²¹ These four considerations were as follows: the effect of the independent variable on the dependent variable

Table 3: Assessment of mindfulness in relationship between perceived stress and quality of life by linear hierarchical regression analysis

Model		B	β	R2	F	p
1	The effect of PS on PAQLQ	-3.012	-0.474	0.225	28.387	0.000**
2	The effect of PS on CAMM	-0.651	-0.501	0.251	32.813	0.000**
3	The effect of CAMM on PAQLQ	1.499	0.307	0.094	10.187	0.002*
4	When controlling CAMM, the effect of PS on PAQLQ	-2.717	-0.427	0.231	14.573	0.000**

Note: PS: Perceives Stress, CAMM: Child and Adolescent Mindfulness Measure, PAQLQ: Pediatric Asthma Quality of Life Questionnaire, *p<.01: statistically significant, **p<.001: statistically significant

Table 4: Assessment of mindfulness in relationship between perceived stress and symptoms by linear hierarchical regression analysis

Model		B	β	R2	F	p
1	The effect of PS on S	-1.454	-0.399	0.159	18.572	0.000**
2	The effect of PS on CAMM	-0.651	-0.501	0.251	32.813	0.000**
3	The effect of CAMM on S	0.717	0.256	0.065	6.86	0.010*
4	When controlling CAMM, the effect of PS on S	-1.318	-0.362	0.163	9.479	0.000**

Note: PS: Perceived Stress, S: Symptoms, CAMM: Child and Adolescent Mindfulness Measure, *p<.05: statistically significant, **p<.001: statistically significant

and the mediator variable should be significant; the effect of the mediator variable on the dependent variable should also be significant. Finally, while controlling the effect of the mediator variable, the significant effect of the independent variable on the dependent variable should disappear. These four aspects were checked, and all analyses were carried out following the linear hierarchical regression analysis steps.

In the last step, the relationship between perceived stress, mindfulness, and quality of life according to asthma control level was examined using 3 different one-way ANOVA analyses.

3. RESULTS

3.1. Demographic data of asthmatic children and parents/caregivers

A total of 100 (31 female / 69 male) children with a mean age and SD of $10 \pm 1,16$ years were included. Of the participants, 51% had controlled asthma according to GINA guidelines.

3.2. The mediation analysis of CAMM in the relationship between perceived stress and PAQLQ

Correlational relationships between the variables were given in Table 2. The correlation between perceived stress and mindfulness ($p<0.01$), quality of life and mindfulness ($p<0.01$), quality of life and perceived stress ($p<0.01$), activity limitation and perceived stress ($p<0.01$), activity limitation and quality of life ($p<0.01$), symptoms and mindfulness ($p<0.05$), symptoms and perceived stress ($p<0.01$), symptoms and quality of life ($p<0.01$), symptoms and activity limitation ($p<0.01$), emotional function and mindfulness ($p<0.01$), emotional function and perceived stress ($p<0.01$), emotional function and quality of life ($p<0.01$), emotional function and activity limitation ($p<0.01$) and finally emotional function and symptoms were significant.

In the first mediation analysis, Linear regression analysis showed that mindfulness did not have a full mediator effect on the relationship between perceived stress and quality of life (Table 3). However, according to Sobel Test analysis, mindfulness has a partial mediator effect on the relationship between perceived stress and quality of life (Sobel z-statistics=2.81; one-tailed=0.000; two-tailed=0.000).

3.3. The mediation analysis of CAMM in the relationship between perceived stress and symptoms

In the second mediation analysis, Linear regression analysis determined that mindfulness did not have a full mediator effect on the relationship between perceived stress and symptoms (Table 4). However, Sobel Test analysis demonstrated that mindfulness had a partial mediator effect on the relationship between perceived stress and symptoms (Sobel z-statistics=-2.38; one-tailed=0.000; two-tailed=0.010).

3.4. The mediation analysis of CAMM in the relationship between perceived stress and emotional function

In the final mediation analysis, Linear regression analysis determined that mindfulness did not have a full mediator effect on the relationship between perceived stress and emotional function (Table 5). However, Sobel Test analysis demonstrated that mindfulness had a partial mediator effect on the relationship between perceived stress and emotional function (Sobel z-statistics=-2.93; one-tailed=0.000; two-tailed=0.000).

3.5. Relationship between asthma control level, perceived stress, CAMM and PAQLQ

One way-ANOVA analysis showed that quality of life significantly differed by asthma control levels ($F_{(2,97)}=11.03$; $p=0.000$) (Table 6). Post-Hoc test determined that quality of life significantly

Table 5: Assessment of mindfulness in relationship between perceived stress and emotional function by linear hierarchical regression analysis

Model		B	β	R2	F	p
1	The effect of PS on EF	-1.131	-0.514	0.265	35.266	0.000*
2	The effect of PS on CAMM	-0.651	-0.501	0.251	32.813	0.000*
3	The effect of CAMM on EF	0.644	0.381	0.145	16.650	0.000*
4	When controlling CAMM, the effect of PS on EF	-0.949	-0.432	0.285	19.329	0.000*

Note: PS: Perceived Stress, EF: Emotional Function, CAMM: Child and Adolescent Mindfulness Measure, *p<.001: statistically significant

Table 6: Comparison of perceived stress, CAMM and PAQLQ according to asthma control level

Scale	Uncontrolled Group (n=13)		Partially-Controlled Group (n=36)		Controlled Group (n=51)		p
	Mean	SD	Mean	SD	Mean	SD	
PS	18.85	3.63	16.78	5.13	15.55	4.52	0.068
CAMM	26.08	6.44	25.56	5.72	28.18	6.16	0.127
PAQLQ	85.31	24.58	112.33	25.17	124.8	30.1	0.000**
AL	15.54	5.01	21.31	4.82	23.53	6.57	0.000**
S	31.69	14.77	47.86	12.70	53.90	17.92	0.000**
EF	38.08	10.43	43.17	10.63	47.37	9.43	0.008*

Note: PS: Perceived Stress, CAMM: Child and Adolescent Mindfulness Measure, PAQLQ: Pediatric Asthma Quality of Life Questionnaire, AL: Activity Limitation, S: Symptoms, EF: Emotional Function, *p<.05: statistically significant, **p<.01: statistically significant

differed between the uncontrolled group and partially controlled group ($p=0.003$), between the uncontrolled group and controlled group ($p=0.000$), and between the partially controlled group and controlled group ($p=0.039$).

There was a significant relationship between asthma control level and activity limitation which is a subscale of quality of life ($F_{(2,97)}=9.94$; $p=0.000$) (Table 6). According to Post-Hoc analysis, this significance was raised from the relationship between the uncontrolled group and the partially controlled group ($p=0.003$) and the relationship between the uncontrolled group and the controlled group ($p=0.000$).

Symptoms also significantly differed by asthma control level ($F_{(2,97)}=10.3$; $p=0.000$) (Table 6). The relationship between the uncontrolled group and the partially controlled group ($p=0.002$) and the relationship between the uncontrolled group and the controlled group ($p=0.000$) made this difference (Table 6).

There was a significant relationship between asthma control level and emotional function ($F_{(2,97)}=5.09$; $p=0.008$) (Table 6). According to Post-Hoc analysis, the difference occurred because of the relationship between the uncontrolled group and the controlled group ($p=0.004$).

4. DISCUSSION

In this study, the authors determined that mindfulness had a partial mediator effect on the relationship between perceived stress and the quality of life, on the relationship between perceived stress and activity limitation, and on the relationship between perceived stress and emotional function. However, there was no significant correlational relationship between mindfulness and symptoms, which is a subscale of the quality of life.

Mindfulness-based practices improve the psychological and physiological health of children and adolescents.²² Mindfulness positively affects health by reducing the stress of chronic diseases.²³ In a study conducted on college students with a mean age of 18.3 years, it was demonstrated that people with a higher level of mindfulness are less likely to have had an asthma diagnosis and less likely to have the symptoms of persistent dry cough and wheezing.²⁴ Besides, a randomized control trial

held in asthmatic adults showed that an 8-week mindfulness-based stress reduction (MBSR) group training program, which has a specific impact on emotional functionality improved the quality of life of patients.²⁵ Another MBSR program, when used as supplementary to routine medical treatments, improved the quality of life of patients with asthma.²⁶ Mindfulness practices regulate physiological stimulations via regulating breathing and slowing heartbeats, hence having a positive impact on the management of the disease.^{27, 28} The reduction in perceived stress, which is associated with mindfulness correlates with the increase in the quality of life of patients with asthma.⁸

Although a significant relationship between mindfulness and activity limitation was shown in some studies²⁵, we could not find such a significant correlation between mindfulness and activity limitation. In our study group, half of the children (51%) had controlled asthma, nearly one-third (36%) had partially controlled asthma, and only 13% of children had uncontrolled asthma. The majority of our patients not having activity limitations may partly explain this discrepancy. Furthermore, the effects of seasonal variations on children's physical activity might be a confounding factor. There are studies showing that asthma control levels improve due to the low activity rates of people with asthma in winter. For example, adults with asthma may think that their asthma control levels are better in winter because they restrict their activities in winter compared to the summer seasons.²⁹ The data collection process of our study coincided with the months of January, February and March. As our participant group attends to school during the winter months, school time coincided with most daily activities.

Asthma symptoms positively correlate with the stress level of asthmatic people.³⁰ There is an inverse relationship between asthma symptoms and mindfulness. Asthma symptoms decrease in parallel with increased mindfulness.²⁴ In our study, it was shown that mindfulness in pediatric asthma had a partial effect on the relationship between perceived stress and symptoms.

People with asthma decrease their emotional functionality due to inefficient coping with stress.³¹ Conversely, it has been observed that the increase in mindfulness levels affects emotional functionality.³² In our study, it was determined that

mindfulness had a partial effect on the relationship between perceived stress and emotional function.

Demographic characteristics of the sample according to asthma control were given in Table 1. In our study, we observed that the perceived stress does not differ according to the asthma control levels, but the perceived stress level decreases hierarchically with the increase of the asthma control level. Literature has shown that the level of stress varies according to asthma control and, consistent with our research, low asthma control, and excessive stress are associated.^{33, 34} In a study, perceived stress increased in direct proportion with both the incidence of asthma and asthma severity.³⁵

Mindfulness level did not change according to asthma control level. Similarly, it has been observed that asthma control has no effect on mindfulness scores in adolescents.³³

Quality of life, activity limitation, symptoms, and emotional function differed according to asthma control. When the level of asthma control increase, the quality of life and emotional function also increase; besides, activity limitation, symptoms scores decreased. A similar study found that when asthma control decreases, quality of life decreases as well.³⁶ People with uncontrolled asthma had a higher activity limitation.³⁷ According to GINA criteria, there is a negative correlation between asthma control and asthma symptoms.³⁸ Moreover, aggression, which is one of the emotional factors known that influence emotional functionality, has been found to decrease with increasing asthma control.³⁹

Our research has some limitations. Compared to other studies, the number of participants was limited, and 70% of the sample was 9 or 10 years old. The study design was cross-sectional. Therefore, mindfulness may not have a full mediator effect on the relationships in question. Although perceived stress decreases when asthma control increases, this model was not significant. It may also be related to the low number of participants. Thus, further research might investigate the same relationships with a higher number of participants and the design might be longitudinal. Moreover, studying activity limitation via equally asthma control group subclasses might occur crucial consequences. It may also be advisable to add a control group to the study to make the results more reliable on a scientific scale.

CONCLUSION

In conclusion, our study found that there is a relationship between perceived stress, mindfulness and the quality of life in pediatric asthma. Our results suggest that mindfulness has a partial mediator impact on the relationship between perceived stress, and quality of life, on the relationship between perceived stress and activity limitation which is the subscale of quality of life, and on the relationship between perceived stress and emotional function which is the quality of life's subscale. Adding mindfulness-based practices to the current medical treatments might reduce perceived stress and increase the quality of life via reducing activity limitation, and increasing emotional function.

Hakem Değerlendirmesi: Dış bağımsız.

Yazar Katkıları: Çalışma Konsepti/Tasarım- Y.Ö.U., M.C.U.; Veri Toplama- Y.Ö.U., M.C.U.; Veri Analizi/Yorumlama- M.C.U.; Yazı Taslağı- M.C.U.; İçeriğin Eleştirel İncelemesi- Y.Ö.U.; Son Onay ve Sorumluluk- Y.Ö.U., M.C.U.

Çıkar Çatışması: Yazarlar çıkar çatışması beyan etmemişlerdir.

Finansal Destek: Yazarlar finansal destek beyan etmemişlerdir.

Peer Review: Externally peer-reviewed.

Author Contributions: Conception/Design of Study- Y.Ö.U., M.C.U.; Data Acquisition- Y.Ö.U., M.C.U.; Data Analysis/Interpretation- M.C.U.; Drafting Manuscript- M.C.U.; Critical Revision of Manuscript- Y.Ö.U.; Final Approval and Accountability- Y.Ö.U., M.C.U.

Conflict of Interest: Authors declared no conflict of interest.

Financial Disclosure: Authors declared no financial support.

REFERENCES

1. Masoli, M, Fabian, D, Holt, S, Beasley, R. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy*, 2004, 59.5: 469-478.
2. The Global Initiative for Asthma (GINA). Global strategy for asthma management and prevention. 2018 Update.
3. Lai, CK, Beasley, R, Crane, J, Foliaki, S, Shah, J, Weiland, S, & ISAAC Phase Three Study Group. Global variation in the prevalence and severity of asthma symptoms: phase three of the international study of asthma and allergies in childhood (ISAAC). *Thorax* 2009; 64(6), 476-483.
4. Kurt, E, Metintas, S, Başıyigit, I, Bulut, I, Coşkun, E, Dabak, S, et al. Prevalence and risk factors of allergies in Turkey: Results of a multicentric cross-sectional study in children. *Pediatric Allergy and Immunology* 2007; 18(7), 566-574.
5. Blackman, JA, & Gurka, MJ. Developmental and behavioral comorbidities of asthma in children. *Journal of Developmental & Behavioral Pediatrics* 2007; 28(2), 92-99.
6. Grizenko, N, Osmanliu, E, Fortier, MÈ, & Joober, R. Increased risk of asthma in children with ADHD: Role of prematurity and maternal stress during pregnancy. *Journal of the Canadian Academy of Child and Adolescent Psychiatry* 2015; 24(2), 109.
7. Katon, W, Lozano, P, Russo, J, McCauley, E, Richardson, L, & Bush, T. The prevalence of DSM-IV anxiety and depressive disorders in youth with asthma compared with controls. *Journal of Adolescent Health* 2007; 41(5), 455-463.
8. Kimura, T, Yokoyama, A, Kohno, N, Nakamura, H, & Eboshida, A. Perceived stress, severity of asthma and quality of life in young adults with asthma. *Allergology International* 2009; 58(1), 71-79.
9. Ramratnam, S, Jaffee, K, Visness, C, & Gern, JE. (2017). Exposure to early life maternal stress and depression and risk of asthma. In C26. Asthma in infants and children. *American Thoracic Society* 2009; A5088-A5088.
10. Rona, RJ, Smeeton, NC, Amigo, H, & Vargas, C. Do psychological distress and somatization contribute to misattribution of asthma? A Chilean study. *Journal of Psychosomatic Research* 2007; 62(1), 23-30.

11. Bellin, MH, Kub, J, Frick, KD, Bollinger, ME, Tsoukleris, M, Walker, et al. Stress and quality of life in caregivers of inner-city minority children with poorly controlled asthma. *Journal of Pediatric Health Care* 2013; 27(2), 127-134.
12. Boran, P, Tokuç, G, Pişgin, B, & Öktem, S. Assessment of quality of life in asthmatic Turkish children. *Turkish Journal of Pediatrics* 2008; 50(1).
13. Türk Toraks Derneği, astım tanı ve tedavi rehberi güncellemesi. *Turkish Thoracic Journal* 2016.
14. Kabat-Zinn, J. Mindfulness-based interventions in context: Past, present and future. *Clinical Psychology: Science and Practice* 2003; 10(2), 144-156.
15. Snoeren, F, & Hoefnagels, C. Measuring perceived social support and perceived stress among primary school children in the Netherlands. *Child Indicators Research* 2014; 7(3), 473-486.
16. Oral, T & Ersan, C. Çocuklarda (8-11 yaş) Algılanan Stres Ölçeği'nin Türkçeye uyarılma çalışması. *Selçuk Üniversitesi Edebiyat Fakültesi Dergisi* 2017; (37), 419-428.
17. Greco, L, Baer, RA, & Smith, GT. Assessing mindfulness in children and adolescents: Development and validation of the child and adolescent mindfulness measure (CAMM). *Psychological Assessment* 2011; 23(3), 606-614.
18. Sünbül, ZA. Psychometric evaluation of child and adolescent mindfulness measure (CAMM) with Turkish sample. *Online Submission* 2018; 7(2), 56-59.
19. Juniper, EF, Guyatt, GH, Feeny, DH, Ferrie, PJ, Griffith, LE, & Townsend, M. Measuring quality of life in children with asthma. *Quality of life research* 1996; 5(1), 35-46.
20. Yüksel, H, Yılmaz, O, Kırmaz, C, & Eser, E. Validity and reliability of the Turkish translation of the pediatric asthma quality of life questionnaire. *The Turkish Journal of Pediatrics* 2009; 51(2), 154.
21. Baron, RM, & Kenny, DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic and statistical considerations. *Journal of Personality and Social Psychology* 1986; 51, 1173-1182.
22. Bränström, R, Duncan, LG, & Moskowitz, JT. The association between dispositional mindfulness, psychological well-being and perceived health in a Swedish population-based sample. *British Journal of Health Psychology* 2011; 16(2), 300-316.
23. Creswell, JD, & Lindsay, EK. How does mindfulness training affect health? A mindfulness stress buffering account. *Current Directions in Psychological Science* 2014; 23(6), 401-407.
24. Shi, L, Liang, D, Gao, Y, Huang, J, Nolan, C, Mulvaney, A, et al. Mindfulness and asthma symptoms: A study among college students. *Journal of Asthma* 2018; 55(1), 101-105.
25. Pbert, L, Madison, JM, Druker, S, Olenzki, N, Magner, R, Reed, G, et al. Effect of mindfulness training on asthma quality of life and lung function: A randomised controlled trial. *Thorax* 2012; 67(9), 769-776.
26. Nadiri, M, & Khanpour, F. The effect of teaching mindfulness-based stress reduction on the quality of life of patients with asthma. *Depiction of Health* 2019; 10(2), 144-156.
27. Delgado, LC, Guerra, P, Perakakis, P, Vera, MN, Reyes del Paso, G, & Vila, J. Treating chronic worry: Psychological and physiological effects of a training programme based on mindfulness. *Behaviour Research and Therapy* 2010; 48(9), 873-882.
28. Vlemincx, E, Vigo, D, Vansteenwegen, D, Van den Bergh, O, & Van Diest, I. Do not worry, be mindful: Effects of induced worry and mindfulness on respiratory variability in a nonanxious population. *International Journal of Psychophysiology* 2013; 87(2), 147-151.
29. Bacon, SL, Lemiere, C, Moullec, G, Ninot, G, Pepin, V, & Lavoie, KL. Association between patterns of leisure time physical activity and asthma control in adult patients. *BMJ Open Respiratory Research* 2015; 2(1).
30. Luria, CJ, Sitarik, AR, Havstad, S, Wegienka, GR, Kim, H, Zoratti, EM, et al. Association between asthma symptom scores and increased perceived stress and trait anxiety in asthmatic adolescents. *Journal of Allergy and Clinical Immunology* 2016; 137(2).
31. Lewandowska, K, Specjalski, K, Jassem, E, & Słomiński, JM. Style of coping with stress and emotional functioning in patients with asthma. *Advances in Respiratory Medicine* 2009; 77(1), 31-36.
32. Borders, A, Earleywine, M, & Jajodia, A. Could mindfulness decrease anger, hostility and aggression by decreasing rumination? *Aggressive Behavior: Official Journal of the International Society for Research on Aggression* 2010; 36(1), 28-44.
33. Cillessen, L, van de Ven, MO, & Karremans, J. C. The role of trait mindfulness in quality of life and asthma control among adolescents with asthma. *Journal of Psychosomatic Research* 2017; 99, 143-148.
34. Kishore, N. The impact of severe housing stress on child asthma control and pediatric asthma caregiver quality of life (PACQOL) (Doctoral Dissertation). 2014.
35. Rod, NH, Kristensen, TS, Lange, P, Prescott, E, & Diderichsen, F. Perceived stress and risk of adult-onset asthma and other atopic disorders: A longitudinal cohort study. *Allergy* 2012; 67(11), 1408-1414.
36. Li, Z, Thompson, LA, Gross, HE, Shenkman, EA, Reeve, BB, DeWalt, DA, et al. Longitudinal associations among asthma control, sleep problems and health-related quality of life in children with asthma: A report from the PROMIS® pediatric asthma study. *Sleep Medicine* 2016; 20, 41-50.
37. Haselkorn, T, Jones, CA, Chen, H, Rhoa, M, Miller, DP, Peters, SP, et al. Relationship between asthma control and activity limitation: Insights from the REACT study. *Journal of Allergy and Clinical Immunology* 2008; 121(2), S160.
38. The Global Initiative for Asthma (GINA). A pocket guide for health professionals (Updated 2019) 2019.
39. Coban, H, & Aydemir, Y. The relationship between allergy and asthma control, quality of life and emotional status in patients with asthma: A cross-sectional study. *Allergy, Asthma & Clinical Immunology* 2014; 10(1), 67.