



Investigation of cytotoxic and genotoxic effects of olive leaf extract on colon cancer cells and normal cell lines

Emre Öztürk¹, Fatma Çalık^{1*}, Derya Ulusoy¹

¹Department of Molecular Biology and Genetics, Erzurum Technical University, 25050 Erzurum, Turkey.

Cite: Öztürk E, Çalık F, Ulusoy D. Investigation of cytotoxic and genotoxic effects of olive leaf extract on colon cancer cells and normal cell lines. *Eurasian Mol Biochem Sci* 2022;1(2): 25-30.

Received: 19 September 2022, Accepted: 22 October 2022

DOI: 10.54672/ejms.2022.11

Abstract

Colon cancer is a type of cancer that occurs when cells in the mucous layer membrane that surrounds the inner surface of the large intestine sections multiply unevenly. This study aimed to investigate olive leaf extract's cytotoxic and genotoxic effects on HTC-116 colon cancer cell lines and HDF healthy cell lines. In the study, olive leaf extract was prepared in methanol. The passaged HTC-116 and HDF cell lines were then incubated in Dulbecco's Modified Eagle's Medium (DMEM). Olive leaf extract extracted in methanol was applied to these cell lines at 3, 10, 20, 50, and 80 µg/ml and left for incubation for 24 hours. Then, the cytotoxic effect was determined by MTT analysis. The genotoxic effect of olive leaf extracts applied to HTC-116 and HDF cell lines after a 24-hour incubation period was determined by Hoechst stain. DNA damage was visualized with a DAPI filter on a camera-attached trinocular fluorescence microscope 30 minutes after staining with Hoechst. The MTT analysis revealed that the cytotoxic effects of olive leaf extract applied to the HTC-116 cell line were close to each group after the 24-hour incubation period. As a result of the examination of the genotoxic activity with Hoechst DNA staining in colon cancer cells (HCT-116), olive leaf extract at various concentrations treated to the cell line had a minor genotoxic effect at low doses (3, 10, and 20 µg/ml), while at high concentrations (50 and 80 µg/ml), the genotoxic effects of the applications were determined. In examining the genotoxic impact of Hoechst DNA staining in the HDF cell line, no genotoxic effect was observed at low doses (3, 10, and 20 µg/ml) of olive leaf extract applied to the cell line at various doses. The little genotoxic effects of high doses (50 and 80 µg/ml) were also observed. Olive leaf contains phenolic substances and has antioxidant, antimicrobial, anti-inflammatory, antiatherogenic, anticarcinogenic and antiviral effects. **Keywords:** In vitro culture, plant regeneration, callus, hormones.

Keywords: Olive leaf extract, colon cancer, cytotoxicity, genotoxicity

Introduction

Cancer is a disease that involve the proliferation of cell that have lost their genetic stability and cell cycle

control through mutation [1]. More than 14 million new cancer cases were reported worldwide in 2012 [2], raising a need to further develop treatments and preventive strategies. Cancer is a disease that mainly occurs with aging, so this disease related to better living conditions and increased life expectancy will continue

* Corresponding Author: Fatma ÇALIK,
Department of Molecular Biology and Genetics,
Erzurum Technical University, 25050 Erzurum, Turkey.
E-mail: fatma.calik25@erzurum.edu.tr



with increasing prevalence today. Today, there are factors other than age that contribute to the development of cancer. There is strong scientific support that the traditional Mediterranean diet (MD) protects against some cancers [3,4]. MD plays a vital role in reducing age-related and non-communicable diseases such as cancer with its anti-inflammatory activity. However, the effects of MD on inflammation have not been fully elucidated [5,6]. MD has shown a protective role in cancer as a whole [6]. However, it is important to understand whether any beneficial effects attributed to MD are due to a particular component of the diet rather than the entire diet. As an example, polyphenol bioactive compounds have shown particular promise.

Olive tree leaves (*Olea europaea*) are an alternative medicine agent widely used in traditional medicine in the Mediterranean region [7]. The bioactive properties of this Leaf have formed a basis for its use as an antioxidant, anti-hypertensive, anti-atherogenic, anti-inflammatory, hypoglycemic and hypocholesterolemic agent [7]. As a result, olive leaf extract may have more significant potential than Extra virgin olive oil (EVOO) for improving health outcomes. During EVOO processing, leaves may be unintentionally left in the mix if separation methods are inadequate [8]. OLE components not detected in Olive Fruit oil include flavonoids such as luteolin and apigenin, which show anti-cancer properties [9,10].

In this study, we investigated the effect of olive leaf extract on human colon cancer cells. It was aimed to investigate the cytotoxic and genotoxic effects of olive leaf extract on HT-116 cell line and Human Dermal Fibroblast (HDF) cell lines.

Materials and Methods

Chemicals: HTC-116 and HDF cell lines were taken from YUTAM cancer laboratories and used for examinations. DMEM (Dulbecco's Modified Eagle's Medium), PBS, Penicillin, and L-glutamine were used

in cell passages and treatment of oil leaf extract. Trypsin EDTA and PBS were used for the passage of cells.

Oil Leaf extract preparation: The leaves were ground with a grinder. They were then left to dissolve in methanol, and extraction was completed. Then, a rotary evaporator separated the extract from the solvent (Laborota 4001, Heidolph). The temperature of the water bath in the rotary evaporator was set at 40°C and the rotation frequency at 60 rpm. The pressure in the condenser was adjusted with a vacuum valve to evaporate the solvent faster. The evaporating solvent was concentrated via a condenser. Finally, olive leaf extract was obtained to be used in future studies.

Cell viability analysis: The viability of Oil leaf extract treated cells was measured using flow cytometry according to the manufacturer's instructions (Ecotech Biotechnology, Türkiye). HCT-116 and HDF cells were seeded in 96-well plates at 1000 cells per well and treated for 24h at concentrations of 3, 10, 20, 50, and 80 µg/mL of Oil leaf extract diluted in the medium. After incubation for the indicated time, the MTT solution has added to the wells and allowed to be incubated for at least 3 hours in darkness. The cell number and viability were measured by the colorimetric absorbance of cells (Thermo Fischer, USA). The absorbance values were used for the cell viability assay.

Genotoxicity analysis: For preparing Hoechst dye, 20 µl of Hoechst stock dye was added to 50 ml of PBS. For Hoechst staining, the HCT-116 and HDF cell lines were seeded in 24 well-plates and the concentration of Oil leaf extract 3, 10, 20, 50, and 80 µg/mL incubated the cells and then, the cells were fixed with 4% paraformaldehyde in phosphate-buffered saline at 4 °C for 30 min. the cells were washed with phosphate-buffered solution (PBS), and samples were incubated with 1 mM Hoechst 33258 fluorescent dye (Sigma-

Aldrich®, USA) for 5 min at room temperature. Nuclear abnormalities were observed under the fluorescent microscope (Leica® DM IL LED, excitation/emission wavelength = 365/420 nm) on a total of 1000 cells in each well. The scored nuclear alterations (NAs) were divided into the following categories: lobed (L), notched (N), and micronuclei (MN) [11].

Results

Cell viability results: MTT analysis was performed to observe whether olive leaf extract has cytotoxic effects on both healthy human fibroblast cell line and colon cancer cell line. Cells were treated with olive leaf extract after 24 hours of incubation.

Cytotoxicity was determined after incubation with the olive leaf extract at concentrations of 3, 10, 20, 50 and 80 µg/ml to colon cancer cells (HCT-116) and human dermal fibroblast cells (HDF) for 24 hours. While it was observed that the cytotoxic effects of olive leaf extract concentrations were close to each other in these applications, the most effective cytotoxicity value was found in the group administered at a dose of 20 µg/ml. It was determined that the cytotoxicity of 50 and 80 µg/ml concentrations was less than the other concentrations. To determine the cytotoxic activity of olive leaf extract in normal cells, olive leaf extract was applied to HDF cells for 24 hours. After the incubation period, no cytotoxic effect was detected in any of the doses of olive leaf extract in the MTT analysis. However, it was determined that olive leaf extract at a concentration of 20 µg/ml showed a proliferative effect (Table 1, Fig 2 and 3).

Table 1. MTT (Cell viability) values after 24 hours of incubation with olive leaf extract treated to HCT-116 and HDF cell lines

Group	HCT-116		HDF	
	Mean	S.D.	Mean	S.D.
Control	0.174	0.010	0.175	0.050
3 µg/mL	0.154	0.006	0.174	0.022
10 µg/mL	0.157	0.015	0.187	0.077
20 µg/mL	0.153	0.008	0.203	0.006
50 µg/mL	0.162	0.016	0.193	0.032
80 µg/mL	0.160	0.011	0.179	0.036



Figure 1. Process of Olive leaf methanol extract preparation; 1: Olive leaves, 2: Homogeneous extract, 3: Evaporator device, 4: Planning by scraping the dried extract, 5: Extract transferred to test tube.




Figure 2. Illustration of inverted Microscope after the application of olive leaf extract at various concentrations to HCT-116 cell lines.




Figure 3. Illustration of inverted Microscope after the application of olive leaf extract at various concentrations to HDF cell lines.

Genotoxic analysis: In evaluating genotoxic activity with Hoechst DNA staining, a small amount of genotoxic effect of olive leaf extract applied to colon cancer cell line at various concentrations was determined at low doses. In contrast, the genotoxic effects of applications at high concentrations (50 and 80 µg/ml) were determined. In the evaluation of genotoxic activity, no genotoxic effect was observed at

low doses of olive leaf extract at various doses applied to the human dermal fibroblast cell line. In contrast, genotoxic effects were determined at high doses (50 and 80 µg/ml) (Fig. 4 and 5).




Figure 4. Fluorescent microscope images of the HDF cell line after hoechst dye application




Figure 5. Fluorescent microscope images of the HCT-116 cell line after hoechst dye application

Discussion

Colon cancer is the third most common type of cancer in the world and in Turkey, and ranks third in cancer-related deaths. It is known that most of the colon cancer starts with the abnormal growth of healthy epithelial cells in the mucous layer of the colon or rectum [12]. In this study, it was aimed to investigate the cytotoxic and genotoxic effects of olive leaf extract on colon cancer cells.

The emergence of colon cancer arises with the interaction of many hereditary, environmental and genetic factors. For these reasons, the susceptibility to mutations and excessive consumption of red meat, insufficient intake of vitamins, bile acids, and mineral intakes, HNPCC is a hereditary type of colon cancer observed at young ages (before the age of 50) and caused an average of 4% of all colon cancer types. This variety is considered genetically predisposed [13]. According to previous research, one of the most important factors in the emergence of colon cancer is the presence of a family history of colon cancer. More than 90% of colon cancer cases were determined accidentally, without any particular symptoms. In the remaining colon cancer cases, it has been determined that there is a high degree of genetic factors. It is known that many such critical genetic factors come together and cause colon cancer [13].

In this study, we examined the effects of olive leaf extract on cancer cell viability in the treatment of colon cancer. As a result of the study showed that olive leaf extract showed cytotoxic effects at 20 µg/ml but not cytotoxic at other concentrations on the HCT-116 cell line. Additionally, olive leaf extract was not found to have a toxic effect on HDF cells. Therefore, it was thought that olive leaf extract could accelerate cell death in cancer cell lines. Previous researches were reported that olive leaf extract have cytotoxic effects on many different cancer cells including breast and colon cancers [14,15]. Also, Barbaro et al. [16] reported that the antitumor activity of olive leaf and oleuropein may be associated with reactive oxygens species (ROS).




Antigenotoxic agents often show expected therapeutic effects that may effectively control cancer [17]. As a result of the study, it was determined a significant number of nuclear abnormalities in the higher concentration of treated colon cancer cells. This feature of olive leaf extract may be explained by its capacity to act as a potent free radical scavenger [18,19,20,21].

This preliminary study aims to assess the effectiveness of methanol extraction from olive leaves and investigate olive leaf extracts as anticancer agents. This study's limitations include being limited to one cancer cell line, and the molecular mechanisms underlying activity were not investigated. Moreover, the individual compounds responsible for the anti-pancreatic cancer activity have not yet been identified. However, this study provides a platform for further research to evaluate the molecular mechanisms involved in the anticancer activity of the phenolic compounds of olive leaf extract in colon cancer cells.

Declaration of Interest: No potential conflict of interest relevant to this article was reported.

Authors' Contributions: EÖ, FÇ, and DU contributed to the study conception and design. EÖ, FÇ, and DU contribution to laboratory work. Literature research (EÖ and FÇ), Writing the article (EÖ and FÇ). All authors read and approved the final manuscript. EÖ; Emre Öztürk, FÇ; Fatma Çalık, DU; Derya Ulusoy.

ORCID:

Emre Öztürk  0000-0002-5847-0721
Fatma Çalık  0000-0003-1548-8689
Derya Ulusoy  0000-0001-7772-075X

References

1. Hanahan D., Weinberg R.A. Hallmarks of cancer: The next generation. *Cell*. 2011; 144:646–674. doi: 10.1016/j.cell.2011.02.013.
2. Ferlay J., Soerjomataram I., Dikshit R., Eser S., Mathers C., Rebelo M., Parkin D.M., Forman D.D., Bray F. Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *Int. J. Cancer*. 2014;136 doi: 10.1002/ijc.29210.
3. Filomeno M., Bosetti C., Bidoli E., Levi F., Serraino D., Montella M., La Vecchia C., Tavani A. Mediterranean diet and risk of endometrial cancer: A pooled analysis of three Italian case-control studies. *Br. J. Cancer*. 2015;112:1816–1821.

4. Schwingshackl L., Hoffmann G. Does a Mediterranean-Type Diet Reduce Cancer Risk? *Curr. Nutr. Rep.* 2015;5:9–17.
5. Ostan R., Lanzarini C., Pini E., Scurti M., Vianello D., Bertarelli C., Fabbri C., Izzi M., Palmas G., Biondi F., et al. Inflammaging and cancer: A challenge for the Mediterranean diet. *Nutrients.* 2015;7:2589–2621. doi: 10.3390/nu7042589.
6. Renna M., Rinaldi V.A., Gonnella M. The Mediterranean Diet between traditional foods and human health: The culinary example of Puglia (Southern Italy) *Int. J. Gastron. Food Sci.* 2015;2:63–71.
7. El S.N., Karakaya S. Olive tree (*Olea europaea*) leaves: Potential beneficial effects on human health. *Nutr. Rev.* 2009;67:632–638.
8. Mihailova A., Abbado D., Pedentchouk N. Differences in n-alkane profiles between olives and olive leaves as potential indicators for the assessment of olive leaf presence in virgin olive oils. *Eur. J. Lipid Sci. Technol.* 2015;117:1480–1485.
9. Pratheeshkumar P., Son Y.-O., Budhraj A., Wang X., Ding S., Wang L., Hitron A., Lee J.-C., Kim D., Divya S.P., et al. Luteolin inhibits human prostate tumor growth by suppressing vascular endothelial growth factor receptor 2-mediated angiogenesis. *PLoS ONE.* 2012;7:513.
10. Li F., Ye L., Lin S., Leung L.K. Dietary flavones and flavonones display differential effects on aromatase (CYP19) transcription in the breast cancer cells MCF-7. *Mol. Cell. Endocrinol.* 2011;344:51–58.
11. Sisman T., Askin H., Turkez H., Ozkan H., Incekara U., Colak S (2015) Determination of nuclear abnormalities in peripheral erythrocytes of the frog *Pelophylax ridibundus* (Anura: Ranidae) sampled from Karasu River Basin (Turkey) for pollution impacts. *LimnoFish* 1(2):75–81.
12. Özer, B. Kolon Kanseri Tedavisi İçin Yeni Nanoterapötik Modalitelerin Geliştirilmesi. Doktora Tezi. Trakya Üniversitesi Fen Bilimleri Enstitüsü Biyoteknoloji Ve Genetik Anabilim Dalı, 2021. Edirne.
13. NALKIRAN, İ. (2014). Kolon kanserinde MnSOD ve GPX1 gen anlatımlarının incelenmesi/ Examination of MnSOD and GPX1 gene expressions in colon cancer (Master Thesis).
14. Cárdeno A, Sánchez-Hidalgo M, Rosillo MA, Alarcón de la Lastra C. Oleuropein, a secoiridoid derived from olive tree, inhibits the proliferation of human colorectal cancer cell through downregulation of HIF-1 α . *Nutr. Cancer.* 2013;65:147-156.
15. Hassan ZK, Elamin MH, Omer SA, Daghestani MH, Al-Olayan ES, Elobeid MA, Virk P. Oleuropein induces apoptosis via the p53 pathway in breast cancer cells. *Asian Pac J Cancer Prev.* 2014;14:6739-6742.
16. Barbaro B, Toietta G, Maggio R, Arciello M, Tarocchi M, Galli A, Balsano C. Effects of the olive-derived polyphenol oleuropein on human health. *Int J Mol Sci.* 2014;15:18508-18524.
17. Tunca B, Tezcan G, Cecener G, Egeli U, Ak S, Malyer H, Tumen G, Bilir A. *Olea europaea* leaf extract alters microRNA expression in human glioblastoma cells. *Journal of cancer research and clinical oncology*, 138(11): 1831-1844, 2012.
18. Bektay, M. Y., Güler, E. M., Gökçe, M., & KIZILTAŞ, M. V. (2021). Investigation of the genotoxic, cytotoxic, apoptotic, and oxidant effects of olive leaf extracts on liver cancer cell lines. *Turkish Journal of Pharmaceutical Sciences*, 18(6), 781.
19. Kara, A., Akman, S., Ozkanlar, S., Tozoglu, U., Kalkan, Y., Canakci, C. F., & Tozoglu, S. (2013). Immune modulatory and antioxidant effects of melatonin in experimental periodontitis in rats. *Free Radical Biology and Medicine*, 55, 21-26.
20. Arabacı, T., Kermen, E., Ozkanlar, S., Köse, O., Kara, A., Kızıldağ, A., ... & Ibişoğlu, E. (2015). Therapeutic effects of melatonin on alveolar bone resorption after experimental periodontitis in rats: a biochemical and immunohistochemical study. *Journal of periodontology*, 86(7), 874-881.
21. Ozkanlar, S., Kara, A., Sengul, E., Simsek, N., Karadeniz, A., & Kurt, N. (2016). Melatonin modulates the immune system response and inflammation in diabetic rats experimentally-induced by alloxan. *Hormone and Metabolic Research*, 48(02), 137-144.