



Non-linear mixed Jordan triple 1- $*$ -product on von Neumann algebras

Abdul Nadim Khan^{*1} , Mohd Arif Raza¹ , Husain Alhazmi² 

¹Department of Mathematics, College of Science & Arts- Rabigh, King Abdulaziz University, Saudi Arabia

²Department of Mathematics, Faculty of Science, King Abdulaziz University, Jeddah-21589, Saudi Arabia

Abstract

It is shown that if M and N are two von Neumann algebras, one of which has no central abelian projection with $\psi : M \rightarrow N$ satisfying mixed Jordan triple 1- $*$ -product, i.e.,

$$\psi(A \circ B \bullet C) = \psi(A) \circ \psi(B) \bullet \psi(C)$$

for all $A, B, C \in M$, then there exists a bijective map $\Psi : M \rightarrow N$ such that $\Psi(A) = \psi(I)\psi(A)$ with $\psi(I)^2 = I$, whenever $\psi(I)$ is central, and there exist a central projection $\mathfrak{P} \in M$ such that the restriction of ψ to $M\mathfrak{P}$ is a linear $*$ -isomorphism, and to $M(I - \mathfrak{P})$ is a conjugate linear $*$ -isomorphism.

Mathematics Subject Classification (2020). 47B48, 46L10

Keywords. prime ring, semiprime ring, normal ring, involution, left $*$ -centralizer, Jordan left $*$ -centralizer, generalized derivation

1. Notations and introduction

Let M be a von Neumann algebra and $A, B \in M$. We express $A \bullet_{\lambda} B = AB + \lambda BA^*$, the Jordan λ - $*$ -product. For $\lambda = \pm 1$, we say Jordan 1- $*$ -product and Jordan (-1) - $*$ -product, respectively. Traditionally, numerous algebraists were already committed to analyse those mappings that aren't necessarily additive preserved Jordan $*$ -products on various algebras. The study of non-linear preserving problems is one of the premier areas in matrix theory as well as operator theory. A variety of research objectives on certain algebras such as von Neumann algebras, operator algebras, prime $*$ -algebras, etc were discussed in depth [2, 3, 7–11, 14–16] and references therein. The first implementation of this theory was presented by Šemrl [17]. In addition, with the relation to quadratic functionals, the Jordan (-1) - $*$ -product was introduced and studied by him. In [1], Bai and Du revealed that the sum of linear and conjugate linear $*$ -isomorphisms would be any bijective map on von Neumann algebras without central abelian projections, which preserved the Jordan (-1) - $*$ -product. Quite few generalizations throughout the last result can be found [4, 6, 7, 11] done by plenty of authors.

*Corresponding Author.

Email addresses: abdulnadimkhan@gmail.com (A. N. Khan), arifraza03@gmail.com (M. A. Raza), hsalhazmi@kau.edu.sa (H. Alhazmi)

Received: 11.06.2023; Accepted: 24.03.2024

Throughout this line of questioning, recently Huo *et al.* [5] extended the above-mentioned interpellation for Jordan triple η -product. Specifically, he stated that: “Assume that ψ is a bijection between two von Neumann algebras that is not necessarily linear, of which one has abelian projections which is not central with $\psi(I) = I$ and having the Jordan triple η -*-product. If η is not real, then ψ is a linear *-isomorphism, and if η is real, then ψ is the sum of a linear *-isomorphism and a conjugate linear *-isomorphism”. Additionally, they also addressed a conjecture that whether this result is relevant without $\psi(I) = I$. In 2017, Li and Lu [8] provided the affirmative response to this problem and developed the consequence on von Neumann algebras for Jordan’s triple 1-*-product, of which one has abelian projections which is not central. In this article, we also provide a constructive response to the above problem but not only dismantle the presumption of $\psi(I) = I$, we demonstrate the result in a somewhat broader sense by considering mixed Jordan 1-*-product which is defined as for any $A, B, C \in M$,

$$A \circ B \bullet C = (AB + BA) \bullet C = ABC + BAC + CB^*A^* + CA^*B^*.$$

Within this manuscript, we are primarily interested in exploring how non-linear maps are formed on von Neumann algebras satisfying mixed Jordan triple 1-*-product i.e., $\psi(A \circ B \bullet C) = \psi(A) \circ \psi(B) \bullet \psi(C)$ for all $A, B, C \in M$. Over few years some significant work drawn an attention of researchers has been consecrated to the evaluation of mixed Lie and Jordan triple products and derivations ([12, 18–20]). Such studies reported above encourage us to prove the following:

Theorem 1.1. *Let M and N be two von Neumann algebras, one of which has no central abelian projection. Define a map $\psi : M \rightarrow N$ such that*

$$\psi(A \circ B \bullet C) = \psi(A) \circ \psi(B) \bullet \psi(C)$$

for all $A, B, C \in M$. If $\psi(I)$ is central, then there exists a bijective map $\Psi : M \rightarrow N$ such that $\Psi(A) = \psi(I)\psi(A)$ with $\psi(I)^2 = I$ and there exists a central projection $\mathfrak{P} \in M$ such that the restriction of ψ to $M\mathfrak{P}$ is a linear *-isomorphism and the restriction of ψ to $M(I - \mathfrak{P})$ is a conjugate linear *-isomorphism.

We systematize the proof of aforementioned result in two sections. Section 2 presents some preliminary notions and useful lemmas that are essential to show ψ is additive. In Section 3, we shall provide numerous constructive remarks and lemmas to elaborate the assertion of Theorem 1.1.

2. Additivity of ψ

Theorem 2.1. *Let M and N be two von Neumann algebras and define a bijective map $\psi : M \rightarrow N$ such that*

$$\psi(A \circ B \bullet C) = \psi(A) \circ \psi(B) \bullet \psi(C)$$

for all $A, B, C \in M$. Then ψ is additive.

Proof. Take into account that $\mathfrak{P}_1 \in M$ and $\mathfrak{P}_2 = I - \mathfrak{P}_1$ are projections, whereas I is an unit element of M . We write $M_{jk} = \mathfrak{P}_j M \mathfrak{P}_k$ for $j, k = 1, 2$. Then by Peire’s decomposition of M , we have $M = M_{11} \oplus M_{12} \oplus M_{21} \oplus M_{22}$. It should be noted that any operator $A \in M$ can be written as $A = A_{11} + A_{12} + A_{21} + A_{22}$.

In view of the approximately facts, the verification of the theorem is given within the presentation of the following lemmas:

Lemma 2.2. $\psi(0) = 0$.

Proof. Due to ψ being surjective, there is $A \in M$ such that $\psi(A) = 0$. Thus

$$\psi(0) = \psi(0 \circ 0 \bullet A) = \psi(0) \circ \psi(0) \bullet \psi(A) = 0.$$

□

Lemma 2.3. *Let $A_{12} \in M_{12}$ and $A_{21} \in M_{21}$. Then $\psi(A_{12} + A_{21}) = \psi(A_{12}) + \psi(A_{21})$.*

Proof. Let $\Phi = (A_{12} + A_{21}) - \psi^{-1}(\psi(A_{12}) + \psi(A_{21}))$. Then, we have

$$\begin{aligned} \psi(A_{12} + A_{21}) \circ \psi(\mathfrak{P}_2) \bullet \psi(\mathfrak{P}_1) &= \psi((A_{12} + A_{21}) \circ \mathfrak{P}_2 \bullet \mathfrak{P}_1) \\ &= \psi(A_{12} \circ \mathfrak{P}_2 \bullet \mathfrak{P}_1) + \psi(A_{21} \circ \mathfrak{P}_2 \bullet \mathfrak{P}_1) \\ &= \psi(A_{12}) \circ \psi(\mathfrak{P}_2) \bullet \psi(\mathfrak{P}_1) + \psi(A_{21}) \circ \psi(\mathfrak{P}_2) \bullet \psi(\mathfrak{P}_1) \\ &= (\psi(A_{12}) + \psi(A_{21})) \circ \psi(\mathfrak{P}_2) \bullet \psi(\mathfrak{P}_1). \end{aligned}$$

Apply ψ^{-1} on both sides of above expression. This gives $\Phi \circ \mathfrak{P}_2 \bullet \mathfrak{P}_1 = 0$, which yields $\Phi_{21} = 0$. Similarly, we can show that $\Phi_{12} = 0$ by replacing \mathfrak{P}_2 by \mathfrak{P}_1 and \mathfrak{P}_1 by \mathfrak{P}_2 , respectively. Next, we have

$$\begin{aligned} \psi(I) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet \psi(A_{12} + A_{21}) &= \psi(I \circ (\mathfrak{P}_1 - \mathfrak{P}_2) \bullet (A_{12} + A_{21})) \\ &= \psi(I \circ (\mathfrak{P}_1 - \mathfrak{P}_2) \bullet A_{12}) + \psi(I \circ (\mathfrak{P}_1 - \mathfrak{P}_2) \bullet A_{21}) \\ &= \psi(I) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet \psi(A_{12}) \\ &\quad + \psi(I) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet \psi(A_{21}) \\ &= \psi(I) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet (\psi(A_{12}) + \psi(A_{21})). \end{aligned}$$

Again, impose ψ^{-1} in last relation, we get $I \circ (\mathfrak{P}_1 - \mathfrak{P}_2) \bullet \Phi = 0$. This further implies $\Phi_{11} = \Phi_{22} = 0$. Thus $\Phi = 0$ i.e.,

$$\psi(A_{12} + A_{21}) = \psi(A_{12}) + \psi(A_{21}).$$

□

Lemma 2.4. *For any $A_{11} \in M_{11}, A_{12} \in M_{12}$ and $A_{21} \in M_{21}$,*

- (i) $\psi(A_{11} + A_{12} + A_{21}) = \psi(A_{11}) + \psi(A_{12}) + \psi(A_{21})$;
- (ii) $\psi(A_{12} + A_{21} + A_{22}) = \psi(A_{12}) + \psi(A_{21}) + \psi(A_{22})$.

Proof. Let $\Theta = (A_{11} + A_{12} + A_{21}) - \psi^{-1}(\psi(A_{11}) + \psi(A_{12}) + \psi(A_{21}))$. Then by Lemma 2.3, we have

$$\begin{aligned} \psi(A_{11} + A_{12} + A_{21}) \circ \psi(\mathfrak{P}_1) \bullet \psi(\mathfrak{P}_2) &= \psi((A_{11} + A_{12} + A_{21}) \circ \mathfrak{P}_1 \bullet \mathfrak{P}_2) \\ &= \psi(A_{11} \circ \mathfrak{P}_1 \bullet \mathfrak{P}_2) + \psi(A_{12} \circ \mathfrak{P}_1 \bullet \mathfrak{P}_2) \\ &\quad + \psi(A_{21} \circ \mathfrak{P}_1 \bullet \mathfrak{P}_2) \\ &= \psi(A_{11}) \circ \psi(\mathfrak{P}_1) \bullet \psi(\mathfrak{P}_2) + \psi(A_{12}) \circ \psi(\mathfrak{P}_1) \\ &\quad \bullet \psi(\mathfrak{P}_2) + \psi(A_{21}) \circ \psi(\mathfrak{P}_1) \bullet \psi(\mathfrak{P}_2) \\ &= (\psi(A_{11}) + \psi(A_{12}) + \psi(A_{21})) \circ \psi(\mathfrak{P}_1) \bullet \psi(\mathfrak{P}_2). \end{aligned}$$

The last expression yields $\Theta \circ \mathfrak{P}_1 \bullet \mathfrak{P}_2 = 0$, and hence $\Theta_{12} = 0$. Similarly, we can get $\Theta_{21} = 0$. Now, we only need to show $\Theta_{11} = \Theta_{22} = 0$. It follows from the hypothesis and

Lemma 2.3 that

$$\begin{aligned}
& \psi\left(\frac{I}{2}\right) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet \psi(A_{11} + A_{12} + A_{21}) \\
&= \psi\left(\frac{I}{2} \circ (\mathfrak{P}_1 - \mathfrak{P}_2) \bullet (A_{11} + A_{12} + A_{21})\right) \\
&= \psi\left(\frac{I}{2} \circ (\mathfrak{P}_1 - \mathfrak{P}_2) \bullet A_{11}\right) + \psi\left(\frac{I}{2} \circ (\mathfrak{P}_1 - \mathfrak{P}_2) \bullet A_{12}\right) \\
&\quad + \psi\left(\frac{I}{2} \circ (\mathfrak{P}_1 - \mathfrak{P}_2) \bullet A_{21}\right) \\
&= \psi\left(\frac{I}{2}\right) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet \psi(A_{11}) + \psi\left(\frac{I}{2}\right) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet \psi(A_{12}) \\
&\quad + \psi\left(\frac{I}{2}\right) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet \psi(A_{21}) \\
&= \psi\left(\frac{I}{2}\right) \circ \psi(\mathfrak{P}_1 - \mathfrak{P}_2) \bullet (\psi(A_{11}) + \psi(A_{12}) + \psi(A_{21})).
\end{aligned}$$

Reasoning as above, we obtain $\Theta_{11} = \Theta_{22} = 0$, and hence

$$\psi(A_{11} + A_{12} + A_{21}) = \psi(A_{11}) + \psi(A_{12}) + \psi(A_{21}).$$

Similarly, we can show

$$\psi(A_{12} + A_{21} + A_{22}) = \psi(A_{12}) + \psi(A_{21}) + \psi(A_{22}).$$

This completes the proof. \square

Lemma 2.5. For any $A_{ij} \in M_{ij}$, $1 \leq i, j \leq 2$, we have

$$\psi\left(\sum_{i,j=1}^2 A_{ij}\right) = \sum_{i,j=1}^2 \psi(A_{ij}).$$

Proof. Assume that $\nabla = \sum_{i,j=1}^2 A_{ij} - \psi^{-1}\left(\sum_{i,j=1}^2 \psi(A_{ij})\right)$. In view of Lemma 2.4(i) and $\mathfrak{P}_1 \circ I \bullet A_{22} = 0$, we have

$$\begin{aligned}
\psi(\mathfrak{P}_1) \circ \psi(I) \bullet \psi\left(\sum_{i,j=1}^2 A_{ij}\right) &= \psi(\mathfrak{P}_1 \circ I \bullet \sum_{i,j=1}^2 A_{ij}) \\
&= \psi(\mathfrak{P}_1 \circ I \bullet A_{11}) + \psi(\mathfrak{P}_1 \circ I \bullet A_{12}) \\
&\quad + \psi(\mathfrak{P}_1 \circ I \bullet A_{21}) + \psi(\mathfrak{P}_1 \circ I \bullet A_{22}) \\
&= \psi(\mathfrak{P}_1) \circ \psi(I) \bullet \psi(A_{11}) + \psi(\mathfrak{P}_1) \circ \psi(I) \bullet \psi(A_{12}) \\
&\quad + \psi(\mathfrak{P}_1) \circ \psi(I) \bullet \psi(A_{21}) + \psi(\mathfrak{P}_1) \circ \psi(I) \bullet \psi(A_{22}) \\
&= \psi(\mathfrak{P}_1) \circ \psi(I) \bullet \sum_{i,j=1}^2 \psi(A_{ij}).
\end{aligned}$$

Apply ψ^{-1} on both sides of above expression which yields $\mathfrak{P}_1 \circ I \bullet \nabla = 0$, and hence $\nabla_{11} = \nabla_{12} = \nabla_{21} = 0$. We can show in similar manner that $\nabla_{22} = 0$. Thus $\nabla = 0$ i.e.,

$$\psi\left(\sum_{i,j=1}^2 A_{ij}\right) = \sum_{i,j=1}^2 \psi(A_{ij}).$$

\square

Lemma 2.6. For any $A_{ij}, B_{ij} \in M_{ij}$ with $i \neq j$, $\psi(A_{ij} + B_{ij}) = \psi(A_{ij}) + \psi(B_{ij})$.

Proof. Since $\frac{I}{2} \circ (\mathfrak{P}_i + A_{ij}) \bullet (\mathfrak{P}_j + B_{ij}) = A_{ij} + B_{ij} + A_{ij}^* + B_{ij}A_{ij}^*$, so it follows from Lemma 2.4 that

$$\begin{aligned}
 \psi(A_{ij} + B_{ij}) + \psi(A_{ij}^*) + \psi(B_{ij}A_{ij}^*) &= \psi\left(\frac{I}{2} \circ (\mathfrak{P}_i + A_{ij}) \bullet (\mathfrak{P}_j + B_{ij})\right) \\
 &= \psi\left(\frac{I}{2}\right) \circ \psi(\mathfrak{P}_i + A_{ij}) \bullet \psi(\mathfrak{P}_j + B_{ij}) \\
 &= \psi\left(\frac{I}{2}\right) \circ (\psi(\mathfrak{P}_i) + \psi(A_{ij})) \bullet (\psi(\mathfrak{P}_j) + \psi(B_{ij})) \\
 &= \psi\left(\frac{I}{2}\right) \circ \psi(\mathfrak{P}_i) \bullet \psi(\mathfrak{P}_j) + \psi\left(\frac{I}{2}\right) \circ \psi(\mathfrak{P}_i) \bullet \psi(B_{ij}) \\
 &\quad + \psi\left(\frac{I}{2}\right) \circ \psi(A_{ij}) \bullet \psi(\mathfrak{P}_j) \\
 &\quad + \psi\left(\frac{I}{2}\right) \circ \psi(A_{ij}) \bullet \psi(B_{ij}) \\
 &= \psi\left(\frac{I}{2} \circ \mathfrak{P}_i \bullet \mathfrak{P}_j\right) + \psi\left(\frac{I}{2}\right) \circ \mathfrak{P}_i \bullet B_{ij} \\
 &\quad + \psi\left(\frac{I}{2} \circ A_{ij} \bullet \mathfrak{P}_j\right) + \psi\left(\frac{I}{2} \circ A_{ij} \bullet B_{ij}\right) \\
 &= \psi(B_{ij}) + \psi(A_{ij} + A_{ij}^*) + \psi(B_{ij}A_{ij}^*) \\
 &= \psi(A_{ij}) + \psi(B_{ij}) + \psi(A_{ij}^*) + \psi(B_{ij}A_{ij}^*).
 \end{aligned}$$

Thus

$$\psi(A_{ij} + B_{ij}) = \psi(A_{ij}) + \psi(B_{ij}).$$

□

Lemma 2.7. For any $A_{ii}, B_{ii} \in M_{ii}$, $\psi(A_{ii} + B_{ii}) = \psi(A_{ii}) + \psi(B_{ii})$.

Proof. Suppose $\Pi = (A_{ii} + B_{ii}) - \psi^{-1}(\psi(A_{ii}) + \psi(B_{ii}))$. It is easy to find

$$\begin{aligned}
 \psi(\mathfrak{P}_j) \circ \psi(I) \bullet \psi(A_{ii} + B_{ii}) &= \psi(\mathfrak{P}_j \circ I \bullet (A_{ii} + B_{ii})) \\
 &= \psi(\mathfrak{P}_j \circ I \bullet A_{ii}) + \psi(\mathfrak{P}_j \circ I \bullet B_{ii}) \\
 &= \psi(\mathfrak{P}_j) \circ \psi(I) \bullet \psi(A_{ii}) + \psi(\mathfrak{P}_j) \circ \psi(I) \bullet \psi(B_{ii}) \\
 &= \psi(\mathfrak{P}_j) \circ (\psi(A_{ii}) + \psi(B_{ii})) \bullet \psi(\mathfrak{P}_j).
 \end{aligned}$$

From above, we have $\mathfrak{P}_j \circ I \bullet \Pi = 0$. This yields $\Pi_{ij} = \Pi_{ji} = \Pi_{jj} = 0$. Next, according to Lemma 2.5 and Lemma 2.6, for any $C_{ij} \in M_{ij}$ with $i \neq j$, we have

$$\begin{aligned}
 \psi(\mathfrak{P}_i \circ (A_{ii} + B_{ii}) \bullet C_{ij}) &= \psi(A_{ii}C_{ij} + A_{ii}C_{ij} + B_{ii}C_{ij} + B_{ii}C_{ij}) \\
 &= \psi(A_{ii}C_{ij} + A_{ii}C_{ij}) + \psi(B_{ii}C_{ij} + B_{ii}C_{ij}) \\
 &= \psi(\mathfrak{P}_i \circ A_{ii} \bullet C_{ij}) + \psi(\mathfrak{P}_i \circ B_{ii} \bullet C_{ij}) \\
 &= \psi(\mathfrak{P}_i) \circ \psi(A_{ii}) \bullet \psi(C_{ij}) + \psi(\mathfrak{P}_i) \circ \psi(B_{ii}) \bullet \psi(C_{ij}) \\
 &= \psi(\mathfrak{P}_i) \circ (\psi(A_{ii}) + \psi(B_{ii})) \bullet \psi(C_{ij})
 \end{aligned}$$

On the other hand,

$$\psi(\mathfrak{P}_i \circ (A_{ii} + B_{ii}) \bullet C_{ij}) = \psi(\mathfrak{P}_i) \circ \psi(A_{ii} + B_{ii}) \bullet \psi(C_{ij}).$$

Hence $\mathfrak{P}_i \circ \Pi \bullet C_{ij} = 0$. This gives $\Pi_{ii} = 0$. Thus $\Pi = 0$ i.e.,

$$\psi(A_{ii} + B_{ii}) = \psi(A_{ii}) + \psi(B_{ii}).$$

□

Lemma 2.8. ψ is an additive map.

Proof. It follows from Lemmas 2.2-2.7 that ψ is additive. □

3. Proof of Theorem 1.1

While we proceed upon our scientific findings, we could present some information, including some few conceptual details. The unital von Neumann algebra M is indeed a weakly closed, self-adjoint operator algebra in the Hilbert space H . The collection $Z(M) = \{S \in M : ST = TS \text{ for all } T \in M\}$ is referred as the center of M . If $\mathfrak{P} \in Z(M)$ or $\mathfrak{P}M\mathfrak{P}$ abelian, then the projection \mathfrak{P} is called the central abelian projection. Aware that perhaps the central carrier of A , denoted by \overline{A} , seems to be the smallest \mathfrak{P} central projection that meets $\mathfrak{P}A = A$. The central carrier of A can be viewed as the projection onto the closed subspace spanned by $\{BA(x) : B \in M, x \in H\}$. The core of A , denoted by \underline{A} , is $\sup\{S \in Z(M) : S = S^*, S \leq A\}$, if A is self-adjoint. If \mathfrak{P} is a projection, then it is obvious that $\underline{\mathfrak{P}}$ is the largest central Q projection that satisfies $Q \leq \mathfrak{P}$. If $\underline{\mathfrak{P}} = 0$, then the projection \mathfrak{P} is said to be core-free. It is straightforward to see it now $\underline{\mathfrak{P}} = 0$ if and only if $\overline{(I - \mathfrak{P})} = I$. Following remarks are critical for the proof of our main result:

Remark 3.1. [13, Lemma 4] “If M is a von Neumann algebra with no central abelian projection $\mathfrak{P} \in M$, then there exists a projection $\mathfrak{P} \in M$ such that $\underline{\mathfrak{P}} = 0$ and $\overline{\mathfrak{P}} = I$.”

Remark 3.2. [8, Lemma 2.2] “Let M be a von Neumann algebra on a Hilbert space H . Let A be an operator in M and $\mathfrak{P} \in M$ is a projection with $\overline{\mathfrak{P}} = I$. If $AB\mathfrak{P} = 0$ for all $B \in M$, then $A = 0$. Consequently, if $\mathfrak{Z} \in Z(M)$, then $\mathfrak{Z}\mathfrak{P} = 0$ implies $\mathfrak{Z} = 0$.”

Remark 3.3. [8, Lemma 2.3] “Let M be a von Neumann algebra and $A \in M$. Then $AB + BA^* = 0$ for all $B \in M$ implies that $A = -A^* \in Z(M)$.”

We can see from Theorem 2.1, ψ would be an additive map. Throughout the succeeding arguments, the unit elements of the algebra M and N weren't differentiated and we'll see within next proof that this does not impact our argument. We'll demonstrate that theorem progressively through implementing:

Lemma 3.4. $2I = \psi(I)^2 + (\psi(I)^*)^2$.

Proof. Let $A \in M$ such that $\psi(A) = I$. Then it follows from the additivity of ψ that

$$4I = 4\psi(A) = \psi(I \circ I \bullet A) = \psi(I) \circ \psi(I) \bullet I = 2(\psi(I)^2 + (\psi(I)^*)^2). \quad (3.1)$$

This implies $2I = \psi(I)^2 + (\psi(I)^*)^2$. \square

Lemma 3.5. Let $\Theta_{\mathfrak{P}} = \frac{1}{2}(\psi(I)\psi(\mathfrak{P}) + \psi(I)^*\psi(\mathfrak{P})^*)$, where $\mathfrak{P} \in M$ is a projection. Then $\Theta_{\mathfrak{P}}$ is a projection such that $\psi(\mathfrak{P}) = \psi(I)\Theta_{\mathfrak{P}}$.

Proof. From the hypothesis, we have

$$\begin{aligned} 4\psi(\mathfrak{P}) &= \psi(I \circ \mathfrak{P} \bullet I) \\ &= \psi(I) \circ \psi(\mathfrak{P}) \bullet \psi(I) \\ &= 2\psi(I)\psi(\mathfrak{P}) \bullet \psi(I) \\ &= 2\psi(I)(\psi(I)\psi(\mathfrak{P}) + \psi(I)^*\psi(\mathfrak{P})^*) \\ &= 4\psi(I)\Theta_{\mathfrak{P}}. \end{aligned}$$

Also

$$\begin{aligned} 4\psi(\mathfrak{P}) &= \psi(I \circ \mathfrak{P} \bullet \mathfrak{P}) \\ &= \psi(I) \circ \psi(\mathfrak{P}) \bullet \psi(\mathfrak{P}) \\ &= 2\psi(I)\psi(\mathfrak{P}) \bullet \psi(\mathfrak{P}) \\ &= 2\psi(\mathfrak{P})^2\psi(I) + 2\psi(\mathfrak{P})\psi(\mathfrak{P})^*\psi(I)^* \\ &= 2\psi(\mathfrak{P})(\psi(I)\psi(\mathfrak{P}) + \psi(I)^*\psi(\mathfrak{P})^*) \\ &= 4\psi(\mathfrak{P})\Theta_{\mathfrak{P}} = 4\psi(I)\Theta_{\mathfrak{P}}^2. \end{aligned}$$

It follows from the last two relations that

$$\psi(\mathfrak{P}) = \psi(I)\Theta_{\mathfrak{P}}^2. \quad (3.2)$$

Since $\Theta_{\mathfrak{P}}$ is self-adjoint, so we have

$$\psi(\mathfrak{P})^* = \psi(I)^*\Theta_{\mathfrak{P}}^2. \quad (3.3)$$

Multiply (3.2) by $\psi(I)$ and (3.3) by $\psi(I)^*$, and add the so obtained relations, we get

$$\psi(I)\psi(\mathfrak{P}) + \psi(I)^*\psi(\mathfrak{P})^* = (\psi(I)^2 + (\psi(I)^*)^2)\Theta_{\mathfrak{P}}^2. \quad (3.4)$$

Observe from Lemma 3.4 that $2\Theta_{\mathfrak{P}} = 2\Theta_{\mathfrak{P}}^2$. Therefore $\Theta_{\mathfrak{P}}$ is a projection. \square

Lemma 3.6. *For any $A_{12} \in M_{12}$ and a projection $\mathfrak{P} \in M$, $\psi(A_{12}) = \Theta_{\mathfrak{P}}\psi(A_{12}) + \psi(A_{12})\Theta_{\mathfrak{P}}$.*

Proof. It follows from Lemma 3.5 that

$$\begin{aligned} 2\psi(A_{12}) &= \psi(I \circ \mathfrak{P} \bullet A_{12}) \\ &= \psi(I) \circ \psi(\mathfrak{P}) \bullet \psi(A_{12}) \\ &= 2\psi(I)\psi(\mathfrak{P})\psi(A_{12}) + 2\psi(A_{12})\psi(I)^*\psi(\mathfrak{P})^* \\ &= 2\psi(I)^2\Theta_{\mathfrak{P}}\psi(A_{12}) + 2(\psi(I)^2)^*\psi(A_{12})^*\Theta_{\mathfrak{P}}. \end{aligned}$$

Multiply both sides of above equation by $\Theta_{\mathfrak{P}}$, we get $\Theta_{\mathfrak{P}}\psi(A_{12})\Theta_{\mathfrak{P}} = 0$. Similarly, if we multiply above expression from left and right by $I - \Theta_{\mathfrak{P}}$, we obtain $(I - \Theta_{\mathfrak{P}})\psi(A_{12})(I - \Theta_{\mathfrak{P}}) = 0$. Therefore,

$$\psi(A_{12}) = \Theta_{\mathfrak{P}}\psi(A_{12}) + \psi(A_{12})\Theta_{\mathfrak{P}}.$$

Hence the proof. \square

Lemma 3.7. $\psi(I)^2 = I$.

Proof. By the hypothesis, we can choose a projection $\mathfrak{P}' \in N$, where N has no central abelian projection, such that $\mathfrak{P}' = 0$ and $\overline{\mathfrak{P}'} = I$. Let $\mathcal{N} \in N$ such that $\mathcal{N} = \mathfrak{P}'\mathcal{N}(I - \mathfrak{P}')$. Assume that $\mathfrak{P} = \frac{1}{2}(\psi^{-1}(I)\psi^{-1}(\mathfrak{P}') + \psi^{-1}(I)^*\psi^{-1}(\mathfrak{P}')^*)$. It is clear from Lemma 3.5 and 3.6 that \mathfrak{P} is a projection and $\psi^{-1}(\mathcal{N}) = \mathfrak{P}\psi^{-1}(\mathcal{N}) + \psi^{-1}(\mathcal{N})\mathfrak{P}$. Now, it follows that

$$\begin{aligned} \psi(\mathfrak{P}) &= \frac{1}{2}\psi(\psi^{-1}(I)\psi^{-1}(\mathfrak{P}') + \psi^{-1}(I)^*\psi^{-1}(\mathfrak{P}')^*) \\ &= \frac{1}{4}\psi(\psi^{-1}(I) \circ \psi^{-1}(\mathfrak{P}') \bullet I) \\ &= \frac{1}{4}(I \circ \mathfrak{P}' \bullet \psi(I)) = \psi(I)\mathfrak{P}'. \end{aligned}$$

Also, we have

$$\begin{aligned} \mathcal{N} &= \psi(\mathfrak{P}\psi^{-1}(\mathcal{N}) + \psi^{-1}(\mathcal{N})\mathfrak{P}) \\ &= \frac{1}{2}\psi(I \circ \mathfrak{P} \bullet \psi^{-1}(\mathcal{N})) \\ &= \frac{1}{2}(\psi(I) \circ \psi(\mathfrak{P}) \bullet \mathcal{N}) \\ &= \psi(I)\psi(\mathfrak{P})\mathcal{N} + \psi(I)^*\mathcal{N}\psi(\mathfrak{P})^* \\ &= \psi(I)^2\mathfrak{P}'\mathcal{N} + (\psi(I)^2)^*\mathcal{N}\mathfrak{P}' \\ &= \psi(I)^2\mathcal{N}. \end{aligned}$$

This gives $(I - \psi(I)^2)\mathcal{N} = 0$. Thus, $(I - \psi(I)^2)\mathfrak{P}'\mathcal{N}(I - \mathfrak{P}') = 0$ and since $\overline{(I - \mathfrak{P}')} = I$, in view of Remark 3.1, we obtain $(I - \psi(I)^2)\mathfrak{P}' = 0$. Note that $(I - \psi(I)^2) \in Z(N)$ and $\overline{\mathfrak{P}'} = I$, then again by Remark 3.1, we have $I - \psi(I)^2 = 0$ i.e., $\psi(I)^2 = I$. This completes the proof. \square

Now, for any $A \in M$, define a map $\Psi : M \rightarrow N$ by $\Psi(A) = \psi(I)\psi(A)$. Then Ψ has the following characteristics:

Lemma 3.8. (a) Ψ is an additive bijective map satisfying

$$\Psi(A \circ B \bullet C) = \Psi(A) \circ \Psi(B) \bullet \Psi(C) \quad \text{for all } A, B, C \in M;$$

(b) $\Psi(I) = I$;

(c) $\Psi(A^*) = \Psi(A)^*$ for all $A \in M$;

(d) \mathfrak{P} is a projection in M iff $\Psi(\mathfrak{P})$ is a projection in N .

Proof. For any $A, B \in M$, we have

$$\Psi(A + B) = \psi(I)\psi(A + B) = \psi(I)\psi(A) + \psi(I)\psi(B) = \Psi(A) + \Psi(B).$$

On the other hand, for any $A, B, C \in M$, we have

$$\begin{aligned} \Psi(A \circ B \bullet C) &= \psi(I)\psi(A \circ B \bullet C) \\ &= \psi(I)(\psi(A) \circ \psi(B) \bullet \psi(C)) \\ &= \psi(I)((\psi(A)\psi(B) + \psi(B)\psi(A)) \bullet \psi(C)) \\ &= (\psi(I)\psi(A)\psi(B) + \psi(I)\psi(B)\psi(A))\psi(C) \\ &\quad + (\psi(C)(\psi(I)\psi(B)^*\psi(A)^* + \psi(I)\psi(A)^*\psi(B)^*)). \end{aligned}$$

An application of Lemma 3.7 gives

$$\begin{aligned} \Psi(A \circ B \bullet C) &= (\psi(I)\psi(A)\psi(I)\psi(B) + \psi(I)\psi(B)\psi(I)\psi(A))\psi(I)\psi(C) \\ &\quad + (\psi(I)\psi(C)(\psi(I)^*\psi(B)^*\psi(I)^*\psi(A)^* + \psi(I)^*\psi(A)^*\psi(I)^*\psi(B)^*)) \\ &= (\psi(I)\psi(A)\psi(I)\psi(B) + \psi(I)\psi(B)\psi(I)\psi(A))\psi(I)\psi(C) \\ &\quad + (\psi(I)\psi(C)((\psi(I)\psi(B))^*(\psi(I)\psi(A))^* + (\psi(I)\psi(A))^*(\psi(I)\psi(B))^*)) \\ &= (\Psi(A)\Psi(B) + \Psi(B)\Psi(A))\Psi(C) + \Psi(C)(\Psi(B)^*\Psi(A)^* + \Psi(A)^*\Psi(B)^*) \\ &= \Psi(A) \circ \Psi(B) \bullet \Psi(C) \end{aligned}$$

This completes the proof.

(b) It follows directly from hypothesis and Lemma 3.7.

(c) By the hypothesis, we have

$$2(\Psi(A) + \Psi(A^*)) = 2\Psi(A + A^*) = \Psi(I \circ A \bullet I) = I \circ \Psi(A) \bullet I = 2(\Psi(A) + \Psi(A)^*).$$

Above relation yields $\Psi(A^*) = \Psi(A)^*$ for all $A \in M$.

(d) Since $\Psi(A) = \psi(I)\psi(A)$ for all $A \in M$, so for $A = \mathfrak{P}$ and from Lemma 3.5, we have $\Psi(\mathfrak{P}) = \psi(I)\psi(\mathfrak{P}) = \psi(I)^2\Theta_{\mathfrak{P}} = \Theta_{\mathfrak{P}}$. As we know $\Theta_{\mathfrak{P}}$ is a projection. Thus $\Psi(\mathfrak{P})$ is also, as we asserted. \square

As N has no central abelian projections, it follows from Lemma 3.4 that there exists a projection $Q_1 \in N$ such that $\underline{Q_1} = 0$ and $\overline{Q_1} = I$. Then by Lemma 3.8 (d), $\mathfrak{P}_1 = \Psi^{-1}(Q_1)$ is a projection in M . We denote $A_{ij} = \mathfrak{P}_i M \mathfrak{P}_j$ and $B_{ij} = \mathfrak{P}_i N \mathfrak{P}_j$, respectively. Keep it into mind, we now prove the following:

Lemma 3.9. For any $A_{ij} \in M_{ij}$ and $B_{ij} \in N_{ij}$, $1 \leq i, j \leq 2$, we have $\Psi(A_{ij}) = B_{ij}$.

Proof. First we prove for $i = 1, j = 2$. It follows from hypothesis that

$$\begin{aligned} 2\Psi(A_{12}) &= \Psi(I \circ \mathfrak{P}_1 \bullet A_{12}) \\ &= I \circ Q_1 \bullet \Psi(A_{12}) \\ &= 2Q_1\Psi(A_{12}) + 2\Psi(A_{12})Q_1. \end{aligned}$$

Multiply above relation, first by Q_1 on both sides and then by $Q_2 = I - Q_1$, we obtain $\Psi(A_{12}) = B_{12} + B_{21}$ for some $B_{12} \in N_{12}$ and $B_{21} \in N_{21}$. Next, we prove that $B_{21} = 0$. Observe that

$$\begin{aligned} 0 &= \Psi(I \circ A_{12} \bullet \mathfrak{P}_1) \\ &= I \circ \Psi(A_{12}) \bullet Q_1 \\ &= 2(B_{21} + B_{21}^*). \end{aligned}$$

This gives $B_{21} = 0$, and hence $\Psi(B_{12}) \subseteq M_{12}$. Since Ψ is a bijection, so we can easily obtain $\Psi(B_{12}) = M_{12}$. Similarly, we can show $\Psi(B_{21}) = M_{21}$. \square

Lemma 3.10. *For any $A_{ii} \in M_{ii}$ and $B_{ii} \in N_{ii}$, we have $\Psi(A_{ii}) \subseteq B_{ii}$.*

Proof. For $j \neq i$, we have

$$\begin{aligned} 0 &= \Psi(I \circ \mathfrak{P}_j \bullet A_{ii}) \\ &= I \circ Q_j \bullet \Psi(A_{ii}) \\ &= 2(Q_j \Psi(A_{ii}) + \Psi(A_{ii}) Q_j). \end{aligned}$$

This yields $Q_i \Psi(A_{ii}) Q_i = \Psi(A_{ii}) \subseteq B_{ii}$. \square

Lemma 3.11. *For any $A_{ij}, B_{ij} \in M_{ij}$, $i \leq i, j \leq 2$, we have*

- (a) $\Psi(A_{11}B_{12}) = \Psi(A_{11})\Psi(B_{12})$ and $\Psi(A_{22}B_{21}) = \Psi(A_{22})\Psi(B_{21})$;
- (b) $\Psi(A_{12}B_{21}) = \Psi(A_{12})\Psi(B_{21})$ and $\Psi(A_{21}B_{12}) = \Psi(A_{21})\Psi(B_{12})$;
- (c) $\Psi(A_{11}B_{11}) = \Psi(A_{11})\Psi(B_{11})$ and $\Psi(A_{22}B_{22}) = \Psi(A_{22})\Psi(B_{22})$;
- (d) $\Psi(A_{12}B_{22}) = \Psi(A_{12})\Psi(B_{22})$ and $\Psi(A_{21}B_{11}) = \Psi(A_{21})\Psi(B_{11})$.

Proof. (a) It follows from Lemma 3.9 and 3.10 that $\Psi(B_{12}A_{11}^*) = \Psi(B_{12})\Psi(A_{11})^* = 0$. Thus

$$\begin{aligned} 2\Psi(A_{11}B_{12}) + 2\Psi(B_{12}A_{11}^*) &= \Psi(I \circ A_{11} \bullet B_{12}) \\ &= I \circ \Psi(A_{11}) \bullet \Psi(B_{12}) \\ &= 2\Psi(A_{11})\Psi(B_{12}) + 2\Psi(B_{12})\Psi(A_{11})^*. \end{aligned}$$

This implies $\Psi(A_{11}B_{12}) = \Psi(A_{11})\Psi(B_{12})$. Similarly, we can show $\Psi(A_{22}B_{21}) = \Psi(A_{22})\Psi(B_{21})$. Next, to show (b), see from Lemma 3.9 that $\Psi(B_{21})\Psi(A_{12})^* = 0$. Therefore,

$$2\Psi(A_{12}B_{21}) = \Psi(A_{12} \circ I \bullet B_{21}) = 2\Psi(A_{12})\Psi(B_{21}).$$

Hence, $\Psi(A_{12}B_{21}) = \Psi(A_{12})\Psi(B_{21})$. Equivalently, one can easily show $\Psi(A_{21}B_{12}) = \Psi(A_{21})\Psi(B_{12})$. Now, we establish (c). Let $X_{12} \in N_{12}$ such that $C_{12} = \Psi^{-1}(X_{12}) \in M_{12}$ from Lemma 3.9. It follows from (a) that

$$\Psi(A_{11}B_{11})X_{12} = \Psi(A_{11}B_{11}C_{12}) = \Psi(A_{11})\Psi(B_{11}C_{12}) = \Psi(A_{11})\Psi(B_{11})X_{12}$$

for all $X_{12} \in M_{12}$. Since $\overline{Q_2} = I$, it follows from Remark 3.1 and 3.2 that $\Psi(A_{11}B_{11}) = \Psi(A_{11})\Psi(B_{11})$. Similarly, we can show $\Psi(A_{22}B_{22}) = \Psi(A_{22})\Psi(B_{22})$. Finally, to prove (d), we see from Lemma 3.9 that $E_{21} = \Psi^{-1}(Y_{21}) \in M_{21}$ for any $Y_{21} \in N_{21}$. So

$$\Psi(A_{12}B_{22})Y_{21} = \Psi(A_{12}B_{22}E_{21}) = \Psi(A_{12})\Psi(B_{22}E_{21}) = \Psi(A_{12})\Psi(B_{22})Y_{21}.$$

Reasoning as above, we obtain $\Psi(A_{12}B_{22}) = \Psi(A_{12})\Psi(B_{22})$. Similarly, we can have $\Psi(A_{21}B_{11}) = \Psi(A_{21})\Psi(B_{11})$. \square

Lemma 3.12. Ψ is a \mathbb{R} -linear *-isomorphism.

Proof. Since we know from Lemma 3.8 that Ψ is additive, so it follows from Lemma 3.11 that

$$\begin{aligned}
\Psi(AB) &= \Psi(A_{11}B_{11} + A_{11}B_{12} + A_{12}B_{12} + A_{12}B_{22} \\
&\quad + A_{21}B_{11} + A_{21}B_{12} + A_{22}B_{21} + A_{22}B_{22}) \\
&= \Psi(A_{11}B_{11}) + \Psi(A_{11}B_{12}) + \Psi(A_{12}B_{12}) + \Psi(A_{12}B_{22}) \\
&\quad + \Psi(A_{21}B_{11}) + \Psi(A_{21}B_{12}) + \Psi(A_{22}B_{21}) + \Psi(A_{22}B_{22}) \\
&= \Psi(A_{11})\Psi(B_{11}) + \Psi(A_{11})\Psi(B_{12}) + \Psi(A_{12})\Psi(B_{12}) + \Psi(A_{12})\Psi(B_{22}) \\
&\quad + \Psi(A_{21})\Psi(B_{11}) + \Psi(A_{21})\Psi(B_{12}) + \Psi(A_{22})\Psi(B_{21}) + \Psi(A_{22})\Psi(B_{22}) \\
&= \Psi(A)\Psi(B)
\end{aligned}$$

for all $A, B \in M$. Therefore, Ψ is an isomorphism, and hence $*$ -isomorphism by Lemma 3.8(c). Now we show Ψ is \mathbb{R} -linear. Thus, for every $\eta \in \mathbb{R}$, there exist two rational sequences $\{r_n\}, \{s_n\}$ such that $r_n \leq \eta \leq s_n$ and $\lim r_n = \lim s_n = \eta$ when $n \rightarrow \infty$. It is clear that Ψ preserves positive elements, then Ψ preserves order. So, by the additivity of Ψ , we have

$$r_n I = \Psi(r_n I) \leq \Psi(\eta I) \leq \Psi(s_n I) = s_n I.$$

Hence,

$$\Psi(\eta I) = \eta I$$

for $\eta \in \mathbb{R}$. It means that Ψ is \mathbb{R} -linear. Thereby the proof is completed. \square

Lemma 3.13. *The restriction of Ψ to $M\mathfrak{P}$ is linear and restriction to $M(I - \mathfrak{P})$ is conjugate linear.*

Proof. By Lemma 3.12, $\Psi(iI)^2 = \Psi((iI)^2) = -\Psi(I) = -I$. Also by Lemma 3.8(c), $\Psi(iI)^* = \Psi((iI)^*) = -\Psi(iI)$. Let $F = \frac{I - i\Psi(iI)}{2}$. Then it is easy to verify that F is a central projection in M . Let $\mathfrak{P} = \Psi^{-1}(F)$. Then by Lemma 3.8(d), \mathfrak{P} is a central projection in N . Moreover, for $A \in N$, there hold

$$\Psi(iA\mathfrak{P}) = \Psi(A)\Psi(\mathfrak{P})\Psi(iI) = i\Psi(A)\Psi(\mathfrak{P})(2F - I),$$

and

$$\Psi(iA(I - \mathfrak{P})) = \Psi(A)\Psi(I - \mathfrak{P})\Psi(iI) = -i\Psi(A)(I - F) = -i\Psi(A(I - \mathfrak{P})).$$

That is, the restriction of Ψ to $M\mathfrak{P}$ is linear and restriction to $M(I - \mathfrak{P})$ is conjugate linear. This together with Lemmas 3.8, 3.11 and 3.12 completes the proof of Theorem 1.1. \square

Acknowledgment. The authors wish to give their thanks to the referees for their helpful comments and suggestions that make much improvement of the paper. This work was supported by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, under grant No. **(G-212-662-1441)**. The authors, therefore, gratefully acknowledge DSR technical and financial support.

References

- [1] Z. Bai and S. Du, *Maps preserving products $XY - YX^*$ on von Neumann algebras*, J. Math. Anal. Appl. **386**, 103-109, 2012.
- [2] Z. Bai and S. Du, *Multiplicative Lie isomorphism between prime rings*, Comm. Algebra **36**, 1626-1633, 2008.
- [3] Z. Bai and S. Du, *Multiplicative $*$ -Lie isomorphism between factors*, J. Math. Anal. Appl. **346**, 327-335, 2008.

- [4] L. Dai and F. Lu, *Nonlinear maps preserving Jordan *-products*, J. Math. Anal. Appl. **409**, 180-188, 2014.
- [5] D. Huo, B. Zheng and H. Liu, *Nonlinear maps preserving Jordan triple η -*-products*, J. Math. Anal. App. **430** (2), 830-844, 2015.
- [6] D. Huo, B. Zheng, J. Xu and H. Liu, *Nonlinear mappings preserving Jordan multiple *-product on factor von Neumann algebras*, Linear Multilinear Algebra, **63** (5), 1026-1036, 2015.
- [7] P. Ji and Z. Liu, *Additivity of Jordan maps on standard Jordan operator algebras*, Linear Algebra Appl., **430** (1), 335-343, 2009.
- [8] C. Li and F. Lu, *Nonlinear maps preserving the Jordan triple 1-*-product on von Neumann algebras*, Complex Anal. Oper. Theory, **11**, 109-117, 2017.
- [9] C. Li, F. Lu and X. Fang, *Nonlinear mappings preserving product $XY + YX^*$ on factor von Neumann algebras*, Linear Algebra Appl. **438** (5), 2339-2345, 2013.
- [10] C. Li, F. Lu and T. Wang, *Nonlinear maps preserving the Jordan triple *-product on von Neumann algebras*, Ann. Func. Anal. **7**, 496-507, 2016.
- [11] F. Lu, *Additivity of Jordan maps on standard operator algebras*, Linear Algebra Appl. **357** (1-3), 123-131, 2002.
- [12] L. Yaoxian and Z. Jianhua, *Nonlinear mixed Lie triple derivation on factor von Neumann algebras*, Acta Math. Sin. Chinese Ser. **62** (1), 13-24, 2019.
- [13] C.R. Miers, *Lie homomorphisms of operator algebras*, Pacific. J. Math. **38** (3), 717-735, 1971.
- [14] W. S. Martindale III, *When are multiplicative mappings additive?*, Proc. Amer. Math. Soc. **21**, 695-698, 1969.
- [15] A. Taghavi, V. Darvish and H. Rohi, *Additivity of maps preserving products $AP \pm PA^*$ on C^* -algebras*, Math. Slovaca, **67**, 213-220, 2017.
- [16] A. Taghavi, H. Rohi and V. Darvish, *Additivity of maps preserving Jordan η_* -products on C^* -algebras*, Bull. Iranian Math. Soc. **41**, 107-116, 2015.
- [17] P. Šmerl, *Quadratic and quasi-quadratic functionals*, Proc. Amer. Math. Soc. **119**, 1105-1113, 1993.
- [18] Z. Yang and Y. Zhang, *Nonlinear maps preserving the second mixed Lie triple products on factor von Neumann algebras*, Linear Multilinear Algebra, **68**(2), 377-390, 2020.
- [19] Z. Yang and Y. Zhang, *Nonlinear maps preserving mixed Lie triple products on factor von Neumann algebras*, Ann. Funct. Anal. **10**, 325-336, 2019.
- [20] Y. Zhou, Z. Yang and J. Zhang, *Nonlinear mixed Lie triple derivation on prime *-rings*, Comm. Algebra, **47**, 4791-4796, 2019.