



# International Journal of Secondary Metabolite

Volume: 5 Number: 1  
January 2018

ISSN-e: 2148-6905

Journal homepage: <http://www.ijate.net/>

<http://dergipark.gov.tr/ijsm>

## The Effect of Increasing Mycorrhiza Applications on Nutrition of Pak Choi (*Brassica rapa* L. subsp. *chinensis* L.) Plant

Sevinc Adiloglu, Funda Eryilmaz Acikgoz, Funda Irmak Yilmaz, Yusuf Solmaz, Aydin Adiloglu

**To cite this article:** Adiloglu, S., Eryilmaz Acikgoz, F., Irmak Yilmaz, F., Solmaz, Y., Adiloglu, A. (2018). The Effect of Increasing Mycorrhiza Applications on Nutrition of Pak Choi (*Brassica rapa* L. subsp. *chinensis* L.) Plant. *International Journal of Secondary Metabolite*, 5(1), 27-33. DOI: [10.21448/ijsm.351753](https://doi.org/10.21448/ijsm.351753)

**To link to this article:** <http://www.ijate.net/index.php/ijsm/issue/archive>  
<http://dergipark.gov.tr/ijsm>

This article may be used for research, teaching, and private study purposes.

Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Authors alone are responsible for the contents of their articles. The journal owns the copyright of the articles.

The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of the research material.

Full Terms & Conditions of access and use can be found at  
<http://ijate.net/index.php/ijsm/about>

## The Effect of Increasing Mycorrhiza Applications on Nutrition of Pak Choi (*Brassica rapa* L. subsp. *chinensis* L.) Plant

Sevinc Adiloglu<sup>1\*</sup>, Funda Eryilmaz Acikgoz<sup>2</sup>, Funda Irmak Yilmaz<sup>3</sup>,  
Yusuf Solmaz<sup>1</sup>, Aydin Adiloglu<sup>1</sup>

<sup>1</sup>Department of Soil Science and Plant Nutrition, Namik Kemal University, Tekirdag, Turkey

<sup>2</sup>Department of Plant and Animal Production, Namik Kemal University, Tekirdag, Turkey

<sup>3</sup>Department of Soil Science and Plant Nutrition, Ordu University, Ordu, Turkey

**Abstract:** The study was done to determine the effect of increasing mycorrhiza application on some macro and micro nutrient element contents of pak choi (*Brassica rapa* L. subsp. *chinensis* L.) plant. According to the experiment results, important increases in some macro and micro nutrient element contents of pak choi plant were determined with increasing mycorrhiza applications. The contents were determined as P (0.38 %, 0.42 %, 0.45 %, 0.49 % and 0.51 %), K (4.01 %, 4.30 %, 4.41 %, 4.56 % and 4.70 %), Ca (1.83 %, 2.01 %, 2.06 %, 2.20 % and 2.36 %), Mg (0.14 %, 0.15 %, 0.15 %, 0.16 % and 0.16 %), Fe (309 mgkg<sup>-1</sup>, 417 mgkg<sup>-1</sup>, 678 mgkg<sup>-1</sup>, 1009 mgkg<sup>-1</sup> and 1696 mgkg<sup>-1</sup>), Cu (5.49 mgkg<sup>-1</sup>, 6.10 mgkg<sup>-1</sup>, 6.53 mgkg<sup>-1</sup>, 7.05 mgkg<sup>-1</sup> and 7.63 mgkg<sup>-1</sup>), Mn (45.90 mgkg<sup>-1</sup>, 52.23 mgkg<sup>-1</sup>, 60.20 mgkg<sup>-1</sup>, 70.40 mgkg<sup>-1</sup> and 80.00 mgkg<sup>-1</sup>) and Zn (32.23 mgkg<sup>-1</sup>, 35.40 mgkg<sup>-1</sup>, 37.00 mgkg<sup>-1</sup>, 40.70 mgkg<sup>-1</sup> and 46.86 mgkg<sup>-1</sup>) at I. dose, (control): 0 ml plant<sup>-1</sup>, II. dose: 15 ml plant<sup>-1</sup>, III. dose: 20 ml plant<sup>-1</sup>, IV. dose: 30 ml plant<sup>-1</sup> and V. dose: 40 ml plant<sup>-1</sup>, respectively. These P, K, Ca and Mg contents increases were determined significant at the level of P<0.05, statistically. The highest nutrient element contents of pak choi plant were obtained V. dose: 40 ml plant<sup>-1</sup> applications for P, K, Ca, Mg, Fe, Cu, Mn and Zn nutrient elements.

### ARTICLE HISTORY

Received: 11 July, 2017

Revised: 07 November, 2017

Accepted: 11 November, 2017

### KEYWORDS

Mycorrhiza,  
Macro and micro element,  
Pak choi (*Brassica rapa* L.  
subsp. *chinensis* L.),  
Exotic vegetable,

### 1. Introduction

As a highly rated leafy variety of vegetables and a marvelous food alternative, brassicas is grown for its enlarged, edible, terminal buds; and is preferably eaten almost everywhere in the world as well [1]. This green vegetable was made known around the world by the efforts of the travelers and immigrants [2-4].

\*Corresponding Author E-mail: [sadiloglu@hotmail.com](mailto:sadiloglu@hotmail.com)

Pak choi name was given syn. *Brassica chinensis* L. (1759), *Brassica campestris* L. subsp. *chinensis* (L.) [5], *Brassica rapa* L. subsp. *chinensis* (L.) [6] in China. This vegetable which has been known since the 5<sup>th</sup> A. D. and grown widely in China and Taiwan is classified within the group which is identified as Chinese vegetables [7, 8]. Its leaves are edible and it has 50-60 days of vegetation approximately [9, 10].

A greenhouse experiment was made for the effect of increasing mycorrhiza application on the some nutrient element contents of maize plant (*Zea mays* L.) [11]. According to the research results, N, K, Fe and Cu contents of maize plant increased with increasing mycorrhiza application.

The effect of salt (0 and 100 mg Na Cl kg<sup>-1</sup>) and increasing zinc applications (0, 25, 50 mg Zn kg<sup>-1</sup>) application on phosphorus and zinc uptake of maize (*Zea mays* L.) plant with mycorrhiza and non-mycorrhiza conditions. At the end of the experiment it was determined that mycorrhiza inoculated applications provided a significant increase in fresh weight, dry weight, phosphorus and zinc contents compared to the non-mycorrhiza applications [12].

The research was done to determine the effect of increasing mycorrhiza application on some macro and micro nutrient element contents of pak choi plant.

## 2. Material and Methods

Using high tunnel cold greenhouse covered by polyetilen (PE) with UV additive which belongs to Namik Kemal University, Vocational College of Technical Sciences, Plant and Animal Production Department, the experiments were carried out autumn in Tekirdag city (40°98' N, 27°48' E) Turkey in 2016.

Research was designed as three replications according to randomized block experimental design. The white mini variety of Pak Choi (Zengarden Firm) was used for the research (Figure 1). Seeds were sown in multi-celled trays filled with peat (Klasmann-Deilmann, Potground H, Germany) in October. Some properties of the used peat are: 160-260 mg l<sup>-1</sup> N, 180-280 mg l<sup>-1</sup> P<sub>2</sub>O<sub>5</sub>, 200-150 mg l<sup>-1</sup> K<sub>2</sub>O, 80-150 mg l<sup>-1</sup> Mg, pH: 6, 70 % organic matter and 35 % C. When the seedlings became 2 to 3 true leaves (21<sup>th</sup> days for pak choi after seed sowing) they were planted to pre-prepared places in high tunnel cold greenhouse with 10x10 cm intervals and 10 plants in each parcel.

Then five mycorrhiza doses (I. dose, (control): 0 mlplant<sup>-1</sup>, II. dose: 15 mlplant<sup>-1</sup>, III. dose: 20 mlplant<sup>-1</sup>, IV. dose: 30 mlplant<sup>-1</sup> and V. dose: 40 mlplant<sup>-1</sup>) were applied before a month from harvesting time and plants were harvested 54 days after seed sowing. Some macro and micro nutrient elements (P, K, Ca, Mg, Fe, Cu, Mn and Zn) contents of plants were determined via ICP-OES instrument [13]. Then experiment analysis results were evaluated SPSS 21 statistically program. ANOVA variance analysis was done and Duncan multiple comparison tests were done on this research results.

Some chemical properties of the soil samples which used in this experiment it can be seen in Table 1. The some climate data measured inside the tunnel during the growing of the plants it can be seen in Table 2. Since there were no diseases and pests, no pesticides was used during the growing period.



**Figure 1.** A general view from pak choi experiment and post-harvested plant (Original).

According to the Table 1, pH value alkaline, salt less, low organic matter content, little lime, medium available phosphorus, exchangeable K, Ca and Mg sufficient, also, micro element (Mn, Cu, Fe and Zn) contents are sufficient of experiment area soil

**Table 1.** Some chemical properties of soil samples

Soil properties	Results
pH	8.01
Salinity (%)	0.07
CaCO <sub>3</sub> (%)	2.74
Organic matter (%)	1.35
Ca (%)	0.54
P (mgkg <sup>-1</sup> )	36.40
K (mgkg <sup>-1</sup> )	253.80
Mg (mgkg <sup>-1</sup> )	473.10
Mn (mgkg <sup>-1</sup> )	5.68
Cu (mgkg <sup>-1</sup> )	0.81
Fe (mgkg <sup>-1</sup> )	7.43
Zn (mgkg <sup>-1</sup> )	0.97

**Table 2.** Average some climate data in unheated greenhouse during the experiment period.

Month	Average temperature (°C)	Maximum temperature (°C)	Minimum temperature (°C)	Average humidity (%)
October	16.50	19.03	14.01	87
November	12.30	15.01	9.60	89
December	7.40	10.50	4.30	90

### 3. Results and Discussion

The effect of increasing mycorrhiza application on some macro nutrient element content of pak choi plant is given in Table 3. According to the Table 3, phosphor content of pak choi plant was obtained 0.38 % and 0.51 % for I. dose and V. dose, respectively. Potassium content of pak choi plant was obtained 4.01 % and 4.70 % for I. dose and V. dose, respectively.

Calcium content of pak choi plant was obtained 1.83 % and 2.36 % for I. dose and V. dose, respectively. Magnesium content of pak choi plant was obtained 0.14 % and 0.16 % for I. dose and V. dose, respectively.

P, K, Ca and Mg contents of pak choi plant increased with increasing mycorrhiza application (Table 3). These increases were found statistically significant at the level of  $p < 0.05$ . These results were found to be some earlier researchers [11, 14].

**Table 3.** The effect of increasing mycorrhiza application on some macro nutrient element contents of pak choi plant, (%)

Mycorrhiza doses	P	K	Ca	Mg
I.dose (Control)	0.38±0.31 <b>c</b>	4.01±0.18 <b>d</b>	1.83±0.77 <b>d</b>	0.14±0.07 <b>a</b>
II. dose	0.42±0.13 <b>b</b>	4.30±0.30 <b>c</b>	2.01±0.29 <b>c</b>	0.15±0.21 <b>ab</b>
III. dose	0.45±0.08 <b>b</b>	4.41±0.51 <b>bc</b>	2.06±0.11 <b>c</b>	0.15±0.13 <b>ab</b>
IV. dose	0.49±0.09 <b>a</b>	4.56±0.73 <b>ab</b>	2.20±0.99 <b>b</b>	0.16±0.15 <b>a</b>
V. dose	0.51±0.06 <b>a</b>	4.70±0.59 <b>a</b>	2.36±0.62 <b>a</b>	0.16±0.80 <b>a</b>

\*: values are average of three replications,  
 \*\*: each parameter evaluated individually,  
 \*\*\*:  $p < 0.05$

The effect of increasing mycorrhiza application on some micro nutrient element content of pak choi plant is given in Table 4. According to the Table 4, iron content of pak choi plant was obtained 309 mgkg<sup>-1</sup> and 1696 mgkg<sup>-1</sup> for I. dose and V. dose, respectively. Copper content of pak choi plant was obtained 5.49 mgkg<sup>-1</sup> and 7.63 mgkg<sup>-1</sup> for I. dose and V. dose, respectively.

Zinc content of pak choi plant was obtained 32.23 mgkg<sup>-1</sup> and 46.86 mgkg<sup>-1</sup> for I. dose and V. dose, respectively. Manganese content of pak choi plant was obtained 45.90 mgkg<sup>-1</sup> and 80.00 mgkg<sup>-1</sup> for I. dose and V. dose, respectively.

Fe, Cu, Zn and Mn contents of pak choi plant increased with increasing mycorrhiza application according to the Table 4. These increases were found statistically significant at the level of  $p < 0.05$ . These results were found to be some earlier researchers [15-17].

**Table 4.** The effect of increasing mycorrhiza application on some micro nutrient element contents of pak choi plant, mgkg<sup>-1</sup>

Mycorrhiza doses	Fe	Cu	Zn	Mn
I.dose (Control)	309±4.3d	5.49±0.02d	32.23±0.10d	45.90±0.4d
II. dose	417±3.9d	6.10±0.02c	35.40±0.10c	52.23±0.2d
III. dose	678±5.5c	6.53±0.01c	37.00±0.02c	60.20±0.4c
IV. dose	1009±19.3b	7.05±0.08b	40.70±0.10b	70.40±0.1b
V. dose	1696±4.9a	7.63±0.04a	46.86±0.30a	80.00±0.5a

\*: values are average of three replications,

\*\* : each parameter evaluated individually,

\*\*\*: p<0.05

Mycorrhiza could also enhance crop quality not only by enrichment in macronutrients (like P) [18, 19] and in micronutrients [20-22]. Mycorrhiza yield mineral element contents are decreasing, compromising the nutritional value of food. The use of beneficial bacteria results in an increase in the concentration or availability of mineral elements in the production. Micro nutrient is regarded as a promising strategy to overcome malnutrition in terms of nutrition [20, 23]. Beneficial bacteria use, macro and micronutrient element affects the soil intake positively [24, 25]. According to Gianinazzi and Gollotte, 2010 [26] increasing mineral nutrient and water uptake by plants. And promote plant growth while reducing fertilizer requirement.

#### 4. Conclusion

According to this greenhouse experiment results, increasing doses of mycorrhiza application was increased some macro (N, P, K, Ca and Mg) and micro (Fe, Cu, Zn and Mn) element contents of pak choi plant. These increases were found to be statistically significant at the level of p<0.05. This result is very important nutrition and quality of pak choi plant. Because, excess chemical fertilizers application to the agricultural soils was destroyed soil fertility and quality. Therefore, mycorrhiza, another different microbial fertilizers, other organic fertilizers and organic materials should be applied to the agricultural soils for quality plant production and sustainable soil fertility. Also, according to the earlier research result, mycorrhiza applications were improved of some physical, chemical and biological properties of the agricultural soils.

#### Acknowledgement

The abstract of this paper has been presented in The 3rd International Symposium on Eurasian Biodiversity (SEAB 2017).

#### 5. References

- [1] Tirasoglu, E., Cevik, U., Ertugrul, B., Apaydin, G., Baltas, H., Ertugrul, M. (2005). Determination of trace elements in cole (*Brassica oleraceae* var. *acephale*) at Trabzon region in Turkey. *J. Quantitative, Spectroscopy, Radiative Transfer*, 94, 181-187.
- [2] Nieuwhof, M. (1969). *Cole Crops*. Leonard Hill, London, 102-104.
- [3] Balkaya, A., Yanmaz, R. (2005). Promising kale (*Brassica oleracea* var. *acephala*) populations from Black Sea region, Turkey. *New Zealand Journal of Crop and Horticultural Science*, 33, 1-7.
- [4] Adiloglu, S., Eryilmaz Acikgoz, F., Adiloglu, A. (2015). The effect of increasing doses of N application on some agronomic characteristics, vitamin C, protein and mineral content

- of mibuna (*Brassica rapa* var. *nipposinica*) and mizuna (*Brassica rapa* var. *japonica*) plants. *Journal of Agricultural Faculty, Uludag University*, 29 (2), 1-11.
- [5] Makino, T. (1912). Observations on the flora of Japan. *Botanical Magazine of Toky.*, 23, 93-102.
- [6] Hanelt, P. (1986). *Formal and informal classifications of the infraspecific variability of cultivated plants-advantages and limitations*. In: Styles, B.T. (ed.), *Infraspecific Classification of Wild and Cultivated Plants*, 139-156. Oxford.
- [7] Dixon, G.R. (2007). *Vegetable brassicas and related crucifers*. CAB International North American Office, 875 Massachusetts Avenue, 7th Floor, Cambridge, MA 02139 USA., 327.
- [8] Eryilmaz Acikgoz, F. (2016). Seasonal variations on quality parameters of Pak Choi (*Brassica rapa* L. subsp. *chinensis* L.). *Advances in Crop Science and Technology*, 4, 4.
- [9] Opena, R.T., Kuo C.G., Yoon, J.Y. (1988). Breeding and seed production of Chinese cabbage in the tropics and subtropics. *AVRDC, Shanhua, Tairan, Tech Bull.*, 17, 92-95.
- [10] Larkcom, J. (2008). *Oriental vegetables*. Frances Lincoln Ltd., London, UK.
- [11] Karatekin, M. (2014). *Effect of the applications mycorrhizal and bacterial (Bacillus sp 189) inoculation and sulfur on phosphorus usefulness of a calcareous soil*. Selcuk University Graduate School of Natural and Applied Sciences, Dept. of Soil Science, MSc. Thesis.
- [12] Sonmez, F., Cıg, F., Erman, M., Tufenkci, S. (2013). Effects of zinc, salt and mycorrhiza applications on the development and the phosphorus and zinc uptake of maize. *Yuzuncuyil University, Journal of Agricultural Sciences*, 23(1), 1-9.
- [13] Kacar, B., Inal, A. (2010). *Plant Analysis*. Nobel Publish., Ankara, 849, 659.
- [14] Koide, R.T. (1991). Nutrient supply, nutrient demand and plant response to mycorrhizal infection. *New Phytologist*, 117, 365- 386.
- [15] Arancon, N.Q., Edwards, C.A., Bierman, P., Welch, C., Metzger, J.D. (2004). The influence of vermicompost applications to strawberries on growth and yield. *Bioresource Technology*, 93, 145-153.
- [16] Smith, S., Read, D.J. (2008). *Mycorrhizal Symbiosis*. Academic Press, San Diego, CA.
- [17] Kucukyumuk, Z., Gultekin, M., Erdal, I. (2014). Effects of vermicompost and mycorrhiza on plant growth and mineral nutrition in pepper. *Süleyman Demirel University, Journal of Agricultural Faculty*, 9 (1), 51-58.
- [18] Karandashov, V., Bucher, M. (2005). Symbiotic phosphate transport in arbuscular mycorrhizas. *Trends Plant Sci.*, 10, 22-29.
- [19] Veresoglou, S.D., Chen, B., Rillig, M.C. (2012). Arbuscular mycorrhiza and soil nitrogen cycling. *Soil Biol., Biochem*, 46, 53-62.
- [20] White, P.J., Broadley, M.R. (2009). Biofortification of crops with seven mineral elements often lacking in human diets e iron, zinc, copper, calcium, magnesium, selenium and iodine. *New Phytol.*, 182, 49-84.
- [21] Antunes, P.M., Franken, P., Schwarz, D., Rillig, M.C., Cosme, M., Scott, M., & Hart, M.M. (2012). *Linking soil biodiversity and human health: do arbuscular mycorrhizal fungi contribute to food nutrition*. In: Wall, D.H. et al. (Eds.), *Soil Ecology and Ecosystem Services*. Oxford University Press, Oxford, UK, 153-172.
- [22] He, X., Kazuhide Nara, K. (2007). Element biofortification: can mycorrhizas potentially offer a more effective and sustainable pathway to curb human malnutrition? *Trends Plant Sci.*, 12, 331-333.

- [23] Pellegrino E., Bedini S. (2014). Enhancing ecosystem services in sustainable agriculture: Bio fertilization and bio fortification of chickpea (*Cicer arietinum* L.) by arbuscular mycorrhiza fungi. *Soil Biology & Biochemistry*, 68, 429-439.
- [24] Weber, E., Saxena, M.C., George, E., & Marschner, H. (1993). Role of vesicular-arbuscular mycorrhiza in the mineral nutrition of chickpea (*Cicer arietinum* L.) grown in Northern Syria. *Field Crops Research*, 32 (1-2), 115-128.
- [25] Gianinazzi, S., Trouvelot, A., Gianinazzi-Pearson, V. (1990). Role and use of mycorrhizas in horticultural crop production. *Advances Horticultural Science*, 4, 25-30.
- [26] Gianinazzi, S., Gollotte, A. (2010) Agroecology: the key role of arbuscular mycorrhizas in ecosystem services. *Mycorrhiza*, 20, 519-530.