

Experimental and Theoretical Investigation of Molecular Structure and Vibrational Frequencies of 3-Cyano-7-hydroxycoumarin by Density Functional Theory

Adnan SAĞLAM*¹

¹Bartın University, Engineering Faculty, Electrical Electronical Engineering Department, 74100, Bartın

(Alınış / Received: 04.05.2017, Kabul / Accepted: 19.07.2017, Online Yayınlanma / Published Online: 15.08.2017)

Keywords

3C7HC,
Vibrational frequencies,
IR,
Raman spectra,
DFT

Abstract: In this research, geometrical structure and vibrational spectra of 3-cyano-7-hydroxycoumarin (3C7HC) were studied experimentally and theoretically. Experimentally, FT-IR (infrared) and FT-R (Raman) spectra of 3C7HC molecule were respectively obtained in the range of 4000-400 cm⁻¹ and 3500-100 cm⁻¹. Theoretically, the optimized geometric structures, vibrational spectra and corresponding assignments according to two possible configurations of the 3C7HC molecule were investigated using the density functional method (B3LYP) with 6-311++G (d,p) basis set. When analyzes were performed, C_s symmetry property of 3C7HC molecule was used. It was seen that the experimental and calculated data of the 3C7HC molecule were highly compatible with each other.

Yoğunluk Fonksiyonel Teorisi (DFT) ile 3-Cyano-7-hydroxycoumarin molekülünün Titreşim Frekansı ve Moleküler Yapısı Üzerine Teorik ve Deneysel Bir Çalışma

Anahtar Kelimeler

3C7HC,
Titreşim frekansı,
IR,
Raman spektrumu,
DFT

Özet: Bu çalışmada 3-siyano-7-hidroksikoumarin (3C7HC) molekülünün geometrik yapısı ve titreşim spektrumu deneysel ve teorik olarak incelendi. Deneysel olarak 3C7HC molekülünün FT-IR (infrared) ve FT-R (Raman) spektrumu, sırasıyla 4000-400 cm⁻¹ ve 3500-100 cm⁻¹ aralığında elde edildi. Teorik olarak, 3C7HC molekülünün iki muhtemel konfigürasyonuna göre optimize geometrik yapıları, titreşim spektrumları ve bunlara karşılık gelen atamalar, yoğunluk fonksiyon teorisi (DFT) ile B3LYP 6-311++G(d,p) temel parametreleri kullanılarak yapıldı. Analizlerde, 3C7HC molekülünün C_s simetri özelliği kullanıldı. 3C7HC molekülünün deneysel ve hesaplanan verilerinin birbirleri ile oldukça uyumlu olduğu görüldü.

1. Introduction

3-Cyano-7-hydroxycoumarin is known as 3-Cyanoumbelliferone, 7-hydroxy-2-oxo-1-benzopyran-3-carbonitrile and 7-hydroxy-2-oxochromene-3-carbonitrile, respectively. It is the derivative molecule of Coumarine, having a chemical formula of C₁₀H₅NO₃. Coumarine derivatives exhibit various biological activities and therapeutic properties. As a result of their presence in the chemical structures, they have been extensively studied in biomedical and industrial areas [1-14]. In recent years, the vibrational spectra and geometrical structures of Coumarine molecules have been experimentally and theoretically studied by many researchers. DFT method has been widely used in investigation of electronic molecular system. Density functional theory is based on the assumption that the electron density of an electronic system's energy can be defined using probability terms [15-

14]. In the literature, no study was found on experimental and theoretical investigation of 3C7HC molecule.

In this study, experimental and theoretical conformations of 3C7HC molecule were carried out. The vibrational frequencies and geometrical structure of 3C7HC molecule were calculated with DFT/B3LYP method using 6-311++G(d,p) basis set. The calculated geometrical parameters and vibrational frequencies were compared with experimental results and previous studies.

2. Material and Method

2.1. Experimental details

3-cyano-7-hydroxycoumarin with 97% purity was purchased commercially from Sigma-Aldrich

*Corresponding author: adnansaglam@bartin.edu.tr

Company USA and used in pure form without further purification. The 3C7HC sample is found as powder at room temperature. 3C7HC was pelleted with KBr at room temperature using approximately 10 MPa pressure. The FT-IR spectrum of the 3C7HC was obtained within 4000-100 cm^{-1} range with the Perkin Elmer Spectrometer. FTR spectrum of 3C7HC was obtained within the range of 3200-100 cm^{-1} with Jasco NRS-3100 Raman Spectrometer.

2.2. Computational method

Optimized geometric structures, vibrational frequencies, zero-point energies and dipole moments according to two different conformation of 3C7HC were calculated with B3LYP 6-311++G(d,p) basis parameters of DFT method. The computations were made with Gaussian 09 [25] and Gauss-View software [26]. While the vibrational frequencies of 3C7HC were calculated theoretically, the 0.9614 scale was used [27].

3. Results and Discussion

3.1. Molecular geometry

3C7HC molecule has a planar geometric structure represented by the Cs symmetry which consists of identity E and a mirror plane σ elements. In this study, two probable conformers of optimized geometrical structures of the 3C7HC molecule with hydroxyl group (O-H) oriented in line with C-O bonding were investigated. When O-H is farther away

from the oxygen atom of the pyrone ring of 3C7HC, the conformer is called as rotamer (Rot1). When O-H is closer to the oxygen atom, the conformer is called as rotamer2 (Rot2). The optimized geometrical structures of the Rot1 and Rot2 conformers of 3C7HC molecule are shown in Fig. 1. Orientation angles of O-H for conformers of 3C7HC are 110.6° for Rot1 and 249.6° for Rot2.




Figure 1. Rot1 and Rot2 orientations of the 3C7HC molecule.

In the case of Rot1 and Rot2 of the 3C7HC molecule the zero-point, relative energy and dipole moments are given in Table 1.

The relative energy between two conformers is considerably low and it has a value of ~ 0.32 kcal/mol. As indicated by zero-point energies and dipole moments of the molecule, Rot1 conformer is more stable. Experimental [28, 29] and computed geometric parameters of conformers of 3C7HC molecule were compared in Table 2. The largest differences in computed and experimental parameters were obtained as Rot1 (0.04 Å), Rot2 (0.042 Å) for the bond lengths, and Rot1 (1.7°), Rot2 (1.8°) for the bond angles.

Table 1. Zero-point, relative energy and dipole moment of 3C7HC

Conformers	Zero-point energy [Hartree/particle]	Relative energy [kcal/mole]	Dipole moment [Debye]
Rot1	-664.537840		8.0795
Rot2	-664.537338	0.314799	9.7508

Table 2. Rot1 and Rot2 conformers of 3C7HC

Parameters	Exp. ^a	Calculated		Parameters	Exp. ^a	Calculated	
		Rot1	Rot2			Rot1	Rot2
Bond lengths (Å)				Bond angles (°)			
O1-C2	1.367	1.399	1.397	O1-C2-C3	118.2	118	118.1
C1-C9	1.382	1.359	1.36	O1-C2-O11	115.9	115.2	115.3
C2-C3	1.433	1.47	1.471	C2-C3-C4	120.6	121.1	121.1
C2-O11	1.224	1.197	1.197	C2-C3-C12	-	117.5	117.5
C3-C4	1.341	1.365	1.364	C3-C12-N13	178.5 ^b	178.3	178.2
C3-C12	1.430 ^b	1.425	1.425	C3-C4-C10	121	121.2	121.1
C12-N13	1.145 ^b	1.155	1.155	C3-C4-H15	-	119.6	119.6
C4-C10	1.424	1.424	1.425	C4-C10-C9	118.2	117.7	117.8
C4-H15	-	1.085	1.085	C10-C5-C6	120.7	121.1	120.9
C10-C5	1.411	1.411	1.408	C10-C5-H16	-	119	119.1
C10-C9	1.368	1.408	1.41	C5-C6-C7	120.7	119.5	119.6
C5-C6	1.393	1.377	1.38	C5-C6-H17	-	121.7	120.4
C5-H16	-	1.084	1.084	C6-C7-C8	120.6	120.9	120.9
C6-C7	1.396	1.408	1.408	C6-C7-O14	-	116.6	122.2
C6-H17	-	1.082	1.085	C7-O14-H18	-	110.6	110.5
C7-C8	1.398	1.393	1.393	C7-C8-C9	118	118.7	118.7
C7-O14	1.351	1.356	1.357	C7-C8-H19	-	122	120.8
O14-H18	-	0.964	0.964	C8-C9-C10	123	121.7	121.8
C8-C9	1.376	1.39	1.387	C8-C9-O1	117	117.2	117.3
C8-H19	-	1.084	1.082	C2-O1-C9	121.9	123.6	123.7
Rot1 angle = 110.6°				Rot2 angle = 249.6°			

^aTaken from Ref.[28] ^bRef.[29]

3.2. Vibrational analysis

3C7HC molecule has 19 atoms and is in the Cs symmetry. 19 atoms' three cartesian displacements were provided with 57 internal modes that is;

$$\Gamma_{\text{inter.}} = 38A' + 19A'' \quad (1)$$

In the event of translation ($\Gamma_{\text{trans.}} = A' + 2A''$) and rotation ($\Gamma_{\text{rot.}} = 2A' + A''$) from Cs point group table, vibrational modes are expressed as;

$$\Gamma_{\text{vib.}} = \Gamma_{57} - \Gamma_{\text{trans.}} - \Gamma_{\text{rot.}} = 35A' + 16A'' \quad (2)$$

In infrared (FT-R) and Raman (FT-R) all vibrational values are active. In the Cs symmetry, the 2N-3 of the vibrations is in plane and the N-3 of the vibrations is out of plane [30]. As a result, 35 out of 51 vibrational frequencies of the 3C7HC molecule are in plane and 16 of them are out of plane. Out of plane vibrations of the molecule are represented with A'' symmetries, while planar vibrations are represented with A' symmetries. This is clearly indicated in visual examination of vibrations performed with Gauss-View visualization program.

FT-IR and FT-R spectra of 3C7HC molecule are given in Fig.2 and Fig. 3. Observed and computed vibrational frequencies, Raman activities, infrared intensities, in plane -out of plane assignments are given in Table 3. Correlation values of vibrational frequencies for both conformers are given in the Table. The obtained results, normal coordinate analysis and the vibrational frequencies are significantly consistent with the results of previous studies.




Figure 2. FT-IR spectrum of 3C7HC molecule.

3.2.1. C-H vibrations

CH stretching vibrations in the aromatic molecular structures widely occur in the form of weak multiple bands within 3100-3000 cm⁻¹ interval. This region generally represents C-H stretching vibrations [31]. In this region the characteristics of substituent have no significant effect on the bands and four C-H stretching vibrations occur in 3C7HC molecule. In this study, aromatic C-H stretching vibrations of 3C7HC were obtained as 3090, 3043 cm⁻¹ in FT-IR and

3089 cm⁻¹ in FT-R. In plane C-H bending vibrations are related with C-C stretching vibrations and the vibrations of aromatic molecular structures demonstrated in 1500-1100 cm⁻¹ region [32]. C-H in plane bending vibrations of 3C7HC were obtained as 1611, 1566, 1505, 1460, 1385, 1374, 1318, 1272, 1230, 1203, 1150, 1123, 1066, 976, 953 cm⁻¹ at FT-IR and 1608, 1596, 1557, 1459, 1374, 1314, 1273, 1233, 1160, 1127, 1068, 980 cm⁻¹ at FT-R. The absorption bands stemming from C-H out of plane bending vibrations generally occurred within 1000-750 cm⁻¹ region [33, 34]. The observed bands 885, 825, 765 cm⁻¹ in FT-IR and 741 cm⁻¹ in FT-R were assigned to C-H out of plane bending vibrations of 3C7HC.




Figure 3. FT-R spectrum of 3C7HC molecule.

3.2.2. O-H vibrations

O-H bending and stretching vibrations can be identified with stretching vibration of the frequency band that is related to the hydrogen bonding extent. Because the oxygen atom is lighter than the hydrogen atom, the stagnation of the molecular structure had no effect on the O-H stretching vibration. O-H stretching vibrations are demonstrated by strong absorption bands in 3700-3584 cm⁻¹ region [16]. In this study, the O-H stretching vibration was obtained as 3690 cm⁻¹ in FT-IR. Generally, O-H in plane bending stretching vibrations are found with C-H in plane and CC stretching vibrations. O-H in plane bending stretching vibrations demonstrated in 1617-1114 cm⁻¹ region [16, 37]. In plane bending stretching vibrations were obtained as 1566, 1505, 1460, 1385, 1374, 1318, 1274, 1230, 1150, 1123, 1066, 976 cm⁻¹ at FT-IR and 1596, 1557, 1459, 1374, 1314, 1273, 1233, 1160, 1127, 1068, 980 cm⁻¹ at FT-R. O-H out of plane bending vibrations demonstrated as in 450-350 cm⁻¹ region [35-38]. Out of plane bending vibration was obtained as 359 cm⁻¹ at FT-R. It has been verified that the experimental and calculated O-H vibrations are in accordance with the previous studies.

3.2.3. CO vibrations

CO stretching vibrations are called carbonyl groups and have a decisive feature. C=O stretching vibrations are widely studied with FT-IR and FT-R. The C=O stretching vibrations are demonstrated in 1780-1700

cm⁻¹ region [38-40]. In this research, C=O stretching vibration of 3C7HC molecule was obtained as 1705 cm⁻¹ at FT-IR and 1704 cm⁻¹ at FT-R. C-O stretching vibrations of pyrone rings of coumarins demonstrated in 1250-850 cm⁻¹ region [38-40]. C-O stretching vibrations were obtained as 1272,1150, 1123, 976, 953 cm⁻¹ at FT-IR and 1273, 1160, 1127, 980 cm⁻¹ at FT-R. C-O in plane bending vibrations was obtained as 652 cm⁻¹ at FT-IR and 650 cm⁻¹ at FT-R. C-OH stretching vibrations were obtained as 1460, 1272, 1203 cm⁻¹ at FT-IR and 1459, 1273, 1215 cm⁻¹ at FT-R. All C-OH vibrations are significantly consistent with the previous studies [16,38-40].

3.2.4. CN vibrations

There is a limited number of absorbing groups in the area where C≡N stretching is, so C≡N stretching restraint is quite easy. The bands showing the absorption vibration of cyano groups are quite dense within the range 2240-2220 cm⁻¹ and the stretching vibrations in this region show a characteristic feature of C≡N. The amplitude of C≡N stretching vibration is demonstrated as a significantly strong band 2223 cm⁻¹ in FT-IR and 2224 cm⁻¹ in FT-R for 4-methy-2-cyanobiphenyl [15,39] and 2229 cm⁻¹ at FT-IR and 2221 cm⁻¹ at FT-R for 3-cyano-4-methylcoumarin [15].

Table 3. Experimental and computed vibrational frequencies of Rot1 and Rot2 conformers of 3C7HC

Symmetry	Assignments	Exp. freq.(cm ⁻¹)		Calculated frequencies (cm ⁻¹)					
		IR	R	Rot1			Rot2		
				DFT	IR int. (km/mole)	R act. (Å ⁴ /amu)	DFT	IR int. (km/mole)	R act. (Å ⁴ /amu)
A ¹	v(OH)	3700	-	3676	121	122	3680	141	249
A ¹	v(CH)	3090	3089	3085	1	1	3093	2	104
A ¹	v(CH)	-	-	3063	3	2	3063	5	159
A ¹	v(CH)	-	-	3059	4	114	3057	2	51
A ¹	v(CH)	3043	-	3056	1	4	3045	7	110
A ¹	v(CN)	2235	2339	2245	45	1072	2245	44	1078
A ¹	v(C=O)	1705	1704	1752	569	95	1752	560	96
A ¹	v(C=C)+v(CC)	1611	1608	1591	405	152	1596	480	332
A ¹	v(C=C)+v(CC)+δ(CH)+δ(OH)	1566	1596	1580	281	530	1580	68	339
A ¹	v(C=C)-v(CC)+δ(CH)+δ(OH)	1505	557	1534	96	420	1532	261	533
A ¹	v(CC)+v(C-OH)+δ(CH)+δ(OH)	1460	1459	1478	37	21	1475	66	1
A ¹	v(C=C)+v(CC)+δ(CH)+δ(OH)	1385	-	1416	64	57	1421	21	58
A ¹	v(C=C)+v(CC)+δ(OH)+δ(CH)	1374	1374	1338	68	147	1337	10	97
A ¹	v(CC)+δ(CH)+δ(OH)	1318	1314	1332	5	102	1332	23	48
A ¹	v(CO)+v(C-OH)+δ(CH)+δ(OH)	1272	1273	1288	16	13	1282	158	18
A ¹	v(CC)+δ(CH)+δ(OH)	1230	1233	1237	58	63	1235	191	108
A ¹	v(CC)+v(C-OH)+δ(CH)	1203	1215	1225	157	58	1222	21	20
A ¹	v(CC)+v(CO)+δ(CH)+δ(OH)	-	1160	1164	69	20	1176	40	82
A ¹	δ(CH)+δ(OH)	1150	-	1143	307	15	1149	4	6
A ¹	v(CC)+v(CO)+δ(CH)+δ(OH)	1123	1127	1125	28	4	1113	103	2
A ¹	δ(CH)	1066	1068	1095	23	4	1105	144	9
A ¹	v(CO)+δ(CH)+δ(OH)	976	980	973	53	89	973	116	83
A ¹	v(CO)+δ(CH)	953	-	941	21	14	943	12	10
A ^u	γ(CH)	-	-	936	9	1	927	19	2
A ^u	γ(CH)	885	-	916	9	2	899	0	1
A ¹	δ(ring)	-	859	834	1	7	833	1	7
A ^u	γ(CH)	825	-	813	46	0	827	26	0
A ¹	δ(ring)+v(C-OH)	-	-	802	3	0	803	5	0
A ^u	γ(ring)+γ(CH)	765	-	793	10	0	768	29	0
A ^u	γ(ring)	740	741	729	17	3	727	17	3
A ¹	δ(ring)+δ(CN)	-	-	722	4	49	719	1	52
A ^u	γ(ring)	697	-	694	1	0	684	3	0
A ¹	δ(ring)+δ(CN)+δ(CO)	652	650	630	5	5	631	5	4
A ^u	γ(ring)	618	620	613	5	1	609	11	1
A ¹	δ(ring)	-	-	598	1	2	597	1	1
A ¹	δ(ring)+δ(CN)	554	553	548	22	1	549	17	2
A ^u	γ(ring)+γ(CN)	526	516	514	0	3	514	0	3
A ¹	δ(ring)+δ(C-OH)	482	479	445	16	10	446	1	10
A ^u	γ(ring)	460	446	440	10	0	443	2	0
A ¹	δ(ring)+δ(CN)	-	427	432	3	6	431	6	6
A ¹	δ(ring)+δ(C-OH)	-	-	408	7	7	408	3	8
A ^u	γ(CH)+γ(ring)	-	-	400	3	1	397	1	2
A ^u	γ(OH)	-	359	362	115	2	343	112	2
A ¹	δ(ring)	-	-	336	2	9	336	10	8
A ^u	γ(ring)	-	294	267	0	1	268	3	1
A ¹	ρ(ring)	-	-	248	4	2	249	0	2
A ^u	γ(ring)	-	-	238	0	1	237	1	1
A ^u	γ(ring)	-	163	166	7	0	165	5	0
A ¹	ρ(CN)	-	115	126	3	3	126	3	3
A ^u	w(ring)	-	-	87	2	0	87	0	0
A ^u	w(ring)	-	-	58	0	1	58	1	1

R ot1 R² = 0,9991

Rot2 R² = 0,9991

v:stretching,δ:bending,γ:out of plane bending, ρ: rocking,w:wagging.

In this research, the stretching vibration was observed as 2235 cm^{-1} at FT-IR and 2239 cm^{-1} at FT-R. $\text{C}\equiv\text{N}$ in plane bending vibrations are demonstrated in $617\text{-}437\text{ cm}^{-1}$ region [15]. In plane $\text{C}\equiv\text{N}$ bending vibrations were obtained as $652, 554\text{ cm}^{-1}$ at FT-IR and $650, 553, 427\text{ cm}^{-1}$ at FT-R, whereas out of plane $\text{C}\equiv\text{N}$ bending vibration was obtained as 526 cm^{-1} at FT-IR and 516 cm^{-1} at FT-R. $\text{C}\equiv\text{N}$ rocking vibration was obtained as 115 cm^{-1} at FT-R.

3.2.5. CC Ring vibrations

CC stretching vibrations are significantly dominant in spectra of coumarin derivative molecules and it has the characteristic of the aromatic ring. CC stretching vibrations of aromatic structures are demonstrated strongly within $1624\text{-}726\text{ cm}^{-1}$ range [16]. In this study, stretching vibrations for 3C7HC molecule were obtained as $1611, 1566, 1505, 1460, 1385, 1374, 1318, 1227, 1203, 1123\text{ cm}^{-1}$ at FT-IR and $1608, 1596, 1557, 1459\text{ cm}^{-1}, 1374, 1314, 1233, 1150, 1127\text{ cm}^{-1}$ at FT-R. In plane -out of plane bending ring vibrations of 3C7HC molecule are shown in Table 3. It has been verified that these values obtained are consistent with the previous studies [16, 40].

4. Discussion and Conclusion

In this research, FT-IR and FT-R spectra were obtained and vibrational assignments are shown in detail for the first time for 3C7HC. The differences in observed and scaled vibrational frequencies were found to be significantly low. FT-IR and FT-R spectra of 3C7HC were studied by using B3LYP 6-311++G(d,p) parameter of density functional theory. Vibrational frequencies of optimized molecular structures, geometry parameters, relative energy and dipole moment, corresponding vibrational assignments, FT-IR intensities and FT-R activities of the two different conformers of 3C7HC were obtained using DFT method. The calculated geometrical parameters and vibrational frequencies of these conformers were found to be consistent with the experimental data.

Acknowledgment

The present study was financially supported by Bartın University BAP with Project no: BAP-2012-1-44. The author acknowledges the help of Assoc. Prof. Dr. Halil OTURAK for the Gaussian 09W software.

References

- [1] Symeonidis, T., Chamilos, M., Hadjipavlou-Litina, D. J., Kallitsakis, M., Litinas, K. E. 2009. Synthesis of hydroxycoumarins and hydroxybenzo[f] or [h]coumarins as lipid peroxidation inhibitors. *Bioorg. Med. Chem. Lett.*, 19(4) (2009), 1139-1142.
- [2] Venugopala, K. N., Jayashree, B. S. 2008. Microwave-induced synthesis of Schiff bases of aminothiazolylbromocoumarins as antibacterials. *Indian J. Pharm. Sci.*, 70(1) (2008), 88-91.
- [3] Campos-Toimil, M., Orallo, F., Santana, L., Uriarte, E. 2002. Synthesis and vasorelaxant activity of new coumarin and furocoumarinderivatives. *Bioorg. Med. Chem. Lett.*, 12(5) (2002), 783-786.
- [4] Satyanarayana, S. V., Sreevani, P., Sivakumar, A., Vijayakumar, V. 2008. Synthesis and antimicrobial activity of new Schiff bases containing coumarin moiety and their spectral characterization. *Arkivoc*, 17(2008), 221-233.
- [5] Hwu, J. R., Singha, R., Hong, S. C., Chang, Y. H., Das, A. R., Vliegen, I., De Clercq, E., Neyts, J. 2008. Synthesis of new benzimidazole-coumarin conjugates as anti-hepatitis C virus agents. *Antiviral Res.*, 77(2) (2008), 157-162.
- [6] Daly, A. K. 2009. Pharmacogenomics of anticoagulants: steps toward personal dosage. *Genome Med.*, 1(1), 10-14.
- [7] Anderson, D. M., Shelley, S., Crick, N., Buraglio, M. 2002. No effect of the novel antidiabetic agent nateglinide on the pharmacokinetics and anticoagulant properties of warfarin in healthy volunteers. *J. Clin. Pharmacol.*, 42(12)(2002), 1358-1365.
- [8] Chattha, F. A., Munawar, M. A., Nisa, M., Ashraf, M., Kousar, S., Arshad, S. 2015. Potential antibacterial activity of coumarin and coumarin-3-acetic acid derivatives. *Pakistan Journal of Pharmaceutical Science*, 28(3) (2015), 819-823.
- [9] Timonen, J. M., Nieminen, R. M., Sareila, O., Goulas, A., Moilanen, L. J., Haukka, M., Vainiotalo, P., Moilanen, E., Aulaskari, P. H. 2011. Synthesis and anti-inflammatory effects of a series of novel 7-hydroxycoumarin derivatives. *European Journal of Medicinal Chemistry*, 46(2011), 3845-3850
- [10] Bhattacharyya, S. S., Paul, S., Mandal, S. K., Banerjee, A., Boujedaini, N., Khuda-Bukhsh, A. R. 2009. A synthetic coumarin (4-methyl-7-hydroxycoumarin) has anti-cancer potentials against DMBA-induced skin cancer in mice. *European Journal of Pharmacology*, 614(2009), 128-136.
- [11] KinzaAslam K., Khosa M. K., Jahan, N., Nosheen, S. 2010. Synthesis and applications of coumarin. *Pakistan Journal of Pharmaceutical Science*, 23(4) (2010), 449-454.
- [12] Maheswara, M., Siddaiah, V., Damu, G. L., Rao, Y. K., Rao, C. V. 2006. A solvent-free synthesis of coumarins via Pechmann condensation using heterogeneous catalyst. *Journal of Molecular Catalysis A Chemical*, 255(2006), 49-52.

- [13] Zabrandik, M.1992. The production and application of fluorescent brightening agents. John Wiley & Sons, New York.
- [14] Divyesh, P., Kumari, P., Patel, N. 2010. Synthesis characterization and biological evaluation of some thiazolidinone derivatives as antimicrobial agents. *Journal of Chemical and Pharmaceutical Research*, 2(5) (2010), 84-91.
- [15] Udaya Sri, N., Chaitanya, K., Prasad, M. V. S., Veeraiah, V., Veeraiah, A. 2012. FT-IR, FT-Raman and UV-Vis spectra and DFT calculations of 3-cyano-4-methylcoumarin. *Spectrochimica Acta Part A, Molecular and Biomolecular Spectroscopy*, 97(2012), 728-736.
- [16] Sebastian, S., Sylvestre, S., Jayarajan, D., Amalanathan, M., Oudayakumar, K., Gnanapoongothai, T., Jayavarthanam, T. 2013. Molecular structure, normal coordinate analysis, harmonic vibrational frequencies, natural bond orbital, TD-DFT calculations and biological activity analysis of antioxidant drug 7-hydroxycoumarin. *Spectrochimica Acta Part A, Molecular and Biomolecular Spectroscopy*, 101(2013), 370-381.
- [17] Sortur, V., Yenagi, J., Tonannavar, J., Jadhav, V. B., Kulkarni, M. V. 2006. Fourier transform-infrared and Raman spectra, ab initio calculations and assignments for 6-methyl-4-bromomethylcoumarin. *Spectrochimica Acta A, Molecular and Biomolecular Spectroscopy*, 64(2006), 301-307.
- [18] Dereli, Ö., 2016. Molecular Structure and Spectral (FT-IR, Raman) Investigations of 3-Aminocoumarin. *Optics and Spectroscopy Journal*, 120(2016), 690-700.
- [19] Arjunan, V., Sakiladevi, S., Marchewka, M, K., Mohan, S. 2013. FTIR, FT-Raman, FT-NMR and quantum chemical investigations of 3-acetylcoumarin. *Spectrochimica Acta A, Molecular and Biomolecular Spectroscopy*, 109(2013), 79-89.
- [20] Tonannavar, J., Yenagi, J., Sortur, V., Jadhav, V. B., Kulkarni, M. V. 2010. Vibrational spectra, normal modes, ab initio and DFT calculations for 6-Chloro- and 7-Chloro-4-bromomethylcoumarins. *Spectrochimica Acta A*, 77(2010), 351-358.
- [21] Warde, U., Sekar N. 2017. Solvatochromic benzo[h] coumarins: Synthesis, solvatochromism. *Optical Materials Journal*, 72 (2017), 346-358.
- [22] Sarıkaya, E. K., Dereli, Ö., Erdoğdu, Y., Güllüoğlu, M.T. 2012. Molecular structure and vibrational spectra of 7-Etoxycoumarin by density functional method. *Journal of Molecular Structure*, 1049(2012) 220-226.
- [23] Joseph, L., Sajan, D., Reshmy, R., ArunSasi, B. S., Erdoğdu, Y., Kurien Thomas, K. 2012. Vibrational spectra, structural conformations, scaled quantum chemical calculations and NBO analysis of 3-acetyl-7-methoxycoumarin. *Spectrochimica Acta Part A, Molecular and Biomolecular Spectroscopy*, 99 (2012) 234-247.
- [24] Moghanian, H., Mobinikhaledi, A., Monjezi, R. 2013. Synthesis, spectroscopy (vibrational, NMR and UV-vis) studies, HOMO-LUMO and NBO analysis of 8-formyl-7-hydroxy-4-methylcoumarin by ab initio calculations. *Journal of Molecular Structure* 1052(2013), 135-145.
- [25] Frisch, M. J., Trucks, G. W., Schlegel, H.B., Scuseria, G. E., Robb, M.A., Cheeseman, J. R., Montgomery Jr, J. A., Vreven, T., Kudin, K. N., Burant, J. C., Millam, J. M., Iyengar, S. S., Tomasi, J., Barone, V., Mennucci, B., Cossi, M., Scalmani, G., Rega, N., Petersson, G. A., Nakatsuji, H., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Klene, M., Li, X., Knox, J. E., Hratchian, H. P., Cross, J. B., C., Adamo, Jaramillo, J., Gomperts, R., Stratmann, R. E., Yazyev, O., Austin, A. J., Cammi, R., Pomelli, C., Ochterski, J. W., Ayala, P. Y., Morokuma K., Voth, G. A., Salvador, P., Dannenberg, J. J., Zakrzewski, V. G., Dapprich, S., Daniels, A. D., Strain, M. C., Farkas, O., Malick, D. K., Foresman, J.B., Ortiz, J.V., Cui, Q., Baboul, A.G., Foresman, J. B., Ortiz, J.V., Cui, Q., Baboul, A.G., Clifford, S., Rabuck, A.D., Raghavachari, K., Cioslowski, J., Stefanov, B. B., Liu, G., Liashenko, A., Piskorz, P., Komaromi, I., Martin, R. L., Fox, D. J., Keith, T., Al-Laham, M. A., Peng, C. Y., Nanayakkara, A., M. Challacombe, Gill, P. M. W., Johnson, B., Chen, W. Wong, M. W., Gonzalez, C., Pople, J. A. 2003. GAUSSIAN 03, Revision C.02, Gaussian Inc., Pittsburgh, PA.
- [26] Frish, A., Nielsen, A. B., Holder, A. J. 2001. Gauss View User Manual, Gaussian Inc. Pittsburg, PA.
- [27] Young, D. C. 2001. Computational Chemistry. A Practical Guide for Applying Techniques to Real-World Problems, (Electronics). John Wiley & Sons, New York.
- [28] Ueno, K. 1985. Structures of umbelliferone and 7-ethoxycoumarin. *Acta Crystallographica*, C41 (1985), 1786-1789.
- [29] Vijayalakshmi, L., Parthasarathi, V., Varu, B., Shah, A. 2001. 3-Cyano-4-[2-(4-methyl-thiophenyl)-ethenyl]-2H-1-benzo-pyran-2-one. *Acta Crystallographica*, E57 (2001), 0436-438.
- [30] Wilson, E.B., Decius, J.C., Cross, P.C. 1980. *Molecular vibrations*, Dover Pub., New York.
- [31] Varsanyi, G., Szoke S. 1969. *Vibrational spectra of benzene derivatives*. Academic press, New York.
- [32] Du, X., Guo C., Hansell, E., Doyle, P. S., Caffrey, C. R., Holler, T. P., McKerrow, J. H., Cohen, F. E. 2002. Synthesis and structure-activity

- relationship study of potent trypanocidalthiosemicarbazone inhibitors of the trypanosomal cysteine protease cruzain. *J. Med. Chem.*, 45(13) (2002), 2695-2707.
- [33] Smith, B. C. 1998. *Infrared spectral interpretation: A Systematic Approach*. CRC Press, Boca Raton, Florida.
- [34] Socrates, G. 2004. *Infrared and Raman characteristic group frequencies: tables and charts*, Wiley.
- [35] Vein, D. L., Colthup, N.B., Fateley, W.G., Grasselli, J. 1991. *The Handbook of IR and Raman characteristic frequencies of organic molecules*. Academic Press, New York.
- [36] Colthup, N. B., Daly, L. H., Wiberly, S. E. 1990. *Introduction to Infrared and Raman spectroscopy*, Academic Press, New York.
- [37] Uçun, F., Güçlü, V., Sağlam, A. 2008. Ab initio Hartree-Fock and density functional theory study on molecular structures, energies, and vibrational frequencies of conformations of 2-hydroxy-3-nitropyridine and 3-hydroxy-2-nitropyridine. *Spectrochimica Acta Part A, Molecular and Biomolecular Spectroscopy*, 70(2008), 524-531.
- [38] Sajan, D., Erdoğan, Y., Reshmy, R., Dereli, Ö., Kurien Thomas, K. K., Hubert Joe, I. 2011. DFT-based molecular modeling, NBO analysis and vibrational spectroscopic study of 3-(bromoacetyl)coumarin. *Spectrochimica Acta Part A, Molecular and Biomolecular Spectroscopy*, 82(2011), 118-125.
- [39] Sebastian, S., Sundaraganesan, N., Karthikeyan, B., Srinivasan, V. 2011. Quantum mechanical study of the structure and spectroscopic (FT-IR, FT-Raman, ¹³C, ¹H and UV), first order hyperpolarizabilities, NBO and TD-DFT analysis of the 4-methyl-2-cyanobiphenyl *Spectrochimica Acta A, Molecular and Biomolecular Spectroscopy*, 78(2011), 590-600.
- [40] Erdoğan, Y. 2013. Investigations of FT-IR, FT-Raman, FT-NMR spectra and quantum chemical computations of Esculetin molecule. *Spectrochimica Acta A, Molecular and Biomolecular Spectroscopy*, 106(2013), 25-33.