



Biological Activities of Extracts and Isolated Calotropin from *Gomphocarpus Purpurascens* Leaves: A Computational and Experimental Study

Tsegu Kiros^{1*}, Seid Mohammed², Aman Dekebo^{3,4*} and Yadessa Melaku^{3,4*}

¹Haramaya University, Chemistry Department, Dire Dawa, P.O.Box. 138, Ethiopia

²Adama Science and Technology University, Department of Applied Biology, Adama,
P.O.Box. 1888, Ethiopia

³Adama Science and Technology University, Department of Applied Chemistry, Adama,
P.O.Box. 1888, Ethiopia

⁴ Adama Science and Technology University, Institute of Pharmaceutical Sciences, Ethiopia

*Correspondence: kirosorg@gmail.com; amandekeb@gmail.com; yadessamelaku2010@gmail.com

SUPPLEMENTARY MATERIAL

91-2110 (GPL-01).1.fid
1H




Figure S1: ¹H NMR (400 MHz, CD₃OD, δ in ppm) spectrum of calotropin

91-2110 (GPL-01).2.fid
C13
— 208.72
— 175.88




Figure S2: ^{13}C NMR (100 MHz, CD_3OD) spectrum of calotropin

91-2110 (GPL-01).3.fid
Dept-135




Figure S3: DEPT-135 NMR (100 MHz, CD₃OD) spectrum of calotropin