








Research Article

## HIGH PREVALENCE OF TRANSFERABLE INTEGRON-ASSOCIATED DRUG RESISTANCE IN *Escherichia coli* STRAINS ISOLATED FROM BLOOD CULTURES IN A UNIVERSITY HOSPITAL IN TRABZON, TURKEY

 Ahu REİS <sup>1</sup>,  Osman Birol ÖZGÜMÜŞ <sup>2</sup>,  Erva RAKICI <sup>2</sup>,  İnci DURUKAN <sup>1</sup>,  
 Celal Kurtuluş BURUK <sup>1</sup>,  Gülçin BAYRAMOĞLU <sup>1</sup>,  Ali Osman KILIÇ <sup>1</sup>

<sup>1</sup> Department of Medical Microbiology, Faculty of Medicine, Karadeniz Technical University, Trabzon, TURKIYE

<sup>2</sup> Department of Medical Microbiology, Faculty of Medicine, Recep Tayyip Erdogan University, Rize, TURKIYE

\*Correspondence: [akilic@ktu.edu.tr](mailto:akilic@ktu.edu.tr)

### ABSTRACT

**Objective:** This study aimed to determine the carriage of transferable integron-associated drug resistance in *Escherichia coli* (*E. coli*) strains isolated from blood cultures.

**Materials and Methods:** A total of 111 *E. coli* isolates were included in this study. Antimicrobial susceptibility testing of the isolates against 17 antibiotics was performed using an automated microbiology system. Integron-specific polymerase chain reactions (PCR) were used to detect the presence of integrons. The antibiotic resistance gene cassettes in the variable regions of integrons were analyzed by DNA sequencing. Plasmid transfer assays were performed using the broth mating method. The clonal relationships among integron-carrying strains were evaluated by pulsed-field gel electrophoresis (PFGE).

**Results:** Resistance rates to antibiotics ranged from 0.9% to 63%. Thirty-eight strains carried gene cassettes encoding *dfrA7*, *dfr17-aadA5*, *dfrV*, *dfrA1-aadA1*, and *dfrA12-aadA2*. Seven strains possessed class 2 integrons with gene arrays *dfrA1-sat2-aadA1* and *dfrA1-sat2-aadA30*. Twenty-two integron-carrying isolates harbored conjugative resistance plasmids, three of which were identified as belonging to the IncN group. Two strains with class 1 integrons, isolated from different clinics, exhibited similar patterns in the PFGE analysis.

**Conclusion:** Approximately 50% of *E. coli* isolates from blood cultures at our hospital were found to carry integron-associated transferable drug resistance, suggesting their potential role in the horizontal dissemination of resistance genes. Further research is needed to understand the prevalence of *E. coli* strains of blood origin and the role of integrons and gene cassette arrays in the spread of resistance.

**Keywords:** *Escherichia coli*, Gene Cassette, Integron, Conjugation, Plasmid

Received: 01 October 2024  
Revised: 02 December 2024  
Accepted: 04 December 2024  
Published: 22 December 2024



Copyright: © 2024 by the authors. Published by Aydın Adnan Menderes University, Faculty of Medicine and Faculty of Dentistry. This is an open access article under the Creative Commons Attribution Non-Commercial 4.0 International (CC BY-NC 4.0) License.

---

## INTRODUCTION

Bloodstream infections (BSIs) are among the most significant nosocomial infections, contributing to increased mortality and morbidity due to rising antimicrobial resistance (1). *Escherichia coli*, an opportunistic pathogen, is capable of causing a wide range of infections, from those in the gastrointestinal tract to extraintestinal infections, such as those affecting the central nervous system, urinary tract, and bloodstream. Globally, it is a leading cause of bloodstream infections and exhibits resistance to many drugs (2). While most bacterial infections are treatable with antibiotics, *E. coli* and other pathogens can rapidly develop antibiotic resistance, making infections more difficult to manage and potentially life-threatening (3). The excessive and inappropriate use of antimicrobials accelerates the emergence and spread of resistance. Therefore, the timely and effective use of antimicrobial drugs is essential in preventing resistance (4). Bacteria can develop resistance to antibiotics through genetic mechanisms involving mobile genetic elements (MGEs), such as plasmids, transposons (Tn), insertion sequences (IS), and integrons. Integrons, especially those located on plasmids or transposons, play a critical role in disseminating antimicrobial resistance genes, particularly in gram-negative bacteria. These integrons carry drug-resistant genes and facilitate the exchange of genetic elements. They are classified based on the sequence of *intI* gene, with classes 1 and 2 being the most prevalent (5).

The class 1 integrons consist of two conserved regions namely 5' (5'CS) and 3' (3'CS), and a variable region (VR) containing gene cassettes. The 5'CS includes the integrase gene (*intI*), which encodes the tyrosine recombinase enzyme, the recombination site (*attI*), and a promoter (P) region involved in the expression of inserted genes or gene cassettes. The 3'CS contains the defective quaternary ammonium resistance gene (*qacEΔ1*) and the *sulI* gene, which mediates resistance to sulfonamides. The variable region, located between these two conserved regions, contains the *attC* site, where various antibiotic resistance cassettes are integrated via recombination. Unlike class 1 integrons, class 2 integrons lack *sul* genes and typically include resistance gene cassettes such as *dfra1*, *sat2*, and *aadA1*, which confer resistance to aminoglycosides and sulfamethoxazole (6).

Although several studies in Turkey have investigated integron-associated drug resistance in *E. coli* isolates from urine samples (7, 11), very few reports are available (12) on the *E. coli* isolates from blood samples.

This study aimed to identify transferable integron-associated drug resistance in *E. coli* strains isolated from patients in various clinics.

## MATERIALS AND METHODS

### *Sampling, isolation and identification of bacterial strains*

A total of 111 *E. coli* strains were isolated from hospitalized patients at Karadeniz Technical University's Medical Microbiology Laboratory between January and December 2014. The bacterial strains were cultured using 5% sheep blood agar and EMB agar. Conventional biochemical methods and MALDI-TOF MS (Microflex™, Bruker, Germany) were employed to confirm species-level identification.

### *Antimicrobial susceptibility testing*

Antimicrobial susceptibility testing was performed using the Phoenix ID/AST automated microbiology system (Becton Dickinson Diagnostic Instrument Systems, Sparks, MD, USA), following the manufacturer's instructions. Antibiotic susceptibility test results were interpreted according to Clinical and Laboratory Standards Institute (CLSI) criteria. The isolates were tested for the following antibiotics: amikacin (AK), ampicillin/sulbactam (SAM), aztreonam (ATM), cefazolin (CZ), cefepime (FEP), cefoperazone/sulbactam (CES), cefoksitin (FOX), ceftazidime (CAZ), ceftriaxone (CRO), ciprofloxacin (CIP), ertapenem (ERT), gentamicin (CN), imipenem (IPM), levofloxacin (LEV), meropenem (MEM), piperacillin tazobactam (TZP) and trimethoprim/sulfamethoxazole (SXT). The susceptibility of isolates containing class 1 and class 2 integrons was determined using the disc diffusion method in accordance with CLSI criteria (13). Antibiotic discs (Oxoid, UK) containing ampicillin (10 µg), streptomycin (10 µg), tetracycline (30 µg), nalidixic acid (30 µg), and chloramphenicol (30 µg) were used for this purpose.

### ***DNA isolation***

DNA templates for polymerase chain reaction (PCR) were prepared using a modified boiling method based on the protocol by Perez et al (14). After washing, the supernatant was used as the DNA template for PCR analyses.

### ***Detection of integron gene cassettes and integrase genes***

To detect integron gene cassettes in *E. coli* isolates, specific primers were used: For class 1 integrons: *intI*-1F (GGTCAAGGATCTGGATTTGG) and *intI*-1R (ACATGCGTGTAATCATCGTC). For class 2 integrons: *intI*-2F (CACGGATATGCGACAAAAGGT) and *intI*-2R (GTAGCAAACGAGTGACGAAATG). To amplify the variable regions of the integrons, the following primers were used: For class 1 integrons: 5'-CS (GGCATCCAAGCAGCAAG) and 3'-CS (AAGCAGACTTGACCTGA). For class 2 integrons: hep51 (GATGCCATCGCAAGTACGAG) and hep74 (CGGGATCCCGGACGGATGCACGATTTGTA). PCR reaction conditions and compositions followed previously described methods (15).

### ***DNA sequencing and data analysis***

PCR products from class 1 and class 2 integrons were purified using the High Pure PCR Clean-up Micro Kit (Roche, Germany). Cloning was performed with the pGEM-T Easy Vector System (Promega, USA) and T4 DNA Ligase enzyme, following the manufacturer's guidelines as described by Sambrook et al. (16). After incubation, blue-white colony screening was used for selection. Plasmid DNA from selected white colonies was isolated using the alkaline lysis method (17). Recombinant plasmids carrying amplicons of class 1 and class 2 integrons were sequenced on an ABI 3130 Genetic Analyzer (Applied Biosystems, USA) using SP6 and T7 promoter primers. DNA sequence data were analyzed using BLAST (<http://www.ncbi.nlm.nih.gov/BLAST>) and CLUSTALW2 (<http://www.ebi.ac.uk/Tools/msa/clustalw2>) for sequence alignment and comparisons (18).

### ***Conjugation experiment***

Transfer experiments of plasmids from integron-containing isolates were performed based on the broth mating principle, which was previously described (19). *Escherichia coli* isolates containing integrons were used as donors, and *E. coli* J53-2 (met pro Rif<sup>R</sup>) as the recipient cells. Broth cultures of both donor and recipient were mixed in equal volumes (1:1) and incubated for 20 hours at 37°C without shaking. Transconjugants were selected on LB agar supplemented with rifampicin (300 µg/mL), to which the recipient cells are resistant, and either trimethoprim (25 µg/mL) or streptomycin (100 µg/mL) to which the donor cells are resistant.

### ***Isolation of plasmid DNA***

A colony was picked from transconjugant plates and sub-cultured in 3 mL of LB broth at 150 rpm shaking at 37°C for 16-18 hours. A 1.5-ml aliquot of the overnight culture was centrifuged at 12,000 rpm to pellet the bacteria. The cells were lysed, and plasmid DNA was isolated using the standard alkali-lysis method. Finally, the pellet was dried at 37°C and resuspended in 30-40 µl of 1X TE buffer containing 0.5-1 µg/mL of RNase. Plasmid DNA was analyzed by electrophoresis, and visualized under a UV transilluminator (20).

### ***PCR-based replicon typing***

To identify the incompatibility (Inc) groups that contribute significantly to the dissemination of resistance genes, plasmids isolated from transconjugants were subjected to replicon typing using specific primers as described by Götz et al. (21).

### ***Pulsed-field gel electrophoresis (PFGE)***

Clonal relationship among 58 *E. coli* isolates containing integrase genes (*intI-1* and *intI-2*) belonging to class 1 and class 2 integrons were determined using the PFGE method, with modifications based on the studies of Durmaz et al. (22). PFGE was performed with the CHEF DR-III (Bio-Rad, USA) system using a 1% ultrapure agarose gel in 2L of 0.5X TBE. The gel was stained in a solution containing 10 mg/mL ethidium bromide for 30

minutes. Gel visualization and photography were conducted using the VersaDoc™ system, and images were analyzed with GelJ software, version 2.


### Statistical analysis

Statistical analyses were performed using Pearson's Chi-square test and Fisher's Exact test with the Statistical Package for the Social Sciences (SPSS) version 23 for Windows. A p-value of < 0.05 was considered statistically significant.

## RESULTS

### Antimicrobial resistance phenotypes of *E. coli* isolates


Antimicrobial susceptibility tests for 111 *E. coli* isolates from blood cultures were performed using the Phoenix™ 100 device. The resistance rates of the isolates to 17 antibiotics are shown in Figure 1.



**Figure 1.** Total antimicrobial resistance rates of 111 clinical *Escherichia coli* isolates.

The highest resistance percentage was observed for SAM 63% (n=70), while the lowest was for ERT 0.9% (n=1). Resistance rates to other antibiotics were as follows: ATM 44.1% (n=49), CZ 58.5% (n=65), FEP 40.5% (n=45), CES 22.5% (n=25), FOX 18.1% (n=20), CAZ 36% (n=40), CRO 48.6% (n=54), CIP 36% (n=40), CN 25.2% (n=28), LEV 35.1% (n=39), TZP 18.9% (n=21), and SXT 54% (n=60). None of the isolates was resistant to AK, IPM, and MEM (Fig. 1). In a subset of 45 *E. coli* isolates with class 1 and class 2 integron gene cassettes, tested using the

disc diffusion method, resistance to ampicillin, streptomycin, and tetracycline was found to be 66.6% (n=30), while resistance to nalidixic acid was 64.4% (n=29). The lowest resistance rate was 26.6% (n=12) for chloramphenicol (Figure 2).



**Figure 2.** Total antimicrobial resistance rates of *E. coli* isolates containing class 1 and class 2 integrons.

Statistical analysis revealed significant differences in antimicrobial resistance between strains with and without class 1 and class 2 integrase genes for SAM, ATM, CZ, FEP, CAZ, CRO, CIP, and SXT ( $p < 0.05$ ). However, no significant differences were observed for CES, FOX, ERT, CN, LEV, and TZP antibiotics ( $p > 0.05$ ). Statistical analysis could not be conducted for AK, IPM, and MEM, as no resistant strains were found in either group.

### PCR for integrons

A total of 111 *E. coli* isolates were screened for the presence of class 1 and class 2 integrase genes. Of these, 51 isolates (46%) were positive for class 1 integrase, and seven isolates (6%) were positive for class 2 integrase. PCR analysis of the variable region of class 1 integrons showed that 38 isolates (34%) contained class 1 integrons, while 13 isolates (11.7%) carried empty integrons. Class 2 integrons were detected in 7 isolates (6%) (Supplementary Table 1). Gene cassette arrays of 45 *E. coli* isolates with integrons were determined by DNA sequencing. Among 27 of the 38 isolates with class 1 integrons, five gene cassette arrays were identified: *dfrA7*, *dfr17-aadA5*, *dfrV*, *dfrA1-aadA1* ve *dfrA12-aadA2*, associated with trimethoprim and streptomycin-

spectinomycin resistances. These arrays ranged from 800-1800 bp in length. Gene cassette arrays could not be sequenced for 11 class 1 integrons due to technical difficulties of the unknown nature (TEC002, TEC005, TEC010, TEC015, TEC020, TEC043, TEC055, TEC057, TEC072, TEC078 and TEC091). For the seven isolates with class 2 integrons, two gene cassette arrays were identified: *dfrA1-sat2-aadA1* and *dfrA1-sat2-aadA30*, approximately 2500 bp in length. Sequencing was unsuccessful for three class 2 integrons (TEC012, TEC033, TEC081) (Supplementary Table 1).


### ***Conjugation experiments***

Plasmid transfer studies were executed on 45 *E. coli* isolates with integrons. Among these, 22 isolates (48.8%) carried conjugative plasmids larger than 100 kb. Conjugation frequency ranged from  $10^{-9}$  to  $10^{-8}$ . Transconjugants were validated genotypically using previously established PCR methods. PCR assays revealed that three plasmids (pTEC010, pTEC015 and pTEC075) belonged to the IncN group. No Inc P, IncQ, or IncW group plasmids were detected in any of the transconjugants (Supplementary Table 1).

### ***Pulsed-field gel electrophoresis analysis***

PFGE dendrogram analysis of 50 *E. coli* strains with integrons, their antibiotic resistance patterns, and sources shown in Figure 3. These strains were divided into 39 distinct groups, some of which were further subdivided into subgroups (A, B, C). Eleven subgroups (4, 7, 9, 10, 12, 13, 14, 16, 25, 35, 37) showed similarity rates above 80%. Subgroups 13 showed similarity rates exceeding 90%. Notably, strains TEC010 and TEC015 all carrying class 1 integrons, demonstrated high clonal similarity despite significant differences in their clinical origins and antibiotic resistance patterns. Eight strains were excluded from the analysis because the DNA pattern quality obtained as a result of PFGE experiments was not sufficient for dendrogram analysis.





**Figure 3.** Dendrogram analysis of *E. coli* isolates containing class 1 and class 2 integrons

## DISCUSSION

Infections caused by Gram-negative bacteria, such as septicemia, are significant contributors to mortality and morbidity. The isolation of bacteria, particularly from blood cultures, often indicates a severe invasive infection requiring immediate antimicrobial therapy. Among these, *E. coli* is one of the most commonly isolated pathogens from blood cultures. The World Health Organization (WHO) has identified *E. coli* as a priority pathogen of concern due to its role in the global development of antimicrobial resistance, especially infections affecting the blood and urinary tract. Antibiotic resistance presents a considerable challenge to public health. Accurate detection of resistant bacteria and determination of antimicrobial resistance, particularly in clinical laboratories, are therefore essential for enabling clinicians to administer effective antibiotic treatments (23, 24).

In our study, the antibiotic resistance patterns of *E. coli* from blood cultures were analyzed against 17 different antibiotics. Resistance rates were highest against SAM, CZ, CRO, and SXT, as also reported in the previous studies (25, 26). As a matter of fact, these antibiotics have historically been used extensively in the empirical treatment of sepsis and other infections (23). Our findings align with those of Kreidl et al. (27), who reported resistance rates of 71.8% against aminopenicillins and 37.5% against CIP. Furthermore, all *E. coli* isolates in our study were sensitive to AK and carbapenems (IPM and MEM), consistent with findings by Anvarinejad et al. (28).

Timely and accurate diagnosis and treatment are essential for mitigating the severe effects of sepsis and reducing mortality. Identifying common bacterial pathogens and their resistance patterns is crucial for guiding appropriate antimicrobial therapy. Ampicillin, nalidixic acid, SXT, and tetracyclines, widely used in both clinical and veterinary medicine (28), showed significant resistance in this study. Notably, strains with class 1 and class 2 integrons exhibited the highest resistance rates. These classes of integrons contain genes conferring resistance to trimethoprim and streptomycin. Class 1 integrons, in particular, harbor the *sul1* gene encoding sulfonamide resistance adjacent to the 3'-conserved region (29). Integrons are genetic elements that facilitate the capture, insertion, and exchange of antibiotic resistance gene cassettes, thereby playing a significant role in the dissemination of resistance genes, particularly among Gram-negative bacteria (30). The high prevalence of trimethoprim and streptomycin resistance in this study are thought to be resulted in the integrons containing various antibiotic gene cassette arrays.

Multidrug resistance in Enterobacterales is often associated with antimicrobial resistance genes carried by integrons (31). In Türkiye, numerous studies have focused on integrons in bacteria from clinical (32), animal (33), and aquatic (34) sources. Although there are some epidemiological studies in clinical bacteria, especially in urine samples (9), research on the molecular characterization of *E. coli* strains from blood samples remains limited. In this study, 46% (111/51) of isolates carried the *intI1* gene, with 38 isolates (34%) containing class 1 integrons, and 13 isolates (11.7%) having empty integrons. Additionally, 6% (111/7) of isolates carried the *intI2* gene, and all of these belong to class 2 integrons.

The frequency of the *intI1* gene (34%) observed in this study is comparable to findings by Liu et al. (35) (41.33%) but lower than results reported by Abdel-Rhman et al. (36) (98.33%) and Liu et al. (35) (67.39 %). The lower prevalence of class 1 integrons may be attributed to alternative resistance mechanisms, such as of beta-lactamase enzyme production, efflux pumps overexpression, reduced bacterial outer membrane permeability, and intrinsic resistance factors (37). Class 2 integrons were identified at a rate of 6%, consistent with Ahumado-Santos et al. (38), who reported a low prevalence (1.4%).

The predominance of class 1 integrons over class 2 integrons in our isolates is consistent with another study (11). Empty integrons found in 13 isolates may represent bacteria containing empty integrons are probably due to the presence of a very large gene cassette and the inability to amplify these gene cassettes (39, 40). We believe that the presence of empty integrons in bacteria could make these isolates have the potential to become multi-drug resistant by capturing antibiotic resistance genes from other bacteria in the hospital environment and spreading to the other bacteria of both commensal and clinical origin.

The presence of *dfrA* genes encoding trimethoprim resistance and *aad/aac* genes encoding aminoglycoside resistance has been widely reported (41, 42). Gene cassette arrays such as *dfrA1*, *dfrA5*, *aadA1*, and *aadA5* are common in clinical bacteria, as reported by Cicek et al. (8) and Sandalli et al. (7). The *dfrA17/aadA5* gene array, which confers resistance to trimethoprim and streptomycin, is often plasmid-mediated and associated with the proliferation of resistant bacterial isolates in clinical settings (43). In our study, 22 isolates (48.8%) containing class 1 and class 2 integrons harbored conjugative plasmids (>100 kb) confirmed through mating assays. These plasmids, particularly of the IncP and IncN groups, are known for their broad host range and role in spreading multidrug resistance in clinical settings (44).

The clonal relationships of integron-containing *E. coli* isolates were evaluated using PFGE. While PFGE effectively differentiated subgroups, it showed limited correlation with antibiotic resistance profiles, suggesting the influence of other genetic factors. Notably, isolates from different clinical origins and resistance profiles exhibited over 90% similarity in PFGE analyses, while clonally unrelated isolates from the same source displayed similar resistance profiles (Fig. 3). Also, TEC078 and TEC087 isolates, which were isolated from the same clinical source and had almost the same antibiotic resistance profiles, were determined as clonally

unrelated in different branches. As a result, PFGE patterns determined in *E. coli* isolates containing integrons and similar gene cassettes showed little clonal association.

## CONCLUSION

This study is among the few to investigate the prevalence of integrons in *E. coli* isolates from blood samples in Türkiye. The transfer of antibiotic resistance genes via integrons and plasmids poses significant challenges for managing infectious diseases. Continuous surveillance of antimicrobial resistance in invasive *E. coli* isolates is essential to monitor trends and guide effective treatment strategies. Understanding the mechanisms of resistance, including the role of integrons, can inform the development of targeted antimicrobial therapies. Further research is needed to explore the interplay between integrons, gene cassette arrays, and resistance mechanisms in bloodstream infections.

## Acknowledgments

We sincerely thank the Scientific Research Project Unit of Karadeniz Technical University for their support of this study (Grant number: 5335/2015).

## Authorship contributions

AR, OBÖ and AOK designed the study; AR, CKB and GB collected *E. coli* isolates and carried out phenotypic identification; AR, ER and İD performed the molecular assay; AR, OBÖ, ER and AOK prepared and revised the manuscript. All authors gave the final approval of the version to be published.

## Data availability statement

The authors state that the data supporting the study's results can be found in the article. Additionally, the raw data can be obtained from the corresponding author upon a reasonable request.

## Declaration of competing interest

The authors declare that they have no conflict of interest.

## Ethics

This study was approved by Karadeniz Technical University Faculty of Medicine Scientific Research Ethics Committee (Date: 12.10.2015, Decision no:3, Reference number 2015/117).

## Funding

The present study was supported by the Scientific Research Project Unit of Karadeniz Technical University (Grant number: 5335/2015).

## REFERENCES

1. Delle RD, Pezzotti P, Fontana C, Altieri A, Minelli S, Mariotti B, Cerretti R, Leoni D, Andreoni M, Sarmati L. An in-depth analysis of nosocomial bloodstream infections due to Gram-negative bacilli: clinical features, microbiological characteristics and predictors of mortality in a 1 year, prospective study in a large tertiary care Italian hospital. *Infectious Diseases* 2019; 51(1): 12-22.
2. Hastak P, Cummins ML, Gottlieb T, Cheong E, Merlino J, Myers GS, Chowdhury PR. Genomic profiling of *Escherichia coli* isolates from bacteraemia patients: a 3-year cohort study of isolates collected at a Sydney teaching hospital. *Microbial genomics* 2020; 6(5).
3. Lipworth S, Vihta KD, Chau K, Barker L, George S, Kavanagh J, Davies T, Vaughan A, Andersson M, Jeffery K, Oakley S, Morgan M, Hopkins S, Peto T, Crook DW, Walker AS, Stoesser N. Ten-year longitudinal molecular epidemiology study of *Escherichia coli* and *Klebsiella* species bloodstream infections in Oxfordshire, UK. *Genome Medicine*; 13(1): 1-13.
4. Aricò MO, Valletta E, Caselli D. Appropriate Use of Antibiotic and Principles of Antimicrobial Stewardship in Children. *Children* 2023; 10(4): 740.
5. Boucher Y, Labbate M, Koenig JE, Stokes HW. Integrons: mobilizable platforms that promote genetic diversity in bacteria. *Trends Microbiol.* 2007; 15:301-9.
6. Barlow RS, Gobius KS. Diverse class 2 integrons in bacteria from beef cattle sources. *Journal of Antimicrobial Chemotherapy* 2006; 58: 1133-1138.
7. Sandalli C, Buruk CK, Sancaktar M, Ozgumus OB. Prevalence of integrons and a new *dfrA17* variant in Gram-negative bacilli which cause community-acquired infections. *Microbiol Immunol* 2010; 54: 164-169. <https://doi.org/10.1111/j.1348-0421.2010.00197.x>
8. Cicek CA, Ozgumus OB, Saral A, Sandalli C. Antimicrobial resistance patterns and integron carriage of *Escherichia coli* isolates causing community-acquired infections in Turkey. *Annals of laboratory medicine* 2014; 34(2): 139-44.
9. Cicek CA, Sandalli C, Budak EE, Yağmur G, Çizmeçi Z, Ak S, Balcı PÖ, Şay Coşkun SU, Ay Altıntop Y, Fırat M, Sarı F, Çalışkan A, Yıldız N Sancaktar M, Özgümüş OB. İdrar Kültürlerinden İzole Edilen *Escherichia coli* Suşlarında Sınıf 1 ve Sınıf 2 İntegron Gen Kasetlerinin Karakterizasyonu: Çok Merkezli Bir Çalışma. *Mikrobiyol Bul* 2016; 50(2): 175-185.
10. Düzgün AÖ, Okumuş F, Saral A, Çiçek AÇ, Cinemre S. Determination of antibiotic resistance genes and virulence factors in *Escherichia coli* isolated from Turkish patients with urinary tract infection. *Revista da Sociedade Brasileira de Medicina Tropical* 2019; 52, e20180499.
11. Cicek CA, Şemen V, Ejder NA, Gündoğdu DZU, Kalcan SK, Köse FT, Özgümüş OB. Molecular epidemiological analysis of integron gene cassettes and tetA/tetB/tetD gene associations in *Escherichia coli* strains producing extended-spectrum  $\beta$ -lactamase (ESBL) in urine cultures. *Advances in Clinical and Experimental Medicine* 2022; 31(1): 71-79.
12. Öner SZ, Karaday E, Çalışkan A, Demir M, Şenol H, Kaleli İ. Integron distribution and relationship to antimicrobial resistance in *E. coli* isolated from blood culture. *Indian Journal of Medical Microbiology* 2024; 48, 100554.
13. Wayne, PA. National Committee for Clinical Laboratory Standards. Performance standards for antimicrobial susceptibility testing. Tenth Informational supplement, M100-S10 (M2): National Committee for Clinical Laboratory Standards, 2000.
14. Perez-Perez FJ, Hanson ND. Detection of plasmid-mediated AmpC beta-lactamase genes in clinical isolates by using multiplex PCR. *J Clin Microbiol* 2002; 40(6): 2153-62.
15. Machado E, Canto'n R, Baquero F, Gala'n JC, Rolla'n A, Peixe L, Coque TM. Integron Content of Extended-Spectrum-Lactamase-Producing *Escherichia coli* Strains over 12 Years in a Single Hospital in Madrid, Spain. *Antimicrobial Agents and Chemotherapy* 2005; 49(5): 1823-1829.
16. Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual* (No. Ed. 2). Cold spring harbor laboratory press
17. Kado CI, Liu ST. Rapid procedure for detection and isolation of large and small plasmids. *J Bacteriol* 1981; 145(3): 1365-73.

18. Rohlf FJ (1993). NSYS-PC: Numerical Taxonomy and Multivariate Analysis System. New York: Exeter Software.
19. Mokracka J, Koczura R, Pawłowski K, Kaznowski A. Resistance patterns and integron cassette arrays of Enterobacter cloacae complex strains of human origin. Journal of Medical Microbiology 2011; 60: 737-743.
20. Majumdar S, Chatterjee A. Ammonium acetate induced isolation of plasmid DNA (2014) Online International Interdisciplinary Research Journal ISSN2249-9598, Volume-IV
21. Götz A, Pukall R, Smit E, Tietze E, Prager R, Tschäpe H, Van Elsas JD, Smalla K. Detection and characterization of broad-host-range plasmids in environmental bacteria by PCR. Applied and environmental microbiology 1996; 62(7): 2621-2628.
22. Durmaz R, Otlu B, Koksall F. The optimization of a rapid pulsed-field gel electrophoresis protocol for the typing of *Acinetobacter baumannii*, *Escherichia coli* and *Klebsiella* spp. Jpn J Infect Dis 2009; 62: 372-377.
23. Ibrahim DR, Dodd CE, Stekel DJ, Meshioye RT, Diggle M, Lister M, Hobman JL. Multidrug-Resistant ESBL-Producing *E. coli* in Clinical Samples from the UK. Antibiotics 2023; 12(1): 169.
24. Birru M, Woldemariam M, Manilal A, Aklilu A, Tsalla T, Mitiku A, Gezmu T. Bacterial profile, antimicrobial susceptibility patterns, and associated factors among bloodstream infection suspected patients attending Arba Minch General Hospital, Ethiopia. Scientific reports 2021; 11(1): 1-13.
25. Duran H, Çeken N, Atik B. *Escherichia coli* ve *Klebsiella pneumoniae* Türlerinde Antibiyotik Direnci ne Durumda? Yoğun Bakım Ünitesinden Beş Yıllık Analiz. Fırat Tıp Dergisi/Firat Med J 2022; 27(2): 116-120
26. Son T, Manh ND, Trung NT, Quyen DT, Meyer CG, Phuong NTK, Hoan PQ, Sang VV, Nurjadi D, Velavan T, Bang MH, Song LH. Molecular detection of blaCTX-M gene to predict phenotypic cephalosporin resistance and clinical outcome of *Escherichia coli* bloodstream infections in Vietnam. Annals of clinical microbiology and antimicrobials 2021; 20(1): 1-9.
27. Kreidl P, Kirchner T, Fille M, Heller I, Lass-Flörl C, Orth-Höller D. Antibiotic resistance of blood cultures in regional and tertiary hospital settings of Tyrol, Austria (2006-2015): Impacts & trends. PloS one 2019; 14(10): e0223467.
28. Anvarinejad M, Khalifeh M, Mardaneh J, Pouladfar G, Dehyadegari MA. Bloodstream infections in the south of Iran: microbiological profile and antibiotic-resistance patterns of isolated bacteria. Crescent J Med Biol Sci 2021; 8(4): 312-318.
29. Hu LF, Chang X, Ye Y, Wang ZX, Shao YB, Shi W, Li X, Li JB. *Stenotrophomonas maltophilia* resistance to trimethoprim/sulfamethoxazole mediated by acquisition of sul and dfrA genes in a plasmid-mediated class 1 integron. Int J Antimicrob Agents 2011; 37(3): 230-4.
30. Sacristán C, Esperón F, Herrera-León S, Iglesias I, Neves E, Nogal V, Muñoz MJ, de la Torre A. Virulence genes, antibiotic resistance and integrons in *Escherichia coli* strains isolated from synanthropic birds from Spain. Avian Pathol 2014; 43(2): 172-5.
31. Zhu W, Wang T, Zhu Y, Xiao L, Liu W, Wei Q. Two-dimensional PCR for detecting class 1, 2 and 3 integrons. Heliyon 2022; 8(11): e11844.
32. Başkan C, Sırıken B, Tüfekci EF, Kılınç Ç, Ertürk Ö, Erol İ (2022) Presence of quorum sensing system, virulence genes, biofilm formation and relationship among them and class 1 integron in carbapenem-resistant clinical *Pseudomonas aeruginosa* isolates. Archives of Microbiology 2022; 204(8): 464.
33. Rakici E, Altunisik A, Sahin K, Ozgumus OB. Determination and molecular analysis of antibiotic resistance in Gram-negative enteric bacteria isolated from *Pelophylax* sp. in the Eastern Black Sea Region. Acta Veterinaria Hungarica 2021; 69(3): 223-233.
34. Sivri N, Sandalli C, Ozgumus OB, Colakoglu F, Dogan D. Antibiotic resistance profiles of enteric bacteria isolated from Kucukcekmece Lagoon (Istanbul-Turkey). Turkish Journal of Fisheries and Aquatic Sciences 2012; 12(3).
35. Liu W, Zhao F, Chen J. Analysis of drug resistance genes of integrons in clinical isolates of *Escherichia coli* from elderly bloodstream infections. Cellular and Molecular Biology 2022; 68(6): 67-72.
36. Abdel-Rhman SH, Elbargisy RM, Rizk DE. Characterization of integrons and quinolone resistance in clinical *Escherichia coli* isolates in Mansoura City, Egypt. International Journal of Microbiology 2021.
37. Dehkordi MK, Halaji M, Nouri S. Prevalence of class 1 integron in *Escherichia coli* isolated from animal sources in Iran: a systematic review and meta-analysis. Tropical Medicine and Health 2020; 48:16.
38. Ahumada-Santos YP, Báez-Flores ME, Díaz-Camacho SP, de Jesús Uribe-Beltrán M, Eslava-Campos CA, Parra-Unda JR, Delgado-Vargas F. Association of phylogenetic distribution and presence of integrons with multidrug resistance in *Escherichia coli* clinical isolates from children with diarrhoea. Journal of Infection and Public Health 2020; 13(5): 767-772.
39. Partridge SR, Tsafnat G, Coiera E, Iredell JR. Gene cassettes and cassette arrays in mobile resistance integrons. FEMS Microbiol Rev 2009; 33(4): 757-784.

- 
40. Ndi OL, Barton MD. Incidence of class 1 integron and other antibiotic resistance determinants in *Aeromonas* spp. from rainbow trout farms in Australia. *J Fish Dis* 2011; 34(8): 589-99.
  41. Szmolka A, Nagy B. Multi drug resistant commensal *Escherichia coli* in animals and its impact for public health. *Frontier in microbiology* 2013; 4: 258. <https://doi.org/10.3389/fmicb.2013.00258>
  42. Düzgün AÖ, Saral A (2018) Investigation of Antibiotic Resistance Genes and Class 1 Integron in Multi-Drug Resistant *P. aeruginosa* and *E. coli* Strains. *Cumhuriyet Sci. J.* 2018; Vol. 39-4: 1063-1068.
  43. Chen J, Su Z, Liu Y, Wang S, Dai X, Li Y, Peng S, Shao Q, Zhang H, Wen P, Yu J, Huang X, Xu H. Identification and characterization of class 1 integrons among *Pseudomonas aeruginosa* isolates from patients in Zhenjiang, China. *International Journal of Infectious Diseases* 2009; 13(6): 717-721.
  44. Schlüter A, Szczepanowski R, Pühler A, Top EM. Genomics of IncP-1antibiotic resistance plasmids isolated from wastewater treatment plants provides evidence for a widely accessible drug resistance gene pool. *FEMS Microbiol Rev* 2007; 31: 449-477.