



Effects of acetlysalicylic acid with indole-3-acetic acid on rooting and pigmentation in *Amygdalus L.*

Gulcin Beker AKBULUT¹ and Emel YİĞİT^{2,*}

¹Tunceli University, Tunceli Vocational School, Department of Organic Agriculture, Tunceli/TURKEY

²Inonu University, Science and Art Faculty, Department of Biology, 44280 Malatya/TURKEY

Received: 05.04.2013; Accepted: 03.07.2013

Abstract. Vegetative propagation is a key step, playing an important role in the succesful production of elite clones. The use of plant hormanes can increase the rroting capacity of cuttings. In this experiment, we investigated whether exogenously applied acetylsalicylic acid (ASA) with indole-3-acetic acit (IAA) (50, 100 mg/L) through the rooting medium could increase effects on *Amygdalus* spp or not. In the experiment, one year old semihardwood shoot cuttings were used. The highest callus formation was determined on the applied cuttings with 50 mg/L ASA with IAA in all groups. Rooting among the application groups in *A. lycioides* was observed around 10% in 50 mg/L ASA with IAA treatment group. Pigmentation contents in cuttings were found to be important in a time-dependent fashion.

Keywords: Rooting, acetylsalicylic acid, indole-3-acetic acid,callus

1. INTRODUCTION

Many species and cultivars of *Amygdalus* are produced for their edible fruit or as garden ornamentals, and a few species are also grown for their edible seeds [1]. *Prunus amygdalus* seedlings have been used for production since many centuries [2]. *P.amygdalus* is a nut tree, which has been grown in large areas in the world due to having climatic tolerance and having many microclimates suitable for almond growth [3]. In Turkey, the production budding and/or grafting are not common; and instead seed has been usually used for production [4]. The initiation and development of adventitious roots are vital processes for successful clonal propagation by cuttings.

Although rooting of several *Prunus* species has been reported, including apricot [5], cherry [6], peach [7], and plum [8], *in vitro* rooting of almond has proven difficult [9]. Reports describing rooting of adult almond explants are limited, and have primarily focused on hard shell cultivars grown in Europe [10, 11]. In Australia and the USA, paper shell types such as ‘Nonpareil’ are the main cultivars grown. Limited success has so far been achieved for the *in vitro* induction of adventitious roots with the premier paper shell cultivar ‘Nonpareil’ [12]. Nicotra and Pellegrini (1989) were reported that adventitious roots values given emitted higher than 30% only in 7 varieties (M 51, M 49, Ferragnes, M55, Genco, M50 and Texas) of almond’s 26 varieties [13].

* Corresponding author. *Email address:* emel-yigit@windowslive.com

Cuttings from some species, clones, or developmental phases within clones are competent for adventitious root initiation, while others are not [14]. Adventitious root formation is a requirement in any successful vegetative propagation program, many ecologically and economically important hardwood tree species have a low genetic or physiological capacity for adventitious root formation and are considered recalcitrant to routine commercial-scale vegetative propagation. However, successful propagation of difficult-to-root species can be achieved if the type of cutting (hardwood, softwood, or root) date of collection (seasonal growth development), stock plant or cutting manipulation (pruning, wounding etc.), rooting treatment (auxin type and concentration, rooting media) and greenhouse parameters (mist bed system, supplemental, lighting, temperature, etc.) are carefully considered [15].

Stimulation of root initiation, the second was the first practical use to be made of growth regulators [16]. Furthermore, the stimulation of adventitious roots by phenolic compounds has been reported for a number of plant species. Catechol, *p*-hydroxybenzoic acid, pyrogallol and salicylic acid stimulated root initiation in mung bean cuttings [17].

Salicylic acid (SA), a common plant phenolic compound, influences numerous physiological and biochemical process in plants. These include adventitious root initiation [18, 19], inhibition of ethylene biosynthesis [20], disease resistance [21], salt and osmotic stress [19, 22], ozone [23], growth and photosynthesis [22, 24], heat production [25], chilling tolerance [26]. Thus more and more attention was paid to the action of SA, and as a result, SA has been proposed as a new plant hormone [27]. In tissue culture applications have emerged for SA or its artificial analogue acetylsalicylic acid (ASA) which has been found to be more effective in some systems [28]. Auxins have been shown to have the greatest effect on rooting [29-31]. Nevertheless, auxins have failed to promote root initiation or they had only a slight rooting effect in the case of hard-to-root olive cultivars [32, 33]. Treatment of cuttings with acetylsalicylic acid in combination with IAA can be of value in the propagation of woody plants [17].

In this study, we investigated the effects of ASA with IAA (50, 100 mg/L) on rooting and pigmentation in three species of *Amygdalus* (*A. trichamygdalus*, *A. orientalis* and *A. lycioides*).

MATERIALS AND METHODS

Preparation of the plant samples

One year old shoot cuttings of ten-year-old plants of *Amygdalus trichamygdalus*, *Amygdalus orientalis* and *Amygdalus lycioides* taken in November and December 2002 from Gündüzbey country of Malatya were used. The cuttings were prepared, 20-25 cm in length with 8-10 buds. The cuttings were washed for 24 h with tap water. Basal ends of cuttings were soaked in 0, 50, 100, mg/L ASA+IAA dissolved in water for 24 hours. The treatments had three replicates which have ten cuttings each. The cuttings were grown in controlled climate chamber at 24±2°C, 60±5% relative humidity, a photoperiod of 16 h, and a light intensity of 12.000 lux. Samples were grown perlite and with three-repetitions by using Hoagland solution [34]. The callus was recorded two months after planting.

Pigment extraction and determination

In stem cuttings were analyzed chlorophyll a (Chl a), chlorophyll b (Chl b), total chlorophyll and carotenoid (Car) contents after ASA+IAA treatment. During the extraction of the pigments, methods suggested by De-Kok and Graham (1989) were used [35]. The absorbance values of the extraction process, were determined (UV-1201V Shimadzu spectrophotometer) at 470, 645 and 662, nm according to Lichtenthaler and Welburn (1983) and total rates of Chl a, Chl b, total Chl and Car were calculated [36].

Statistical analysis

Statistical analysis were performed using SPSS 10.0 for Windows. Duncan's Multiple Range Test was employed to determine the statistical significance of differences among the means [37].

RESULTS

Application of ASA with IAA caused a significant increase in callus rate of *A. lycioides* in November and in December. At the same time a rooting of approximately 10% was occurred in the group applied 50 mg/L of ASA with IAA in November. As seen in Table 1, compared to control ASA with IAA treatment caused a marked decrease in callus rate of *A. orientalis* in November. On the other hand in *A. trichamygdalus* no such variations were recorded in November and in December (Table 1).

Table 1 Callus rate in the cuttings of *A. trichamygdalus*, *A. orientalis* and *A. lycioides* on a monthly basis.

			Callus rate (%)	Rooting rate (%)
November 2002	<i>A. trichamygdalus</i>	Control	47.20±2.45 ^b	ND
		50 mg/L ASA+IAA	54.85±5.01 ^a	ND
		100 mg/L ASA+IAA	46.97±2.63 ^b	ND
	<i>A. orientalis</i>	Control	90.00±2.55 ^a	ND
		50 mg/L ASA+IAA	59.87±3.76 ^b	ND
		100 mg/L ASA+IAA	56.70±2.90 ^b	ND
	<i>A. lycioides</i>	Control	87.78±4.72 ^b	ND
		50 mg/L ASA+IAA	100.00±0.00 ^a	9.86±0.09
		100 mg/L ASA+IAA	93.63±5.53 ^b	ND
December 2002	<i>A. trichamygdalus</i>	Control	40.00±0.05 ^b	ND
		50 mg/L ASA+IAA	50.00±0.05 ^a	ND
		100 mg/L ASA+IAA	40.00±0.05 ^b	ND
	<i>A. orientalis</i>	Control	40.00±0.05 ^b	ND
		50 mg/L ASA+IAA	50.00±0.05 ^a	ND
		100 mg/L ASA+IAA	40.00±0.05 ^b	ND
	<i>A. lycioides</i>	Control	83.33±5.77 ^b	ND
		50 mg/L ASA+IAA	93.33±5.77 ^a	ND
		100 mg/L ASA+IAA	80.00±3.64 ^b	ND

n=3

Within same column, means followed by letters are significantly different from each other ($P<0.05$) according to Duncan's test

ND=Not determined

Photosynthetic pigment content (Chl a, Chl b, total Chl and Car) were presented in Table 2. The highest Chl a amount was observed in *A. orientalis* groups which were treated with 50 mg/L ASA with IAA during the month of November as 5.08 µg/g fresh weight (FW) and December (6.20 µg/g FW). The highest Chl b amount, on the other hand, was determined in the same groups treated with 100 mg/L ASA with IAA during the month of December. In the species of *A. trichamygdalus* and *A. orientalis* applied 100 mg/L of ASA+IAA, the total chlorophyll amount increased noticeably. The highest carotenoid content was recorded in the groups applied 50 mg/L of ASA with IAA as 2.68 µg/g FW and 3.83 µg/g FW in November and December months respectively (Table 2).

Effects of acetlysalicylic acid with indole-3-acetic acid

Table 2 Pigment content ($\mu\text{g.g}^{-1}$ fresh weight) in the cuttings of *A. trichamygdalus*, *A. orientalis* and *A. lycioides* on a monthly basis.

			Chl a	Chl b	Total chlorophyll	Carotenoid
November 2002	<i>A. trichamygdalus</i>	Control	4.13±0.42 ^b	3.99±1.79 ^a	8.12±2.05 ^b	1.14±0.46 ^b
		50mg/L ASA+IAA	4.09±1.55 ^b	3.77±1.34 ^b	7.86±2.87 ^c	0.97±0.33 ^c
		100 mg/L ASA+IAA	4.78±0.94 ^a	4.01±1.42 ^a	8.79±2.33 ^a	1.46±0.60 ^a
	<i>A. orientalis</i>	Control	3.86±0.55 ^c	2.27±0.42 ^c	6.13±0.38 ^c	1.84±0.27 ^b
		50 mg/L ASA+IAA	5.08±1.23 ^a	3.04±0.41 ^b	8.12±1.34 ^b	2.68±1.28 ^a
		100 mg/L ASA+IAA	4.95±0.03 ^b	4.70±1.31 ^a	9.65±1.31 ^a	1.01±0.10 ^c
	<i>A. lycioides</i>	Control	2.97±0.15 ^b	2.87±0.24 ^a	5.84±0.34 ^a	1.95±0.06 ^c
		50 mg/L ASA+IAA	2.80±0.16 ^c	2.23±0.22 ^c	5.03±0.16 ^c	2.04±0.06 ^b
		100 mg/L ASA+IAA	3.21±0.48 ^a	2.56±1.10 ^b	5.77±0.68 ^b	2.22±0.08 ^a
December 2002	<i>A. trichamygdalus</i>	Control	4.00±0.49 ^c	3.13±1.53 ^c	7.13±1.75 ^c	1.49±0.51 ^b
		50 mg/L ASA+IAA	4.59±1.43 ^b	3.38±1.01 ^b	7.97±2.15 ^b	1.83±1.26 ^a
		100 mg/L ASA+IAA	4.87±0.64 ^a	4.35±1.35 ^a	9.22±1.86 ^a	1.24±0.47 ^c
	<i>A. orientalis</i>	Control	3.54±0.54 ^c	2.46±0.55 ^c	6.00±0.44 ^c	2.05±0.19 ^b
		50 mg/L ASA+IAA	6.20±0.20 ^a	3.12±0.39 ^b	9.32±0.20 ^b	3.83±0.16 ^a
		100 mg/L ASA+IAA	4.95±0.04 ^b	4.89±0.09 ^a	9.84±0.13 ^a	0.93±0.04 ^c
	<i>A. lycioides</i>	Control	2.54±0.27 ^c	1.84±1.13 ^b	4.38±1.08 ^c	1.96±0.05 ^c
		50 mg/L ASA+IAA	3.03±0.32 ^a	2.07±0.29 ^a	5.10±0.21 ^a	2.04±0.11 ^b
		100 mg/L ASA+IAA	2.83±0.53 ^b	2.12±0.86 ^a	4.95±0.99 ^b	2.29±0.21 ^a

n=3

Within same column, means followed by letters are significantly different from each other ($P<0.05$) according to Duncan's test

DISCUSSION

Many researches reported that rooting process in *Amygdalus* propagation was very difficult. Due to this reason alternative rooting forms was studied by the use of cutting exudation solution [38] or indicated plant growth regulators as indol-butyric acid (IBA) and naphthalene acetic acid (NAA), nevertheless it is necessary to determine the best concentration of these products specifically for *Amygdalus* L. [39, 40]. Antonopoulou et al. (2005) reported that the effect of vitamin riboflavin (B₂) on *in vitro* of the almond x peach hybrid clone GF 667 did not stimulate adventitious rooting of the explants and rooting was very low in comparison with the control treatment. Our results were determined difficult of rooting of *Amygdalus* spp. However, in this study, in ASA with IAA application groups were observed high callus formation and rarely root formation [41].

In addition to auxins, phenolic compounds either alone or in combination of auxins [17, 42] have also stimulated adventitious root formation in cuttings of several species. Singh (1993) [43] found that SA stimulated root formation in young shoots of ornamental plants and Li and Li (1995) reported the formation of adventitious roots on hypocotyl cuttings of mung beans [44]. SA had synergistically acted with IAA and promoted the root formation in mung bean cuttings, but had no effect on *Acer* cuttings [17]. SA combined with NAA synergistically promoted the root

number and root lengths of the cuttings of several *Populus* spp. Combined application of IBA with various concentration of SA significantly promoted the rooting of cuttings with respect to untreated ones. But the rooting was markedly low compared to effect of IBA (5 g l⁻¹) [45]. Although this effect was seemed to be in relation with the clonal differences and cutting time rather than concentration and treatment method [46]. In our study, reported that a rooting of approximately 10% was occurred in the group applied 50 mg/L of ASA+IAA in November in *A. lycioides*. The best callus formation was observed on the applied cuttings with 50 ppm ASA with IAA in all application groups (Table 1).

The effect of SA application was expected in view of earlier studies that showed increased or decreased photosynthetic pigments following SA application, depending on type of species or cultivar. For instance, Chandra and Bhatt (1998) observed that an increasing or decreasing effect of SA on chlorophyll content of cowpea (*Vigna unguiculata*) depends on the genotype [47]. In another study treatment with SA increased pigment contents in maize [22] and wheat [48] grown under normal or stress conditions [19]. Radwan et al. (2006) were reported that in all SA treatments, chlorophyll a/b ratio was higher than healthy controls [49]. Considering phenols effect, the lower and the moderate concentrations of salicylic acid (50 and 100 mg/L) recorded the highest values of chlorophyll a, b and carotenoids [50]. In the species of *A. trichamygdalus* and *A. orientalis* applied 100 mg/L of ASA with IAA, the total chlorophyll amount increased noticeably. The highest carotenoid amount was recorded in the groups applied 50 mg/L of ASA with IAA as 2.68 µg/g FW and 3.83 µg/g FW in November and December months respectively (Table 2). Our results obtained from this work agree with similar findings of other researchers [19, 46, 47].

In conclusion, exogenously applied ASA can influence a number of biological processes in plants [27]. The present research, in one year old shoot cuttings of *Amygdalus* spp. described effects on rooting and pigment content of ASA with IAA. In shoot cuttings of *A. lycioides* was seen to enhance adventitious rooting of ASA with IAA at 50 mg/L. The results confirmed that ASA with IAA promoted to callusing in application groups. Pigmentation changes occurred in application groups, but there was no correlation between pigmentation rates and callusing. In the results of study callus were formed high proportion (%93-100) in treated groups, unfortunately no or very few new rooting (10%) was observed.

REFERENCES

- [1] Lingdi, L. and Bartholomew, B., 48. *Amygdalus Linnaeus*, Sp. Pl. 1: 472. 1753, Flora of China, 9 (2003) 391-395.
- [2] Edstromand, J.P. and Viveros, M., Almond production manual. In: division of agricultural and natural resources, Micke, W.C. (ed.). Univ. of California Publication, (1996) 3364.
- [3] Dokuzoguz, M. ve Gulcan, R., Almond growth and obstacles, Tubitac Publications (1979) 80.
- [4] Ercisli, S., A short review of the fruit germplasm resources of Turkey, Genetic Resources and Crop Evolution, 51 (2004) 419-435.
- [5] Perez-Tornero, O., Egea, J., Vanoostende, A. and Burgos L., Assessment of factors affecting adventitious shoot regeneration from in vitro cultured leaves of apricot, Plant Science, 158 (2000) 61-70.
- [6] Hammatt, N. and Grant N.J., Micropropagation of mature British wild cherry, Plant Cell Tissue and Organ Culture, 47 (1997) 103-110.
- [7] Hammerschlag, F.A., Bauchan, G.R. and Scorza, R., Factors influencing in vitro multiplication and rooting of peach cultivars, Plant Cell Tissue and Organ Culture, 8 (1987) 235-242.
- [8] Aier, N.B., Sharma, S.D., Micropropagation in some plum cultivars, Fruit Science Reports, 17 (1990) 57-63.
- [9] Kester, D.E and Gradziel, T.M. and Grassely C., Almonds. *Acta Horticulturae*, 290 (1990) 699-758.
- [10] Caboni, E. and Damiano, C., Rooting in two Almond genotypes, Plant Science, 96 (1994) 163-165.
- [11] Caboni, E., Tonelli, M.G., Lauri, P., Lacovacci, P., Kevers, C., Damiano, C. and Gaspar T., Biochemical aspects of Almond microcuttings related to in vitro rooting ability, *Biologia Plantarum*, 39 (1997) 91-97.
- [12] Ainsley, P.J., Collins, G.G. and Sedgley M., In vitro rooting of Almond (*Prunus dulcis* Mill.), *In vitro Cellular & Developmental Biology-Plant*, 37 (2001) 778-785.
- [13] Nicotra, A. and Pellegrini, M., Almond rootstock breeding for easy propagation. *Options méditerranéennes, Série Séminaires*, 5 (1989) 51-60.
- [14] Steven, T.L., Alan, G.S. and Wesley, P.H., Differential gene expression in response to auxin treatment in the wild type and Rac, an adventitious rooting-incompetent mutant of tobacco, *Plant Physiology*, 114 (1997) 1197-1206.
- [15] Pijut, P.M. and Espinosa, A.C., Development of a rooted cutting propagation method for *Prunus serotina*, *The International Plant Propagators' Society Combined Proceedings*, 54 (2005) 129-131.
- [16] Hartmann, H.T. and Kester, D.E., *Plant Propagation: Principles and Practices*. Prentice-Hall, Englewood Cliffs, New Jersey 1975.

- [17] Kling, G.J. and Meyer, M.M., Effect of phenolic compounds and indole acetic acid and adventitious root initiation in *Phaseolus aureus*, *Acer saccharinum* and *Acer griseum*, Horticultural Science, 18 (1983) 352-354.
- [18] Kang, S.M., Jung, H.Y., Kang, Y.M., Dae-Jin, Y., Bahk, J.D., Yang, J.K. and Choi, M.S., Effects of methyljasmonate and salicylic acid on the production of tropane alkaloids and the expression of PMT and H6H in adventitious root cultures of *Scopolia parviflora*, Plant Science, 166 (2004) 745-751.
- [19] Arfan, M., Athar, H.R. and Ashraf, M., Does exogenous application of salicylic acid through the rooting medium modulate growth and photosynthetic capacity in two differently adapted spring wheat cultivars under salt stress, Journal of Plant Physiology, 164 (2007) 685-694.
- [20] Leslie, C.A. and Romani, R.J. Inhibition of ethylene biosynthesis by salicylic acid, Plant Physiology, 88 (1988) 833-837.
- [21] Mandal, S., Nirupama, M. and Adinpunya, M., Salicylic acid-induced resistance to *Fusarium oxysporum* f. sp. *Lycopersici* in tomato, Plant Physiology and Biochemistry, 47 (2009) 642-649.
- [22] Khodary, S.E.A., Effect of salicylic acid on growth, photosynthesis and carbohydrate metabolism in salt stressed maize plants, International Journal of Agriculture and Biology, 6 (2004) 5-8.
- [23] Koch, J.R., Creelman, R.A., Eshita, S.M., Seskar, M., Mullet, J.E. and Davis, K.R., Ozone sensitivity in hybrid poplar correlates with insensitivity to both salicylic acid and Jasmonic Acid. The Role of Programmed Cell Death in Lesion Formation, Plant Physiology, 123 (2000) 487-496.
- [24] El-Tayeb, M.A., Response of barley grains to the interactive effect of salinity and salicylic acid, Plant Growth Regulation, 45 (2005) 215-224.
- [25] Raskin, I., Skubatz, H., Tang, W. and Meeuse, B.J.D., Salicylic acid levels in thermogenic and non-thermogenic plants, Annals of Botany, 66 (1990) 369-373.
- [26] Senaratna, T., Touchell, D., Bunn, E. and Dixon K., Acetylsalicylic acid (Aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plants. Plant Growth Regulation, 30 (2000) 157-161.
- [27] Raskin I., Role of salicylic acid in plants, Annual Review of Plant Physiology and Plant Molecular Biology, 43 (1992) 439-463.
- [28] Lopez-Delgado, H., Dat, J.F, Foyer, C.H. and Scott, I.M., Induction of thermotolerance in potato microplants by acetylsalicylic acid and H₂O₂, Journal of Experimental Botany, Vol. April 49:321 (1998) 713-720.
- [29] Hartmann, H.T., Kester, D.E., Davies, F.T. and Geneve R.L., Plant Propagation Principles and Practices, 5th ed. Prentice-Hall, New Jersey. 2001.
- [30] Kelen, M. and Ozkan G., Relationships between rooting ability and changes of endogenous IAA and ABA during the rooting of hardwood cuttings of some grapevine rootstocks, Eur. J. Hort. Sci., 68 (2003) 8-13.
- [31] Negash, L., Vegetative propagation of the threatened African wild olive [*Olea europaea* L. subsp. *cuspidata* (Wall. ex DC.) Cifffieri]. New Forest, 26 (2003) 137-146.

Effects of acetlysalicylic acid with indole-3-acetic acid

- [32] Wiesman, Z. and Lavee, S., Enhancement of stimulatory effects on rooting of olive cultivar stem cuttings, *Sci Hort*, 62 (1995a) 189-198.
- [33] Denaxa, N.K., Vemmos, S.N. and Roussos, P.A., The role of endogenous carbohydrates and seasonal variation in rooting ability of cuttings of an easy and a hard to root olive cultivars (*Olea europaea* L.), *Scientia Horticulturae* 143 (2012) 9-28.
- [34] Hoagland, D.R. and Arnon, D.I. The water culture method for growing plants without soil, *California Agr Expt Sta Circ*, 37 (1938).
- [35] De-Kok, L. and Graham M., Levels of pigments, soluble proteins, amino acids and sulphhydryl compounds in foliar tissue of *Arabidopsis thaliana* during dark induced and natural senescence, *Plant Physiology and Biochemistry*, 27 (1989) 203-209.
- [36] Lichtenthaler K. and Welburn A.R., Determination of total carotenoids and chlorophylls a and b of leaf extracts in different solvents, *Botanisches Institut der Univeristat, Kaiserstran ße 12, Postfach*, (1983).
- [37] Duncan, D.B., Multiple Range and Multiple F tests. *Biometrics*, 11:1 (1955).
- [38] Mneney, E.E. and Mantell, S.E. Effect of deliberate wounding and natural cashew gum extracts in improving rooting and resilience of cashew cuttings post-excision, *Research & Training Newsletter, Dar es Salam*, 18 (2003) 8-11.
- [39] Martin, K.P. Plant regeneration through direct somatic embryogenesis on seed coat explants of Cashew (*Anacardium occidentale* L.), *Scientia Horticulturae, Wageningen*, 98 (2003) 299-304.
- [40] Lopes, R.L., Cavalcante, I.H.L., Oliveira, I.V.D.M., and Martins A.B.G., Indol butyric acid levels on cashew cloning by air-layering process, *Revista Brasileira de Fruticultura, Jaboticabal-SP*, 27 (2005) 517-518.
- [41] Antonopoulou, C., Dimassi, K., Therios, I., Chatzissavvidis, C. and Tsirakoglou, V., Inhibitory effect of riboflavin (Vitamin B₂) on the in vitro rooting and nutrient concentration of explants of peach rootstock GF 677 (*Prunus amygdalus* x *P. persica*), *Scientia Horticulturae*, 106 (2005) 268-272.
- [42] Nandi, S.K., Rikhari, H.C., Nadeem, M. and Palni, L.M.S., *Physiology and Molecular Biology of Plants*, 3 (1997) 15.
- [43] Singh, S.P., Effect of non-auxinic chemicals on root formation in some ornamental plant cuttings, *Advances in Horticultural Science*, 3 (1993) 207-210.
- [44] Li, L. and Li, L., Effects of recorcinol and salicylic acid on the formation of adventitious Roots on hypocotyl cutting of *Vigna radiata*, *Journal of Topical an Subtropical Botany*, 3 (1995) 67-71.
- [45] Isfendiyaroglu, M. and Ozeker, E., Rooting of *Olea europaea* 'Domat' cuttings by auxin and salicylic acid treatments, *Pakistan Journal of Botany*, 40 (2008) 1135-1141.
- [46] Bojarczuk, T. and Jankiewicz, S.L., Influence of phenolic substances on rooting of softwood cuttings of *Populus alba* L., *P. canescens* Sm., *Acta Agrobotanica*, 28 (1975) 121-129.

[47] Chandra, A. and Bhatt, R.K., Biochemical and physiological response to salicylic acid in relation to the systemic acquired resistance, *Photosynthetica*, 45 (1998) 255-258.

[48] Singh, B. and Usha, K., Salicylic acid induced physiological and biochemical changes in wheat seedlings under water stress, *Plant Growth Regulation*, 39 (2003) 137-141.

[49] Radwan, D.E.M., Fayez, K.A., Mahmoud, S.Y., Hamad, A. and Lu, G., Salicylic acid alleviates growth inhibition and oxidative stress caused by zucchini yellow mosaic virus infection in *Cucurbita pepo* leaves, *Physiological and Molecular Plant Pathology*, 69 (2006) 172-181.

[50] Amin, A.A., Rashad, El-S H.M., El-Abagy, H.M.H., Physiological effect of indole-3-butyric acid and salicylic acid on growth, yield and chemical constituents of onion plants, *Journal of Applied Sciences Research*, 3 (2007) 1554-1563.