

The geometry of tangent conjugate connections

Adara M. Blaga^{*} and Mircea Crasmareanu[†]

Abstract

The notion of conjugate connection is introduced in the almost tangent geometry and its properties are studied from a global point of view. Two variants for this type of connections are also considered in order to find the linear connections making parallel a given almost tangent structure.

2000 AMS Classification: 53C15; 53C05; 53C10; 53C07.

Keywords: almost tangent structure; conjugate linear connection.

Received 22/02/2013 : Accepted 25/03/2014 Doi : 10.15672/HJMS.2015449431

Introduction

Let F be a tensor field of $(1, 1)$ -type on a given smooth manifold M . An interesting object in the geometry of pair (M, F) is provided by the class of F -linear connections i.e. linear connections ∇ making F parallel: $\nabla F = 0$. In order to determine this class, in [9] is introduced the notion of F -conjugate connection associated to a fixed (non-necessary F -connection) ∇ . By denoting $\nabla^{(F)}$ this F -conjugate connection we have studied the geometry of $(M, F, \nabla, \nabla^{(F)})$ until now for two cases: almost complex structures in [1] and almost product structures in [2].

The present work is devoted to another remarkable type of tensor fields of $(1, 1)$ -type, namely *almost tangent structures*. These structures were introduced by Clark and Bruckheimer [5] and Eliopoulos [10] around 1960 and have been investigated by several authors, see [3], [6]-[8], [16], [18]. As it is well-known, the tangent bundle of a manifold carries a canonical integrable almost tangent structure, hence the name. This tangent structure plays an important rôle in the Lagrangian description of analytical mechanics, [7]-[8], [12].

Recall that we are interested in the class of J -linear connections since, according to [15, p. 120], the existence of a symmetric (torsion-free) one in this class implies the

^{*}Department of Mathematics and Computer Science, West University of Timișoara, Bld. V. Pârvan nr. 4, 300223, Timișoara, România
Email: adarablaga@yahoo.com

[†]Faculty of Mathematics, University "Al. I. Cuza", 700506, Iași, România
Email: mcrasm@uaic.ro

integrability of J in the sense of G -structures as is discussed below; for example, J -linear connections of Levi-Civita type are studied in [11]. An important difference between the former structures (almost complex, almost product) and the later (almost tangent) is given by the fact that an almost tangent structure J is a degenerate tensor field due to its nilpotence $J^2 = 0$, see the following Section. An example where this difference is obvious is the duality property $(\nabla^{(F)})^{(F)} = \nabla$ which holds for a non-degenerate F while for almost tangent structures we have ii) of our Proposition 2.1.

The content of paper is as follows. After a short survey in almost tangent geometry we introduce the tangent conjugate connection $\nabla^{(J)}$ in Section 2 following the pattern of [1]-[2]. Its properties are studied following the same way as in the cited papers; for example the difference $\nabla^{(J)} - \nabla$ is expressed again in terms of two tensor fields of $(1, 2)$ -types called *structural* and *virtual* tensor fields. We study also the behavior of the tangent conjugate connections for a family of anti-commuting almost tangent structures. In the last two Sections we generalize $\nabla^{(J)}$, firstly through an exponential process and secondly with a general tensor field of $(1, 2)$ -type.

1. Almost tangent geometry revisited

Let M be a smooth, m -dimensional real manifold for which we denote: $C^\infty(M)$ -the real algebra of smooth real functions on M , $\Gamma(TM)$ -the Lie algebra of vector fields on M , $T_s^r(M)$ -the $C^\infty(M)$ -module of tensor fields of (r, s) -type on M . An element of $T_1^1(M)$ is usually called *vector 1-form* or *affinor*.

Recall the concept of almost tangent geometry:

1.1. Definition. $J \in T_1^1(M)$ is called *almost tangent structure* on M if it has constant rank and:

$$ImJ = \ker J. \quad (1.1)$$

The pair (M, J) is called *almost tangent manifold*.

The name is motivated by the fact that (1.1) implies the nilpotence $J^2 = 0$ exactly as the natural tangent structure of tangent bundles. Denoting $rankJ = n$ it results $m = 2n$. If in addition, we suppose that J is integrable i.e.:

$$N_J(X, Y) := [JX, JY] - J[JX, Y] - J[X, JY] + J^2[X, Y] = 0 \quad (1.2)$$

then J is called *tangent structure* and (M, J) is called *tangent manifold*.

From [17, p. 3246] we get some features of tangent manifolds:

(i) the distribution $ImJ (= \ker J)$ defines a foliation denoted $V(M)$ and called *the vertical distribution*.

1.2. Example. $M = \mathbb{R}^2$, $J_e(x, y) = (0, x)$ is a tangent structure with $\ker J_e$ the Y -axis, hence the name. The subscript e comes from "Euclidean".

(ii) there exists an atlas on M with local coordinates $(x, y) = (x^i, y^i)_{1 \leq i \leq n}$ such that $J = \frac{\partial}{\partial y^i} \otimes dx^i$ i.e.:

$$J \left(\frac{\partial}{\partial x^i} \right) = \frac{\partial}{\partial y^i}, \quad J \left(\frac{\partial}{\partial y^i} \right) = 0. \quad (1.3)$$

We call *canonical coordinates* the above (x, y) and the change of canonical coordinates $(x, y) \rightarrow (\tilde{x}, \tilde{y})$ is given by:

$$\begin{cases} \tilde{x}^i = \tilde{x}^i(x) \\ \tilde{y}^i = \frac{\partial \tilde{x}^i}{\partial x^a} y^a + B^i(x). \end{cases} \quad (1.4)$$

It results an alternative description in terms of G -structures. Namely, a tangent structure is a G -structure with:

$$G = \left\{ C = \begin{pmatrix} A & O_n \\ B & A \end{pmatrix} \in GL(2n, \mathbb{R}); A \in GL(n, \mathbb{R}), B \in gl(n, \mathbb{R}) \right\} \quad (1.5)$$

and G is the invariance group of matrix $J = \begin{pmatrix} O_n & O_n \\ I_n & O_n \end{pmatrix}$ i.e. $C \in G$ if and only if $C \cdot J = J \cdot C$.

The natural almost tangent structure J of $M = TN$ is an example of tangent structure having exactly the expression (1.3) if (x^i) are the coordinates on N and (y^i) are the coordinates in the fibers of $TN \rightarrow N$. Also, J_e of Example 1.2 has the above expression (1.3) with $n = 1$, whence it is integrable. A third class of examples is obtained by duality: if J is an (integrable) endomorphism with $J^2 = 0$ then its dual $J^* : \Gamma(T^*M) \rightarrow \Gamma(T^*M)$, given by $J^* \alpha := \alpha \circ J$ for $\alpha \in \Gamma(T^*M)$, is (integrable) endomorphism with $(J^*)^2 = 0$.

2. Basic properties of tangent conjugate connections

Let ∇ be a linear connection on the almost tangent manifold (M, J) and define the *tangent conjugate connection* of ∇ by:

$$\nabla^{(J)} := \nabla - J \circ \nabla J. \quad (2.1)$$

Remark that $\nabla^{(J)}$ coincides with ∇ if and only if $\nabla J \subseteq \ker J = \text{Im} J$ which means the inclusion $\nabla(\Gamma(TM) \times \ker J) \subseteq \ker J = \text{Im} J$, in particular if ∇ is a J -linear connection; for another case see i) of Proposition 2.3. For any $X, Y \in \Gamma(TM)$ we get:

$$\nabla_X^{(J)} Y = \nabla_X Y - J(\nabla_X JY). \quad (2.2)$$

A first set of properties for this linear connection are given by:

2.1. Proposition. *The tangent conjugate connection $\nabla^{(J)}$ satisfies:*

- i) $\nabla^{(J)} J = \nabla J$, which means that ∇ and $\nabla^{(J)}$ are simultaneous J -linear connections or not;
- ii) $\nabla^{2(J)} := (\nabla^{(J)})^{(J)} = 2\nabla^{(J)} - \nabla$; more generally $\nabla^{n(J)} = n\nabla^{(J)} - (n-1)\nabla$ for $n \in \mathbb{N}^*$;
- iii) its torsion is $T_{\nabla^{(J)}} = T_{\nabla} - J \circ d^{\nabla} J$ where d^{∇} is the exterior covariant derivative induced by ∇ , namely $(d^{\nabla} J)(X, Y) := (\nabla_X J)Y - (\nabla_Y J)X$;
- iv) its curvature is

$$\begin{aligned} R_{\nabla^{(J)}}(X, Y, Z) &= R_{\nabla}(X, Y, Z) - \nabla_X J(\nabla_Y JZ) + \nabla_Y J(\nabla_X JZ) - \\ &\quad - J[\nabla_X J(\nabla_Y Z) - \nabla_Y J(\nabla_X Z) - \nabla_{[X, Y]} JZ]. \end{aligned} \quad (2.3)$$

In particular:

$$R_{\nabla^{(J)}}(X, Y, JZ) = R_{\nabla}(X, Y, JZ) - J[\nabla_X J(\nabla_Y JZ) - \nabla_Y J(\nabla_X JZ)]. \quad (2.4)$$

Proof The general part of ii) follows by induction while for iii) a direct calculus yields $T_{\nabla^{(J)}}(X, Y) = T_{\nabla}(X, Y) - J(\nabla_X JY - \nabla_Y JX)$. \square

Let $f : M \rightarrow M$ be a *tangentomorphism*, that is an automorphism of the G -structure defined by J :

$$f_* \circ J = J \circ f_*. \quad (2.5)$$

Recall that f is an *affine transformation* for ∇ if for any $X, Y \in \Gamma(TM)$:

$$f_*(\nabla_X Y) = \nabla_{f_* X} f_* Y. \quad (2.6)$$

These notions are connected by:

2.2. Proposition. *If the tangentomorphism f is an affine transformation for ∇ then f is also affine transformation for $\nabla^{(J)}$.*

Proof We have:

$$\begin{aligned} f_*(\nabla_X^{(J)}Y) &= f_*(\nabla_X Y) - (f_* \circ J)(\nabla_X JY) = \nabla_{f_*X} f_*Y - J(f_*(\nabla_X JY)) = \\ &= \nabla_{f_*X} f_*Y - J((\nabla_{f_*X} f_*(JY))) = \nabla_{f_*X} f_*Y - J((\nabla_{f_*X} J(f_*Y))) = \nabla_{f_*X}^{(J)} f_*Y \end{aligned}$$

which yields the conclusion. \square

A second class of properties for the tangent conjugate connection is provided by:

2.3. Proposition. *i) If J is ∇ -recurrent i.e. $\nabla J = \eta \otimes J$ for η a 1-form, then $\nabla^{(J)} = \nabla$.
ii) If ∇ is symmetric and $\nabla J = \eta \otimes I$ then $\nabla^{(J)} = \nabla - \eta \otimes J$ and $\nabla^{(J)}$ is a quarter-symmetric connection.*

Proof i) In this case we have $J \circ \nabla J = 0$.

ii) Recall after [1, p. 122] that the quarter-symmetry means the existence of a 1-form π and a tensor field F of (1,1)-type such that $T_{\nabla^{(J)}} = F \wedge \pi := F \otimes \pi - \pi \otimes F$. From Proposition 2.1 we have $T_{\nabla^{(J)}}(X, Y) = T_{\nabla}(X, Y) - \eta(X)JY + \eta(Y)JX$, and the hypothesis $T_{\nabla} = 0$ yields the previous equation with $F = J$ and $\pi = \eta$. \square

2.4. Example. Let N be a smooth n -dimensional manifold and $M = TN$ its tangent bundle; hence $m = 2n$. Let $\{x^i; 1 \leq i \leq n\}$ be a local system of coordinates on N and consider its lift to M given by $\{x^i, y^i; 1 \leq i \leq n\}$ with y^i the coordinates on the fibres of TN . The canonical almost tangent structure J of M has the local expression (1.3) and it is integrable. Fix a general linear connection ∇ on M with local Christoffel symbols Γ as follows:

$$\left\{ \begin{array}{l} \nabla \frac{\partial}{\partial x^i} \frac{\partial}{\partial x^j} = \Gamma_{ij}^{(1)k} \frac{\partial}{\partial x^k} + \Gamma_{ij}^{(2)k} \frac{\partial}{\partial y^k} \\ \nabla \frac{\partial}{\partial x^i} \frac{\partial}{\partial y^j} = \Gamma_{ij}^{(3)k} \frac{\partial}{\partial x^k} + \Gamma_{ij}^{(4)k} \frac{\partial}{\partial y^k} \\ \nabla \frac{\partial}{\partial y^i} \frac{\partial}{\partial x^j} = \Gamma_{ij}^{(5)k} \frac{\partial}{\partial x^k} + \Gamma_{ij}^{(6)k} \frac{\partial}{\partial y^k} \\ \nabla \frac{\partial}{\partial y^i} \frac{\partial}{\partial y^j} = \Gamma_{ij}^{(7)k} \frac{\partial}{\partial x^k} + \Gamma_{ij}^{(8)k} \frac{\partial}{\partial y^k} \end{array} \right. \quad (2.7)$$

Then its tangent conjugate connection has the expression:

$$\left\{ \begin{array}{l} \nabla \frac{\partial}{\partial x^i} \frac{\partial}{\partial x^j} = \Gamma_{ij}^{(1)k} \frac{\partial}{\partial x^k} + \left(\Gamma_{ij}^{(2)k} - \Gamma_{ij}^{(3)k} \right) \frac{\partial}{\partial y^k} \\ \nabla \frac{\partial}{\partial x^i} \frac{\partial}{\partial y^j} = \Gamma_{ij}^{(3)k} \frac{\partial}{\partial x^k} + \Gamma_{ij}^{(4)k} \frac{\partial}{\partial y^k} \\ \nabla \frac{\partial}{\partial y^i} \frac{\partial}{\partial x^j} = \Gamma_{ij}^{(5)k} \frac{\partial}{\partial x^k} + \left(\Gamma_{ij}^{(6)k} - \Gamma_{ij}^{(7)k} \right) \frac{\partial}{\partial y^k} \\ \nabla \frac{\partial}{\partial y^i} \frac{\partial}{\partial y^j} = \Gamma_{ij}^{(7)k} \frac{\partial}{\partial x^k} + \Gamma_{ij}^{(8)k} \frac{\partial}{\partial y^k} \end{array} \right. \quad (2.8)$$

A special case is important in applications: the initial connection ∇ is called *distinguished* or *d-connection* if it preserves the linear structure of the fibres of M which means that:

$$\Gamma^{(2)} = \Gamma^{(3)} = \Gamma^{(6)} = \Gamma^{(7)} = 0. \quad (2.9)$$

It results that ∇ is a J -connection and then its tangent conjugate connection is $\nabla^{(J)} = \nabla$.

3. The structural and the virtual tensor fields

Remark that the tangent conjugate connection $\nabla^{(J)}$ of ∇ can be written in another form as:

$$\nabla^{(J)} = \nabla + C_{\nabla}^J - B_{\nabla}^J \quad (3.1)$$

where:

$$\begin{cases} C_{\nabla}^J(X, Y) := \frac{1}{2}[(\nabla_{JX}J)Y + (\nabla_XJ)JY] \\ B_{\nabla}^J(X, Y) := \frac{1}{2}[(\nabla_{JX}J)Y - (\nabla_XJ)JY]. \end{cases} \quad (3.2)$$

which we call respectively, the *structural* and the *virtual tensor field* of ∇ . We obtain also the following expressions for them:

$$\begin{cases} C_{\nabla}^J(X, Y) = \frac{1}{2}[\nabla_{JX}JY - J(\nabla_{JX}Y + \nabla_XJY)] \\ B_{\nabla}^J(X, Y) = \frac{1}{2}[\nabla_{JX}JY - J(\nabla_{JX}Y - \nabla_XJY)]. \end{cases} \quad (3.3)$$

We notice that they satisfy the following properties:

$$\begin{cases} C_{\nabla}^J(JX, Y) = C_{\nabla}^J(X, JY) = -\frac{1}{2}J(\nabla_{JX}JY); & C_{\nabla}^J(JX, JY) = 0 \\ B_{\nabla}^J(JX, Y) = -B_{\nabla}^J(X, JY) = \frac{1}{2}J(\nabla_{JX}JY); & B_{\nabla}^J(JX, JY) = 0 \\ C_{\nabla}^J(JX, Y) = -B_{\nabla}^J(JX, Y) \end{cases} \quad (3.4)$$

and the skew-symmetry (3.4₂) means that $B_{\nabla}^J(J, \cdot)$ is a vectorial 2-form. Another important property is that these tensor fields are invariant with respect to J -conjugation of linear connections:

$$C_{\nabla^{(J)}}^J = C_{\nabla}^J; \quad B_{\nabla^{(J)}}^J = B_{\nabla}^J. \quad (3.5)$$

With respect to the invariance of these associated tensor fields under projective changes we get that only C^J is invariant:

3.1. Proposition. *Let ∇ and ∇' be two linear projectively equivalent connections:*

$$\nabla' = \nabla + \eta \otimes I + I \otimes \eta \quad (3.6)$$

for η a 1-form. Then $C_{\nabla'}^J = C_{\nabla}^J$ and $B_{\nabla'}^J = B_{\nabla}^J + J \otimes (\eta \circ J)$ while the tangent conjugate connection $\nabla'^{(J)}$ of ∇' satisfies:

$$\nabla'^{(J)} = \nabla^{(J)} + \eta \otimes I + I \otimes \eta - J \otimes (\eta \circ J) \quad (3.7)$$

and so it is not invariant under projective equivalence.

Proof Follows from a direct computation. \square

4. Invariant distributions

Let $\mathcal{D} \subset TM$ be a fixed distribution considered as a vector subbundle of TM . As usually, we denote by $\Gamma(\mathcal{D})$ its $C^\infty(M)$ -module of sections.

4.1. Definition. i) \mathcal{D} is called *J-invariant* if $X \in \Gamma(\mathcal{D})$ implies $JX \in \Gamma(\mathcal{D})$.
ii) The linear connection ∇ *restricts to \mathcal{D}* if $Y \in \Gamma(\mathcal{D})$ implies $\nabla_X Y \in \Gamma(\mathcal{D})$ for any $X \in \Gamma(TM)$.

4.2. Example. The distribution $\mathcal{D}_J = \ker J = \text{Im} J$ is *J-invariant*.

If ∇ restricts to \mathcal{D} then it may be considered as a connection in the vector bundle \mathcal{D} . From this fact, a connection which restricts to \mathcal{D} is called sometimes *adapted to \mathcal{D}* .

4.3. Proposition. *If the distribution \mathcal{D} is J -invariant and the linear connection ∇ restricts to \mathcal{D} then $\nabla^{(J)}$ also restricts to \mathcal{D} .*

Proof Fix $Y \in \Gamma(\mathcal{D})$. Then $JY \in \Gamma(\mathcal{D})$ and for any $X \in \Gamma(TM)$ we have $\nabla_X Y, \nabla_X JY \in \Gamma(\mathcal{D})$. Therefore, $J(\nabla_X JY) \in \Gamma(\mathcal{D})$ and so $\nabla_X^{(J)} Y = \nabla_X Y - J(\nabla_X JY) \in \Gamma(\mathcal{D})$. \square

4.4. Example. Returning to Example 4.2 we have that $\nabla_X = \nabla_X^{(J)}$ on $\mathcal{D}_J = \ker J = \text{Im} J$.

A more general notion like restricting to a distribution is that of geodesically invariance [4, p. 118]. The distribution \mathcal{D} is ∇ -geodesically invariant if for every geodesic $\gamma : [a, b] \rightarrow M$ of ∇ with $\dot{\gamma}(a) \in \mathcal{D}_{\gamma(a)}$ it follows $\dot{\gamma}(t) \in \mathcal{D}_{\gamma(t)}$ for any $t \in [a, b]$. The cited book gives a necessary and sufficient condition for a distribution \mathcal{D} to be ∇ -geodesically invariant: for any $X, Y \in \Gamma(\mathcal{D})$, the symmetric product $\langle X : Y \rangle_\nabla := \nabla_X Y + \nabla_Y X$ to belong to $\Gamma(\mathcal{D})$ or equivalently, for any $X \in \Gamma(\mathcal{D})$ to have $\nabla_X X \in \Gamma(\mathcal{D})$.

A direct computation gives:

$$\langle \cdot : \cdot \rangle_{\nabla^{(J)}} = \langle \cdot : \cdot \rangle_\nabla - J \circ d^\nabla J \quad (4.1)$$

and then the ∇ -geodesically invariance and $\nabla^{(J)}$ -geodesically invariance for \mathcal{D} coincides if and only if $J \circ d^\nabla J$ is zero on $\mathcal{D} \times \mathcal{D}$. In particular, \mathcal{D}_J is ∇ -geodesically invariant if and only if is $\nabla^{(J)}$ -geodesically invariant.

5. Affine combination of tangent conjugate connections

In what follows we shall see what happens to the tangent conjugate connection for families of almost tangent structures. Let J_1, J_2 be two almost tangent structures; conditions for their simultaneous integrability are given in [13]-[14]. Then for any $a, b \in \mathbb{R}$ the tensor field $J_{ab} := aJ_1 + bJ_2$ is an almost tangent structure if and only if $J_1 J_2 = -J_2 J_1$. Then its tangent conjugate connection is given by:

$$\nabla_X^{(J_{ab})} Y = a^2 \nabla_X^{(J_1)} Y + b^2 \nabla_X^{(J_2)} Y + (1 - a^2 - b^2) \nabla_X Y - ab[J_1(\nabla_X J_2 Y) + J_2(\nabla_X J_1 Y)]. \quad (5.1)$$

5.1. Proposition. *Let ∇ be a linear connection and J_1 and J_2 two anti-commuting almost tangent structures. If (∇, J_1, J_2) is a mixed-recurrent structure i.e. $\nabla J_i = \eta \otimes J_j$ for $i \neq j$ then ∇ is the average of the two tangent conjugate connections:*

$$\nabla = \frac{1}{2}[\nabla^{(J_1)} + \nabla^{(J_2)}] \quad (5.2)$$

and $\nabla^{(J_{ab})}$ is an affine combination of them:

$$\nabla^{(J_{ab})} = \frac{1 + a^2 - b^2}{2} \nabla^{(J_1)} + \frac{1 - a^2 + b^2}{2} \nabla^{(J_2)}. \quad (5.3)$$

Proof Applying J_i to $\nabla_X J_i Y - J_i(\nabla_X Y) = \eta(X) J_j Y$ with $i \neq j$ and the anti-commuting hypothesis we obtain:

$$J_1(\nabla_X J_1 Y) = -J_2(\nabla_X J_2 Y). \quad (5.4)$$

Summing the expression of the tangent conjugate connections we get (5.2) and from a previous computation, the relation (5.3). \square

6. Exponential tangent conjugate connections

For θ a real number we define the *exponential tangent conjugate connection* of ∇ as:

$$\nabla^{(J,\theta)} := \nabla - \exp(-\theta J) \circ \nabla \circ \exp(\theta J) \quad (6.1)$$

where $\exp(\pm\theta J) := \cos(\theta) \cdot I \pm \sin(\theta) \cdot J$. Explicitly we get:

$$\nabla^{(J,\theta)} = \sin^2(\theta)\nabla - \frac{1}{2}\sin(2\theta)\nabla J + \sin^2(\theta)J \circ \nabla J = 2\sin^2(\theta)\nabla - \frac{1}{2}\sin(2\theta)\nabla J - \sin^2(\theta)\nabla^{(J)} \quad (6.2)$$

and then:

$$\nabla^{(J,\theta)} J = \sin^2(\theta)\nabla J + \frac{1}{2}\sin(2\theta)J \circ \nabla J. \quad (6.3)$$

It follows:

6.1. Proposition. *Let ∇ be a symmetric linear connection.*

i) *If J is ∇ -recurrent with η the 1-form of recurrence then:*

$$\nabla^{(J,\theta)} = \sin^2(\theta)\nabla - \frac{1}{2}\sin(2\theta) \cdot \eta \otimes J \quad (6.4)$$

and $\nabla^{(J,\theta)}$ is a quarter-symmetric connection.

ii) *If $\nabla J = \eta \otimes I$ then:*

$$\nabla^{(J,\theta)} = \sin^2(\theta)\nabla - \sin(\theta) \cdot \eta \otimes \exp(-\theta J) \quad (6.5)$$

and:

$$T_{\nabla^{(J,\theta)}} = \sin(\theta) \otimes \exp(-\theta J) \wedge \eta. \quad (6.6)$$

Proof i) Follows from the fact that the hypothesis implies $J \circ \nabla J = 0$. The quarter-symmetry elements are $F = J$ and $\pi = \sin(\theta) \cos(\theta) \cdot \eta$.

ii) From $\cos(\theta) \cdot \eta \otimes I - \sin(\theta) \cdot \eta \otimes J = \eta \otimes \exp(-\theta J)$ we get:

$$T_{\nabla^{(J,\theta)}} = -\sin(\theta) \cdot [\eta \otimes \exp(-\theta J) - \exp(-\theta J) \otimes \eta]. \quad \square$$

7. Generalized tangent conjugate connections

In this section we present a natural generalization of the tangent conjugate connection.

7.1. Definition. A *generalized tangent conjugate connection* of ∇ is:

$$\nabla^{(J,C)} = \nabla^{(J)} + C \quad (7.1)$$

with $C \in T_2^1(M)$ an arbitrary (1,2)-tensor field.

Let us search for tensor fields C such that the duality $(\nabla^{(J,C)})^{(J,C)} = 2\nabla^{(J,C)} - \nabla$ holds as is given by Proposition 2.1. It results that we are interested in finding solutions C to the equation:

$$J(C(X, JY)) = 2C(X, Y) \quad (7.2)$$

for all $X, Y \in \Gamma(TM)$ and let us remark that: i) $C_0 = 0$ is a particular solution of (7.2);

ii) applying J to (7.2) gives that $ImC \subseteq \ker J = ImJ$. Then returning to (7.2) it follows from the left-hand-side that C_0 is the unique solution of (7.2).

Also, we have:

$$\nabla^{(J,C)} J = \nabla^{(J)} J + C(\cdot, J\cdot) - J \circ C \quad (7.3)$$

and then:

i) $\nabla^{(J,C)} J = \nabla J$ as in i) of Proposition 2.1 if and only if: $C(\cdot, J\cdot) = J \circ C(\cdot, \cdot)$,

ii) $\nabla^{(J,C)}$ is a J -linear connection if and only if:

$$\nabla J + C(\cdot, J\cdot) = J \circ C. \quad (7.4)$$

Acknowledgement. The first author acknowledges the support by the research grant PN-II-ID-PCE-2011-3-0921.

References

- [1] A. M. Blaga; M. Crasmareanu, *The geometry of complex conjugate connections*, Hacet. J. Math. Stat., 41(2012), no. 1, 119-126. MR2976917
- [2] A. M. Blaga; M. Crasmareanu, *The geometry of product conjugate connections*, An. Stiint. Univ. Al. I. Cuza Iași Mat. (N. S.), 59(2013), no. 1, 73-84. MR3098381
- [3] F. Brickell; R. S. Clark, *Integrable almost tangent structures*, J. Diff. Geom., 9(1974), 557-563. MR0348666 (50 #1163)
- [4] F. Bullo; A. D. Lewis, *Geometric control of mechanical systems. Modeling, analysis, and design for simple mechanical control systems*, Texts in Applied Mathematics, 49, Springer-Verlag, New York, 2005. MR2099139 (2005h:70030)
- [5] R. S. Clark; M. Bruckheimer, *Sur les structures presque tangents*, C. R. A. S. Paris, 251(1960), 627-629. MR0115181 (22 #5983)
- [6] R. S. Clark; D. S. Goel, *On the geometry of an almost tangent manifold*, Tensor, 24(1972), 243-252. MR0326613 (48 #4956)
- [7] M. Crampin, *Defining Euler-Lagrange fields in terms of almost tangent structures*, Phys. Lett. A, 95(1983), no. 9, 466-468. MR0708702 (84k:58072)
- [8] M. Crampin; G. Thompson, *Affine bundles and integrable almost tangent structures*, Math. Proc. Camb. Phil. Soc., 98(1985), 61-71. MR0789719 (86g:53039)
- [9] V. Cruceanu, *Connexions compatibles avec certaines structures sur un fibré vectoriel banachique*, Czechoslovak Math. J., 24(99)(1974), 126-142. MR0353356 (50 #5840)
- [10] H. A. Eliopoulos, *Structures presque tangents sur les variétés différentiables*, C. R. A. S. Paris, 255(1962), 1563-1565. MR0142078 (25 #5472)
- [11] D. S. Goel, *Selfadjoint metrics on almost tangent manifolds whose Riemannian connection is almost tangent*, Canad. Math. Bull., 17(1974/75), no. 5, 671-674. MR0383288 (52 #4169)
- [12] J. Grifone, *Structure presque-tangente et connexions*, I, II. Ann. Inst. Fourier (Grenoble), 22(1972), no. 1, 3, 287-334, 291-338. MR0336636 (49 #1409), MR0341361 (49 #6112)
- [13] V. Kubát, *Simultaneous integrability of two J -related almost tangent structures*, Comment. Math. Univ. Carolin. 20(1979), no. 3, 461-473. MR0550448 (80m:53034)
- [14] V. Kubát, *On simultaneous integrability of two commuting almost tangent structures*, Comm. Math. Univ. Carolinae, 22(1981), no. 1, 149-160. MR0609943 (82e:53052)
- [15] M. de León; P. R. Rodrigues, *Methods of differential geometry in analytical mechanics*, North-Holland Mathematics Studies, 158, North-Holland Publishing Co., Amsterdam, 1989. MR1021489 (91c:58041)
- [16] G. Thompson; U. Schwardmann, *Almost tangent and cotangent structures in the large*, Trans. Amer. Math. Soc., 327(1991), no. 1, 313-328. MR1012509 (91m:53029)
- [17] I. Vaisman, *Lagrange geometry on tangent manifolds*, Int. J. Math. Math. Sci., 51(2003), 3241-3266. MR2018588 (2004k:53116)
- [18] K. Yano; E. T. Davies, *Differential geometry on almost tangent manifolds*, Ann. Mat. Pura Appl. (4), 103(1975), 131-160. MR0390958 (52 #11781)