



Namık Kemal Üniversitesi
Tekirdağ Ziraat Fakültesi Dergisi
Journal of Tekirdag Agricultural Faculty

An International Journal of all Subjects of Agriculture

Sahibi / Owner

Namık Kemal Üniversitesi Ziraat Fakültesi Adına
On Behalf of Namık Kemal University Agricultural Faculty

Prof.Dr. Ahmet İSTANBULLUOĞLU
Dekan / Dean

Editörler Kurulu / Editorial Board

Başkan / Editor in Chief

Prof.Dr. Selçuk ALBUT
Ziraat Fakültesi Biyosistem Mühendisliği Bölümü
Department Biosystem Engineering, Agricultural Faculty
salbut@nku.edu.tr

Üyeler / Members

Prof.Dr. M. İhsan SOYSAL	Zootekni / Animal Science
Prof.Dr. Bülent EKER	Biyosistem Mühendisliği / Biosystem Engineering
Prof.Dr. Servet VARIŞ	Bahçe Bitkileri / Horticulture
Prof.Dr. Aslı KORKUT	Peyzaj Mimarılığı / Landscape Architecture
Prof.Dr. Temel GENÇTAN	Tarla Bitkileri / Field Crops
Prof.Dr. Müjgan KIVAN	Bitki Koruma / Plant Protection
Prof.Dr. Şefik KURULTAY	Gıda Mühendisliği / Food Engineering
Prof.Dr. Aydın ADİLOĞLU	Toprak Bilimi ve Bitki Besleme / Soil Science and Plant Nutrition
Prof.Dr. Fatih KONUKCU	Biyosistem Mühendisliği / Biosystem Engineering
Doç.Dr. Ömer AZABAĞAOĞLU	Tarım Ekonomisi / Agricultural Economics
Yrd.Doç.Dr. Devrim OSKAY	Tarımsal Biyoteknoloji / Agricultural Biotechnology
Yrd.Doç.Dr. Harun HURMA	Tarım Ekonomisi / Agricultural Economics
Yrd.Doç.Dr. M. Recai DURGUT	Biyosistem Mühendisliği / Biosystem Engineering


İndeksler / Indexing and abstracting



CABI tarafından full-text olarak indekslenmektedir / Included in **CABI**



DOAJ tarafından full-text olarak indekslenmektedir / Included in **DOAJ**



EBSCO tarafından full-text olarak indekslenmektedir / Included in **EBSCO**



FAO AGRIS Veri Tabanında İndekslenmektedir / Indexed by **FAO AGRIS Database**



INDEX COPERNICUS tarafından full-text olarak indekslenmektedir / Included in **INDEX COPERNICUS**



TUBİTAK-ULAKBİM Tarım, Veteriner ve Biyoloji Bilimleri Veri Tabanı (TVBBVT) Tarafından taranmaktadır / Indexed by **TUBİTAK-ULAKBİM** Agriculture, Veterinary and Biological Sciences Database

Yazışma Adresi / Corresponding Address

Tekirdağ Ziraat Fakültesi Dergisi NKÜ Ziraat Fakültesi 59030 TEKİRDAĞ

E-mail: ziratdergi@nku.edu.tr

Web adresi: <http://jotaf.nku.edu.tr>

Tel: +90 282 250 20 07

ISSN: 1302-7050

Danışmanlar Kurulu /Advisory Board

Bahçe Bitkileri / Horticulture

Prof.Dr. Kazım ABAK	Çukurova Üniv. Ziraat Fak. Adana
Prof.Dr. Y.Sabit AĞAOĞLU	Ankara Üniv. Ziraat Fak. Ankara
Prof.Dr. Jim HANCOCK	Michigan State Univ. USA
Prof.Dr. Mustafa PEKMEZCİ	Akdeniz Üniv. Ziraat Fak. Antalya

Bitki Koruma / Plant Protection

Prof.Dr. Mithat DOĞANLAR	Mustafa Kemal Üniv. Ziraat Fak. Hatay
Prof.Dr. Timur DÖKEN	Adnan Menderes Üniv. Ziraat Fak. Aydın
Prof.Dr. Ivanka LECHAVA	Agricultural Univ. Plovdiv-Bulgaria
Dr. Emil POCSAI	Plant Protection Soil Cons. Service Velençe-Hungary

Gıda Mühendisliği / Food Engineering

Prof.Dr. Yaşar HIŞIL	Ege Üniv. Mühendislik Fak. İzmir
Prof.Dr. Fevzi KELEŞ	Atatürk Üniv. Ziraat Fak. Erzurum
Prof.Dr. Atilla YETİŞEMİYEN	Ankara Üniv. Ziraat Fak. Ankara
Prof.Dr. Zhelyazko SIMOV	University of Food Technologies Bulgaria

Peyzaj Mimarlığı / Landscape Architecture

Prof.Dr. Mükerrerem ARSLAN	Ankara Üniv. Ziraat Fak. Ankara
Prof.Dr. Bülent ÖZKAN	Ege Üniv. Ziraat Fak. İzmir
Prof.Dr. Güniz A. KESİM	Düzce Üniv. Orman Fak. Düzce
Prof.Dr. Genoveva TZOLOVA	University of Forestry Bulgaria

Tarla Bitkileri / Field Crops

Prof.Dr. Esvet AÇIKGÖZ	Uludağ Üniv. Ziraat Fak. Bursa
Prof.Dr. Özer KOLSARICI	Ankara Üniv. Ziraat Fak. Ankara
Dr. Nurettin TAHSİN	Agric. Univ. Plovdiv Bulgaria
Prof.Dr. Murat ÖZGEN	Ankara Üniv. Ziraat Fak. Ankara
Doç. Dr. Christina YANCHEVA	Agric. Univ. Plovdiv Bulgaria

Tarım Ekonomisi / Agricultural Economics

Prof.Dr. Faruk EMEKSİZ	Çukurova Üniv. Ziraat Fak. Adana
Prof.Dr. Hasan VURAL	Uludağ Üniv. Ziraat Fak. Bursa
Prof.Dr. Gamze SANER	Ege Üniv. Ziraat Fak. İzmir
Dr. Alberto POMBO	El Colegio de la Frontera Norte, Meksika

Tarım Makineleri / Agricultural Machinery

Prof.Dr. Thefanis GEMTOS	Aristotle Univ. Greece
Prof.Dr. Simon BLACKMORE	The Royal Vet.&Agr. Univ. Denmark
Prof.Dr. Hamdi BİLGİN	Ege Üniv. Ziraat Fak. İzmir
Prof.Dr. Ali İhsan ACAR	Ankara Üniv. Ziraat Fak. Ankara

Tarımsal Yapılar ve Sulama / Farm Structures and Irrigation

Prof.Dr. Ömer ANAPALI	Atatürk Üniv. Ziraat Fak. Erzurum
Prof.Dr. Christos BABAJIMOPOULOS	Aristotle Univ. Greece
Dr. Arie NADLER	Ministry Agr. ARO Israel

Toprak / Soil Science

Prof.Dr. Sait GEZGİN	Selçuk Üniv. Ziraat Fak. Konya
Prof.Dr. Selim KAPUR	Çukurova Üniv. Ziraat Fak. Adana
Prof.Dr. Metin TURAN	Atatürk Üniv. Ziraat Fak. Erzurum
Doç. Dr. Pasquale STEDUTO	FAO Water Division Italy

Zootekni / Animal Science

Prof.Dr. Andreas GEORGIDUS	Aristotle Univ. Greece
Prof.Dr. Ignacy MISZTAL	Breeding and Genetics University of Georgia USA
Prof.Dr. Kristaq KUME	Center for Agricultural Technology Transfer Albania
Dr. Brian KINGHORN	The Ins. of Genetics and Bioinformatics Univ. of New England Australia
Prof.Dr. Ivan STANKOV	Trakia Univ. Dept. Of Animal Sci. Bulgaria
Prof.Dr. Nihat ÖZEN	Akdeniz Üniv. Ziraat Fak. Antalya
Prof.Dr. Jozsef RATKY	Res. Ins. Animal Breed. and Nut. Hungary
Prof.Dr. Naci TÜZEMEN	Atatürk Üniv. Ziraat Fak. Erzurum

İÇİNDEKİLER / CONTENTS

T. Yılmaz, D. Gökçe, F. Şavklı, S. Çeşmeci Engellilerin Üniversite Kampüslerinde Ortak Mekanları Kullanabilmeleri Üzerine Bir Araştırma: Akdeniz Üniversitesi Olbia Kültür Merkezi Örneği A Study On Young Disabled People's Use Of Common Areas in The University Campuses Example Of Olbia Culture Center in Akdeniz University	1-10
K. Demirel, Y. Kavdır Toprak Altına Serilen Su Tutma Bariyer Uygulamaları Toprak Profilindeki Tuz İçeriğini Arttırır mı? Does Application of Water Retention Barrier to Soil Increase Salt Content Within Soil Profile?	11-21
S. Çınar, R. Hatipoğlu, A. Aktaş Çukurova Taban Kesimi Meralarında Yabancı Ot Mücadelesi Üzerine Bir Araştırma Research On Weed Control in Pastures Under Lowland Conditions Of Cukurova	22-26
A. Delice, N. Ekinci, F. F. Özdüven, E. Gür Lapseki'de Yetiştirilen 0900 Ziraat Kiraz Çeşidinin Kalite Özellikleri Ve Ekolojik Faktörler Determinations of Factors That Effect on Quality Properties of 0900 Ziraat Cherry Variety in Lapseki	27-34
M. F. Baran, P. Ülger, B. Kayışoğlu Kanola Hasadında Kullanılan Tablanın Hasat Kayıpları Üzerine Etkisi The Effect of Canola Harvest Header Used in Canola Harvesting on Harvest Losses.....	35-44
M. M. Özgüven Kapalı Alanlarda Kullanılan Bazı Hasat Sonrası Tarım Makinalarının Gürültü Haritalarının İncelenmesi Investigation of Noise Maps for Some Post-Harvest Agricultural Machinery Used Indoor Spaces	45-53
A. Semerci Evaluation of The Changes in The Cost Factors of Sunflower Production in Turkey Ayçiçeği Üretiminde Maliyet Faktörlerindeki Değişimin İncelenmesi (Trakya Bölgesi/Türkiye Örneği)	54-61
F. Coşkun, M. Arıcı, G. Çelikyurt, M. Gülcü Farklı Yöntemler Kullanılarak Üretilen Hardalilerin Bazı Özelliklerinde Depolama Sonunda Meydana Gelen Değişmeler Changes occuring at the end of storage in some properties of hardaliye produced by using different methods	62-67
D. Boyraz, H. Sarı Tekirdağ Değirmenaltı-Muratlı Kavşağı Çevre Yolunu Oluşturan Katenadaki Toprakların Fiziksel Ve Zemin Özelliklerinin Değerlendirilmesi Evaluating the Physical and Ground Conditions of The Soils in The Catena Which Forms Tekirdağ Değirmenaltı-Muratlı Intersection Ringroad	68-78
B. E. Öztürk, B. Kaptan, O. Şimşek Determination of Some Heavy Metals Level in Kashar Cheese Produced in Thrace Region Trakya Bölgesinde Üretilen Kaşar Peynirlerinin Bazı Ağır Metal Düzeylerinin Belirlenmesi	79-83
D. Katar, Y. Arslan, İ. Subaşı Ankara Ekolojik Koşullarında Farklı Ekim Zamanlarının Ketencik (Camelina Sativa (L.) Crantz) Bitkisinin Yağ Oranı Ve Bileşimi Üzerine Olan Etkisinin Belirlenmesi Determination of Effect of Different Sowing Dates on Oil Content and Fatty Acid Composition in Camelina (Camelina sativa (L.) Crantz) under Ankara Ecological Condition	84-90
Y. Mutlu, F. Koc, M. L. Ozduven, L. Coskuntuna Effects of Inoculant Preparation Time and Doses on Fermentation and Aerobic Stability Characteristics of the Second Crop Maize Silages İnokulant Hazırlama Süresi ve Dozunun İkinci Ürün Mısır Silajlarının Fermantasyon ve Aerobik Stabilitate Özellikleri Üzerine Etkileri	91-97
G. Güngör, K. Benli, H. Güngör Marmara Denizi'nde Deniz Ürünleri Pazarlaması: İstanbul İli Sahil Şeridi Örneği Marketing Seafood Products in Marmara Sea: A Case Study Along The Coastal Strip in İstanbul Province	98-108
J. M. Kıyıcı, N. Tüzemen Buzağuların Kovadan Süt İçmeyi Öğrenme Davranışlarının Karşılaştırılması Comparison of Learning Behaviour of Calves Drink Milk From The Bucket	109-114

Determination of Some Heavy Metals Level in Kashar Cheese Produced in Thrace Region*

B. E. Öztürk¹

B. Kaptan¹

O. Şimşek¹

¹Department of Food Engineering, Agricultural Faculty, Namık Kemal University, Tekirdag, Turkey

In this research, 50 samples of kashar cheese produced and sold in Trakya (Thrace) region, were studied for analysis of heavy metals as Lead (Pb), tin (Sn), copper (Cu) and mercury (Hg). Heavy metal contents of samples were determined by Atomic Absorption Spectrophotometer (AAS) after having been burned in a microwave oven. As a result of this study, average heavy metal contents of 50 samples found as; lead (Pb):0.0600 ppm, tin (Sn): 0.0366 ppm, copper (Cu): 0.5036 ppm, mercury (Hg): 0.0214 ppm. The obtained results show that mean concentrations of copper and tin in samples are below average levels whereas some results in lead and mercury are above the average levels.

Keywords: Kashar cheese, lead, tin, copper, mercury

*This study is a part of Bengi Eren Öztürk's master thesis

Trakya Bölgesinde Üretilen Kaşar Peynirlerinin Bazı Ağır Metal Düzeylerinin Belirlenmesi*

Bu çalışmada Trakya Bölgesinde üretilen kaşar peynirlerinin kurşun (Pb), kalay (Sn), bakır (Cu) ve civa (Hg) içeriklerinin belirlenmesi amaçlanmıştır. Bu amaçla bölgeden 50 adet kaşar peyniri örneği toplanmıştır. Örnekler mikrodalga fırında yakıldıktan sonra ağır metal içerikleri Atomic Absorption Spectrophotometer (AAS) cihazı kullanılarak belirlenmiştir. Analiz sonuçlarına göre kaşar peyniri örneklerinde ortalama olarak Pb:0.0600 ppm, Sn: 0.0366 ppm, Cu: 0.5036 ppm ve Hg: 0.0214 ppm düzeylerinde belirlenmiştir. Yapılan analizler sonucunda elde edilen değerler örneklerin ortalama Cu ve Sn içerikleri belirlenen limit değerlerin altında, Pb ve Hg içeriklerinin ise limit değerlerin üstünde olduğu göstermektedir.

Anahtar kelimeler: Kaşar peyniri, kurşun, kalay, bakır, civa

*Bu çalışma Bengi Eren Öztürk'ün yüksek lisans tezinden türetilmiştir

Introduction

Because of their long term toxicological effects, inorganic or aggregated forms of chemical substances represent a severe risk in food and hence feeding. Heavy metals are widespread in environment and it is reported that heavy metal levels in milk and milk products in industrially developed areas is much higher than those in rural areas since metal contamination in food increases rapidly with the increase of environmental pollution due to industrial waste products. Even if the concentration of heavy metals in a living organism is low, it is proved to be more toxic or poisonous than any other metal (John & Howard, 1996). Heavy metals have no beneficial effects, they can cause health problems even in very low concentrations; furthermore toxic effects can be seen when the limit of heavy metals are exceeded (FAO/WHO, 1993). Because of this fact, the maximum legal concentrations of

heavy metals in food and beverages consumed on regular base have been determined and are checked regularly by the legal entities (Şimşek et al., 2000).

Especially, milk and its products are contaminated with heavy metals such as lead, cadmium, antimony, arsenic, copper, mercury, manganese, zinc and tin. Due to the contamination, the metal concentration of consumptive (consumed) cheese depends on the type of cheese, the way, place, tools and equipment of production (Feeley et al., 1972, Moreno-Rojas et al., 1994). Although Kashar is one of the most popular cheeses in Turkey, there is no valid data about the heavy metal contents of Kashar cheese produced in industrially developed Thrace region.

The purpose of the present study is to determine the concentration of some heavy metals in Kashar

cheese samples consumed in Thrace region, besides if the legal tolerance levels are exceeded or not.

Material and Method

The collection of samples

A total of fifty Kashar cheese samples were collected from the retail markets in Thrace region of Turkey (approximately 250 g samples in their original packages). The samples, taken to the laboratory, were homogenized and stored at -20 °C before analysed.

Reagents and chemical standarts

During the experiment, nitric acid (HNO₃ 65%), hydrogen peroxide (H₂O₂ 30%), and multi-element calibration standard-2A (10g/ml, Agilent, Palo Alto, CA) were used. Aqueous standards were prepared with appropriate dilution of a 10 mg/l multi-element solution with ultra pure water and stored at refrigerator.

Wet burning of Samples

The burning processes, which are necessary to eliminate organic compounds and transfer the inorganic elements into soluble phase, were accomplished by using a Mars-5 microwave wet ashing system (CEM Corp., Matthews, NC, USA) and its accessories (Ellen and Van Loon 1990, Fuente et al. 1997, CEM corporation, Nordic Committee on Food Analysis (Anonymous, 1998). Approximately 2 g of sample was weighed into 100 ml teflon vessels. Samples were digested with 5 ml HNO₃ and 2 ml of H₂O₂ in a microwave digestion system. The resulting extracts were cooled and diluted to 10 ml with deionized water (Milli-Q Millipore 18.2 MΩ/cm conductivity). A blank digest was carried out in the same way. All sample solutions were clear. The samples were digested according to the following temperature programme as follows: 2 min. for 400 w, 2 min. for 400 w, 6 min. for 400 w, 5 min. for 400 w, 8 min. for 800 w, 8 min. for vent. After this, the clear solutions were analysed by AAS.

Determination of lead, tin, copper, and mercury

Measurements are made for every single element by using different cathode lamps (Varian 280 Z

Atomic Absorption) in spectrophotometer. Experiments for Pb, Sn and Cu elements are carried at wavelengths of 283.3 nm, 286.3 nm and 324.7 nm, respectively (Anonymous 1988) and the experiment for mercury is carried at wavelength of 253.7 nm (Anonymous 1989).

Statistical Analyses

All data is statistically analysed by using SPSS 10.0 professional Statistics 1999. The experimental data is tested by ANOVA (One way ANOVA randomized complete blocks) and the differences between means are achieved by using the Duncan multiple range test.

Results and Discussion

The data about the heavy metal content of Kashar cheese samples produced and sold in Thrace region, is given in Table 1.

Although the content of Pb was between the range of 0.06-0.04 ppm for 21 samples, it was not at a detectable level for 29 samples (Table 1). According to the Turkish Food Codex, the maximum Pb level in Kashar cheese should be at 0.30 ppm (Anonymous 2002) but according to the results, 3 samples of Kashar cheese had higher Pb content than standard level. The difference in Pb content for all Kashar cheese samples is considered to be significant (p<0.01). The reason for high Pb content might be attributed to unsuitable galvanized or tin-coated metal containers used for milking and transportation of milk. The obtained results indicate that using similar equipment during the production of cheese might affect the Pb content of cheese. More researches about that should be done to reveal the source of this contamination. According to a study done in Italy, the average value of Pb content was determined to be 0.60 pp. and according to another study, the amount of Pb in the cheese samples collected from different regions in Romania was between the range of 0,03 and 0.24 ppm, (Hura, 2002). At a study in Turkey, the average values were found to be 0.14 ppm (for Çecil cheese) and 1.20 ppm (for Çömlek cheese) (Mendil, 2006).

Table 1. The contents of lead, copper, tin and mercury in Kashar cheese samples (ppm)

Samples Number	Pb	Cu	Sn	Hg	Samples Number	Pb	Cu	Sn	Hg
1	0.02 ^a	0.67 ^k	nd	nd	26	0.40 ^g	0.55 ^h	nd	nd
2	nd	0.43 ^e	nd	nd	27	0.10 ^c	0.45 ^e	nd	nd
3	nd	0.62 ^j	0.32 ^d	nd	28	0.05 ^b	0.44 ^e	nd	nd
4	0.30 ^f	0.40 ^d	nd	nd	29	nd	0.45 ^e	nd	0.12 ^f
5	nd	0.51 ^g	nd	nd	30	0.13 ^d	0.50 ^g	0.14 ^b	nd
6	nd	0.32 ^b	nd	0.06 ^{bc}	31	0.02 ^a	0.67 ^k	nd	nd
7	nd	0.48 ^f	0.25 ^c	nd	32	nd	0.43 ^e	nd	nd
8	0.11 ^c	0.60 ⁱ	nd	0.10 ^e	33	nd	0.62 ^j	nd	0.10 ^e
9	nd	0.66 ^k	nd	nd	34	nd	0.40 ^d	0.14 ^b	nd
10	nd	0.45 ^e	nd	nd	35	nd	0.51 ^g	nd	nd
11	0.40 ^g	0.55 ^h	nd	nd	36	nd	0.26 ^a	nd	0.05 ^b
12	0.10 ^c	0.45 ^e	nd	nd	37	nd	0.48 ^f	nd	nd
13	0.05 ^b	0.44 ^e	nd	nd	38	0.11 ^c	0.60 ⁱ	nd	0.10 ^e
14	nd	0.45 ^e	nd	0.12 ^f	39	nd	0.66 ^k	nd	nd
15	0.13 ^d	0.50 ^g	0.14 ^b	nd	40	nd	0.45 ^e	nd	nd
16	0.02 ^a	0.67 ^k	nd	nd	41	0.40 ^g	0.55 ^h	nd	0.08 ^d
17	nd	0.43 ^e	nd	nd	42	0.10 ^c	0.45 ^e	nd	nd
18	nd	0.62 ^j	nd	nd	43	0.05 ^b	0.44 ^e	nd	nd
19	nd	0.40 ^d	nd	nd	44	nd	0.45 ^e	nd	0.12 ^f
20	nd	0.51 ^g	nd	nd	45	0.13 ^d	0.50 ^g	0.14 ^b	nd
21	nd	0.34 ^c	nd	0.07 ^{cd}	46	0.02	0.67 ^k	nd	nd
22	nd	0.48 ^f	0.14 ^b	nd	47	nd	0.43 ^e	nd	nd
23	0.11 ^c	0.60 ⁱ	nd	0.10 ^e	48	0.21 ^e	0.62 ^j	0.56 ^e	0.15 ^g
24	nd	0.66 ^k	nd	nd	49	nd	0.40 ^d	nd	nd
25	nd	0.45 ^e	nd	nd	50	nd	0.51 ^g	nd	nd
		<i>Min.value</i>				nd	0.26	nd	nd
		<i>Max.value</i>				0.40	0.67	0.56	0.15
		<i>Average value</i>				0.06	0.5036	0.0366	0.0214
		<i>Limit value</i>				0.30	1.00	250.00	0.03

Different lower case indicate significant differences among themselves of Kaşar cheese (P<0.01)

nd Not Detectable

Pb values, which we have both identified within the specified limits and the values in general, are under the average literature values. If 100 g cheese is considered to be consumed every day, Pb concentration level in body varies between 0.06 ppm and 1 ppm. For an average adult (60 kg body weight), the provisional tolerable daily intake (PTDI) for lead is 214 mg (FAO/WHO 1999) and in our study, the lead level was not exceed the limit. To minimize excessive Pb intake, a maximum tolerance level is important. This can easily be achieved with good production processes and with the prohibition of the sales of contaminated products (Schwartz, 1994, Debake et al., 2002).

In this study, Cu contents of Kashar cheese were determined as follows: minimum 0.26 ppm, maximum 0.67 ppm, and the average was found to be 0.5036 ppm (Table 1). According to the Turkish Food Codex, the acceptable limit value for Cu content is 1.0 ppm (Anonymous, 2002). The obtained results from all of the samples are below this limit. In terms of Cu content, Kashar cheeses have significant differences ($p < 0.01$). Concerning the amount of Cu that can be found in hard cheeses, such as cheddar, the nutritional risk is even greater when the cheese is exposed to lipid oxidation which is technologically important in terms of discoloration (Jiménez et al. 1984). In some studies, ranging limits of the Cu were reported 0.1-1.3 ppm in Mozzarella cheese (Basile et al., 1978), 1.0 ppm in Manchego cheese (Moreno-Rojas et al. 1994) and 0.2-0.4 ppm in Gouda cheese (IDF, 1992). In this study, the Cu values determined in the samples were similar to the hard cheeses' and are also close to the value, described in the Turkish Food Codex, they are below the limit.

Concerning the Sn content, in the analysis of the Kashar cheese samples collected from different markets, the maximum level was 0,56 ppm whereas the minimum levels could not be found and the average level was found to be as 0.0366 ppm (Table 1). The differences among the values were significant ($p < 0.01$). According to the Turkish Food Codex, the Sn content in Kashar cheese must be 250 ppm as maximum and all samples were

found to be acceptable (Anonymous 2002). Food poisoning due to Sn occurs very rarely and just occurs due to the environmental pollution. Yet, some other complications may occur such as, acute eye and skin irritations, headaches, abdominal pain, nausea and dizziness, severe sweating, shortness of breath, long-term depression, liver damage, immune system impairment, chromosomal anomalies, red blood cell deficiency, brain damage, nervousness, sleep disorders, amnesia (Shills et al., 1994).

The average Hg content in Kashar cheese was found to be as 0.0214 ppm (Table 1). The minimum level of Hg content could not be found whereas the maximum value was found to be 0.15 ppm. The differences in the specified amounts of Hg in cheese samples were found to be significant ($p < 0.01$). According to the Turkish Food Codex, the acceptable amount of Hg in cheese must be as maximum 0.03 ppm (Anonymous 2002). In this study, Hg contents of the 12 samples were above the acceptable limit value.

Conclusion

In this study, Pb, Cu, Sn and Hg contents of Kashar cheese were determined in terms of food safety. While Cu and Sn contents of all cheese samples are under the maximum acceptable limit which has been determined by Turkish Food Codex, Pb and Hg contents of these samples were found to be above this limit. The high levels of Pb and Hg contents in cheese samples may be an important risk factor for the human body and may cause serious health problems. Therefore, the containers used in the production process of cheese must not contain heavy metals. The manufacturers must be informed about this issue so the government has an important role for preventing the environmental pollution. Our results showed that some contaminations occurred depending on the equipment used during the production of the milk and/or production process of the cheese. Further researches are necessary to find out the exact source of contamination.

References

Anonymous 1988. Varian Analytical Methods for Graphite Tube Atomizers, Varian Australia Pty Ltd Mulgrave, Victoria, Australia, Publication No 85 100848-00.

Anonymous 1989. Varian Analytical Methods for Flame Atomizers, Varian Australia Pty Ltd Mulgrave, Victoria, Australia, Publication No 85 100009-00.

Anonymous 1998. Metals: Determination by Atomic Adsorption Spectrophotometry After Wet Digestion in Microwave Oven. In *NMKL Method No. 161*, p. 8, Nordic Committee on Food Analysis, Norway.

- Anonymous 2002. Türk food codex. Communication on determination of maximum levels of some contaminants in foods (pp. 1–198). Ankara: T.C. Tarım ve Köy İşleri Bakanlığı.
- Basile, G., V. Tarallo and P. Violante, 1978. II. Contenuto in metalli pesanti nei formaggi a pasta filata prodotti in Campania. *Bollettino dei Laboratori Chimici Provinciali*, 29, 38–46.
- CEM., 1998. *Mars-5 Microwave Accelerated Reaction System*, pp. 119, CEM Corporation, North Carolina.
- Dabeka, R. W., A. D. McKenzie and K. Pepper, 2002. Lead contamination of raisins sold in Canada. *Food Additives and Contaminants*, 19, 47–54.
- ELLEN, G. and J.W. VAN LOON, 1990. Determination of cadmium and Pb in foods by graphite furnace atomic absorption spectrometry with Zeeman background correction: Test with certified reference materials. *Food Addit. Contam.* 7, 265–273.
- FAO/WHO, 1993. Evaluation of Certain Food Additives and Contaminants. 41st Report of the Joint FAO/WHO Committee on Food Additives. WHO technical report series no 837. World Health Organization, Geneva, Switzerland.
- FAO/WHO, 1999. Joint FAO/WHO Expert Committee On Food Additives, *Summary and Conclusions*, 53rd Meeting, Rome, 1–10 June.
- Feeley, R. M., P. E. Criner, E. W. Murphy, and E. W. Toefer, 1972. Major mineral elements in dairy products. *Research*, 61, 505–510
- FUENTE, M.G., B. CARAZO and M. JUARE, 1997. Determination of major minerals in dairy products digested in closed vessels using microwave heating. *J. Dairy Sci.* 80, 806–811.
- Hura, C., 2002. Chemical contaminants in food and human body, 1990–2000. Ceram Press, Iasi, ISBN 973-8188-01-6.
- IDF, 1992. Trace elements in milk and milk products. *International Dairy Federation Bulletin* (Brussels), No. 278.
- Jime'nez, A. M., M. A. Herrador, and A. G. Asuero, 1984. Elementos traza en alimentos. I. Aspectos metodológicos de su determinación. *Alimentaria*, 152, 107–111.
- John, H. D., G. J. Howard, and C. Worthy, 1996. *Fundamental toxicology for chemists*, UK Royal Society of Chemistry Information Services, Cambridge, pp. 516.
- Mendil, D., 2006. Mineral and trace metal levels in some cheese collected from Turkey. *Food Chemistry*, 96, 532–537.
- Moreno-Rojas, R., M. A. Amaro-Lopez, and G. Zuerera-Cosano, 1994. Copper, iron and zinc variations in Manchego-type cheese during the traditional cheese-making process. *Food Chemistry*, 49, 67–72.
- Anonymous, (161 1998), 1998. Metals: Determination by Atomic Adsorption Spectrophotometry After Wet Digestion in Microwave Oven. In *NMKL Method No. 161*, p. 8, Nordic Committee on Food Analysis, Norway. (literatür listesinde yok)
- Schwartz, J., 1994. Societal benefits of reducing lead exposure. *Environmental Research*, 66, 105–124.
- Shills, M. E., J. A. Olson and M. Shike, 1994. *Modern nutrition in health and Disease*. 8th ed. Phill. Lea & Febiger.
- Şimşek, O., R. Gültekin, Ö. Öksüz, O. and S. Kurultay, 2000. The effect of environmental pollution on the heavy metal content of raw milk. *Nahrung*, 44, 360–363.