

Impacts of changing climate on maize production in the transitional zone of Ghana. A case study of Nkoranza south municipality

Victor Adjei^{1*}

Rosina Kyerematen²

¹Climate Change and Sustainable Development Programme, University of Ghana, Legon

²Department of Animal Biology and Conservation Science, University of Ghana, Legon

*Corresponding Author: victoradjei73@gmail.com

Abstract

This study sought to assess the challenges and opportunities that come with climate change and variability impacts on maize farming in the Nkoranza South Municipality in the Transitional Zone of Ghana. The mixed method approach was used in collecting the data. The results of the study showed that farmers had observed changes in climate in the form of decreasing rainfall, rising air temperatures and seasonal changes in rainfall pattern which were affecting their maize farming operations. The major setbacks within the area were deficit in rainy days and intermittent erratic rainfall affecting maize production. The reduction in rainfall was corroborated by data obtained from Ghana Meteorological Agency (GMet) which indicated that the dry cell between the major and minor rainy seasons was getting wetter and the two seasons were gradually merging whereas the first and last quarters of the year were getting drier for a couple of decades. The situation over the last five years had worsened as the amount of total rainfall had reduced by 22% compared to the 30 year period between 1960 and 1982. The major opportunity available to farmers in the face of changing climate in this agroecological zone was cashew production. About 76.8% of the respondents had diversified into cashew farming.

Keywords: Soil, Physical properties, Bade, Yobe

Received: 02.05.2018



Accepted: 05.10.2018



Published (online): 15.10.2018

Introduction

There is now unequivocal evidence that the climate of the Earth has demonstrably warmed up since the pre-industrial era and that most of the warming over the last 50 years is very likely to have been as a result of increase in Greenhouse Gases (GHGs) concentrations in the atmosphere (Intergovernmental Panel on Climate Change (IPCC), 2007).

Three major physical impacts of climate change according to IPCC (AR4) in Ghana are; temperature rise, changing rainfall regime towards a longer dry season and a vanishing wet season. Owusu *et al.*, (2008) also underscore the fact that there has been a noticeable shift in the rainfall pattern in Ghana towards a longer dry season and the disappearance of short dry spells between major and minor seasons. The Ghanaian agricultural-dependent economy has thus suffered severe economic consequences (Kyekyeku Nti, 2008).

Agriculture is the pillar of Ghana's economy providing employment to the masses in the rural areas and sustaining the vast majority of the population. Only about 0.89% of the country's arable land, equivalent to 23,657 hectares is cultivated under irrigation, benefitting only rice production [Environmental Protection Agency (EPA) 2009; Ministry of Food and Agriculture (MoFA), 2003].

According to Environmental Protection Agency of Ghana (EPA, 2000), maize yields in Ghana are low compared to that of developed countries. The national average yield is about 1.6 mt per hectare. This is primarily due to the inadequate input of fertilisers and non-existing

irrigation facilities in the maize production areas (WABS consulting, 2008).

Among the staples cultivated in the country, maize has been identified as one of the most essential within the grains and cereals family [Statistical, Research and Information Directorate (SRID), 2010]. It is grown on more than, 997,661 hectares across all the various agro-ecological regions in Ghana (SRID, 2010)

Generally, high temperatures have been reported as detrimental to grain yields (Badu-Apraku *et al.*, 1982; Lobell *et al.*, 2008). According to the EPA (2000), the projected percentage drop in maize yield in the transition zone ranged from 0.5 percent in the year 2000 to 6.9 percent by the year 2020 as compared to their level in 2000. The study of Lobell *et al.* (2008) indicates that each day's temperature above 30°C would reduce the final yield by 1% under optimal rain-fed conditions, and by 1.7% under drought conditions. In the transition (where Nkoranza is located) and forest belts, though droughts are not the main drawback, variations in rainfall particularly deficits in the number of rainy days pose new challenges to rural livelihoods (Yaro, 2010). There is a shortening of the farming season in several places as well as a gradual waning of the secondary growing season in the transition zone (Owusu and Waylen, 2009; Yaro *et al.*, 2010).

Materials and Methods

The study was conducted in the Nkoranza South Municipality in the Brong Ahafo Region of Ghana.

Cite this article as : Adjei, V. and Kyerematen, R. (2018). Impacts of changing climate on maize production in the transitional zone of Ghana. A case study of Nkoranza south municipality. Int. J. Agric. Environ. Food Sci., 2(4), 164-170. DOI: 10.31015/jaefs.18028

Available online at : www.jaefs.com - <http://dergipark.gov.tr/jaefs>



The Municipality lies within the wet semi-equatorial region of Ghana. It is located in the transitional zone between the savannah woodland of northern Ghana and the forest of the south. The Municipality lies within longitudes 1°10'W and 1°55'W and latitudes 7°20'N and 7°55'N (Ghana District, 2007). It shares common boundaries with Nkoranza North District to the north, Techiman Municipality to the west, Offinso North to the south and Ejura-Sekyere-Dumase to the south-east (Ghana District, 2007).

The eastern part of the Municipality is mainly characterised by savannah woodland and the southern part is principally marked by forest regrowth, made up of shrubs and grasses. It has a mean annual rainfall level ranging between 800-1200 mm, relative humidity ranging from 55-90% and average annual temperature of 26°C (Ghana District, 2007).

Data Collection

Both primary and secondary data were used in obtaining the data. With regards to primary data, two focus group discussions were organized in two communities (Donkro Nkwanta and Aboasu). Twelve key informants were also interviewed, with at least, one coming from each community. The maize farmers were seven, two sub-chiefs, a meteorologist from GMet- Wenchi branch, an agricultural extension officer from Nkoranza South Municipal Assembly and the 2015 best farmer of the Municipality.

Secondary data consisted of meteorological data (rainfall and temperature) from GMet which covered 1960 to 2014. This data (rainfall) was divided into two trenches, 1960 to 1982 and 1988 to 2014. Some data between 1983 to 1987 were unavailable and so were not used for the analyses. Rainfall data over the last five years (2010 to 2014) as well as temperature data from 1983 to 2014 were analysed in determining current rainfall trends in the study area. Maize production data (2010-2015) was also secured from the assembly of the Municipality for analysis.

Study Design

The study relied on a questionnaire survey from selected farming communities, qualitative interview and focus group discussion (FGD) to acquire essential field data. The questionnaire survey was administered among some selected maize farming communities in the Municipality of which purposive and accidental sampling techniques were used to enhance diversity of the respondents. The purposive technique was used to obtain the farming communities; percentage of male and female farmers as respondents, and selection of key informants. The key informants were made up of the aged, chiefs, a meteorologist and an agricultural officer. Any farmer chanced upon was also given the opportunity to answer the questionnaire (Accidental sampling technique).

The questions included close-ended as well as a few open-ended questions, most of which sought information on respondent's perception on climate change and variability, farm size, variety of maize cultivated, farming system, application of fertilizer and other coping strategies meant to reduce the effects of low yields. The survey was conducted face-to-face with the respondents in different locations and in different communities. The questionnaires were read and translated into Akan (the main local language) for the respondents to answer.

In all, 12 key informants were interviewed to gain information about their views on climate change and variability impacts on maize farming in the Municipality.

Seven of the respondents were maize farmers, three were elders from the chief's palace whose ages were between 65 and 73 years (the aged) and the rest were farmers. At least, one respondent came from each of the five farming communities in the study area. Apart from these key informants, one of the Municipal Agricultural Officers was interviewed. Lastly, a staff of Ghana Meteorological Agency (GMet, Wenchi branch) was also interviewed. The questions in these qualitative interviews were unstructured, but were administered with the help of an interview guide. The themes covered were similar to those in the questionnaire survey, but were more interactive and probing, seeking information on climate change and variability of the study area, adaptation of the farmers etc. with ample opportunity for the informants to provide extensive personal narratives. The interviews were done in the participants' homes or residences and were tape-recorded with their permission and later transcribed.

Two FGDs were organized at the end of the data collection in two communities. Averagely the two FGDs were made up of three females and seven male farmers. The focus groups dealt with the changing climate and variability in the area, its impacts both positive and negative on their livelihoods, their adaptive capabilities to alleviate the challenges posed by the changing climate and economic/financial assistance they received elsewhere (if any) to support themselves. All the deliberations were videoed with the permission of the discussants and later transcribed. The selection was made based on how well they answered questions during the questionnaire administration.

Results and Discussion

Demographic Information on Maize Farmers

Table 1 provides basic information about the maize farmers selected for the questionnaire survey. The respondents were made up of both male 154 (70%) and female maize farmers 66 (30%). The number of respondents was proportional to the population of each community. Sixty percent of the respondents were Bonos, 33.6% were settlers from the northern part of Ghana, 3.6% Asantes and 2.3% representing Fulanis and Ewes. Majority of maize farmers rented lands for farming accounting for 51.8%. Maize farmers who worked on their own lands were few representing 20.5%. Several of the farmers had farmlands of size greater than 2 hectares (ha). About 55.5% of the farmers had farmlands greater than 2 ha. Ninety-seven percent of respondents practised monoculture which was the dominant farming system among the maize farmers within the Municipality with 2.3% of respondents practising intercropping or mixed cropping. Eighty-seven percent of the respondents sowed *aburohoma*, a variety of maize prevalent in the Nkoranza South Municipality. Even though approved seeds like *obaatampa* and *omankwa* were available on the market, most farmers preferred the traditional variety to these new cultivars which were approved to be drought resistant and early maturing hybrids.

The percentage of farmers who applied fertilizers on their crops was 71.4%. About 17.4% of the respondents did not apply fertilizers and the farmers who occasionally applied fertilizers were 10%. Most of the farmers enriched their crops with organic fertilizers.

Twenty-nine percent of the respondents had not received any formal education with only 5.5% of the respondents having attained tertiary level education. Majority of the respondents were Junior High School (JHS)/Middle school



leavers representing 40.9%. Data from the survey indicated that males (8.2%) were better educated than females. About 34.8% of the female farmers had not received formal education while 26.6% of males had no formal education. A study by Khan and Ali (2013) revealed that if producers are educated, it enhances the application of best farming practices in tomato production.

Perception of Climate Change and Variability among Farmers

All farmers interviewed indicated that the pattern of rainfall in the area had changed. Of these, 78% indicated that the nature of the change was erratic. The majority (85.9%) indicated the severity of this episode was embarrassing over the past 5 years (Table 2). This had therefore made their farming operations more precarious due to the fact that there is no irrigation system in the Municipality, bringing farming activities to a near halt whenever the rain failed.

About 65.5% of the farmers admitted that unguided activities of unscrupulous chainsaw operators had led to the cutting down of trees which might have resulted in reduction in rains. On the other hand, 25.5% attributed the changes in precipitation trend to nature itself and 0.5% of respondents believed that the changes had come due to excessive emission of GHGs into the atmosphere. Another 5% of farmers also placed the blame on bush fire whereas 3.2% indicated that they did not know what had caused this change.

When they were further probed about what could be done to reverse the situation of the erratic rainfall, 64.1% responded that there should be afforestation to replace the felled trees. Others (27.7%) had the view that only God could intervene to restore the situation while some farmers (5.5%) called for bush fire control. Farmers in the FGDs disclosed that control of bush burning would allow small plants to replace logged trees although this would take a couple of years to establish. About 2.3% of respondents did not know what could be done to bring the rainfall situation back to normal whereas 0.5% also had the assurance that a ban on burning of fossil fuel would be helpful. This indicates that farmers do have some knowledge about causes of changing climate but the remedies were not within their reach due to population growth and poverty (FGDs).

With regard to temperature, when farmers were asked whether air temperatures had increased, stabilized or decreased, 86.8% *strongly agreed* that air temperatures had currently risen. It emerged from the FGDs that air temperatures had been warmer over the past 10 years and even the two cold periods (June-July and December-January) were now short lived. About 7.7% *agreed* that air temperatures had changed but not appreciably while 5.5% did not know whether temperatures had changed or not (Fig. 1). Another 86.8% of respondents had observed that the current temperatures were higher than the temperatures for the same area over the past 20yrs.

Table 1. Demographic information of farmers

Community		Gender		Farm size			Level of Education			Place of birth		
		M	F	< 2 (ha)	2-4 (ha)	>4 (ha)	None	Basic	SHS	Tertiary	Native	Alien
Aboasu	(35)	28	7	34	32	34	37	46	14	3	31	69
Asoano	(30)	24	6	43	37	20	30	60	7	3	90	10
D/Nkwanta	(95)	57	38	49	20	31	25	59	8	8	47	53
Kyekyawere	(30)	24	6	37	30	33	27	63	10	0	30	70
Kyiradeso	(30)	21	9	50	30	20	33	53	7	7	30	70
Total	(220)	154	66	44%	27%	29%	29%	57%	9%	5%	46%	54%

Table 2. Changing rainfall pattern response on the time of the changes

Changing rainfall pattern	Response on the time of the changes		Total (%)
	1-5years (%)	6-10years (%)	
Yes	85.9	14.1	100.0
Total	85.9	14.1	100.0




Figure 1. Farmers' perception on changing air temperatures

Analysis of Rainfall Data of Nkoranza South Municipality (1960-2015) Depicting the Regime of Climate Change and Variability in the Study Area

Figures 2a and 2b cover the precipitation trend for the study area (Nkoranza South Municipality) from 1960-2015. 1983 to 1987 did not appear in the tables/figures because there were a lot of blank cells within this period.

Precipitation was low in the early 1960s but rose steadily in the mid-1960s, falling suddenly in the 1970s. Rainfall however progressively rose and somewhat stabilized in the mid-1970s and declined again in the early 1980s. There was no clear pattern from 1988 to 2015. It was also obvious that from 1988 to 2015, the rainfall amount hardly crossed 1000mm unlike the period between 1960-1982. This confirms what the farmers said that the current rainfall pattern is rather irregular and unpredictable.

The averages of 1960-1982 were higher than that of the present, 1988-2014 (Table 3b). From the months of January to March, the mean of 1960-82 was greater than that of 1988-2014. The farmers were right that lately, the amount of rainfall in March had drastically reduced as the amount from 1960-82 (Table 3a) was well over 100mm (114.7mm) whereas the average amount for the period between 1988 and 2014 had reduced to 71.0 mm, the reason, most probably, why majority of the farmers sowed in April (79.1%). Between 1988 and 2014 rainfall averages had appreciated from April to June, with a deficit in July, rising again in August. Currently, the amount of rainfall in November has decreased to the detriment of farmers. This means that presently, crops do not receive as much rainfall as compared to 30 years ago. It can also be deduced from Tables 3b and 3c that presently, August is getting wetter and the major and minor seasons are progressively merging. It is also clear that the first and last quarters of the year (January-March as well as October- December) are getting drier over the last couple of decades with the last five years the worst.

The future seems to be rather bleak looking at the trend in 1960-1982 with almost no rains in November and December (2010-2014). The average total amount of the period 2010-2014 showed a reduction of 22% over the past five years with a decline between 1960-82 (100.7) and 1988-2014 (91.9) of 6%. The month of March which was meant for planting of crops now obtained 64.5 mm (3c) of rainfall against 114.7 mm (3a) in the past butting farmers assertion that climate had changed.

Irregularity of Rainfall

On rainfall regularity, the pattern of precipitation in the Municipality is characterised by uncertainty. Based on the above tables, Tables 3a-c show that the amount of rainfall in the month of August increased but figures from 2015 deviated from the trend (Table 3d). The precipitation received in the August, 2015 farming season was 0.00mm. Table 3d also showed a reduction in rainfall amounts between June and August. This phenomenon (irregularity) might have had a negative effect on maize production in 2015 resulting in the lowest yields (1.5mt/ha) of maize recorded for the Municipality in the last six years. A report by EPA (2000) indicates that even though other contributing factors exist, rising temperature and irregularity in precipitation are the major causes of the continuous reduction in maize yields. Annual rainfall for Nkoranza South Municipality since the 2008/9 farming season for the month of March had never risen above 100mm but rose to 113mm in 2015 which confirms that the nature of rains in the

transitional zone is irregular. According to Stanturf *et al.*, (2011) declines in total precipitation and increasing irregular rainfall patterns make farming more hazardous, increasing the likelihood of crop failures and reducing agricultural production.

Impacts of Climate Variability and Change on Maize Production in the Municipality

The major threats of changing climate were rainfall (erratic) and rising air temperatures (FGD). The intensity of precipitation had reduced but rainy days which had declined had had a greater effect on crop yields than amount of rainfall. Over the past six years, the intensity as well as rainy days in the Municipality had varied from one year to another as shown in Table 4. The lowest amount of rainfall over the past six years (2010-2015) was in 2010 (812.3mm) with the highest in 2013 (1487.2). Some of the rainy days within the year were spread out (it rained on several days) while others were not. For instance, in 2010, the rainy days were 85 whereas in 2015, the rainy days were only 44. What can be inferred from the rainfall regime over the past six years is that the rainy days had become more inconsistent than the amount or intensity of rainfall. This confirms a study by Owusu and Waylen (2009) that more rainy days with less amount of rainfall does not have much impact on yields. The reduction in rainfall intensity in the study area was in line with a similar study by Travasso *et al.*, (2009) in Argentina which showed that there had been an observed reduction in the growing season of maize crops by 27 days, consequently causing a reduction in yields when crop yield (maize) was simulated using version 3 of Hadley Center Coupled Model (HadCM3).

There was a relationship between rainy days and maize production in the Municipality. When rainfall spread over more days, crop yields increased but when rainy days were few with large amounts of rain, crop yields dropped. For instance, in 2010, the amount of rainfall (812.3mm) was the least among the six years under review, however, average maize yields in the Municipality was 2.0 mt/ha compared to 2011, which had 1270.8mm of rainfall over 51 rainy days (Table 4). In 2011, the average yield was 1.9 mt/ha yet total cultivated area was bigger (61,231 ha) than that of 2010 (59,616 ha). The farmers who were interviewed hammered on the fact that erratic rainfall had been one of their major setbacks and 2015, for instance, was a terrible year. Rainy days had never gone below 50 days over the past six years but 2015 recorded 44 rainy days which was exceptional (Table 4). The farmers complained that in both the major and the minor seasons of 2015, the rains started late and the cessation too was early. The minor season's onset was somewhere late in September ending in mid-October which might have prevented some of the farmers from planting their seeds most likely resulting in a smaller cultivated area (42,500 ha) yielding 1.5 mt/ha in 2015 (Table 4). Fosu-Mensa (2012) reported that increased variability of precipitation, which is reflected in the high variability in grain yield, was another factor leading to the reduction of crop yields.

Opportunities Associated with Climate Change and Variability in Nkoranza South Municipality

It was observed that over the past 20 years, the livelihood of the people in the Municipality had somewhat changed. According to the farmers (FGDs), some varieties of corn, yam, groundnut and cassava were fast disappearing. Table 4 summarises the benefits farmers had derived from climate



variability and change. Majority of respondents (76.8%) (Table 5) said that cashew production had given them some form of relief with. About 66.9% of the cashew farmers being Bonos while 26.6% (Busanga, Dagarti, Wale, Kasena, Kusase, Gruma, Frafra) were from the three northern regions. Three major reasons respondents attributed to high rate of cashew farming were that it did not demand much time, energy and money. The other alternate form of livelihood that helped farmers adapt to the changing climate was livestock rearing as climate variability did not have any great impact on it (livestock), however, its major problem was pests and diseases (57.1%), followed by theft (33.3%). Most of the animals were reared by farmers from the North Ghana and this represented 78.9% with the Bonos accounting for 21.1%.

Conclusion

The study revealed that changes in the onset and cessation of rain coupled with intermittent dry spells had negative effects on maize farming production in the Nkoranza South Municipality, and this posed a great threat to household incomes as most farmers earned their living from farming. Vulnerability of maize production to climate change and variability in this Municipality depended on the onset of rainfall with respect to their growth stage of the crop. Maize production in the Municipality solely depended on

rainfall hence its variability affected production

Cultivation of cashew has become the sole alternative form of livelihood providing income for households as maize production had become more precarious and unreliable.

The amount of rainfall had reduced drastically over the last five years to the detriment of maize farming. The major and minor seasons were gradually merging as August had become wetter over the last two-three decades with the exception of 2015. January, February, March, October, November and December were becoming progressively drier in the last 5-10 years as compared to three decades ago. This demonstrates clearly that meteorological conditions over the transitional zone would continue to alter negatively at the expense of poor-resourced maize farmers whose farming operations are dependent on rain-fed.

Acknowledgement

We are grateful to the Climate Change and Sustainable Development Programme at the University of Ghana for making it possible for this research to be carried out. We would also like to thank GMet-Wenchi, Nkoranza South Municipal Assembly (Agricultural Department), as well as the chiefs, elders and people of Nkoranza South Municipality for their willingness to take part in the survey and for the valuable information they gave us.




Figure 2a. Rainfall data for Nkoranza South Municipality (1960-1982) (Computed from GMet data)




Figure 2b. Rainfall data of Nkoranza South Municipality (1988-2015) (Computed from GMet data)

**Table 3a.** Average monthly rainfall 1960-1982 for Nkoranza

month	J	F	M	A	M	J	J	A	S	O	N	D	total avg.
mm	10.8	46.0	114.7	129.8	154.0	154.0	116.6	72.7	192.2	169.9	37.0	11.4	100.7

Table 3b. Average monthly rainfall 1988-2014 for Nkoranza

month	J	F	M	A	M	J	J	A	S	O	N	D	total avg.
mm	7.1	35.1	71.0	148.2	159.0	174.0	97.7	79.9	162.1	141.1	20.7	6.8	91.9

Table 3c. Average monthly rainfall for the past 5yrs (2010-2014) for Nkoranza

month	J	F	M	A	M	J	J	A	S	O	N	D	total avg.
mm	5.0	34.9	64.5	121.0	147.5	158.8	80.8	88.5	138.3	90.9	8.1	0.0	78.2

Table 3d. Annual Rainfall Figures for 2015

month	J	F	M	A	M	J	J	A	S	O	N	D	Total avg
mm	0.0	43.9	113.0	35.3	88.3	66.1	50.1	0.0	214.8	160.8	86.5	0.0	71.6

Table 4. Rainfall and Maize production data for Nkoranza South Municipality (2010-2015)

Year	Rainfall intensity (mm)	Rainy days	Average yield (mt/ha)	Production level (mt)	Total cultivated area (ha)
2010	812.3	85.0	2.0	119232.0	59616
2011	1270.8	51.0	1.9	116338.9	61231
2012	1110.7	57.0	2.1	124452.0	59263
2013	1487.2	70.0	2.0	124899.0	62494
2014	1131.3	65.0	1.9	115250.0	60658
2015	949.0	44.0	1.5	83570.0	42500

Source: Computed from Nkoranza South Assembly data, 2015

Table 5. Response on the Time of the Changes Benefits from Climate Change Variability

Tribe	Benefits from climate change variability					Total
	Cashew production	Animal rearing	Trading	Others(oil palm, plantain, vegetable, melon)	Nothing	
Bono	85.0%	3.0%	0.8%	1.5%	9.8%	100.0%
Asante	87.5%	0.0%	0.0%	12.5%	0.0%	100.0%
Settlers from the North	60.8%	20.3%	1.4%	1.4%	16.2%	100.0%
Others(Ewe and Fulani)	80.0%	0.0%	0.0%	0.0%	20.0%	100.0%
Total	76.8%	8.6%	0.9%	1.8%	11.8%	100.0%

References

- Badu-Apraku, B., Hunter, R.B., and Tollenaar, M. (1983). Temperature effects on maize grain filling. *Canadian Journal of Plant Science*, 63: 357-363.
- Environmental Protection Agency, Ghana (EPA) (2000). Climate change vulnerability and adaptation assessment of coastal Zone of Ghana. Final report. Accra: Ministry of Environment Science and Technology.
- Intergovernmental Panel on Climate Change (IPCC), (2007). Climate Change 2007: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change [Parry, Martin L., Canziani, Osvaldo F., Palutikof, Jean P., van der Linden, Paul J., and Hanson, Clair E. (eds.)]. Cambridge University Press, Cambridge, United Kingdom, 1000 pp
- Environmental Protection Agency, Ghana (EPA), (2009). Ghana's national climate change adaptation strategy. Zero draft. Accra: Ministry of Environment Science and Technology.
- Fosu-Mensah, Y.B. (2012). Modelling maize (*Zea mays* L.) productivity and impact of climate change on yield and nutrient utilization in sub-humid Ghana. In: Vlek, P.L.G., Denich, M., Martius, C., Manschadi, A., Bogardi, J. (Eds), Ecology and Development Series No. 87. Cuvillier Verlag, Gottingen. Pp.: 1-171
- Ghana Districts, (2007). District Profiles. <http://www.ghanadistricts.com>
- Khan, H., and Ali, F. (2013). Measurement of Productive Efficiency of Tomato Growers in Peshawar, Pakistan. *Agric. Econ. CZECH*, 2013, 59, No. 8, p. 381-388. ISSN 0139-570 X.
- Kyekeyku Nti, F. (2008). Climate Change Vulnerability and Coping Mechanisms among Farming Communities. MSc Thesis, Department of Agricultural Economics College of Agriculture, University of Ghana, Legon.
- Lobell, D.B., Burke, M.B., Tebaldi, C., Mastrandrea, M.D., Falcon, W.P., and Naylor, R.L. (2008). Prioritizing climate change adaptation and needs for food security in 2030. *Science*, 319, 607-610.
- Ministry of Food and Agriculture (MoFA), (2003). Agriculture in Ghana: facts and figures. Produced by the Statistics, Research and Information Directorate Accra. <http://www.fao.org/docrep/008/a0013e/a0013e05.htm> cited on 12th March, 2013



- Owusu, K., and P. Waylen. (2009). "Trends in spatiotemporal variability in annual rainfall in Ghana (1951–2000)." *Weather* 64(5): 115–120.
- Owusu, K., Waylen, P., and Qiu, Y. (2008). Changing rainfall inputs in the Volta basin: implications for water sharing in Ghana. *GeoJournal*, 71, 201–210.
- Stanturf, J. A., Warren, M.L., Charnley, S., Polasky, S.C., Goodrick, S.L., Arma, F., and Nyarko Y.A., (2011). Ghana climate vulnerability and adaptation assessment. US Agency for International Development, Washington D.C., USA
- Statistics, Research and Information Directorate (SRID), (2010). Agriculture in Ghana (2009). Facts and figures. Ministry of Food and Agriculture. Accra, Ghana. Available from <http://www.mofa.gov.gh/AGRICULTURE%20IN%20GHANA%20%202009.pdf>, (accessed 14th April, 2011).
- Travasso, M.I., Graciela, O.M., and Gabriel, R.R., (2009). Climate change impacts on regional maize yields and possible adaptation measures in Argentina. *Int. J. Global Warming*, Vol. 1, Nos 1/2/3
- WABS Consulting, (2008). Draft Report: Maize value chain study in Ghana: Enhancing efficiency and competitiveness. WABS Consulting Ltd. Accra.
- Yaro, J. A. (2010). The Social Dimensions of Adaptation to Climate Change in Ghana. The World Bank Discussion paper no.15. Washington: The World Bank.
- Yaro, J. A., Dogbe, T. D., Bizikova, L., Bailey, P., Ahiable, G., Yahaya, T., and Salam, K. A. (2010). Development and Climate Change: The Social Dimensions of Adaptation to Climate Change in Ghana. Discussion Paper Number 15, 14th December 2010. Washington, DC; World Bank.