

Laparoscopic cholecystectomy in left-sided gallbladder detected during operation

Ameliyat sırasında saptanan sol yerleşimli safra kesesinde laparoskopik kolesistektomi

Yahya Kemal Çalışkan¹, Fatih Başak², Abdullah Şişik²

¹ Department of General Surgery, University of Health Science, Kanuni Education and Research Hospital, Istanbul, Turkey

² Department of General Surgery, University of Health Science, Umraniye Education and Research Hospital, Istanbul, Turkey

ORCID ID of the author(s)

YKÇ: 0000-0003-1999-1601

FB: 0000-0003-1854-7437

AŞ: 0000-0002-7500-8651

Abstract

The case where the gallbladder was located on the lower face of the left lateral segment of the liver was defined as a left-sided gallbladder. It is a rare hereditary anomaly and can be seen with the right-sided ligamentum teres. Vascular and biliary anomalies can be seen in the cases of placement anomaly. Full dissection of the calot's triangle applied as a standard in laparoscopic cholecystectomy is important for the safety of the cases in anomaly cases. In this study, we aimed to present a laparoscopic cholecystectomy performed in a case of a left-sided gallbladder detected during surgery.

Keywords: Gallbladder, Variation, Laparoscopic cholecystectomy

Öz

Safra kesesinin karaciğerin sol lateral segmentinin alt yüzüne yerleşmesi durumu sol yerleşimli safra kesesi olarak tanımlanmıştır. Nadir görülen kalıtsal bir anomali olup, sağ yerleşimli ligamentum teres ile birlikte görülebilmektedir. Yerleşim anomalisi olgularında damarsal ve safra yolları anomalisi görülebilmektedir. Laparoskopik kolesistektomi ameliyatında standart olarak uygulanan kalot üçgeninin tam diseksiyonu anomali olgularında da ameliyat güvenliği için önemlidir. Bu çalışmada ameliyat esnasında saptanan sol yerleşimli safra kesesi olgusunda yapılan laparoskopik kolesistektomi ameliyatını sunmak amaçlanmıştır.

Anahtar kelimeler: Safra kesesi, Varyasyon, Laparoskopik kolesistektomi

Introduction

A left-sided gallbladder (LSG) is a gallbladder located on the left side of the liver, round ligament, and not on the right side, which is its mundane location. It is first described from Hochstetter in 1856. The reported incidence is between 0.1% and 1.2% [1-4]. The present case report demonstrates a case of LSG identified during laparoscopic cholecystectomy. Gallbladder is located in the lower left side of the left lateral segment of the liver. It is a rare hereditary anomaly and can be optically discerned with a right-sided ligamentum teres. In this study, we aimed to present challenges in a laparoscopic cholecystectomy performed in a case of a LSG detected during surgery.

Corresponding author / Sorumlu yazar:

Yahya Kemal Çalışkan

Address / Adres: Genel Cerrahi Kliniği, Kanuni Eğitim ve Araştırma Hastanesi, Küçükçekmece, İstanbul, Türkiye

e-Mail: yahyakemal@yahoo.com

Informed Consent: The authors stated that the written consent was obtained from the patient presented in the study.

Hasta Onamı: Yazar çalışmada sunulan hastadan yazılı onam alındığını ifade etmiştir.

Conflict of Interest: No conflict of interest was declared by the authors.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemişlerdir.

Financial Disclosure: The authors declared that this study has received no financial support.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

Published: 5/28/2019

Yayın Tarihi: 28.05.2019

Copyright © 2019 The Author(s)

Published by JOSAM

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-NoDerivatives License 4.0 (CC BY-NC-ND 4.0) where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.



Case presentation

A 55-year-old woman presented with abdominal pain and jaundice. She is hospitalized with the pre-diagnosis of cholangitis and choledocholithiasis due to occlusion of choledochus. He was discharged after medical treatment. Magnetic resonance cholangiography revealed no abnormality in the gallbladder or pathology in the biliary tract. The patient was prepared for elective cholecystectomy six weeks later. Operation revealed that the gallbladder was located in the left segment of the left lobe of the liver and left of the ligamentum teres (Figure 1, 2). It was decided to continue cholecystectomy without changing the places of entry. Cystic canal and cystic artery were revealed in the caliper dissection with technique of open window and critical view of safety. No major anomalies were observed in the dissection area. Standard retrograde (infundibulum-first) cholecystectomy was performed laparoscopically. The patient was discharged without any problems after one day. The pathological examination of the specimen was reported as chronic cholecystitis with gall stone.



Figure 1: Intraoperative view of left-sided gallbladder



Figure 2: Gallbladder attached to the left lobe of the liver

Discussion

LSG is a rare anomaly. Vascular and biliary anomalies may be seen in these cases with placement anomaly. In laparoscopic cholecystectomy, for surgical safety, surgeon should complete dissection applied as a standard to reveal anomaly of the calot's triangle. Antegrade cholecystectomy (fundus-first) or, if necessary, conversion should be kept in mind

when the association with the main bile duct is suspicious or in cases of such anomaly [3-5].

Aberrant gallbladder can be of 4 different types: Left-sided; intrahepatic; transverse; Retro settled. These are the most rarely seen gallbladder without situs inversus [5]. LSG can be found in two anatomic variants. The first is the actual LSG where the gallbladder is located in the left lobe of the liver. In this case, there may be subtypes depending on how the cystic channel is incorporated into the biliary tree [4-6]. The cystic channel is connected to the right bile duct (CBD) on the right side. The normal gallbladder bud does not go to the left and right of the left lobe, to the left and right of the left lobe, and to the left of the round ligament. Left cystic canal or left CBD connected. It accompanies failure in the development of the normal structure of the right and right side of the liver canal [7]. Second, the gallbladder is on the left side of the round ligament, but the round ligament is still in the right lobe of the liver because it is still attached to the right liver [6].

The inability to predict the cystic duct or the general bile duct may be challenging, and the selective use of selective intraoperative cholangiography might contribute to the safe laparoscopic management of this unusual problem [7,8].

In conclusion, left-sided gallbladder is a rare operative finding. Preoperative imaging may not detect the anomaly. Intraoperative cholangiography should be performed to detect anomalies associated with biliary trees. When the surgeon is suspicious, conversion to open surgery is recommended to prevent complications.

References

1. Idu M, Jakimowicz J, Iuppa A, Cuschieri A. Hepatobiliary anatomy in patients with transposition of the gallbladder: implications for safe laparoscopic cholecystectomy. *Br J Surg.* 1996;83:1442-3.
2. Nagai M, Kubota K, Kawasaki S, Takayama T, Bandai Y, Makuuchi M. Are left-sided gallbladders really located on the left side? *Ann Surg.* 1997;225:274-80.
3. Maetani Y, Itoh K, Kojima N, et al. Portal vein anomaly associated with deviation of the ligamentum teres to the right and malposition of the gallbladder. *Radiology.* 1998;207:723-8.
4. Hsu SL, Chen TY, Huang TL, et al. Left-sided gallbladder: its clinical significance and imaging presentations. *World J Gastroenterol.* 2007;13:6404-9.
5. Wong LS, Rusby J, Ismail T. Left-sided Gallbladder: A Diagnostic and Surgical Challenge. *ANZ J Surg.* 2001;71:557-8.
6. Kubo S, Lee S, Yamamoto T, Edagawa A, Kinoshita H. Left-sided Gallbladder Associated with Anomalous Branching of the Portal Vein Detected by Sonography. *J Osaka Cty Med Cntr.* 2000;46:95-8.
7. Matsumura N, Tokumura H, Yasumoto A, et al. Laparoscopic cholecystectomy and common bile duct exploration for cholecystocholedocholithiasis with a left-sided gallbladder: report of a case. *Surg Today.* 2009;39:252e255.
8. Hasbahecci M, Erol C, Seker M, Basak F, Alimoglu O. Standard laparoscopic cholecystectomy for malposition of the gallbladder caused by right-sided ligamentum teres. *J Minim Access Surg.* 2013 Oct;9(4):177-9.

The National Library of Medicine (NLM) citation style guide is used in this paper.

Suggested citation: Patrias K. Citing medicine: the NLM style guide for authors, editors, and publishers [Internet]. 2nd ed. Wendling DL, technical editor. Bethesda (MD): National Library of Medicine (US); 2007 [updated 2015 Oct 2; cited Year Month Day]. Available from: <http://www.nlm.nih.gov/citingmedicine>