



**Turkish Small Animal Veterinary Association**  
**14 International Continuing Education Congress**  
**8-9-10 NOVEMBER 2019**  
WOW ISTANBUL HOTELS & CONVENTION CENTER



**PS10**

### **Neurofibroma Case in a Dog**

Aynur DEMİR, Eylem BEKTAŞ BİLGİÇ\* Gülşen SEVİM KARAGÖZOĞLU

Istanbul University-Cerrahpaşa Veterinary Faculty, Surgery Department, Istanbul, TURKEY

#### **Abstract**

Peripheral nerve tumors arise in the cranial nerves, spinal nerve roots, and peripheral nerves. Peripheral nerve tumors [peripheral nerve sheath tumors (PNSTs)] are infrequently reported in dogs and rarely in cats. These tumors occur most commonly in middle-aged to older dogs of medium and large breeds. Peripheral nerve tumors occur mainly in the cranial, spinal and associated peripheral nerves and less commonly in the autonomic nervous system. These tumors arise from Schwann cells, perineurial fibroblast or a combination of these two cell types. PNSTs are also classified as schwannomas, neurofibromas and neurofibrosarcomas. Neurofibroma is a benign nerve-sheath tumor in the peripheral nervous system. In this case a 5-year-old Terrier dog was presented which was brought to our clinic with a complaint of swelling back of its left leg. In the examination a big solid swelling, unrelated to the bone, was seen on the back of the left femur. After the diagnostic procedure the patient was prepared for the operation. The dog was anesthetized and performed an excisional biopsy. As a result of the pathological examination the tumor type was determined as neurofibroma.

**Keywords:** Dog, neurofibroma, peripheral nerve sheath tumor (PNST)

\*Corresponding Author: [xxxxxxxx@xxxx.xxx](mailto:xxxxxxxx@xxxx.xxx)