

Aseptomatik dev üreteral polip: Olgu sunumu

Asymptomatic giant ureteral fibroepithelial polyp: A case report

Sacit Nuri Görgel¹, Uğur Balcı¹, Yusuf Kürşad Özdamar², Cengiz Kara¹

¹Izmir Ataturk Training and Research Hospital, 1st.Urology Clinic, Izmir, Turkey

²Manisa Merkez Efendi State Hospital

Özet

Fibroepitelyal polipler (FEP) üriner kanalın mezodermal dokusundan kaynaklanan iyi huylu tümörlerdir. Fibroepitelyal poliple rin etyolojisi bilinmemektedir. Hematüri ve obstrüktif üriner semptomlar en sık rastlanan bulgulardır. Üreterde dev bir polip olmasına rağmen, biz vakamızda herhangi bir semptom tespit etmedik. Ultrasonografi ile rastlantısal olarak sağ üreterde tespit edilen aseptomatik fibroepitelyal polip odusunu sunuyoruz.

Anahtar Kelimeler: Üreter, fibroepitelyal polip, tedavi, dev

Abstract

Fibroepithelial polyps (FEPs) are benign tumors of the urinary tract, arising from the mesodermal tissue. Etiology of fibroepithelial polyps is unknown. Hematuria and obstructive urinary symptoms are the most common findings. Although there was a giant polyp in the ureter in our case, we did not observe any symptom. We report a case of asymptomatic fibroepithelial polyp of the right ureter incidentally detected with ultrasonography.

Key Words: Ureter, fibroepithelial polyp, treatment, giant

Introduction

Fibroepithelial polyp is a benign neoplasm of mesodermal origin that arises in the wall of the renal pelvis, ureter, bladder or urethra. The etiology of FEP is unclear, congenital factors may play a role in children and chronic infection in adults. The most common symptoms are hematuria and obstructive urinary symptoms. It can be treated by endoscopically or open surgery (1).

Case report

A 73-year-old woman was admitted to our clinic with incidentally detected bladder mass on ultrasonography. Her physical examination was normal. She had history of breast cancer treated with right radical mastectomy and chemotherapy. Hemogram, blood biochemistry and urine analysis were in normal ranges. Abdominal ultrasound examination demonstrated a mass at the right side of the bladder wall. Intravenous urography was normal. (Figure1) Ureteral dilatation, filling defects and hydronephrosis were not detected. Cystoscopic examination was planned. There was a polypoid mass, in white col-

or, about 3 cm long and 1 cm wide, prolapsing into the bladder from the right ureteral orifice with ureteric jet-stream on cystoscopic examination.(Figure 2) Then the ureteroscope was inserted into the right ureter there was a stalked polyp and the polyp was grasped with forceps and excised (Figure 3). After this procedure double J stent was placed for 6 weeks. She was discharged on postoperative 1st day. Histopathologic examination of the tissue specimen revealed a fibroepithelial polyp. A double J stent removed 6 weeks postoperatively. At follow-up, the patient was asymptomatic and had no evidence of disease on cystourethroscopy after three months.

Discussion

Fibroepithelial polyps are considered the most common benign neoplasms of the ureter among other benign lesions such as lymphangiomas, leiomyomas and neurofibromas. They are often smoothly marginated, cylindrical and sessile (2). Because of their histologic organization, FEPs are classified as benign hamartomas; however, malignant degeneration and cystic transformation have also




Figure 1: Intravenous urography of the patient which is normal.

been reported (3).

Fibroepithelial polyps commonly present in adults in the third to fifth decades. Male-to-female ratio of 3:2. In adults, most FEPs occur in the ureter; 62% of these polyps are located in the upper ureter or ureteropelvic junction, 15% are in the renal pelvis, and a small percentage is in the bladder or posterior urethra. In our case fibroepithelial polyp was in the right middle ureter. Fibroepithelial polyps of the lower urinary tract usually occur in the posterior urethra, most often in children. (2) Kara et al reported incidentally detected ureteral fibroepithelial polyps in children (4). Fibroepithelial polyps usually appear as solitary polyps; however, rare cases of multiple and bilateral appearances have been reported (3, 5) The typical length is not more than five centimeters (6).

Although the etiology of FEPs is unclear, they are thought to be either congenital slow-growing lesions that develop as a result of chronic urothelial irritants, such as infection, inflammation, calculi, or obstruction. The most common symptoms of the polyps are hematuria and flank pain (2). Although there was a giant polyp in the right ureter in our case, we did not observe any symptom. The pain is characteristically intermittent and

colicky due to partial obstruction. Urinary frequency, dysuria, and pyuria are other less common findings (2). Sometimes the pain may be severe, torsion of the polyp, resulting in painful ischemia or infarction (7).

Fibroepithelial polyps appear to be long, smooth ureteral filling defects and hydronephrosis on intravenous urography or retrograde urograms (8). In our case ureteral filling defects and hydronephrosis were not detected by intravenous urography. We believe that this giant FEP did not cause hydronephrosis because it did not have an obstructive effect since it was moving in and out with urine efflux. It is important to distinguish FEPs from upper urinary tract carcinomas because management and prognosis can be significantly different. Debruyne et al reported that unnecessary nephroureterectomies were performed with FEP due to an uncertain pre-operative diagnosis (9). However, the definitive diagnosis can only be made by ureterorenoscopy and histopathologic examination

Management of ureteral FEPs include local coagulation by laser, polypectomy by ureteroscopy, and segmental resection with ureteroureterostomy or nephroureterectomy. In the past, treatment of FEP was excision of the polyp and reanastomosis with an open surgery. Recently endoscopic treatment with the ureteroscopes has become more popular. Usually, the polyps are grasped with forceps and resected through the ureteroscope. The holmium: YAG laser is another modality for endoscopic resection (3). Carey and Bird successfully ablated multiple polyps in ureter by using the holmium laser, and removed each polyp from the ureteral wall with grasping forceps




Figure 2: Polypoid mass prolapsing into the bladder from the right ureteral orifice




Figure 3: The mass was about 7 cm long and 1 cm wide

(10). Also ureteral stones are removed concurrently with a basket and a ureteral access sheath is used to facilitate the multiple passes of the ureteroscope and the removal of the polyps and stones from the proximal ureter. Percutaneous antegrade excision should be available for treating polyps in the upper urinary tract (3). Laparoscopic surgery might be preferred when the polyps are too large. When malignancy is suspected or when an endoscopic approach is not adequate, segmental resection of the ureter, ureteroureterostomy, pyeloplasty, or nephroureterectomy may be needed (4) The risk of recurrence, and frequency of follow-up are not clear. But some studies have suggested urine cytology, control ureteroscopy and intravenous urography in the follow up (4).

Conclusion

Fibroepithelial polyps of the ureter are rare benign neoplasm that can mimic other ureteral pathologies and lead to inaccurate diagnosis and treatment. Fibroepithelial polyps are commonly symptomatic and rarely asymptomatic that can be detected incidentally as in our case.

Fibroepithelial polyps appear as ureteral filling defects and hydronephrosis on intravenous urography or retrograde urograms but rarely it can be normal as in presented case. Fibroepithelial polyps should be considered in the diagnosis of ureteral filling defects to prevent aggressive treatments.

Kaynaklar

1. Yeh CC, Lin HC, Chen CC et al. Ureteral fibroepithelial polyp prolapsed into the bladder cavity suspending a bladder stone. J Urol 2001;165(5):1627-8.
2. Williams TR, Wagner BJ, Corse WR, Vestevich JC: Fibroepithelial polyps of the urinary tract. Abdom Imaging 2001; 27:217-21.
3. Lam JS, Bingham BJ, Gupta M: Endoscopic treatment of fibroepithelial polyps of the renal pelvis and ureter. Urology 2003; 62:810-3.
4. Cengiz Kara, Berkan Resorlu, Ural Oğuz, Ali Ünsal Incidentally detected ureteral fibroepithelial polyps in children: is endoscopic treatment of them really necessary? Int Urol Nephrol 2010; 42:1-5
5. Tekdogan UY, Canaklı F, Aslan Y, Han O, Gungor S, Atan A: Bilateral ureteral fibroepithelial polyps and review of the literature. Int J Urol 2005; 12:98-100.
6. Bellin ME, Springer O, Mourey-Gerosa I, et al. CT diagnosis of ureteral fibroepithelial polyps. Eur Radiol. 2002;12(1):125-8.
7. Niu ZB, Yang Y, Hou Y et al (2007) Ureteral polyps: an etiological factor of hydronephrosis in children that should not be ignored. Pediatr Surg Int 23(4):323-6.
8. Yagi S, Kawano Y, Gotanda T, Kitagawa T, Kawahara M, Nakagawa M, Higashi Y: Endoscopic treatment of a long fibroepithelial polyp. Int J Urol 2001; 8:467-9.
9. Debruyne FM, Moonen WA, Daenakindt AA, Delaere KPJ: Fibroepithelial polyp of the ureter. Urology 1980; 16:355-9
10. Carey RI, Bird VG. Endoscopic management of 10 separate fibroepithelial polyps arising in a single ureter. Urology. 2006 Feb;67(2):413-5.

Yazışma / Correspondence

Sacit Nuri Görgel, M.D.

Izmir Atatürk Training and Research Hospital,

1st.Urology Clinic, 35050, Izmir, Turkey

Tel: + 90 505 610 09 93

Fax: + 90 232 243 15 30

E-mail: sngorgel@hotmail.com