



Hesitant Intuitionistic Fuzzy Ideals of Γ -Semirings

Debabrata Mandal*

Department of Mathematics, Raja Peary Mohan College, Uttarpara, Hooghly, India

*Corresponding author, dmandaljumath@gmail.com

ABSTRACT: In this paper, we have defined hesitant intuitionistic fuzzy ideals, hesitant intuitionistic fuzzy bi-ideals and hesitant intuitionistic fuzzy quasi-ideals of a Γ -semiring and obtain some of their related properties. We have discussed some inter-relations between these ideals. We also obtain some characterizations of regular Γ -semiring.

Keywords – Hesitant, Intuitionistic fuzzy ideal, Cartesian product, Regular, Γ -semiring

1. Introduction

The notion of a semiring as first introduced by Vandiver (1934) in 1934 with two associative binary operations where one distributes over the other. But semirings had appeared in earlier studies on the theory of ideals of rings. In structure, semirings lie between semigroups and rings. The results which hold in rings but not in semigroups may hold in semirings, since semiring is a generalization of ring. The study of rings shows that multiplicative structure of ring is an independent of additive structure whereas in semiring multiplicative structure of semiring is not an independent of additive structure of semiring. The additive and the multiplicative structure of a semiring play an important role in determining the structure of a semiring. The theory of rings and theory of semigroups have considerable impact on the development of theory of semirings. Also, semirings has some applications to the theory of automata, formal languages, optimization theory and other branches of applied mathematics. Ideals of semiring play a central role in the structure theory and useful for many purposes. The notion of Γ -semiring was introduced by Rao (1995) as a generalization of Γ -ring as well as of semiring. Γ -semirings also includes ternary semirings and provide algebraic home to nonpositive cones of totally ordered rings.

The theory of fuzzy sets, proposed by Zadeh (1965), has provided a useful mathematical tool for describing the behavior of the systems that are too complex or ill-defined to admit precise mathematical analysis by classical methods and tools. As an important generalization of this notion, Atanassov (1993), introduced the concept of an intuitionistic fuzzy set where besides the degree of membership of each element there was considered a degree of non-membership with (membership value + non-membership value) ≤ 1 . In 2010, Torra (2010) introduced the hesitant fuzzy set which permits the membership degree of an element to a set to be represented by a set of possible values between 0 and 1. The hesitant fuzzy set therefore provides a more accurate representation of peoples hesitancy in stating their preferences over objects than the fuzzy set or its classical extensions. Hesitant fuzzy set theory has been applied to several practical problems, see (Jun and Song, 2014; Rodriguez, 2012; Torra and Narukawa, 2009; Wei 2012; Xia and Xu, 2011; Xia et al., 2013; Xu and Xia, 2011; Zhu et

al., 2012). Jun and Khan (2015) applied notion of hesitant fuzzy sets to semigroups and investigated several properties. Since then many researchers developed this idea.

The main aim of this paper is to combine the notion of intuitionistic fuzzy set and hesitant fuzzy set and apply it in studying several properties of Γ -semirings. We also obtain some characterizations.

2. Preliminaries

Definition 1. (Rao, 1995) Let S and Γ be two additive commutative semigroups with zero. Then S is called a Γ -semiring if there exists a mapping $S \times \Gamma \times S \rightarrow S$ ($((a, \alpha, b) \mapsto a\alpha b$) satisfying the following conditions:

- (i) $(a + b)\alpha c = a\alpha c + b\alpha c$,
- (ii) $a\alpha(b + c) = a\alpha b + a\alpha c$,
- (iii) $a(\alpha + \beta)b = a\alpha b + a\beta b$,
- (iv) $a\alpha(b\beta c) = (a\alpha b)\beta c$.

(v) $0_S \alpha a = 0_S = a\alpha 0_S$,

(vi) $a 0_\Gamma b = 0_S = b 0_\Gamma a$

for all $a, b, c \in S$ and for all $\alpha, \beta \in \Gamma$.

A subset A of a Γ -semiring S is called a left (resp. right) ideal of S if A is closed under addition and $S\Gamma A \subseteq A$ (resp. $A\Gamma S \subseteq A$). A subset A of a Γ -semiring S is called an ideal if it is both left and right ideal of S .

A subset A of a Γ -semiring S is called a quasi-ideal of S if A is closed under addition and $S\Gamma A \cap A\Gamma S \subseteq A$.

A subset A of a Γ -semiring S is called a bi-ideal if A is closed under addition and $A\Gamma S\Gamma A \subseteq A$.

Definition 2. (Zadeh, 1965) A fuzzy subset of a non-empty set X is defined as a function $\mu : X \rightarrow [0,1]$.

Definition 3. (Torra and Narukawa, 2009) Hesitant fuzzy set on S in terms of a function H that when applied to X returns a subset of $[0, 1]$.

Definition 4. (Atanassov, 1983) An intuitionistic fuzzy set defined on a non-empty set X , denoted by $\text{IFS}(X)$. An $\text{IFS}(X)$ is an object having the form $A = \langle \mu_A, \lambda_A \rangle = \{x, \mu_A(x), \lambda_A(x) | x \in X\}$ where the fuzzy sets μ_A and λ_A denote the degree of membership (namely $\mu_A(x)$) and the degree of non-membership (namely $\lambda_A(x)$) of each element $x \in X$ to the set A respectively, and $0 \leq \mu_A(x) + \lambda_A(x) \leq 1$ for all $x \in X$.

Throughout this paper unless otherwise mentioned S denotes the Γ -semiring and for any two set P and Q , we use the following notation:

$$\cap \{P, Q\} = P \cap Q \text{ and } \cup \{P, Q\} = P \cup Q.$$

3. Basic Definitions and Results of Hesitant Intuitionistic Fuzzy Ideals

In this section, the notion of hesitant intuitionistic fuzzy ideals in semiring are introduced and some of their basic properties are investigated.

Definition 5. Let X be a non-empty set. A hesitant intuitionistic fuzzy set A in X is a structure $A = \{ \langle x, \mu_A, \lambda_A \rangle : x \in X \}$ which briefly denoted as $A = \langle \mu_A, \lambda_A \rangle$.

Definition 6. Let $A = \langle \mu_A, \lambda_A \rangle$ be a non empty hesitant intuitionistic fuzzy subset of a Γ -semiring S . Then $\langle \mu_A, \lambda_A \rangle$ is called a hesitant intuitionistic fuzzy left ideal [hesitant intuitionistic fuzzy right ideal] of S if

- (i) $\mu_A(x + y) \supseteq \cap \{\mu_A(x), \mu_A(y)\}$, $\lambda_A(x + y) \subseteq \cup \{\lambda_A(x), \lambda_A(y)\}$ and
(ii) $\mu_A(x\gamma y) \supseteq \mu_A(y)$, $\lambda_A(x\gamma y) \subseteq \lambda_A(y)$
[respectively $\mu_A(x\gamma y) \supseteq \mu_A(x)$, $\lambda_A(x\gamma y) \subseteq \lambda_A(x)$].

for all $x, y \in S$ and $\gamma \in \Gamma$.

A hesitant intuitionistic fuzzy ideal of a Γ -semiring S is a non empty hesitant intuitionistic fuzzy subset of S which is a hesitant intuitionistic fuzzy left ideal as well as a hesitant intuitionistic fuzzy right ideal of S .

Note that if $\langle \mu_A, \lambda_A \rangle$ is a hesitant intuitionistic fuzzy left or right ideal of a Γ -semiring S , then $\mu_A(0) \supseteq \mu_A(x)$ and $\lambda_A(0) \subseteq \lambda_A(x)$ for all $x \in S$.

A hesitant intuitionistic fuzzy right ideal is defined similarly. By a hesitant intuitionistic fuzzy ideal $A = \langle \mu_A, \lambda_A \rangle$, we mean that $A = \langle \mu_A, \lambda_A \rangle$ is both hesitant intuitionistic fuzzy left and hesitant intuitionistic fuzzy right ideal.

Example 1. Let $S = \Gamma =$, the set of all non positive integers. Then S forms a Γ -semiring with usual addition and multiplication of integers. Define $A = \langle \mu_A, \lambda_A \rangle$ be a hesitant intuitionistic fuzzy subset of S as follows

$$\begin{aligned}\mu_A(x) &= [0,1] \text{ if } x = 0 \\ &= 0.2 \cup (0.3,0.8] \text{ if } x \text{ is even} \\ &= [0.4,0.7] \text{ if } x \text{ is odd}\end{aligned}$$

and

$$\begin{aligned}\lambda_A(x) &= \phi \text{ if } x = 0 \\ &= [0.5,0.7] \text{ if } x \text{ is even} \\ &= (0.1,0.8) \text{ if } x \text{ is odd}\end{aligned}$$

The hesitant intuitionistic fuzzy subset $\langle \mu_A, \lambda_A \rangle$ of S defined above is a hesitant intuitionistic fuzzy ideal S .

Throughout this section, we prove results only for hesitant intuitionistic fuzzy left ideals. Similar results can be obtained for hesitant intuitionistic fuzzy right ideals and hesitant intuitionistic fuzzy ideals.

Definition 7. The characteristic hesitant intuitionistic fuzzy set of A is defined to be the structure $\chi_A(x) = \langle x, \mu_{\chi_A}(x), \lambda_{\chi_A}(x) : x \in X \rangle$ where

$$\begin{aligned}\mu_{\chi_A}(x) &= [0,1] \text{ if } x \in A \\ &= \phi \text{ if } x \notin A.\end{aligned}$$

and

$$\begin{aligned}\lambda_{\chi_A}(x) &= \phi \text{ if } x \in A \\ &= [0,1] \text{ if } x \notin A.\end{aligned}$$

Theorem 1. Let I be a non-empty subset of a Γ -semiring S . Then I is a left ideal of S if and only if the characteristic function $\chi_I = \langle \mu_{\chi_I}, \lambda_{\chi_I} \rangle$ is a hesitant intuitionistic fuzzy left ideal of S .

PROOF. Assume that I is a left ideal of S and $x, y \in I$ and $\gamma \in \Gamma$. Suppose χ_I is not a hesitant intuitionistic fuzzy left ideal of S . So, $\mu_{\chi_I}(x + y) \subset \cap \{\mu_{\chi_I}(x), \mu_{\chi_I}(y)\}$ and $\lambda_{\chi_I}(x + y) \supset \cap \{\lambda_{\chi_I}(x), \lambda_{\chi_I}(y)\}$. Since $x, y \in I$, $\mu_{\chi_I}(x) = [0,1]$, $\mu_{\chi_I}(y) = [0,1]$, $\lambda_{\chi_I}(x) = \phi$ and $\lambda_{\chi_I}(y) =$

ϕ ; as a result, $\mu_{\chi_I}(x+y) = \phi$ and $\lambda_{\chi_I} = [0,1]$ i.e. $x+y \notin I$ - contradiction to the fact that I is a left ideal of S . Hence $\mu_{\chi_I}(x+y) \supseteq \cap \{\mu_{\chi_I}(x), \mu_{\chi_I}(y)\}$ and $\lambda_{\chi_I}(x+y) \subseteq \cap \{\lambda_{\chi_I}(x), \lambda_{\chi_I}(y)\}$. Similarly we can show that $\mu_{\chi_I}(x\gamma y) \supseteq \mu_{\chi_I}(y)$, $\lambda_{\chi_I}(x\gamma y) \subseteq \lambda_{\chi_I}(y)$.

Therefore $\chi_I = \langle \mu_{\chi_I}, \lambda_{\chi_I} \rangle$ is a hesitant intuitionistic fuzzy left ideal of S .

Conversely, assume that $\chi_I = \langle \mu_{\chi_I}, \lambda_{\chi_I} \rangle$ is a hesitant intuitionistic fuzzy left ideal of S for any subset I of S . Let $x, y \in I$ and $\gamma \in \Gamma$. Then $\mu_{\chi_I}(x) = [0,1]$, $\mu_{\chi_I}(y) = [0,1]$, $\lambda_{\chi_I}(x) = \phi$ and $\lambda_{\chi_I}(y) = \phi$. Now $\mu_{\chi_I}(x+y) \supseteq \cap \{\mu_{\chi_I}(x), \mu_{\chi_I}(y)\} = [0,1]$, $\lambda_{\chi_I}(x+y) \subseteq \cup \{\lambda_{\chi_I}(x), \lambda_{\chi_I}(y)\} = \phi$ and $\mu_{\chi_I}(x\gamma y) \supseteq \mu_{\chi_I}(y) = [0,1]$, $\lambda_{\chi_I}(x\gamma y) \subseteq \lambda_{\chi_I}(y) = \phi$. This implies $x+y, x\gamma y \in I$.

Hence I is a left ideal of S .

Definition 8. Let $A = \langle \mu_A, \lambda_A \rangle$ and $B = \langle \mu_B, \lambda_B \rangle$ be two hesitant intuitionistic fuzzy sets of a Γ -semiring S . Define intersection of A and B by

$$A \cap B = \langle \mu_A, \lambda_A \rangle \cap \langle \mu_B, \lambda_B \rangle = \langle \mu_A \cap \mu_B, \lambda_A \cup \lambda_B \rangle.$$

Proposition 1. Intersection of a non-empty collection of hesitant intuitionistic fuzzy left ideals is a hesitant intuitionistic fuzzy left ideal of S .

PROOF. Assume that $A_i = \langle \mu_{A_i}, \lambda_{A_i} \rangle : i \in I$ be a non-empty family of ideals of S . Let $x, y \in S$ and $\gamma \in \Gamma$. Then

$$\begin{aligned} (\cap_{i \in I} \mu_{A_i})(x+y) &= \cap_{i \in I} \{\mu_{A_i}(x+y)\} \supseteq \cap_{i \in I} \{\cap \{\mu_{A_i}(x), \mu_{A_i}(y)\}\} \\ &= \cap \{\cap_{i \in I} \mu_{A_i}(x), \cap_{i \in I} \mu_{A_i}(y)\} = \cap \{(\cap_{i \in I} \mu_{A_i})(x), (\cap_{i \in I} \mu_{A_i})(y)\}. \end{aligned}$$

$$\begin{aligned} (\cup_{i \in I} \lambda_{A_i})(x+y) &= \cup_{i \in I} \{\lambda_{A_i}(x+y)\} \subseteq \cup_{i \in I} \{\cup \{\lambda_{A_i}(x), \lambda_{A_i}(y)\}\} \\ &= \cup \{\cup_{i \in I} \lambda_{A_i}(x), \cup_{i \in I} \lambda_{A_i}(y)\} = \cup \{(\cup_{i \in I} \lambda_{A_i})(x), (\cup_{i \in I} \lambda_{A_i})(y)\}. \end{aligned}$$

Again

$$(\cap_{i \in I} \mu_{A_i})(x\gamma y) = \cap_{i \in I} \{\mu_{A_i}(x\gamma y)\} \supseteq \cap_{i \in I} \{\mu_{A_i}(y)\} = (\cap_{i \in I} \mu_{A_i})(y).$$

$$(\cup_{i \in I} \lambda_{A_i})(x\gamma y) = \cup_{i \in I} \{\lambda_{A_i}(x\gamma y)\} \subseteq \cup_{i \in I} \{\lambda_{A_i}(y)\} = (\cup_{i \in I} \lambda_{A_i})(y).$$

Hence $\cap_{i \in I} A_i = \langle \cap_{i \in I} \mu_{A_i}, \cup_{i \in I} \lambda_{A_i} \rangle : i \in I$ is a hesitant intuitionistic fuzzy left ideal of S .

Proposition 2. Let $f: R \rightarrow S$ be a morphism of Γ -semirings [2] and $A = \langle \mu_A, \lambda_A \rangle$ be a hesitant intuitionistic fuzzy left ideal of S , then $f^{-1}(A)$ is a hesitant intuitionistic fuzzy left ideal of R where $f^{-1}(A)(x) = \langle f^{-1}(\mu_A)(x), f^{-1}(\lambda_A)(x) \rangle = \langle \mu_A(f(x)), \lambda_A(f(x)) \rangle$.

PROOF. Suppose $f: R \rightarrow S$ be a morphism of Γ -semirings. Let $A = \langle \mu_A, \lambda_A \rangle$ and $r, s \in R$ and $\gamma \in \Gamma$. Then

$$\begin{aligned} f^{-1}(\mu_A)(r+s) &= \mu_A(f(r+s)) = \mu_A(f(r) + f(s)) \\ &\supseteq \cap \{\mu_A(f(r)), \mu_A(f(s))\} = \cap \{f^{-1}(\mu_A)(r), f^{-1}(\mu_A)(s)\} \end{aligned}$$

$$\begin{aligned} f^{-1}(\lambda_A)(r+s) &= \lambda_A(f(r+s)) = \lambda_A(f(r) + f(s)) \\ &\subseteq \cup \{\lambda_A(f(r)), \lambda_A(f(s))\} = \cup \{(f^{-1}(\lambda_A))(r), (f^{-1}(\lambda_A))(s)\} \end{aligned}$$

Again $(f^{-1}(\mu_A))(r\gamma s) = \mu_A(f(r\gamma s)) = \mu_A(f(r)\gamma f(s)) \supseteq \mu_A(f(s)) = (f^{-1}(\mu_A))(s)$.

$$(f^{-1}(\lambda_A))(r\gamma s) = \lambda_A(f(r\gamma s)) = \lambda_A(f(r)\gamma f(s)) \subseteq \lambda_A(f(s)) = (f^{-1}(\lambda_A))(s).$$

Thus $\langle f^{-1}(\mu_A)(x), f^{-1}(\lambda_A)(x) \rangle$ is a hesitant intuitionistic fuzzy left ideal of R .

Definition 9. Let $A = \langle \mu_A, \lambda_A \rangle$ and $B = \langle \mu_B, \lambda_B \rangle$ be hesitant intuitionistic fuzzy subsets of X . The cartesian product of A and B is defined by

$$\begin{aligned} (A \times B)(x, y) &= (\langle \mu_A, \lambda_A \rangle \times \langle \mu_B, \lambda_B \rangle)(x, y) = (\langle \mu_A \times \mu_B, \lambda_A \times \lambda_B \rangle)(x, y) \\ &= [\cap \{\mu_A(x), \mu_B(y)\}, \cup \{\lambda_A(x), \lambda_B(y)\}] \text{ for all } x, y \in X. \end{aligned}$$

Theorem 2. Let $A = \langle \mu_A, \lambda_A \rangle$ and $B = \langle \mu_B, \lambda_B \rangle$ be hesitant intuitionistic fuzzy left ideals of a Γ -semiring S . Then $A \times B$ is a hesitant intuitionistic fuzzy left ideal of the Γ -semiring $S \times S$.

PROOF. Let $(x_1, x_2), (y_1, y_2) \in S \times S$ and $\gamma \in \Gamma$. Then

$$\begin{aligned} (\mu_A \times \mu_B)((x_1, x_2) + (y_1, y_2)) &= (\mu_A \times \mu_B)(x_1 + y_1, x_2 + y_2) \\ &= \cap \{ \mu_A(x_1 + y_1), \mu_B(x_2 + y_2) \} \\ &\supseteq \cap \{ \cap \{ \mu_A(x_1), \mu_A(y_1) \}, \cap \{ \mu_B(x_2), \mu_B(y_2) \} \} \\ &= \cap \{ \cap \{ \mu_A(x_1), \mu_B(x_2) \}, \cap \{ \mu_A(y_1), \mu_B(y_2) \} \} \\ &= \cap \{ (\mu_A \times \mu_B)(x_1, x_2), (\mu_A \times \mu_B)(y_1, y_2) \} \end{aligned}$$

$$\begin{aligned} (\lambda_A \times \lambda_B)((x_1, x_2) + (y_1, y_2)) &= (\lambda_A \times \lambda_B)(x_1 + y_1, x_2 + y_2) \\ &= \cup \{ \lambda_A(x_1 + y_1), \lambda_B(x_2 + y_2) \} \\ &\subseteq \cup \{ \cup \{ \lambda_A(x_1), \lambda_A(y_1) \}, \cup \{ \lambda_B(x_2), \lambda_B(y_2) \} \} \\ &= \cup \{ \cup \{ \lambda_A(x_1), \lambda_B(x_2) \}, \cup \{ \lambda_A(y_1), \lambda_B(y_2) \} \} \\ &= \cup \{ (\lambda_A \times \lambda_B)(x_1, x_2), (\lambda_A \times \lambda_B)(y_1, y_2) \} \end{aligned}$$

and

$$\begin{aligned} (\mu_A \times \mu_B)((x_1, x_2)\gamma(y_1, y_2)) &= (\mu_A \times \mu_B)(x_1\gamma y_1, x_2\gamma y_2) \\ &= \cap \{ \mu_A(x_1\gamma y_1), \mu_B(x_2\gamma y_2) \} \\ &\supseteq \cap \{ \mu_A(y_1), \mu_B(y_2) \} \\ &= (\mu_A \times \mu_B)(y_1, y_2). \end{aligned}$$

$$\begin{aligned} (\lambda_A \times \lambda_B)((x_1, x_2)\gamma(y_1, y_2)) &= (\lambda_A \times \lambda_B)(x_1\gamma y_1, x_2\gamma y_2) \\ &= \cup \{ \lambda_A(x_1\gamma y_1), \lambda_B(x_2\gamma y_2) \} \\ &\subseteq \cup \{ \lambda_A(y_1), \lambda_B(y_2) \} \\ &= (\lambda_A \times \lambda_B)(y_1, y_2). \end{aligned}$$

Hence $A \times B$ is a hesitant intuitionistic fuzzy left ideal of $S \times S$.

3. Hesitant Intuitionistic Fuzzy Bi-Ideals and Quasi-Ideals

Definition 10. Let $A = \langle \mu_A, \lambda_A \rangle$ and $B = \langle \mu_B, \lambda_B \rangle$ be two hesitant intuitionistic fuzzy sets of a Γ -semiring S . Define composition of A and B by

$$A \circ B = \langle \mu_A, \lambda_A \rangle \circ \langle \mu_B, \lambda_B \rangle = \langle \mu_A \circ \mu_B, \lambda_A \circ \lambda_B \rangle$$

where

$$\begin{aligned} \mu_A \circ \mu_B(x) &= \cup \left[\cap_i \{ \mu_A(a_i), \mu_B(b_i) \} \right] \\ &\quad x = \sum_{i=1}^n a_i \gamma_i b_i \\ &= \phi, \text{ if } x \text{ cannot be expressed as above} \end{aligned}$$

and

$$\begin{aligned} (\lambda_A \circ \lambda_B)(x) &= \cap \left[\cup_i \{ \lambda_A(a_i), \lambda_B(b_i) \} \right] \\ &\quad x = \sum_{i=1}^n a_i \gamma_i b_i \\ &= [0, 1], \text{ if } x \text{ cannot be expressed as above} \end{aligned}$$

where $x, a_i, b_i \in S, \gamma_i \in \Gamma$ and $i = 1, \dots, n$.

Lemma 1. Let $A = \langle \mu_A, \lambda_A \rangle, B = \langle \mu_B, \lambda_B \rangle$ be two hesitant intuitionistic fuzzy ideal of a Γ -semiring S . Then $A \circ B \subseteq A \cap B \subseteq A, B$.

PROOF. Suppose $A = \langle \mu_A, \lambda_A \rangle, B = \langle \mu_B, \lambda_B \rangle$ be two hesitant intuitionistic fuzzy ideal of a Γ -semiring S . Then

$$\begin{aligned}
(\mu_A \circ \mu_B)(x) &= \bigcup \left\{ \bigcap_i \{ \mu_A(a_i), \mu_B(b_i) \} \right\} \\
&\quad x = \sum_{i=1}^n a_i \gamma_i b_i \\
&\quad \text{where } x, a_i, b_i \in S, \gamma_i \in \Gamma \text{ and } i = 1, \dots, n. \\
&\subseteq \bigcup \left\{ \bigcap_i \{ \mu_A(a_i) \} \right\} \\
&\subseteq \bigcup \left\{ \bigcap \left\{ \mu_A \left(\sum_{i=1}^n a_i \gamma_i b_i \right) \right\} \right\} = \mu_A(x)
\end{aligned}$$

$$\begin{aligned}
(\lambda_A \circ \lambda_B)(x) &= \bigcap \left\{ \bigcup_i \{ \lambda_A(a_i), \lambda_B(b_i) \} \right\} \\
&\quad x = \sum_{i=1}^n a_i \gamma_i b_i \\
&\quad \text{where } x, a_i, b_i \in S, \gamma_i \in \Gamma \text{ and } i = 1, \dots, n. \\
&= \bigcap \left\{ \bigcup_i \lambda_A(a_i) \right\} \\
&\supseteq \bigcap \left\{ \bigcup \left\{ \lambda_A \left(\sum_{i=1}^n a_i \gamma_i b_i \right) \right\} \right\} = \lambda_A(x)
\end{aligned}$$

Since this is true for every representation of x , $A \circ B \subseteq A$.

Similarly, we can prove that $A \circ B \subseteq B$.

Therefore $A \circ B \subseteq A \cap B$.

Hence the lemma.

Definition 11. A hesitant intuitionistic fuzzy subset $\langle \mu_A, \lambda_A \rangle$ of a Γ -semiring S is called hesitant intuitionistic fuzzy bi-ideal if for all $x, y, z \in S$ and $\alpha, \beta \in \Gamma$ we have

- (i) $\mu_A(x + y) \supseteq \bigcap \{ \mu_A(x), \mu_A(y) \}$, $\lambda_A(x + y) \subseteq \bigcup \{ \lambda_A(x), \lambda_A(y) \}$
- (ii) $\mu_A(x\alpha y) \supseteq \bigcap \{ \mu_A(x), \mu_A(y) \}$, $\lambda_A(x\alpha y) \subseteq \bigcup \{ \lambda_A(x), \lambda_A(y) \}$
- (iii) $\mu_A(x\alpha y\beta z) \supseteq \bigcap \{ \mu_A(x), \mu_A(z) \}$, $\lambda_A(x\alpha y\beta z) \subseteq \bigcup \{ \lambda_A(x), \lambda_A(z) \}$

Definition 12. A hesitant intuitionistic fuzzy subset $\langle \mu_A, \lambda_A \rangle$ of a Γ -semiring S is called hesitant intuitionistic fuzzy quasi-ideal if for all $x, y \in S$ we have

- (i) $\mu_A(x + y) \supseteq \bigcap \{ \mu_A(x), \mu_A(y) \}$, $\lambda_A(x + y) \subseteq \bigcup \{ \lambda_A(x), \lambda_A(y) \}$
- (ii) $(\mu_A \circ \mu_{\chi_S}) \cap (\mu_{\chi_S} \circ \mu_A) \subseteq \mu_A$, $(\lambda_A \circ \lambda_{\chi_S}) \cup (\lambda_{\chi_S} \circ \lambda_A) \supseteq \lambda_A$.

Theorem 3. A hesitant intuitionistic fuzzy subset $\langle \mu_A, \lambda_A \rangle$ of a Γ -semiring S is a hesitant intuitionistic fuzzy left ideal of S if and only if for all $x, y \in S$, we have

- (i) $\mu_A(x + y) \supseteq \bigcap \{ \mu_A(x), \mu_A(y) \}$, $\lambda_A(x + y) \subseteq \bigcup \{ \lambda_A(x), \lambda_A(y) \}$
- (ii) $\mu_{\chi_S} \circ \mu_A \subseteq \mu_A$, $\lambda_{\chi_S} \circ \lambda_A \supseteq \lambda_A$.

PROOF. Assume that $\langle \mu_A, \lambda_A \rangle$ is a hesitant intuitionistic fuzzy left ideal of S . Then it is sufficient to show that the condition (ii) is satisfied. Let $x \in S$. If x can be expressed as $x = \sum_{i=1}^n a_i \gamma_i b_i$, for $a_i, b_i \in S$, $\gamma_i \in \Gamma$ and $i=1, \dots, n$, then we have

$$\begin{aligned}
(\mu_{\chi_S} \circ \mu_A)(x) &= \bigcup \left[\bigcap_i \{ \mu_{\chi_S}(a_i), \mu_A(b_i) \} \right] \subseteq \bigcup \left[\bigcap_i \{ \mu_A(a_i \gamma_i b_i) \} \right] \\
&\quad x = \sum_{i=1}^n a_i \gamma_i b_i \qquad \qquad \qquad x = \sum_{i=1}^n a_i \gamma_i b_i \\
&\subseteq \bigcup \left[\bigcap \left\{ \mu_A \left(\sum_{i=1}^n a_i \gamma_i b_i \right) \right\} \right] = \mu_A(x). \\
(\lambda_{\chi_S} \circ \lambda_A)(x) &= \bigcap \left[\bigcup_i \{ \lambda_{\chi_S}(a_i), \lambda_A(b_i) \} \right] = \bigcap \left[\bigcup_i \{ \lambda_A(b_i) \} \right] \\
&\quad x = \sum_{i=1}^n a_i \gamma_i b_i \qquad \qquad \qquad x = \sum_{i=1}^n a_i \gamma_i b_i \\
&\geq \bigcap \left[\bigcup_i \{ \lambda_A(a_i \gamma_i b_i) \} \right] \geq \bigcap \left[\bigcup \left\{ \lambda_A \left(\sum_{i=1}^n a_i \gamma_i b_i \right) \right\} \right] = \lambda_A(x).
\end{aligned}$$

This implies that $\mu_{\chi_S} \circ \mu_A \subseteq \mu_A$, $\lambda_{\chi_S} \circ \lambda_A \supseteq \lambda_A$.

Conversely, assume that the given conditions hold. Then it is sufficient to show the second condition of the definition of hesitant intuitionistic fuzzy ideal. Let $x, y \in S$ and $\gamma \in \Gamma$.

Then we have

$$\begin{aligned}\mu_A(x\gamma y) &\supseteq (\mu_{\chi_S} \circ \mu_A)(x\gamma y) = \bigcup_{x\gamma y = \sum_{i=1}^n a_i \gamma_i b_i} [\bigcap \{ \mu_{\chi_S}(a_i), \mu_A(b_i) \}] \\ &\supseteq \mu_A(y) \text{ (since } x\gamma y = x\gamma y \text{)}.\end{aligned}$$

$$\begin{aligned}\lambda_A(x\gamma y) &\subseteq (\lambda_{\chi_S} \circ \lambda_A)(x\gamma y) = \bigcap_{x\gamma y = \sum_{i=1}^n a_i \gamma_i b_i} [\bigcup \{ \lambda_{\chi_S}(a_i), \lambda_A(b_i) \}] \\ &\subseteq \lambda_A(y) \text{ (since } x\gamma y = x\gamma y \text{)}.\end{aligned}$$

Hence $\langle \mu_A, \lambda_A \rangle$ is a hesitant intuitionistic fuzzy left ideal of S .

Theorem 4. Let $A = \langle \mu_A, \lambda_A \rangle$ and $B = \langle \mu_B, \lambda_B \rangle$ be a hesitant intuitionistic fuzzy right ideal and a hesitant intuitionistic fuzzy left ideal of a Γ -semiring S , respectively. Then $A \cap B$ is a hesitant intuitionistic fuzzy quasi-ideal of S .

PROOF. Let x, y be any element of S . Then

$$\begin{aligned}(\mu_A \cap \mu_B)(x + y) &= \bigcap \{ \mu_A(x + y), \mu_B(x + y) \} \\ &\supseteq \bigcap \{ \bigcap \{ \mu_A(x), \mu_A(y) \}, \bigcap \{ \mu_B(x), \mu_B(y) \} \} \\ &= \bigcap \{ \bigcap \{ \mu_A(x), \mu_B(x) \}, \bigcap \{ \mu_A(y), \mu_B(y) \} \} \\ &= \bigcap \{ (\mu_A \cap \mu_B)(x), (\mu_A \cap \mu_B)(y) \}.\end{aligned}$$

$$\begin{aligned}(\lambda_A \cup \lambda_B)(x + y) &= \bigcup \{ \lambda_A(x + y), \lambda_B(x + y) \} \\ &\subseteq \bigcup \{ \bigcup \{ \lambda_A(x), \lambda_A(y) \}, \bigcup \{ \lambda_B(x), \lambda_B(y) \} \} \\ &= \bigcup \{ \bigcup \{ \lambda_A(x), \lambda_B(x) \}, \bigcup \{ \lambda_A(y), \lambda_B(y) \} \} \\ &= \bigcup \{ (\lambda_A \cup \lambda_B)(x), (\lambda_A \cup \lambda_B)(y) \}.\end{aligned}$$

On the other hand, we have

$$((A \cap B) \circ \chi_S) \cap (\chi_S \circ (A \cap B)) \subseteq (A \circ \chi_S) \cap (\chi_S \circ B) \subseteq (A \cap B).$$

This completes the proof.

Lemma 2. Any hesitant intuitionistic fuzzy quasi-ideal of S is a hesitant intuitionistic fuzzy bi-ideal of S .

PROOF. Let $A = \langle \mu_A, \lambda_A \rangle$ be any hesitant intuitionistic fuzzy quasi-ideal of S . It is sufficient to show that $\mu_A(x\alpha y\beta z) \supseteq \bigcap \{ \mu_A(x), \mu_A(z) \}$, $\lambda_A(x\alpha y\beta z) \subseteq \bigcup \{ \lambda_A(x), \lambda_B(z) \}$ and $\mu_A(x\alpha y) \supseteq \bigcap \{ \mu_A(x), \mu_A(y) \}$, $\lambda_A(x\alpha y) \subseteq \bigcup \{ \lambda_A(x), \lambda_B(y) \}$ for all $x, y, z \in S$ and $\alpha, \beta \in \Gamma$.

In fact, by the assumption, we have

$$\begin{aligned}\mu_A(x\alpha y\beta z) &\supseteq ((\mu_A \circ \mu_{\chi_S}) \cap (\mu_{\chi_S} \circ \mu_A))(x\alpha y\beta z) \\ &= \bigcap \{ (\mu_A \circ \mu_{\chi_S})(x\alpha y\beta z), (\mu_{\chi_S} \circ \mu_A)(x\alpha y\beta z) \} \\ &= \bigcap \{ \bigcup_{x\alpha y\beta z = \sum_{i=1}^n a_i \gamma_i b_i} (\bigcap \{ \mu_A(a_i), \mu_{\chi_S}(b_i) \}), \bigcup_{x\alpha y\beta z = \sum_{i=1}^n a_i \gamma_i b_i} (\bigcap \{ \mu_{\chi_S}(a_i), \mu_A(b_i) \}) \} \\ &\supseteq \bigcap \{ \bigcap \{ \mu_A(x), \mu_{\chi_S}(z) \}, \bigcap \{ \mu_{\chi_S}(x), \mu_A(z) \} \} \text{ (since } x\alpha y\beta z = x\alpha y\beta z \text{)} \\ &= \bigcap \{ \mu_A(x), \mu_A(z) \}\end{aligned}$$

(i). $A \cap B \subseteq AoBoA$ for every hesitant intuitionistic fuzzy bi-ideal $A = \langle \mu_A, \lambda_A \rangle$ and every hesitant intuitionistic fuzzy ideal $B = \langle \mu_B, \lambda_B \rangle$ of S .

(ii). $A \cap B \subseteq AoBoA$ for every hesitant intuitionistic fuzzy quasi-ideal $A = \langle \mu_A, \lambda_A \rangle$ and every hesitant intuitionistic fuzzy ideal $B = \langle \mu_B, \lambda_B \rangle$ of S .

PROOF. Suppose S is a regular Γ -semiring and $A = \langle \mu_A, \lambda_A \rangle, B = \langle \mu_B, \lambda_B \rangle$ be any hesitant intuitionistic fuzzy bi-ideal and hesitant intuitionistic fuzzy ideal of S , respectively and x be any element of S . Since S is regular, there exist $a \in S$ and $\alpha, \beta \in \Gamma$ such that $x = x\alpha\beta x$.

$$\begin{aligned} (\mu_A \circ \mu_B \circ \mu_A)(x) &= \bigcup_{x=\sum_{i=1}^n a_i \gamma_i b_i} (\mu_A \circ \mu_B)(a_i, \mu_A(b_i)) \\ &\supseteq \bigcap \{(\mu_A \circ \mu_B)(x\alpha a), \mu_A(x)\} \\ &= \bigcap \left\{ \bigcup_{x\alpha a=\sum_{i=1}^n a_i \gamma_i b_i} \{(\mu_A(a_i), \mu_B(b_i))\}, \mu_A(x) \right\} \\ &\supseteq \bigcap \{ \bigcap \{ \mu_A(x), \mu_B(a\beta x\alpha a), \mu_A(x) \} \text{ (since } x\alpha a = x\alpha\beta x\alpha a) \} \\ &\supseteq \bigcap \{ \mu_A(x), \mu_B(x) \} = (\mu_A \cap \mu_B)(x). \end{aligned}$$

$$\begin{aligned} (\lambda_A \circ \lambda_B \circ \lambda_A)(x) &= \bigcap_{x=\sum_{i=1}^n a_i \gamma_i b_i} \{(\lambda_A \circ \lambda_B)(a_i), \lambda_A(b_i)\} \\ &\subseteq \bigcup \{(\lambda_A \circ \lambda_B)(x\alpha a), \lambda_A(x)\} \\ &= \bigcup \left\{ \bigcap \left(\bigcup_{x\alpha a=\sum_{i=1}^n a_i \gamma_i b_i} \{(\lambda_A(a_i), \lambda_B(b_i))\} \right), \lambda_A(x) \right\} \\ &\subseteq \bigcup \{ \bigcup \{ \lambda_A(x), \lambda_B(a\beta x\alpha a), \lambda_A(x) \} \text{ (since } x\alpha a = x\alpha\beta x\alpha a) \} \\ &\supseteq \bigcup \{ \lambda_A(x), \lambda_B(x) \} = (\lambda_A \cup \lambda_B)(x). \end{aligned}$$

(i) \Rightarrow (ii) follows from Lemma 2.

References

- Atanassov, K. T., 1983. Intuitionistic Fuzzy Sets. *Fuzzy Sets and Systems*, 20: 87-96.
- Dutta, T. K., Sardar, S. K., 2002. On matrix Γ -semirings. *East Journal of Mathematical Sciences*, 7(1): 17-31.
- Jun, Y. B., Khan, M., 2015. Hesitant Fuzzy Bi-Ideals in Semigroups. *Communications of the Korean Mathematical Society*, 30(3): 143-154.
- Jun, Y. B., Song, S. Z., 2014. Hesitant Fuzzy Set Theory Applied to Filters in MTL-Algebras. *Honam Mathematical Journal*, 36(4): 813-830.
- Rao, M. M. K., 1995. Γ -semirings-1. *Southeast Asian Bulletin of Mathematics*, 19: 49-54.
- Rodriguez, R. M., Martinez, L., Herrera, F., 2012. Hesitant Fuzzy Linguistic Term Sets for Decision Making. *IEEE Transactions on Fuzzy Systems*, 20(1): 109-119.
- Rosenfeld, A., 1971. Fuzzy Groups. *Journal of Mathematical Analysis and Applications*, 35: 512-517.
- Torra, V., 2010. Hesitant Fuzzy Sets. *International Journal of Intelligent Systems*, 25: 529-539.
- Torra, V., Narukawa, Y., 2009. On Hesitant Fuzzy Sets and Decision. Eds: Bezdek, J. C., Kacprzyk, J., Ruspini, E. H., Sugeno, M. *The 18th IEEE International Conference on Fuzzy Systems*, pp. 1378-1382, 20-24 August, Jeju Island, South Korea.
- Vandiver, H. S., 1934. Note on A Simple Type of Algebra in which Cancellation Law of Addition Does Not Hold. *Bulletin of the American Mathematical Society*, 40: 914-920.
- Wei, G., 2012. Hesitant Fuzzy Prioritized Operators and Their Application to Multiple Attribute Decision Making. *Knowledge-Based Systems*, 31: 176-182.
- Xia, M., Xu, Z. S., 2011. Hesitant Fuzzy Information Aggregation in Decision Making. *International Journal of Approximate Reasoning*, 52(3): 395-407.
- Xia, M., Xu, Z. S., Chen, N., 2013. Some Hesitant Fuzzy Aggregation Operators with Their Application in Group Decision Making. *Group Decision and Negotiation*, 22: 259-279.
- Xu, Z. S., Xia, M., 2011. Distance and Similarity Measures for Hesitant Fuzzy Sets. *Information Sciences*, 181(11): 2128-2138.
- Zhu, B., Xu, Z., Xia, M., 2012. Hesitant Fuzzy Geometric Bonferroni Means. *Information Sciences*, 205: 72-85.
- Zadeh, L.A., 1965. Fuzzy Sets. *Information and Control*, 8: 338-353.