

SOME INCLUSION RELATIONS BETWEEN VARIOUS SUBCLASSES OF PLANAR HARMONIC MAPPINGS INVOLVING CONFLUENT HYPERGEOMETRIC DISTRIBUTION SERIES

S. PORWAL, §

ABSTRACT. The purpose of the present paper is to establish connections between various subclasses of harmonic univalent functions by applying certain convolution operator involving Confluent Hypergeometric distribution series. To be more precise, we investigate such connections with Goodman-Rønning-type harmonic univalent functions in the open unit disc \mathbb{U} .

Keywords: Harmonic, Univalent functions, Confluent Hypergeometric Distribution Series.

AMS Subject Classification: 30C45.

1. INTRODUCTION

Let \mathcal{A} denote the class of functions f of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1)$$

which are analytic in the open unit disk $\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$ and satisfy the normalization condition $f(0) = f'(0) - 1 = 0$. Recently, Porwal [12] introduced a Poisson distribution series as

$$K(m, z) = z + \sum_{n=2}^{\infty} \frac{m^{n-1}}{(n-1)!} e^{-m} z^n.$$

The Poisson distribution series is a recent topic of study in Geometric Function Theory and co-relates G.F.T. with Probability distribution. It opened up a new and interesting direction of research in G.F.T. After the appearance of this paper some researchers ([1], [2], [15]) investigated hypergeometric distribution series, hypergeometric-type distribution series and confluent hypergeometric distribution series and give some beautiful applications on various subclasses of analytic and harmonic univalent functions. Noteworthy contribution in this direction are given in ([5], [6], [10], [14] and [16]). Very Recently, Porwal and Kumar [15] generalized the Poisson distribution series by introducing confluent

Sri Radhey Lal Arya Inter College, Ehan, Hathras-204101, (U.P.), India.

e-mail: saurabhjcb@rediffmail.com; ORCID: <https://orcid.org/0000-0003-0847-3550>.

§ Manuscript received: December 16, 2017; accepted: April 3, 2018.

TWMS Journal of Applied and Engineering Mathematics, Vol.9, No.4 © Işık University, Department of Mathematics, 2019; all rights reserved.

hypergeometric distribution series in the following way. For this purpose first we recall the following definitions.

The confluent hypergeometric function is given by the power series

$$F(a; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n(1)_n} z^n,$$

where a, c are complex numbers such that $c \neq 0, -1, -2, \dots$ and $(a)_n$ is the Pochhammer symbol defined in terms of the Gamma function, by

$$(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)} = \begin{cases} 1, & \text{if } n = 0 \\ a(a+1)\dots(a+n-1), & \text{if } n \in N = \{1, 2, 3, \dots\} \end{cases}$$

is convergent for all finite value of z .

Now we define for $a, c, m > 0$ such that the series

$$F(a; c; m) = \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n(1)_n} m^n$$

is convergent.

Porwal and Kumar [15] introduce the confluent hypergeometric distribution whose probability mass function is

$$\frac{(a)_n m^n}{(c)_n n! F(a; c; m)},$$

$n = 0, 1, 2, \dots$

If we put $a = c$ then it reduce to the Poisson distribution.

Now, we introduce a new series $I(a; c; m; z)$ whose coefficients are probabilities of confluent hypergeometric distribution

$$I(a; c; m; z) = z + \sum_{n=2}^{\infty} \frac{(a)_{n-1} m^{n-1}}{(c)_{n-1} (n-1)! F(a; c; m)} z^n,$$

where $a, c, m > 0$.

Denote by S_H the class of functions $f = h + \bar{g}$ that are harmonic univalent and sense-preserving in the open unit disc \mathbb{U} for which $f(0) = f_z(0) - 1 = 0$. Then for $f = h + \bar{g} \in S_H$, we may express the analytic functions h and g as

$$h(z) = z + \sum_{n=2}^{\infty} A_n z^n, \quad g(z) = \sum_{n=1}^{\infty} B_n z^n, \quad |B_1| < 1. \tag{2}$$

We also let the subclass S_H^0 of S_H

$$S_H^0 = \{f = h + \bar{g} \in S_H : g'(0) = B_1 = 0\}.$$

The classes S_H^0 and S_H were first studied in [7]. Also, we let $K_H^0, S_H^{*,0}$ and C_H^0 denote the subclasses of S_H^0 of harmonic functions which are, respectively, convex, starlike and close-to-convex in \mathbb{U} . Also, let T_H^0 be the class of sense-preserving, typically real harmonic functions $f = h + \bar{g}$ in S_H . For definitions and properties of these classes, one may refer to [7] or [8], (see also [19]).

For $0 \leq \gamma < 1$, let

$$N_H(\gamma) = \left\{ f \in S_H : \Re \left(\frac{f'(z)}{z'} \right) \geq \gamma, \quad z = re^{i\theta} \in \mathbb{U} \right\},$$

and

$$G_H(\gamma) = \left\{ f \in S_H : \Re \left\{ (1 + e^{i\alpha}) \frac{zf'(z)}{f(z)} - e^{i\alpha} \right\} \geq \gamma, \alpha \in R, z \in \mathbb{U} \right\},$$

where

$$z' = \frac{\partial}{\partial \theta} (z = re^{i\theta}), \quad f'(z) = \frac{\partial}{\partial \theta} f(re^{i\theta}).$$

Define

$$TN_H(\gamma) \equiv N_H(\gamma) \cap T \text{ and } TG_H(\gamma) \equiv G_H(\gamma) \cap T,$$

where T consists of the functions $f = h + \bar{g}$ in S_H so that h and g are of the form

$$h(z) = z - \sum_{n=2}^{\infty} |A_n| z^n, \quad g(z) = \sum_{n=1}^{\infty} |B_n| z^n. \quad (3)$$

The classes $N_H(\gamma)$, $TN_H(\gamma)$, $G_H(\gamma)$ and $TG_H(\gamma)$ were initially introduced and studied, respectively, in ([4], [17]). A function f in $G_H(\gamma)$ is called Goodman-Rønning-type harmonic univalent functions in \mathbb{U} .

Now, for $a_1, c_1, a_2, c_2, m_1, m_2 > 0$, we introduce the operator $\Omega(f) = \Omega \begin{pmatrix} a_1, & c_1, & m_1 \\ a_2, & c_2, & m_2 \end{pmatrix} f(z)$ for $f(z) \in S_H$ as

$$\Omega \begin{pmatrix} a_1, & c_1, & m_1 \\ a_2, & c_2, & m_2 \end{pmatrix} f(z) = I(a_1; c_1; m_1; z) * h(z) + \overline{I(a_2; c_2; m_2; z) * g(z)} = H(z) + \overline{G(z)},$$

where

$$H(z) = z + \sum_{n=2}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} A_n z^n, \quad G(z) = B_1 z + \sum_{n=2}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)! F(a_2; c_2; m_2)} B_n z^n. \quad (4)$$

Throughout this paper, we will frequently use the notation

$$\Omega(f) = \Omega \begin{pmatrix} a_1, & c_1, & m_1 \\ a_2, & c_2, & m_2 \end{pmatrix} f,$$

The purpose of the present paper is an attempt to interconnect between distribution function and Geometric Function Theory. Motivated by results on connections between various subclasses of analytic and harmonic univalent functions by using hypergeometric functions, generalized Bessel functions, Poisson distribution series, hypergeometric distribution series, hypergeometric-type distribution series and confluent hypergeometric distribution series (see [1], [2], [3], [9], [11], [13] and [18]), we establish a number of connections between the classes $G_H(\gamma)$, K_H^0 , $S_H^{*,0}$, C_H^0 and $N_H(\beta)$ by applying the convolution operator Ω .

2. MAIN RESULTS

In order to establish connections between harmonic convex functions and Goodman-Rønning-type harmonic univalent functions, we need following results in Lemma 2.1 and Lemma 2.2.

Lemma 2.1. (see [7], [8]). *If $f = h + \bar{g} \in K_H^0$ where h and g are given by (2) with $B_1 = 0$, then*

$$|A_n| \leq \frac{n+1}{2}, \quad |B_n| \leq \frac{n-1}{2}.$$

Lemma 2.2. (See [17]). Let $f = h + \bar{g}$ be given by (2). If $0 \leq \gamma < 1$ and

$$\sum_{n=2}^{\infty} (2n-1-\gamma) |A_n| + \sum_{n=1}^{\infty} (2n+1+\gamma) |B_n| \leq 1-\gamma, \quad (5)$$

then f is sense-preserving, Goodman-Rønning-type harmonic univalent functions in \mathbb{U} and $f \in G_H(\gamma)$.

Remark 2.1. In [17], it is also shown that $f = h + \bar{g}$ given by (3) is in the family $TG_H(\gamma)$, if and only if the coefficient condition (5) holds. Moreover, if $f \in TG_H(\gamma)$, then

$$|A_n| \leq \frac{1-\gamma}{2n-1-\gamma}, \quad n \geq 2,$$

$$|B_n| \leq \frac{1-\gamma}{2n+1+\gamma}, \quad n \geq 1.$$

Theorem 2.1. Let $a_1, c_1, a_2, c_2, m_1, m_2 > 0$. If for some $\gamma (0 \leq \gamma < 1)$, the inequality

$$\begin{aligned} & \frac{1}{F(a_1; c_1; m_1)} \left\{ 2 \frac{a_1(a_1+1)}{c_1(c_1+1)} m_1^2 F(a_1+2; c_1+2; m_1) + (7-\gamma) \frac{a_1}{c_1} m_1 F(a_1+1; c_1+1; m_1) \right. \\ & \left. + 2(1-\gamma) (F(a_1; c_1; m_1) - 1) \right\} + \frac{1}{F(a_2; c_2; m_2)} \left\{ 2 \frac{a_2(a_2+1)}{c_2(c_2+1)} m_2^2 F(a_2+2; c_2+2; m_2) \right. \\ & \left. + (5+\gamma) \frac{a_2}{c_2} m_2 F(a_2+1; c_2+1; m_2) \right\} \\ & \leq 2(1-\gamma), \end{aligned}$$

is satisfied then $\Omega(K_H^0) \subset G_H(\gamma)$.

Proof. Let $f = h + \bar{g} \in K_H^0$ where h and g are of the form (2) with $B_1 = 0$. We need to show that $\Omega(f) = H + \bar{G} \in G_H(\gamma)$, where H and G defined by (4) with $B_1 = 0$ are analytic functions in \mathbb{U} .

In view of Lemma 2.2, we need to prove that

$$P_1 \leq 1-\gamma,$$

where

$$P_1 = \sum_{n=2}^{\infty} (2n-1-\gamma) \left| \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} A_n \right| + \sum_{n=2}^{\infty} (2n+1+\gamma) \left| \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)! F(a_2; c_2; m_2)} B_n \right|. \quad (6)$$

In view of Lemma 2.1, we have

$$\begin{aligned}
 P_1 &\leq \frac{1}{2} \left[\sum_{n=2}^{\infty} (n+1)(2n-1-\gamma) \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} \right. \\
 &\quad \left. + \sum_{n=2}^{\infty} (n-1)(2n+1+\gamma) \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)! F(a_2; c_2; m_2)} \right] \\
 &= \frac{1}{2} \left[\sum_{n=2}^{\infty} \{2(n-1)(n-2) + (7-\gamma)(n-1) + 2(1-\gamma)\} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} \right. \\
 &\quad \left. + \sum_{n=2}^{\infty} \{2(n-2) + (5+\gamma)\} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-2)! F(a_2; c_2; m_2)} \right] \\
 &= \frac{1}{2} \left[\frac{1}{F(a_1; c_1; m_1)} \left\{ 2 \sum_{n=3}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-3)!} + (7-\gamma) \sum_{n=2}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-2)!} + 2(1-\gamma) \sum_{n=2}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)!} \right\} \right. \\
 &\quad \left. + \frac{1}{F(a_2; c_2; m_2)} \left\{ 2 \sum_{n=3}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-3)!} + (5+\gamma) \sum_{n=2}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-2)!} \right\} \right] \\
 &= \frac{1}{2} \left[\frac{1}{F(a_1; c_1; m_1)} \left\{ 2 \frac{a_1(a_1+1)}{c_1(c_1+1)} m_1^2 F(a_1+2; c_1+2; m_1) + (7-\gamma) \frac{a_1}{c_1} m_1 F(a_1+1; c_1+1; m_1) \right. \right. \\
 &\quad \left. \left. + 2(1-\gamma) (F(a_1; c_1; m_1) - 1) \right\} + \frac{1}{F(a_2; c_2; m_2)} \left\{ 2 \frac{a_2(a_2+1)}{c_2(c_2+1)} m_2^2 F(a_2+2; c_2+2; m_2) \right. \right. \\
 &\quad \left. \left. + (5+\gamma) \frac{a_2}{c_2} m_2 F(a_2+1; c_2+1; m_2) \right\} \right] \\
 &\leq 1 - \gamma,
 \end{aligned}$$

by given hypothesis.

This completes the proof of Theorem 2.1. □

Analogous to Theorem 2.1, we next find conditions of the classes $S_H^{*,0}, C_H^0$ with $G_H(\gamma)$. However we first need the following result which may be found in [7], [8].

Lemma 2.3. *If $f = h + \bar{g} \in S_H^{*,0}$ or C_H^0 , where h and g are given by (2) with $B_1 = 0$, then*

$$|A_n| \leq \frac{(2n+1)(n+1)}{6}, \quad |B_n| \leq \frac{(2n-1)(n-1)}{6}.$$

Theorem 2.2. *Let $a_1, c_1, a_2, c_2, m_1, m_2 > 0$. If for some $\gamma(0 \leq \gamma < 1)$ and the inequality*

$$\begin{aligned}
 &\frac{1}{F(a_1; c_1; m_1)} \left\{ 4 \frac{a_1(a_1+1)(a_1+2)}{c_1(c_1+1)(c_1+2)} m_1^3 F(a_1+3; c_1+3; m_1) + (28-2\gamma) \frac{a_1(a_1+1)}{c_1(c_1+1)} m_1^2 F(a_1+2; c_1+2; m_1) \right. \\
 &\quad \left. + (39-9\gamma) \frac{a_1}{c_1} F(a_1+1; c_1+1; m_1) m_1 + 6(1-\gamma) (F(a_1; c_1; m_1) - 1) \right\} \\
 &\quad + \frac{1}{F(a_2; c_2; m_2)} \left\{ 4 \frac{a_2(a_2+1)(a_2+2)}{c_2(c_2+1)(c_2+2)} m_2^3 F(a_2+3; c_2+3; m_2) \right. \\
 &\quad \left. + (20+2\gamma) \frac{a_2(a_2+1)}{c_2(c_2+1)} m_2^2 F(a_2+2; c_2+2; m_2) \right. \\
 &\quad \left. + (15+3\gamma) \frac{a_2}{c_2} m_2 F(a_2+1; c_2+1; m_2) \right\}
 \end{aligned}$$

$$\leq 6(1 - \gamma),$$

is satisfied, then

$$\Omega(S_H^{*,0}) \subset G_H(\gamma) \text{ and } \Omega(C_H^0) \subset G_H(\gamma).$$

Proof. Let $f = h + \bar{g} \in S_H^{*,0}$ (C_H^0 where h and g are given by (2) with $B_1 = 0$). We need to show that $\Omega(f) = H + \bar{G} \in G_H(\gamma)$, where H and G defined by (4) with $B_1 = 0$ are analytic functions in \mathbb{U} . In view of Lemma 2.2, it is enough to show that $P_1 \leq 1 - \gamma$, where P_1 is given by (6).

In view of Lemma 2.3, we have

$$\begin{aligned} P_1 &\leq \frac{1}{6} \left[\sum_{n=2}^{\infty} (2n+1)(n+1)(2n-1-\gamma) \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} \right. \\ &\quad \left. + \sum_{n=2}^{\infty} (2n-1)(n-1)(2n+1+\gamma) \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)! F(a_2; c_2; m_2)} \right] \\ &= \frac{1}{6} \left[\frac{1}{F(a_1; c_1; m_1)} \left\{ \sum_{n=2}^{\infty} \{4(n-1)(n-2)(n-3) + (28-2\gamma)(n-1)(n-2) \right. \right. \\ &\quad \left. \left. + (39-9\gamma)(n-1) + 6(1-\gamma)\} \right\} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)!} \right. \\ &\quad \left. + \frac{1}{F(a_2; c_2; m_2)} \left\{ \sum_{n=2}^{\infty} \{4(n-2)(n-3) + (20+2\gamma)(n-2) + (15+3\gamma)\} \right\} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-2)!} \right] \\ &= \frac{1}{6} \left[\frac{1}{F(a_1; c_1; m_1)} \left\{ 4 \sum_{n=4}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-4)!} + (28-2\gamma) \sum_{n=3}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-3)!} \right. \right. \\ &\quad \left. \left. + (39-9\gamma) \sum_{n=2}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-2)!} + 6(1-\gamma) \sum_{n=2}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)!} \right\} + \frac{1}{F(a_2; c_2; m_2)} \right. \\ &\quad \left. \left\{ 4 \sum_{n=4}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-4)!} + (20+2\gamma) \sum_{n=3}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-3)!} + (15+3\gamma) \sum_{n=2}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-2)!} \right\} \right] \\ &= \frac{1}{6} \left[\frac{1}{F(a_1; c_1; m_1)} \left\{ 4 \frac{a_1(a_1+1)(a_1+2)}{c_1(c_1+1)(c_1+2)} m_1^3 F(a_1+3; c_1+3; m_1) \right. \right. \\ &\quad \left. \left. + (28-2\gamma) \frac{a_1(a_1+1)}{c_1(c_1+1)} m_1^2 F(a_1+2; c_1+2; m_1) \right. \right. \\ &\quad \left. \left. + (39-9\gamma) \frac{a_1}{c_1} F(a_1+1; c_1+1; m_1) m_1 + 6(1-\gamma) (F(a_1; c_1; m_1) - 1) \right\} \right. \\ &\quad \left. + \frac{1}{F(a_2; c_2; m_2)} \left\{ 4 \frac{a_2(a_2+1)(a_2+2)}{c_2(c_2+1)(c_2+2)} m_2^3 F(a_2+3; c_2+3; m_2) \right. \right. \\ &\quad \left. \left. + (20+2\gamma) \frac{a_2(a_2+1)}{c_2(c_2+1)} m_2^2 F(a_2+2; c_2+2; m_2) + (15+3\gamma) \frac{a_2}{c_2} m_2 F(a_2+1; c_2+1; m_2) \right\} \right] \\ &\leq 1 - \gamma, \end{aligned}$$

follows from the given condition. \square

In order to determine connection between $TN_H(\beta)$ and $G_H(\gamma)$, we need the following result in Lemma 2.4.

Lemma 2.4. (See [4]). Let $f = h + \bar{g}$ where h and g are given by (2) with $B_1 = 0$, and suppose that $0 \leq \beta < 1$. Then

$$f \in TN_H(\beta) \Leftrightarrow \sum_{n=2}^{\infty} n |A_n| + \sum_{n=2}^{\infty} n |B_n| \leq 1 - \beta.$$

Remark 2.2. If $f \in TN_H(\beta)$, then $|A_n| \leq \frac{1-\beta}{n}$, $n \geq 2$ and $|B_n| \leq \frac{1-\beta}{n}$, $n \geq 1$.

Theorem 2.3. If $a_1, c_1, a_2, c_2 > 1, m_1, m_2 > 0$. If for some $\beta(0 \leq \beta < 1)$ and $\gamma(0 \leq \gamma < 1)$ and the inequality

$$\begin{aligned} & (1 - \beta) \left[\frac{1}{F(a_1; c_1; m_1)} \left\{ 2(F(a_1; c_1; m_1) - 1) - \frac{(1 + \gamma)(c_1 - 1)}{m_1(a_1 - 1)} \left(F(a_1 - 1; c_1 - 1; m_1) - 1 - \frac{(a_1 - 1)}{(c_1 - 1)} m_1 \right) \right\} \right. \\ & \left. + 2(F(a_2; c_2; m_2) - 1) + \frac{(1 + \gamma)(c_2 - 1)}{m_2(a_2 - 1)} \left(F(a_2 - 1; c_2 - 1; m_2) - 1 - \frac{(a_2 - 1)}{(c_2 - 1)} m_2 \right) \right] \\ & + (3 + \gamma) |B_1| \\ & \leq 1 - \gamma \end{aligned}$$

is satisfied then

$$\Omega(TN_H(\beta)) \subset G_H(\gamma).$$

Proof. Let $f = h + \bar{g} \in TN_H(\beta)$ where h and g are given by (2). In view of Lemma 2.2, it is enough to show that $P_2 \leq 1 - \gamma$, where P_2 is given by the following expression

$$P_2 = \sum_{n=2}^{\infty} (2n - 1 - \gamma) \left| \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} A_n \right| + \sum_{n=2}^{\infty} (2n + 1 + \gamma) \left| \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)! F(a_2; c_2; m_2)} B_n \right| + (3 + \gamma) |B_1|. \tag{7}$$

Using Remark 2.2, we have

$$\begin{aligned} P_2 & \leq (1 - \beta) \left[\sum_{n=2}^{\infty} \left\{ 2 - \frac{(1 + \gamma)}{n} \right\} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} \right. \\ & \left. + \sum_{n=2}^{\infty} \left\{ 2 + \frac{(1 + \gamma)}{n} \right\} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)! F(a_2; c_2; m_2)} \right] + (3 + \gamma) |B_1| \\ & = (1 - \beta) \left[\frac{1}{F(a_1; c_1; m_1)} \left\{ 2 \sum_{n=2}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)!} - (1 + \gamma) \sum_{n=2}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} n!} \right\} \right. \\ & \left. + \frac{1}{F(a_2; c_2; m_2)} \left\{ 2 \sum_{n=2}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)!} + (1 + \gamma) \sum_{n=2}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} n!} \right\} \right] + (3 + \gamma) |B_1| \\ & = (1 - \beta) \left[\frac{1}{F(a_1; c_1; m_1)} \{ 2(F(a_1; c_1; m_1) - 1) \right. \\ & \left. - \frac{(1 + \gamma)(c_1 - 1)}{m_1(a_1 - 1)} \left(F(a_1 - 1; c_1 - 1; m_1) - 1 - \frac{(a_1 - 1)}{(c_1 - 1)} m_1 \right) \right\} \\ & \left. + 2(F(a_2; c_2; m_2) - 1) + \frac{(1 + \gamma)(c_2 - 1)}{m_2(a_2 - 1)} \left(F(a_2 - 1; c_2 - 1; m_2) - 1 - \frac{(a_2 - 1)}{(c_2 - 1)} m_2 \right) \right] \\ & + (3 + \gamma) |B_1| \\ & \leq 1 - \gamma, \end{aligned}$$

by the given hypothesis. □

In next theorem, we establish connections between $TG_H(\gamma)$ and $G_H(\gamma)$.

Theorem 2.4. Let $a_1, c_1, a_2, c_2, m_1, m_2 > 0$. If for some $\gamma(0 \leq \gamma < 1)$ the inequality

$$\frac{1}{F(a_1; c_1; m_1)} (F(a_1; c_1; m_1) - 1) + \frac{1}{F(a_2; c_2; m_2)} (F(a_2; c_2; m_2) - 1) \leq 1 - \frac{3 + \gamma}{1 - \gamma} |B_1| \quad (8)$$

is satisfied, then $\Omega(TG_H(\gamma)) \subset G_H(\gamma)$.

Proof. Making use of Lemma 2.2, we only need to prove that $P_2 \leq 1 - \gamma$, where P_2 is given by (7). Using Remark 2.1, it follows that

$$\begin{aligned} P_2 &= \sum_{n=2}^{\infty} (2n - 1 - \gamma) \left| \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} A_n \right| \\ &\quad + \sum_{n=2}^{\infty} (2n + 1 + \gamma) \left| \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)! F(a_2; c_2; m_2)} B_n \right| + (3 + \gamma) |B_1| \\ &\leq (1 - \gamma) \left[\sum_{n=2}^{\infty} \frac{(a_1)_{n-1} m_1^{n-1}}{(c_1)_{n-1} (n-1)! F(a_1; c_1; m_1)} + \sum_{n=2}^{\infty} \frac{(a_2)_{n-1} m_2^{n-1}}{(c_2)_{n-1} (n-1)! F(a_2; c_2; m_2)} \right] + (3 + \gamma) |B_1| \\ &= (1 - \gamma) \left[\frac{1}{F(a_1; c_1; m_1)} (F(a_1; c_1; m_1) - 1) + \frac{1}{F(a_2; c_2; m_2)} (F(a_2; c_2; m_2) - 1) \right] + (3 + \gamma) |B_1| \\ &\leq (1 - \gamma), \end{aligned}$$

by the given condition and this completes the proof. \square

In next theorem, we present conditions on the parameters $a_1, c_1, a_2, c_2, m_1, m_2$ and obtain a characterization for operator Ω which maps $TG_H(\gamma)$ on to itself.

Theorem 2.5. If $a_1, c_1, a_2, c_2, m_1, m_2 > 0$ and $\gamma(0 \leq \gamma < 1)$. Then

$$\Omega(TG_H(\gamma)) \subset TG_H(\gamma),$$

if and only if, the condition (8) is satisfied.

Proof. The proof of above theorem is similar to that of Theorem 2.4. Therefore we omits the details involved. \square

Remark 2.3. If we put $a_1 = c_1; a_2 = c_2$ then we obtain the corresponding results for the Poisson distribution series.

Acknowledgement

The author is thankful to the referee for his/her valuable comments and suggestions.

REFERENCES

- [1] Ahmad, M.S., Mehmood, Q., Nazeer, W. and Haq, A.U., (2015), An application of a Hypergeometric distribution series on certain analytic functions, Sci. Int.(Lahore), 27(4), pp. 2989-2992.
- [2] Ahmad, D., (2014), Mapping properties of some convolution operator associated with some special function, M.Phil. Dissertation, C.S.J.M. University, Kanpur, India.
- [3] Ahuja, O.P., (2008), Connections between various subclasses of planar harmonic mappings involving hypergeometric functions, Appl. Math. Comput., 198 (1), pp. 305-316.
- [4] Ahuja, O.P., and Jahangiri, J.M., (2002), Noshiro-type harmonic univalent functions, Sci. Math. Jpn., 6 (2), pp. 253-259.
- [5] Altinkaya, S. and Yalcin, S., (2018), Poisson distribution series for analytic univalent functions, Complex Anal. Oper. Theory, 12(5), pp. 1315-1319.
- [6] Altinkaya, S. and Yalcin, S., (2017), Poisson distribution series for certain subclasses of starlike functions with negative coefficients, Annal. Oradea Univ. Math. Fasci., 24 (2), pp. 5-8.
- [7] Clunie, J. and Sheil-Small, T., (1984), Harmonic univalent functions, Ann. Acad. Sci. Fen. Series A.I. Math., 9, pp. 3-25.

- [8] Duren, P., (2004), Harmonic mappings in the plane, Vol. 156, Cambridge University Press, Cambridge.
- [9] Mondal, S.R. and Swaminathan, A., (2012), Geometric properties of Generalized Bessel functions, Bull. Malays. Math. Sci. Soc., 35(1), pp. 179-194.
- [10] Murugusundaramoorthy, G., Vijaya, K. and Porwal, S., (2016), Some inclusion results of certain subclasses of analytic functions associated with Poisson distribution series, Hacettepe J. Math. Stat., 45(4), pp. 1101-1107.
- [11] Ponnusamy, S. and Rønning, F., (1998), Starlikeness properties for convolution involving hypergeometric series, Ann. Univ. Mariae Curie-Sklodowska, LII, 1(16), pp. 141-155.
- [12] Porwal, S., (2014), An application of a Poisson distribution series on certain analytic functions, J. Complex Anal., Art. ID 984135, pp. 1-3.
- [13] Porwal, S., (2015), Some connections between various subclasses of planar harmonic mappings involving generalized Bessel functions, Afr. Mat., 26(5-6), pp. 997-1008.
- [14] Porwal, S. and Kumar, M., (2016), A unified study on starlike and convex functions associated with Poisson distribution series, Afr. Mat., 27(5-6), pp. 1021-1027.
- [15] Porwal, S. and Kumar, S., (2017), Confluent hypergeometric distribution and its applications on certain classes of univalent functions, Afr. Mat., 28, pp. 1-8.
- [16] Porwal, S. and Srivastava, D., (2017), Harmonic starlikeness and convexity of integral operators generated by Poisson distribution series, Math. Moravica, 21, pp. 51-60.
- [17] Rosy, T., Stephen, B.A., Subramanian, K.G. and Jahangiri, J.M., (2001), Goodman-Rønning-type harmonic univalent functions, Kyungpook Math. J., 41 (1), pp. 45-54.
- [18] Sharma, A.K., Porwal, S. and Dixit, K.K., (2013), Class mappings properties of convolutions involving certain univalent functions associated with hypergeometric functions, Electronic J. Math. Anal. Appl., 1(2), pp. 326-333
- [19] Yalcin, S., Öztürk, M., and Yamankaradeniz, M., (2007), On subclass of Salagean-type harmonic univalent functions, J. Inequal. Pure Appl. Math., 8 (2), Art. 54, pp. 1-9.



Dr. Saurabh Porwal is currently working as a lecturer of mathematics in Sri Radhey Lal Arya Inter College, Ehan, Hathras. Formely, He was assistant professor in the Department of Mathematics, UIET, CSJM University, Kanpur, India. He has published several research papers in various reputed national and international journals.
