

## MEASUREMENT OF PLANARITY IN PRODUCT BIPOLAR FUZZY GRAPHS

SUMERA NAZ<sup>1</sup>, SAMINA ASHRAF<sup>2</sup>, HOSSEIN RASHMANLOU<sup>3</sup>, §

**ABSTRACT.** Bipolar fuzzy set theory provides a basis for bipolar cognitive modeling and multiagent decision analysis, where in some situations, the product operator may be preferred to the min operator, from theoretical and experimental aspects. In this paper, the definition of product bipolar fuzzy graphs (PBFs) in [16] is modified. The concepts of product bipolar fuzzy multigraphs (PBFMGs), product bipolar fuzzy planar graphs (PBFPGs) and product bipolar fuzzy dual graphs (PBFDGs) are introduced and their properties are investigated. Meanwhile, the product bipolar fuzzy planarity value of PBFPG is introduced. The relation between PBFPG and PBFDG is also established. Finally, an application of the proposed concepts is provided.

**Keywords:** Product bipolar fuzzy graph, product bipolar fuzzy multigraph, product bipolar fuzzy planar graph, product bipolar fuzzy dual graph.

**AMS Subject Classification:** 05C99

### 1. INTRODUCTION

In 1736, Euler first introduced the theory of planar graphs, by finding an important formula relating the numbers of vertices, edges and faces of polyhedrons, which can be represented by planar graphs. In real-world applications, planar graphs arise quite naturally such as electric printed circuits, railway or road maps, chemical molecules, etc.

In 1965, Zadeh [20] originally introduced the concept of fuzzy set, characterized by a membership function in  $[0, 1]$ , which is very useful in dealing with uncertainty and vagueness. Yager [19] initiated the concept of fuzzy multisets (fuzzy bag). Bipolar fuzzy sets (BFSs) as a generalization of fuzzy sets were first introduced by Zhang [21, 22]. The range of membership degree of BFSs is  $[-1, 1]$ . In a BFS, the membership degree  $(0, 1]$  of an element indicates that the element satisfies the property, the membership degree  $[-1, 0)$  of an element indicates that the element satisfies the implicit counter-property and the membership degree 0 of an element means that the element is irrelevant to the

---

<sup>1</sup> Department of Mathematics, Minhaj University, Lahore, Pakistan.

e-mail: [nsumeranaz@gmail.com](mailto:nsumeranaz@gmail.com); ORCID: <https://orcid.org/0000-0001-6550-9607>.

<sup>2</sup> Quality Enhancement Cell, Lahore College for Women University, Lahore, Pakistan.

e-mail: [saminaa561@gmail.com](mailto:saminaa561@gmail.com); ORCID: <https://orcid.org/0000-0001-6565-7917>.

<sup>3</sup> Department of Computer Science, University College of Rouzbahan, Sari, Iran.

e-mail: [rashmanlou.1987@gmail.com](mailto:rashmanlou.1987@gmail.com); ORCID: <https://orcid.org/0000-0001-8717-9530>.

§ Manuscript received: January 06, 2017; accepted: February 28, 2017.

TWMS Journal of Applied and Engineering Mathematics, Vol.9, No.2 © Işık University, Department of Mathematics, 2019; all rights reserved.

corresponding property [10]. BFSs have received great attention from researchers and have been applied to many fields, such as artificial intelligence, information science, decision science, economics, medical science, social science, computer science, and neural science.

Graphs are basically the bonding of objects. To emphasis on a real life problem, the objects are being bonded by some relations, such as friendship is the bonding of people. But when the ambiguousness or uncertainty in bonding exists, then the corresponding graph can be modeled as fuzzy graph model. In 1975, Rosenfeld [17] put forth the concept of fuzzy graphs and developed its structure, obtaining analogs of several basic graph theoretical concepts. Further, the concept of fuzzy dual graph was initiated by Jabbar et al. [9]. Akram et al. [1] originally proposed the concept of bipolar fuzzy graphs, and made a lot of studies on this extension of fuzzy graphs [2, 3]. Rashmanlou et al. [16] introduced the concept of product bipolar fuzzy graphs. Naz et al. put forward several new concepts concerning the extended structures of fuzzy graphs and provided their applications in decision making [6, 7, 11, 12, 13, 14]. In modern age, pipelines, subway tunnels, metro lines are essential in a city planning, where routes without crossing are perfect for safety. But, due to lack of space, crossing of such lines is allowed. Since crossing between congested (strong) route and non-congested (weak) route is more safe than the crossing between two congested (strong) routes. The terms ‘strong route’ and ‘weak route’ lead strong edge and weak edge of a fuzzy graph, respectively and the permission of crossing between strong and weak edges leads to the concept of fuzzy planar graph. Recently, fuzzy planar graph and its extensions [4, 5, 15, 18] have been studied. Definition of PBFGs in [16] is valid only for the product of even number of negative-membership values. But for odd number of negative-membership values this definition is invalid. So, in this paper, a modified definition of PBFGs is given.

The rest of the paper is organized as follows: Section 2 reviews basic concepts related to BFSs and PBFGs. Section 3 proposes the concepts of multigraphs, planar graphs and dual graphs under bipolar fuzzy environment based on the product operator. In Section 4, an application of the proposed concepts is provided. Section 5 ends up the paper with some concluding remarks.

## 2. PRELIMINARIES

In this section, we recall some basic concepts which are necessary for this paper.

A graph is a pair of sets  $G = (V, E)$ , satisfying  $E \subseteq V \times V$ . The elements of  $V$  are the vertices and the elements of  $E$  are the edges of the graph  $G$ . For a vertex set  $V$  of  $G$ , define an equivalence relation  $\sim$  on  $V \times V - \{xx \mid x \in V\}$  as follows:  $x_1x_2 \sim x'_1x'_2$  if and only if either  $x_1x_2 = x'_1x'_2$  or  $x_1 = x'_2$  and  $x_2 = x'_1$ . The quotient set obtained in this way is denoted by  $\widetilde{V^2}$ . A graph can be drawn in many different ways. A planar graph is a particular diagram which can be drawn on the plane so that no two edges intersect geometrically except at a vertex at which they are both incident. Any plane drawing of a planar graph  $G$  divides the plane into a set of regions, called faces. In any planar graph, the unbounded face is called an infinite face. If a cycle in a planar graph is a boundary of a face, then it is a facial cycle. A graph is said to be a non-planar if it cannot be drawn without crossing. The minimum number of crossings that can occur when  $G$  is drawn in the plane is called the crossing number  $cr(G)$  of a graph  $G$ .

**Euler’s Formula:** Let  $G$  be a connected planar graph with order  $n$ , size  $m$  and  $f$  faces. Then  $n - m + f = 2$ .

A dual graph of a planar graph  $G$  is constructed as follows: place a vertex in each face of

$G$  and if two faces have an edge  $e$  in common, join the corresponding vertices by an edge  $e'$  crossing only  $e$ .

**Definition 2.1.** [19] A fuzzy multiset  $X$  drawn from non-empty set  $V$  is characterized by a function, ‘count membership’ of  $X$  denoted by  $CM_X$  such that  $CM_X : V \rightarrow Q$ , where  $Q$  is the set of all crisp multisets drawn from the unit interval  $[0, 1]$ . Then for any  $x \in V$ , the value  $CM_X(x)$  is a crisp multiset drawn from  $[0, 1]$ .

**Definition 2.2.** [15] Let  $V$  be a non-empty set and  $\sigma : V \rightarrow [0, 1]$  be a mapping. Let  $\mu = \{(xy, \mu(xy)_k) \mid k = 1, 2, \dots, m_{xy}, xy \in V \times V\}$  be a fuzzy multiset of  $V \times V$  such that  $\mu(xy)_k \leq \sigma(x) \wedge \sigma(y)$  for all  $k = 1, 2, \dots, m_{xy}$ , where  $m_{xy} = \max\{k \mid \mu(xy)_k \neq 0\}$ . Then  $G = (V, \sigma, \mu)$  is called fuzzy multigraph, where  $\sigma(x)$  and  $\mu(xy)_k$  denote the membership value of the vertex  $x$  and the membership value of the edge  $xy$  in  $\mathcal{G}$ , respectively.

**Definition 2.3.** [8, 16] A PBF $G$  of a graph  $G = (V, E)$  is a pair  $\mathcal{G} = (X, Y)$ , where  $X = (\mu_X^P, \mu_X^N)$  is a BFS in  $V$  and  $Y = (\mu_Y^P, \mu_Y^N)$  is a bipolar fuzzy relation on  $\widetilde{V}^2$  such that  $\mu_Y^P(xy) \leq \mu_X^P(x) \times \mu_X^P(y)$ ,  $\mu_Y^N(xy) \geq -(\mu_X^N(x) \times \mu_X^N(y))$  for all  $xy \in \widetilde{V}^2$  and  $\mu_Y^P(xy) = \mu_Y^N(xy) = 0$  for all  $xy \in \widetilde{V}^2 - E$ .

**Definition 2.4.** [4] A bipolar fuzzy multiset  $X$  drawn from non-empty set  $V$  is characterized by two functions: ‘count positive membership’ of  $X$  ( $CM_X^P$ ) and ‘count negative membership’ of  $X$  ( $CM_X^N$ ) given respectively by  $CM_X^P : V \rightarrow Q_1$  and  $CM_X^N : V \rightarrow Q_2$ , where  $Q_1$  and  $Q_2$  are the sets of all crisp multisets drawn from the intervals  $[0, 1]$  and  $[-1, 0]$ , respectively.

### 3. PRODUCT BIPOLAR FUZZY PLANAR GRAPHS

**Definition 3.1.** A PBF $G$  with an underlying set  $V$  is defined to be a pair  $\mathcal{G} = (X, Y)$ , where  $X = (\mu_X^P, \mu_X^N)$  is a BFS in  $V$  and  $Y = (\mu_Y^P, \mu_Y^N)$  is a BFS in  $\widetilde{V}^2$  such that

$$\begin{aligned} \mu_Y^P(xy) &\leq \mu_X^P(x)\mu_X^P(y), \mu_Y^N(xy) \geq -|\mu_X^N(x)||\mu_X^N(y)| \text{ for all } xy \in \widetilde{V}^2 \text{ and} \\ \mu_Y^P(xy) &= \mu_Y^N(xy) = 0 \text{ for all } xy \in \widetilde{V}^2 - E. \end{aligned}$$

Considering above modified definition of PBF $G$ s, we define PBF $M$  $G$  using the concept of bipolar fuzzy multiset. Further based on these concepts, the concept of PBF $P$  $G$  is introduced.

**Definition 3.2.** A PBF $M$  $G$  with an underlying set  $V$  is defined to be a pair  $\mathcal{G} = (X, Y)$ , where  $X = (\mu_X^P, \mu_X^N)$  is a BFS in  $V$  and  $Y = \{(xy, \mu_Y^P(xy)_k, \mu_Y^N(xy)_k) \mid k = 1, 2, \dots, m, xy \in \widetilde{V}^2\}$  is a bipolar fuzzy multiset in  $\widetilde{V}^2$  such that

$$\begin{aligned} \mu_Y^P(xy)_k &\leq \mu_X^P(x)\mu_X^P(y), \mu_Y^N(xy)_k \geq -|\mu_X^N(x)||\mu_X^N(y)| \text{ for all } xy \in \widetilde{V}^2 \\ \text{and } \mu_Y^P(xy)_k &= \mu_Y^N(xy)_k = 0 \text{ for all } xy \in \widetilde{V}^2 - E, \text{ for all } k = 1, 2, \dots, m. \end{aligned}$$

In PBF $M$  $G$   $\mathcal{G}$ ,  $Y$  is called product bipolar fuzzy multiedge set.

**Definition 3.3.** Let  $\mathcal{G} = (X, Y)$  be a PBF $M$  $G$ , where  $Y = \{(xy, \mu_Y^P(xy)_k, \mu_Y^N(xy)_k) \mid k = 1, 2, \dots, m, xy \in \widetilde{V}^2\}$ . The degree of a vertex  $x \in V$  in  $\mathcal{G}$ , is denoted by  $deg_{\mathcal{G}}(x)$  and is defined as  $deg_{\mathcal{G}}(x) = (\sum_{k=1}^m \mu_Y^P(xy)_k, \sum_{k=1}^m \mu_Y^N(xy)_k)$  for all  $y \in V$ .

**Definition 3.4.** Let  $Y = \{(xy, \mu_Y^P(xy)_k, \mu_Y^N(xy)_k) \mid k = 1, 2, \dots, m, xy \in \widetilde{V}^2\}$  be a bipolar fuzzy multiedge set in PBF $M$  $G$   $\mathcal{G}$ . A multiedge  $xy$  of  $\mathcal{G}$  is strong if  $\mu_X^P(x)\mu_X^P(y) \leq 2\mu_Y^P(xy)_k$  and  $-|\mu_X^N(x)||\mu_X^N(y)| \geq 2\mu_Y^N(xy)_k$ ,  $k$  is fixed integer.

**Example 3.1.** Consider a multigraph  $G = (V, E)$ , where  $V = \{v_1, v_2, v_3\}$  and  $E = \{v_1v_2, v_1v_2, v_2v_3, v_1v_3\}$ . Let  $X$  be a BFS on  $V$  and  $Y$  be a bipolar fuzzy multiedge set on  $\widetilde{V}^2$  defined by

$X$	$v_1$	$v_2$	$v_3$	$Y$	$v_1v_2$	$v_1v_2$	$v_2v_3$	$v_3v_1$
$\mu_X^P$	0.6	0.5	0.4	$\mu_Y^P$	0.3	0.2	0.2	0.1
$\mu_X^N$	-0.4	-0.7	-0.3	$\mu_Y^N$	-0.1	-0.1	-0.2	-0.1




FIGURE 1. PBFMG.

By direct calculations, it is easy to see from Fig. 1 that  $\mathcal{G} = (X, Y)$  is a PBFMG. Clearly,  $deg_{\mathcal{G}}(v_1) = (0.6, -0.3)$ ,  $deg_{\mathcal{G}}(v_2) = (0.7, -0.4)$  and  $deg_{\mathcal{G}}(v_3) = (0.3, -0.3)$ . Here  $(\mu_Y^P(v_2v_3), \mu_Y^N(v_2v_3))$  is a strong edge as  $(0.4)(0.5) < 2(0.2)$  and  $-|-0.3| - |-0.7| > 2(-0.2)$ .

**Definition 3.5.** Let  $\mathcal{G} = (X, Y)$  be a PBFMG, where  $Y = \{(xy, \mu_Y^P(xy)_k, \mu_Y^N(xy)_k) \mid k = 1, 2, \dots, m, xy \in \widetilde{V}^2\}$  is a bipolar fuzzy multiedge set. An edge  $xy$  of  $\mathcal{G}$  is effective if  $\mu_X^P(x)\mu_X^P(y) = \mu_Y^P(xy)_k$  and  $-|\mu_X^N(x)||\mu_X^N(y)| = \mu_Y^N(xy)_k$ ,  $k$  is fixed integer.

**Definition 3.6.** A PBFMG  $\mathcal{G} = (X, Y)$  is said to be complete if  $\mu_Y^P(xy)_k = \mu_X^P(x)\mu_X^P(y)$  and  $\mu_Y^N(xy)_k = -|\mu_X^N(x)||\mu_X^N(y)|$  for all  $x, y \in V$  and for all  $k = 1, 2, \dots, m$ .

**Example 3.2.** Consider a multigraph  $G = (V, E)$ , where  $V = \{v_1, v_2\}$  and  $E = \{v_1v_2, v_1v_2, v_1v_2\}$ . Let  $\mathcal{G} = (X, Y)$  be a PBFMG on  $V$ , as shown in Fig. 2.




FIGURE 2. Complete PBFMG.

By routine computations, it is easy to see that it is a complete PBFMG.

Assume that geometric insight for PBFMG has only one crossing between bipolar fuzzy edges  $(uv, \mu_Y^P(uv)_k, \mu_Y^N(uv)_k)$  and  $(wx, \mu_Y^P(wx)_k, \mu_Y^N(wx)_k)$ . If  $(\mu_Y^P(uv)_k, \mu_Y^N(uv)_k) = (1, -1)$  and  $(\mu_Y^P(wx)_k, \mu_Y^N(wx)_k) = (0, 0)$  or  $(\mu_Y^P(uv)_k, \mu_Y^N(uv)_k) = (0, 0)$  and  $(\mu_Y^P(wx)_k, \mu_Y^N(wx)_k) = (1, -1)$ , the PBFMG has no crossing, while if  $(\mu_Y^P(uv)_k, \mu_Y^N(uv)_k) = (1, -1)$  and  $(\mu_Y^P(wx)_k, \mu_Y^N(wx)_k) = (1, -1)$ , then there exists a crossing for the representation of the graph.

**Definition 3.7.** The strength of the bipolar fuzzy edge  $uv$  is defined as

$$I_{uv} = (I_{uv}^P, I_{uv}^N) = \left( \frac{\mu_Y^P(uv)_k}{\mu_X^P(u)\mu_X^P(v)}, \frac{-\mu_Y^N(uv)_k}{-|\mu_X^N(u)||\mu_X^N(v)|} \right).$$

An edge  $uv$  of a PBFMG is strong if  $I_{uv}^P \geq 0.5$  and  $I_{uv}^N \leq -0.5$ . An edge of a PBFMG which is not strong is called weak.

**Definition 3.8.** Let  $\mathcal{G} = (X, Y)$  be a PBFMG, where  $Y$  contains two edges  $(uv, \mu_Y^P(uv)_r, \mu_Y^N(uv)_r)$  and  $(wx, \mu_Y^P(wx)_s, \mu_Y^N(wx)_s)$  intersecting at a point  $C$  ( $r$  and  $s$  are fixed integers). The intersecting value at the point  $C$  is defined as

$$\mathcal{I}_C = (\mathcal{I}_C^P, \mathcal{I}_C^N) = \left( \frac{I_{uv}^P + I_{wx}^P}{2}, \frac{I_{uv}^N + I_{wx}^N}{2} \right).$$

**Definition 3.9.** Let  $\mathcal{G}$  be a PBFMG and let  $C_1, C_2, \dots, C_z$  be the intersecting points between the edges for geometric insight.  $\mathcal{G}$  is called a PBFPG with product bipolar fuzzy planarity value  $f$ , where

$$f = (f^P, f^N) = \left( \frac{1}{1 + \{\mathcal{I}_{C_1}^P + \mathcal{I}_{C_2}^P + \dots + \mathcal{I}_{C_z}^P\}}, \frac{-1}{1 - \{\mathcal{I}_{C_1}^N + \mathcal{I}_{C_2}^N + \dots + \mathcal{I}_{C_z}^N\}} \right).$$

Obviously,  $f$  is bounded and  $0 < f^P \leq 1$ ,  $-1 \leq f^N < 0$ . If geometric insight of a PBFPG has no point of intersection, then its product bipolar fuzzy planarity value is  $(1, -1)$  and this PBFPG has underlying crisp graph as the crisp planar graph.

In a PBFMG, if the number of point of intersections increases, planarity decreases. That is,  $\mathcal{I}_C$  is inversely proportional to the planarity.

**Example 3.3.** Consider a PBFMG  $\mathcal{G}$ , such that  $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$  and  $E = \{v_1v_2, v_2v_3, v_3v_4, v_4v_5, v_5v_6, v_6v_1, v_2v_5, v_2v_6, v_3v_6, v_1v_4\}$ , as shown in Fig. 3.




FIGURE 3. PBFMG.

There are two point of intersections  $C_1$  and  $C_2$  in PBFMG.  $C_1$  is the point of intersection between the edges  $(v_2v_5, 0.2, -0.1)$  and  $(v_3v_6, 0.1, -0.1)$  and  $C_2$  is the point of intersection between the edges  $(v_2v_6, 0.1, -0.2)$  and  $(v_3v_6, 0.1, -0.1)$ . For the edges  $(v_2v_5, 0.2, -0.1)$ ,  $(v_2v_6, 0.1, -0.2)$  and  $(v_3v_6, 0.1, -0.1)$ ,  $I_{v_2v_5} = (0.57, -0.42)$ ,  $I_{v_2v_6} = (0.29, -0.83)$  and  $I_{v_3v_6} = (0.67, -0.36)$ , respectively. For the first point of intersection  $C_1$ ,  $\mathcal{I}_{C_1} = (0.62, -0.39)$ . For the second point of intersection  $C_2$ ,  $\mathcal{I}_{C_2} = (0.48, -0.60)$ . Therefore, the product bipolar fuzzy planarity value for PBFMG is  $f = (f^P, f^N) = (0.48, -0.50)$ .

**Theorem 3.1.** Let  $\mathcal{G}$  be a PBFMG such that each intersecting edge is effective. Then product bipolar fuzzy planarity value  $f$  of  $\mathcal{G}$  is given by

$$f = (f^P, f^N) = \left( \frac{1}{1+n_C}, \frac{-1}{1+n_C} \right)$$

where  $n_C$  is the number of point of intersections between the edges in  $\mathcal{G}$ .

*Proof.* Let  $\mathcal{G}$  be a PBFMG such that each intersecting edge is effective. Let  $C_1, C_2, \dots, C_z$ ,  $z \in Z$  be the point of intersections between the edges in  $\mathcal{G}$ . For each intersecting edge  $uv$  in  $\mathcal{G}$ ,  $I_{uv} = (I_{uv}^P, I_{uv}^N) = \left( \frac{\mu_Y^P(uv)_k}{\mu_X^P(u)\mu_X^P(v)}, \frac{\mu_Y^N(uv)_k}{|\mu_X^N(u)||\mu_X^N(v)|} \right) = (1, -1)$ . Thus for  $C_1$ , the point of intersection between the edges  $uv$  and  $wx$ ,  $\mathcal{I}_{C_1} = (\mathcal{I}_{C_1}^P, \mathcal{I}_{C_1}^N) = \left( \frac{1+1}{2}, \frac{-1-1}{2} \right) = (1, -1)$ . Hence  $\mathcal{I}_{C_\eta} = (1, -1)$  for all  $\eta = 1, 2, \dots, z$ .

Now

$$\begin{aligned} f = (f^P, f^N) &= \left( \frac{1}{1 + \{\mathcal{I}_{C_1}^P + \mathcal{I}_{C_2}^P + \dots + \mathcal{I}_{C_z}^P\}}, \frac{-1}{1 - \{\mathcal{I}_{C_1}^N + \mathcal{I}_{C_2}^N + \dots + \mathcal{I}_{C_z}^N\}} \right) \\ &= \left( \frac{1}{1 + \{1 + 1 + \dots + 1\}}, \frac{-1}{1 - \{-1 - 1, \dots, -1\}} \right) \\ &= \left( \frac{1}{1+n_C}, \frac{-1}{1+n_C} \right), \end{aligned}$$

where  $n_C$  is the number of point of intersections between the edges in  $\mathcal{G}$ .  $\square$

**Definition 3.10.** A PBFPG  $\mathcal{G}$  is said to be strong if the product bipolar fuzzy planarity value  $f = (f^P, f^N)$  of  $\mathcal{G}$  is such that  $f^P \geq 0.5$  and  $f^N \leq -0.5$ .

In Example 3.3, the product bipolar fuzzy planarity value  $f = (f^P, f^N)$  of the PBFPG  $\mathcal{G}$  is such that  $f^P \leq 0.5$  and  $f^N \geq -0.5$ . So,  $\mathcal{G}$  is not strong.

**Corollary 3.1.** Let  $\mathcal{G}$  be a complete PBFMG. Then the product bipolar fuzzy planarity value  $f$  of  $\mathcal{G}$  is given by

$$f = (f^P, f^N) = \left( \frac{1}{1+n_C}, \frac{-1}{1+n_C} \right)$$

where  $n_C$  is the number of point of intersections between the edges in  $\mathcal{G}$ .

**Theorem 3.2.** Let  $\mathcal{G}$  be a strong PBFPG. Then the number of point of intersections between strong edges in  $\mathcal{G}$  is at most one.

*Proof.* Let  $\mathcal{G}$  be a strong PBFPG. Suppose  $\mathcal{G}$  has at least two point of intersections  $C_1$  and  $C_2$  between two strong edges in  $\mathcal{G}$ . For any strong edge  $(uv, \mu_Y^P(uv)_k, \mu_Y^N(uv)_k)$ ,  $\mu_X^P(u)\mu_X^P(v) \leq 2\mu_Y^P(uv)_k$  and  $-|\mu_X^N(u)||\mu_X^N(v)| \geq 2\mu_Y^N(uv)_k$ , that is,  $I_{uv}^P \geq 0.5$  and  $I_{uv}^N \leq -0.5$ . Therefore, if two strong edges  $(uv, \mu_Y^P(uv)_k, \mu_Y^N(uv)_k)$  and  $(wx, \mu_Y^P(wx)_k, \mu_Y^N(wx)_k)$  intersect, then  $\mathcal{I}_{C_1}^P = \frac{I_{uv}^P + I_{wx}^P}{2} \geq 0.5$  and  $\mathcal{I}_{C_1}^N = \frac{I_{uv}^N + I_{wx}^N}{2} \leq -0.5$ . Similarly,  $\mathcal{I}_{C_2}^P \geq 0.5$  and  $\mathcal{I}_{C_2}^N \leq -0.5$ . This implies that,  $1 + \mathcal{I}_{C_1}^P + \mathcal{I}_{C_2}^P \geq 2$  and  $1 - (\mathcal{I}_{C_1}^N + \mathcal{I}_{C_2}^N) \geq 2$ . Therefore,  $f^P = \frac{1}{1 + \{\mathcal{I}_{C_1}^P + \mathcal{I}_{C_2}^P\}} \leq 0.5$  and  $f^N = \frac{-1}{1 - \{\mathcal{I}_{C_1}^N + \mathcal{I}_{C_2}^N\}} \geq -0.5$ , a contradiction, as  $\mathcal{G}$  is a strong PBFPG.  $\square$

A fundamental theorem of PBFPG is given below.

**Theorem 3.3.** A PBFPG  $\mathcal{G}$  does not contain any point of intersection between two strong edges, if  $\mathcal{G}$  has product bipolar fuzzy planarity value  $f = (f^P, f^N)$  such that  $f^P > 0.67$  and  $f^N < -0.67$ .

*Proof.* Let  $\mathcal{G}$  be a PBFPG with product bipolar fuzzy planarity value  $f$ , such that  $f^P > 0.67$  and  $f^N < -0.67$ . Let  $C$  be the point of intersection between two strong bipolar fuzzy edges  $(uv, \mu_Y^P(uv)_k, \mu_Y^N(uv)_k)$  and  $(wx, \mu_Y^P(wx)_k, \mu_Y^N(wx)_k)$ . For any strong edge  $(uv, \mu_Y^P(uv)_k, \mu_Y^N(uv)_k)$ ,  $\mu_X^P(u)\mu_X^P(v) \leq 2\mu_Y^P(uv)_k$  and  $-|\mu_X^P(u)||\mu_X^P(v)| \geq 2\mu_Y^N(uv)_k$ . That is,  $I_{uv}^P \geq 0.5$  and  $I_{uv}^N \leq -0.5$ . Similarly  $I_{wx}^P \geq 0.5$  and  $I_{wx}^N \leq -0.5$ . For the minimum value of  $I_{uv}^P$  and  $I_{wx}^P$ ,  $\mathcal{I}_C^P = \frac{0.5+0.5}{2} = 0.5$  and for the maximum value of  $I_{uv}^N$  and  $I_{wx}^N$ ,  $\mathcal{I}_C^N = \frac{-0.5-0.5}{2} = -0.5$ . So,  $f^P = \frac{1}{1+\mathcal{I}_C^P} \leq 0.67$  and  $f^N = \frac{-1}{1-\mathcal{I}_C^N} \geq -0.67$ , a contradiction. Hence,  $\mathcal{G}$  does not contain any point of intersection between strong edges.  $\square$

We denote a PBFPG with product bipolar fuzzy planarity value  $f = (f^P, f^N)$  such that  $f^P > 0.67$  and  $f^N < -0.67$ , as  $(0.67, -0.67)$ -PBFPG.

Face of a PBFPG is an important feature. Face of a PBFPG is a region bounded by bipolar fuzzy edges. If all the edges in the boundary of a product bipolar fuzzy face have membership value  $(1, -1)$ , then it is a crisp face. If one of such edges is removed or has membership value  $(0, 0)$ , the product bipolar fuzzy face does not exist. A product bipolar fuzzy face and its positive membership and negative membership values are defined below.

**Definition 3.11.** Let  $\mathcal{G} = (X, Y)$  be a PBFPG, with product bipolar fuzzy planarity value  $(1, -1)$ . A region bounded by the set of bipolar fuzzy edges  $E^* \subset E$  of a geometric representation of  $\mathcal{G}$  is said to be a product bipolar fuzzy face of  $\mathcal{G}$ . The positive and negative membership values of the product bipolar fuzzy face are  $\prod\{I_{uv}^P \mid uv \in E^*\}$  and  $-\prod\{|I_{uv}^N| \mid uv \in E^*\}$ .

**Definition 3.12.** A product bipolar fuzzy face is said to be strong if its positive membership value is greater than and equal to 0.5 and negative membership value is less than and equal to  $-0.5$ . A product bipolar fuzzy face which is not strong is called weak. Every PBFPG has an infinite region called outer product bipolar fuzzy face. Other faces are called inner product bipolar fuzzy faces.

**Remark 3.1.** Every edge of a strong product bipolar fuzzy face is a strong bipolar fuzzy edge.

**Example 3.4.** Consider a PBFPG  $\mathcal{G}$  as shown in Fig. 4. Let  $f_1, f_2$  and  $f_3$  be the prod-




FIGURE 4. Faces in PBFPG.

uct bipolar fuzzy faces,  $f_1$  is bounded by the edges  $(v_1v_2, (0.1, -0.1))$ ,  $(v_2v_3, (0.15, -0.2))$ ,  $(v_3v_1, (0.1, -0.3))$ ,  $f_2$  is bounded by the edges  $(v_2v_3, (0.15, -0.2))$ ,  $(v_3v_4, (0.1, -0.7))$ ,  $(v_2v_4, (0.25, -0.2))$  and  $f_3$  is bounded by the edges  $(v_1v_2, (0.1, -0.1))$ ,  $(v_1v_3, (0.1, -0.3))$ ,  $(v_3v_4, (0.1, -0.7))$ ,  $(v_2v_4, (0.25, -0.2))$ . The positive membership and negative membership values

of product bipolar fuzzy faces  $f_1, f_2$  and  $f_3$  are  $(0.42, -0.29), (0.67, -0.6)$  and  $(0.28, -0.25)$ , respectively. Here  $f_2$  is a strong product bipolar fuzzy face and  $f_1, f_3$  are weak product bipolar fuzzy faces.

Now we define dual of  $(0.67, -0.67)$ -PBFPFG.

**Definition 3.13.** Let  $\mathcal{G}$  be a  $(0.67, -0.67)$ -PBFPFG, where  $Y = \{(xy, \mu_Y^P(xy)_k, \mu_Y^N(xy)_k) \mid k = 1, 2, \dots, m, xy \in \widetilde{V}^2\}$  is a bipolar fuzzy multiset on  $\widetilde{V}^2$ . Let  $f_1, f_2, \dots, f_t$  be the strong product bipolar fuzzy faces of  $\mathcal{G}$ . A PBFPFG  $\mathcal{G}' = (X', Y')$  such that  $V' = \{x_i, i = 1, 2, \dots, t\}$ , and the vertex  $x_i$  of  $\mathcal{G}'$  is considered for the face  $f_i$  of  $\mathcal{G}$  is said to be a PBFDDG of  $\mathcal{G}$ . The positive membership values of vertices are given by the mapping  $\mu_{X'}^P : V' \rightarrow [0, 1]$  such that  $\mu_{X'}^P(x_i) = \prod\{\mu_Y^P(uv)_k, k = 1, 2, \dots, m \mid uv \text{ is an edge of the boundary of the strong product bipolar fuzzy face } f_i\}$  and negative membership values of vertices are given by the mapping  $\mu_{X'}^N : V' \rightarrow [-1, 0]$  such that  $\mu_{X'}^N(x_i) = -\prod\{|\mu_Y^N(uv)_k|, k = 1, 2, \dots, m \mid uv \text{ is an edge of the boundary of the strong product bipolar fuzzy face } f_i\}$ .

There may exist at least two common edges between two faces  $f_i$  and  $f_j$  of  $\mathcal{G}$ . So, there may exist at least two edges between two vertices  $x_i$  and  $x_j$  in PBFDDG  $\mathcal{G}'$ . The positive membership values and negative membership values of the bipolar fuzzy edges of the PBFDDG are  $\mu_{Y'}^P(uv)_k^l = \mu_{Y'}^P(x_i x_j)_l$  and  $\mu_{Y'}^N(uv)_k^l = \mu_{Y'}^N(x_i x_j)_l$ , where  $(uv)^l$  is an edge in the boundary between two strong product bipolar fuzzy faces  $f_i$  and  $f_j$ ,  $l = 1, 2, \dots, s$ , where  $s$  is the number of edges between  $x_i$  and  $x_j$  or the number of common edges in the boundary between  $f_i$  and  $f_j$ .

**Example 3.5.** Consider a planar graph  $G = (V, E)$ , where  $V = \{v_1, v_2, v_3, v_4, v_5\}$  and  $E = \{v_1v_2, v_2v_3, v_3v_4, v_4v_5, v_5v_1, v_1v_4\}$ . Let  $\mathcal{G} = (X, Y)$  be a PBFDDG on  $V$ , as shown in Fig. 5. Clearly, it is a  $(0.67, -0.67)$ -PBFPFG with three strong product bipolar fuzzy faces  $f_1, f_2$




FIGURE 5. PBFDDG.

and  $f_3$ . Product bipolar fuzzy face  $f_1$  is bounded by edges  $(v_1v_2, 0.7, -0.3), (v_2v_3, 0.6, -0.3), (v_3v_4, 0.3, -0.5)$  and  $(v_1v_4, 0.4, -0.6)$ .  $f_2$  is bounded by edges  $(v_1v_4, 0.4, -0.6), (v_4v_5, 0.15, -0.2)$



and  $(v_5v_1, 0.2, -0.2)$ .  $f_3$  is bounded by edges  $(v_1v_2, 0.7, -0.3)$ ,  $(v_2v_3, 0.6, -0.3)$ ,  $(v_3v_4, 0.3, -0.5)$ ,  $(v_4v_5, 0.15, -0.2)$  and  $(v_5v_1, 0.2, -0.2)$ . We represent the vertices of PBFDG by small white circles and the edges by dashed lines. For each strong product bipolar fuzzy face, we consider a vertex for the PBFDG. Thus the vertex set  $V' = \{x_1, x_2, x_3\}$ , where the vertex  $x_i$  is taken corresponding to the strong product bipolar fuzzy face  $f_i$ ,  $i = 1, 2, 3$ . Therefore

$$\begin{aligned} \mu_{X'}^P(x_1) &= (0.7)(0.6)(0.3)(0.4) = 0.05, \mu_{X'}^N(x_1) = -|-0.3| - 0.3| - 0.5| - 0.6| = -0.03, \\ \mu_{X'}^P(x_2) &= (0.4)(0.2)(0.15) = 0.01, \mu_{X'}^N(x_2) = -|-0.6| - 0.2| - 0.2| = -0.02, \\ \mu_{X'}^P(x_3) &= (0.7)(0.6)(0.3)(0.15)(0.2) = 0.004, \mu_{X'}^N(x_3) = -|-0.3| - 0.3| - 0.5| - 0.2| - 0.2| = \\ &= -0.002. \end{aligned}$$

There are three common edges  $v_1v_2$ ,  $v_2v_3$  and  $v_3v_4$  between the faces  $f_1$  and  $f_3$  in  $\mathcal{G}$ . Therefore, there exist three edges between the vertices  $x_1$  and  $x_3$ , in PBFDG of  $\mathcal{G}$ . Positive and negative membership values of the edges of PBFDG are given by

$$\begin{aligned} \mu_{Y'}^P(x_1x_2) &= \mu_Y^P(v_1v_4) = 0.4, \mu_{Y'}^N(x_1x_2) = \mu_Y^N(v_1v_4) = -0.6, \\ \mu_{Y'}^P(x_2x_3) &= \mu_Y^P(v_1v_5) = 0.2, \mu_{Y'}^N(x_2x_3) = \mu_Y^N(v_1v_5) = -0.2, \\ \mu_{Y'}^P(x_2x_3) &= \mu_Y^P(v_4v_5) = 0.15, \mu_{Y'}^N(x_2x_3) = \mu_Y^N(v_4v_5) = -0.2, \\ \mu_{Y'}^P(x_1x_3) &= \mu_Y^P(v_1v_2) = 0.7, \mu_{Y'}^N(x_1x_3) = \mu_Y^N(v_1v_2) = -0.3, \\ \mu_{Y'}^P(x_1x_3) &= \mu_Y^P(v_2v_3) = 0.6, \mu_{Y'}^N(x_1x_3) = \mu_Y^N(v_2v_3) = -0.3, \\ \mu_{Y'}^P(x_1x_3) &= \mu_Y^P(v_3v_4) = 0.3, \mu_{Y'}^N(x_1x_3) = \mu_Y^N(v_3v_4) = -0.5. \end{aligned}$$

So, the edge set of PBFDG is

$$Y' = \{(x_1x_2, 0.4, -0.6), (x_2x_3, 0.2, -0.2), (x_2x_3, 0.15, -0.2), (x_1x_3, 0.7, -0.3), (x_1x_3, 0.6, -0.3), (x_1x_3, 0.3, -0.5)\}.$$

We state the following theorems without their proofs.

**Theorem 3.4.** Let  $\mathcal{G}$  be a  $(0.67, -0.67)$ -PBFPG without weak edges, with  $n$  vertices,  $m$  bipolar fuzzy edges and  $f$  strong faces, and let  $\mathcal{G}'$  be a PBFDG of  $\mathcal{G}$  with  $n'$  vertices,  $m'$  edges and  $f'$  faces, then  $n' = f$ ,  $m' = m$  and  $f' = n$ .

**Theorem 3.5.** Let  $\mathcal{G}'$  be a PBFDG of a  $(0.67, -0.67)$ -PBFPG  $\mathcal{G}$ . The number of strong product bipolar fuzzy faces in  $\mathcal{G}'$  is less than or equal to the number of vertices of  $\mathcal{G}$ .

**Definition 3.14.** A homomorphism  $h : \mathcal{G}_1 \rightarrow \mathcal{G}_2$  of two PBFPGs  $\mathcal{G}_1$  and  $\mathcal{G}_2$  is a mapping  $h : V_1 \rightarrow V_2$  satisfying the following conditions:

- (a):  $\mu_{X_1}^P(x_1) \leq \mu_{X_2}^P(h(x_1)), \mu_{X_1}^N(x_1) \geq \mu_{X_2}^N(h(x_1)),$
- (b):  $\mu_{Y_1}^P(x_1y_1) \leq \mu_{Y_2}^P(h(x_1)h(y_1)), \mu_{Y_1}^N(x_1y_1) \geq \mu_{Y_2}^N(h(x_1)h(y_1))$   
for all  $x_1 \in V_1, x_1y_1 \in \widetilde{V_1^2}$ .

**Definition 3.15.** An isomorphism  $h : \mathcal{G}_1 \rightarrow \mathcal{G}_2$  of two PBFPGs  $\mathcal{G}_1$  and  $\mathcal{G}_2$  is a bijective mapping  $h : V_1 \rightarrow V_2$  satisfying the following conditions:

- (c):  $\mu_{X_1}^P(x_1) = \mu_{X_2}^P(h(x_1)), \mu_{X_1}^N(x_1) = \mu_{X_2}^N(h(x_1)),$
- (d):  $\mu_{Y_1}^P(x_1y_1) = \mu_{Y_2}^P(h(x_1)h(y_1)), \mu_{Y_1}^N(x_1y_1) = \mu_{Y_2}^N(h(x_1)h(y_1))$   
for all  $x_1 \in V_1, x_1y_1 \in \widetilde{V_1^2}$ .

**Definition 3.16.** A weak isomorphism  $h : \mathcal{G}_1 \rightarrow \mathcal{G}_2$  of two PBFPGs  $\mathcal{G}_1$  and  $\mathcal{G}_2$  is a bijective mapping  $h : V_1 \rightarrow V_2$  satisfying the following conditions:

- (e):  $h$  is homomorphism,

(f):  $\mu_{X_1}^P(x_1) = \mu_{X_2}^P(h(x_1)), \mu_{X_1}^N(x_1) = \mu_{X_2}^N(h(x_1))$  for all  $x_1 \in V_1$ .

**Definition 3.17.** A co-weak isomorphism  $h : \mathcal{G}_1 \rightarrow \mathcal{G}_2$  of two PBFPGs  $\mathcal{G}_1$  and  $\mathcal{G}_2$  is a bijective mapping  $h : V_1 \rightarrow V_2$  satisfying the following conditions:

(g):  $h$  is homomorphism,

(h):  $\mu_{Y_1}^P(x_1y_1) = \mu_{Y_2}^P(h(x_1)h(y_1)), \mu_{Y_1}^N(x_1y_1) = \mu_{Y_2}^N(h(x_1)h(y_1))$  for all  $x_1y_1 \in \widetilde{V}_1^2$ .

As isomorphism between PBFPGs is an equivalence relation, if there exists an isomorphism between two PBFPGs and one is PBFPG, then the other will also be PBFPG. This result is given below.

**Theorem 3.6.** Let  $h : \mathcal{G} \rightarrow \mathcal{H}$  be an isomorphism from a PBFPG  $\mathcal{G}$  to a PBFPG  $\mathcal{H}$ . Then  $\mathcal{H}$  can be drawn as PBFPG with same planarity value of  $\mathcal{G}$ .

*Proof.* Let  $h : \mathcal{G} \rightarrow \mathcal{H}$  be an isomorphism. Since isomorphism preserves size and order of PBFPGs. So, the size and order of  $\mathcal{H}$  will be equal to size and order of  $\mathcal{G}$  and drawing of  $\mathcal{G}$  and  $\mathcal{H}$  is same. Thus the number of intersections between edges and product bipolar fuzzy planarity value of  $\mathcal{H}$  will be same as  $\mathcal{G}$ . Hence  $\mathcal{H}$  can be drawn as PBFPG with same product bipolar fuzzy planarity value as of  $\mathcal{G}$ .  $\square$

Using above Theorem, we can immediately prove the following results.

**Theorem 3.7.** Two isomorphic PBFPGs have the same product bipolar fuzzy planarity values.

**Theorem 3.8.** Let  $\mathcal{G}_1$  and  $\mathcal{G}_2$  be two weak isomorphic PBFPGs with product bipolar fuzzy planarity values  $f_1 = (f_1^P, f_1^N)$  and  $f_2 = (f_2^P, f_2^N)$ , respectively. If the edge positive membership and negative membership values of corresponding intersecting edges are same. Then  $(f_1^P, f_1^N) = (f_2^P, f_2^N)$ .

**Theorem 3.9.** Let  $\mathcal{G}_1$  and  $\mathcal{G}_2$  be two co-weak isomorphic PBFPGs with product bipolar fuzzy planarity values  $f_1 = (f_1^P, f_1^N)$  and  $f_2 = (f_2^P, f_2^N)$ , respectively. If the minimum of positive membership values and maximum of negative membership values of the end vertices of corresponding intersecting edges are same. Then  $(f_1^P, f_1^N) = (f_2^P, f_2^N)$ .

#### 4. APPLICATION OF PBFPGs

We consider a road network as shown in Fig. 6. Each city in the network may be referred as vertex and each road between any two cities may be called as edge. This graph does not contain loops and multiple edges. It is well known that, the length of a road between any two cities is a crisp quantity but vehicle travel time or vehicle capacity on a road network is fuzzy. In this network, the membership values of vertices are representing the degree of capacity of vehicles of a city belongs to a network of 5 cities. The degree of capacity of vehicles of a city is defined in terms of its positive membership and negative membership. Positive membership degree can be depicted as how much capacity, vehicles of a city possess and negative membership can be depicted as how much capacity is lost by the vehicles of a city. The membership values of edges of this graph show the capacity of vehicles on the road joining any two cities. The positive and negative membership degree of edges can be interpreted as the percentage of increasing and decreasing capacity of vehicles on the road between any two cities.

Due to crossing, the vehicle capacity or vehicle travel time decreases, in order to reach from one city to another city and so traveling cost decreases. But we construct the roads in such a way that the number of crossing decreases, that is, the planarity value increases, because generally, as the crossing road increases, vehicle capacity increases, due to the




FIGURE 6. PBFPG of a road network.

increase of crowdedness on roads. That’s why, the measurement of product bipolar fuzzy planarity value is important.

There are only two crossings, at the points  $C_1$  and  $C_2$ , between the roads  $AC$  and  $BE$ , and  $AD$  and  $BE$ , respectively, in this road network of Fig. 6. For the roads  $AC$ ,  $BE$  and  $AD$ ,  $I_{AC} = (0.36, -0.14)$ ,  $I_{BE} = (0.33, -0.18)$  and  $I_{AD} = (0.24, -0.50)$ , respectively. For first crossing at  $C_1$ ,  $\mathcal{I}_{C_1} = (0.35, -0.16)$  and for second crossing at  $C_2$ ,  $\mathcal{I}_{C_2} = (0.29, -0.34)$ . Therefore, the product bipolar fuzzy planarity value  $f$  of above road network is  $(0.61, -0.67)$ . So, the roads are not to much crowded and due to road crossing between cities vehicle travel time saves and traveling cost decreases.

### 5. CONCLUSIONS

The bipolar fuzzy models give more precision, flexibility and compatibility to the system as compared to the classical and fuzzy models. In this paper, we have initiated the concepts of multigraphs, planar graphs and dual graphs under bipolar fuzzy environment based on the product operator. We have also introduced the product bipolar fuzzy planarity value of PBFPGs. If the product bipolar fuzzy planarity value of a PBFPG is  $(1, -1)$ , that is, if there exists no crossing between edges, then the PBFPG is same as crisp planar graph. Therefore, the product bipolar fuzzy planarity value measures the amount of planarity in a PBFPG. In further work, it is necessary and meaningful to extend the concept of PBFPG to product intuitionistic fuzzy planar graphs.

### REFERENCES

- [1] Akram, M., (2011), Bipolar fuzzy graphs, *Information Sciences*, 181(24), pp. 5548-5564.
- [2] Akram, M. and Dudek, W. A., (2012), Regular bipolar fuzzy graphs, *Neural Computing and Applications*, 21(1), pp. 197-205.
- [3] Akram, M., (2013), Bipolar fuzzy graphs with applications, *Knowledge-Based Systems*, 39, pp. 1-8.
- [4] Akram, M., Samanta, S. and Pal, M., (2017), Application of bipolar fuzzy sets in planar graphs, *International Journal of Applied and Computational Mathematics*, 3(2), pp. 773-785.
- [5] Alshehri, N. and Akram, M., (2014), Intuitionistic fuzzy planar graphs, *Discrete Dynamics in Nature and Society*, pp. 1-9.
- [6] Ashraf, S., Naz, S. and Kerre, E. E., (2018), Dombi fuzzy graphs, *Fuzzy Information and Engineering*, 10(1), pp. 58-79.
- [7] Ashraf, S., Naz, S., Rashmanlou, H. and Malik, M. A., (2017), Regularity of graphs in single-valued neutrosophic environment, *Journal of Intelligent & Fuzzy Systems*, 33(1), pp. 529-542.
- [8] Ghorai, G. and Pal, M., (2017), Certain types of product bipolar fuzzy graphs, *International Journal of Applied and Computational Mathematics*, 3(2), pp. 605-619.

- [9] Jabbar, N. A., Naoom, J. H. and Ouda, E. H., (2009), Fuzzy dual graph, J Al-Nahrain Univ, 12(4), pp. 168-171.
- [10] Lee, K. M., (2000), Bipolar-valued fuzzy sets and their basic operations, In Proc. Int. Conf. on Intelligent Technologies, Bangkok, Thailand, pp. 307-312.
- [11] Naz, S., Ashraf, S. and Karaaslan, F., (2018), Energy of a bipolar fuzzy graph and its application in decision making, Italian Journal of Pure and Applied Mathematics, 40, pp. 339-352.
- [12] Naz, S., and Malik, M. A., (2018), Single valued neutrosophic line graphs, TWMS Journal of Applied and Engineering Mathematics, 8(2), pp. 483-494.
- [13] Naz, S., Rashmanlou, H. and Malik, M. A., (2017), Operations on single-valued neutrosophic graphs with application, Journal of Intelligent & Fuzzy Systems, 32(3), pp. 2137-2151.
- [14] Naz, S., Malik, M. A. and Rashmanlou, H., (2018), Hypergraphs and transversals of hypergraphs in interval-valued intuitionistic fuzzy setting, The Journal of Multiple-Valued Logic and Soft Computing, 30 (4-6), pp. 399-417.
- [15] Pal, A., Samanta, S. and Pal, M., (2013), Concept of fuzzy planar graphs, In Science and Information Conference (SAI) IEEE, pp. 557-563.
- [16] Rashmanlou, H., Samanta, S., Pal, M. and Borzooei, R. A., (2015), A study on bipolar fuzzy graphs, Journal of Intelligent & Fuzzy Systems, 28(2), pp. 571-580.
- [17] Rosenfeld, A., (1975), Fuzzy graphs, Fuzzy Sets and their Applications (L. A. Zadeh, K. S. Fu, M. Shimura, Eds.) Academic Press, New York, pp. 77-95.
- [18] Samanta, S., Pal, M. and Pal, A., (2014), New concepts of fuzzy planar graph, International Journal of Advanced Research in Artificial Intelligence, 3(1), pp. 52-59.
- [19] Yager, R. R., (1986), On the theory of bags, International Journal of General System, 13(1), pp. 23-37.
- [20] Zadeh, L. A., (1965), Fuzzy sets, Information and Control, 8(3), pp. 338-353.
- [21] Zhang, W-R., (1994), Bipolar fuzzy sets and relations: a computational framework for cognitive modeling and multiagent decision analysis, Proceedings of IEEE conference, pp. 305-309.
- [22] Zhang, W-R., (1998), Bipolar fuzzy sets, Proceedings of FUZZ-IEEE, pp. 835-840.

**Sumera Naz** for the photography and short autobiography, see TWMS J. App. Eng. Math., V.8, N.2, 2018.



**Samina Ashraf** received her Ph.D. degree from Lahore University of Management Sciences, Pakistan. Currently, she is working as a Director of Quality Enhancement Cell, Lahore College for Women University, Lahore, Pakistan. Her research interests include fuzzy logic, fuzzy mathematics and decision making. She has published 40 research articles in international peer-reviewed journals.



**Hossein Rashmanlou** received his Ph.D. degree from Islamic Azad University, Central Tehran Branch, Tehran, Iran. Currently, he is working as a post doc researcher in Department of Mathematics, University of Mazandaran, Babolsar, Iran. His research interests include fuzzy algebra, fuzzy logic, fuzzy graphs and its applications in different sciences.