

APPLICATION OF THE HOMOTOPY PERTURBATION METHOD (HPM) AND VARIATIONAL ITERATION METHOD (VIM) TO GAS DYNAMIC EQUATION

M. KAZEMI-SANGSEREKI¹, D. D. GANJI¹, M. GORGI¹ §

ABSTRACT. A gas-dynamic control system is one where the path of an object in flight is controlled by either the generation or redirection of gas flow out of an orifice rather than with the traditional movable control surfaces. In this paper considering Hes Homotopy perturbation and Variational iteration methods are calculated the homogeneous gas dynamics equations. The exact (analytic) solution of the equation is calculated in the form of a series with easily computable components like to Adomians decomposition method. The Comparison of result of HPM and VIM with Adomians decomposition method show that they are agreement with them, and HPM, VIM can solve large class of nonlinear problems.

Keywords: Gas dynamics equation, Adomians Decomposition method, Homotopy Perturbation Method (HPM), Variational Iteration Method (VIM).

AMS Subject Classification: 35A15

1. INTRODUCTION

The recent years have seen significant development in the use of various methods for the numerical and analytic solution of the partial differential equation. The aim of the present paper is to implement the homotopy perturbation and variational iteration methods [1220] for the homogeneous gas dynamics equations to an approximation solution. The solution of this method is derived in the form of power series with easy computable components.

Over the last decades several analytical/approximate methods have been developed to solve linear and nonlinear ordinary and partial differential equations. Some of these techniques include variational iteration method (VIM) (He [16,17]; Ganji et al., [12,14,15];), homotopy perturbation method (HPM) (He [2,8,9,10,11]; Ganji et al., [5,6,7,12,14,15];) etc. Linear and Nonlinear phenomena play an important role in various fields of science and engineering. Most models of real life problems are still very difficult to solve. Therefore, approximate analytical solutions such as homotopy perturbation method [2-15] were

¹ Babol Noshiravani University of Technology, Department of Mechanical Engineering, P.O. Box 484, Babol, Iran,
e-mail: sangsereki@gmail.com
e-mail: ddg_davood@yahoo.com
e-mail: gorji@nit.ac.ir

§ Manuscript received 26 February 2011.

TWMS Journal of Applied and Engineering Mathematics Vol.1 No.1 © Işık University, Department of Mathematics 2011; all rights reserved.

introduced. This method is the most effective and convenient ones for both linear and nonlinear equations.

Perturbation method is based on assuming a small parameter. The majority of nonlinear problems, especially those having strong nonlinearity, have no small parameters at all and the approximate solutions obtained by the perturbation methods, in most cases, are valid only for small values of the small parameter. Generally, the perturbation solutions are uniformly valid as long as a scientific system parameter is small. However, we cannot rely fully on the approximations, because there is no criterion on which the small parameter should exist. Thus, it is essential to check the validity of the approximations numerically and/or experimentally. To overcome these difficulties, HPM have been proposed recently.

Recently, He (1999) proposed a variational iteration method based on the use of restricted variations and correction functionals which has found a wide application for the solution of nonlinear ordinary and partial differential equations. This method does not require the presence of small parameters in the differential equation, and provides the solution (or an approximation to it) as a sequence of iterates. The method does not require that the nonlinearities be differentiable with respect to the dependent variable and its derivatives.

In this paper we will apply the homotopy perturbation method and variational iteration method to gas dynamics equation and comparing the result of HPM, VIM with ADM [21-23]

2. BASIC IDEA OF HOMOTOPY-PERTURBATION METHOD

To explain this method, let us consider the following function:

$$A(u) - f(r) = 0, \quad r \in \Omega \quad (1)$$

With the boundary conditions of:

$$B(u, \frac{\partial u}{\partial n}) = 0, \quad r \in \Gamma, \quad (2)$$

Where $A, B, f(r)$ and Γ are a general differential operator, a boundary operator, a known analytical function and the boundary of the domain Ω , respectively. Generally speaking the operator A can be divided into a linear part L and a nonlinear part $N(u)$. Eq.(1) can therefore, be written as:

$$L(u) + N(u) - f(r) = 0 \quad (3)$$

By the homotopy technique, we construct a homotopy

$u(r, p) : \Omega \times [0, 1] \rightarrow \mathbb{R}$ Which satisfies:

$$H(v, p) = (1 - p)[L(v) - L(u_0)] + p[A(v) - f(r)] = 0, \quad (4)$$

$$p \in [0, 1], \quad r \in \Omega,$$

or

$$H(v, p) = L(v) - L(u_0) + pL(u_0) + p[N(u) - f(r)] = 0, \quad (5)$$

where $p \in [0, 1]$ is an embedding parameter, while u_0 is an initial approximation of Eq. (1), which satisfies the boundary conditions. Obviously, from Eqs. (4) and (5) we will have:

$$H(v, 0) = L(v) - L(u_0) = 0, \quad (6)$$

$$H(v, 1) = A(v) - f(r) = 0, \tag{7}$$

The changing process of p from zero to unity is just that of $v(r, p)$ from u_0 to $u(r)$. In topology, this is called deformation, while $L(v) - L(u_0)$ and $A(v) - f(r)$ are called homotopy. According to the HPM, we can first use the embedding parameter p as a small parameter, and assume that the solutions of Eqs. (4), (5) can be written as a power series in p :

$$v = v_0 + pv_1 + p^2v_2 + \dots, \tag{8}$$

Setting $p = 1$ yields in the approximate solution of Eq. (4) to:

$$u = \lim_{p \rightarrow 1} v = v_0 + v_1 + v_2 + \dots, \tag{9}$$

The combination of the perturbation method and homotopy method is called the HPM, which eliminates the drawbacks of the traditional perturbation methods while keeping all its advantage. The series (9) is convergent for most cases. However, the convergent rate depends on the nonlinear operator $A(v)$. Moreover, He (1999) made the following suggestions:

- . The second derivative of $N(v)$ with respect to v must be small because the parameter may be relatively large, i.e. $p \rightarrow 1$.
- . The norm of $L^{-1} \frac{\partial N}{\partial v}$ must be smaller than one so that the series converges.

3. BASIC IDEA OF VARIATIONAL ITERATION METHOD:

To clarify the basic ideas of VIM, we consider the following differential equation:

$$Lu + Nu = g(t), \tag{10}$$

Where L is a linear operator, N is a nonlinear operator and $g(t)$ is a homogeneous term. According to VIM, we can write down a correction functional as follows:

$$u_{n+1}(t) = u_n(t) + \int_0^t \lambda(Lu_n(\tau) + N\tilde{u}(\tau) - g(\tau))d\tau \tag{11}$$

Where λ is a general lagrangian multiplier which can be identified optimally via the variational theorem, The subscript n indicates the n th approximation and u_n is considered as a restricted variation, i.e. $\delta\tilde{u} = 0$.

4. THE GAS DYNAMIC EQUATION IN GENERAL:

The classical gas dynamic equation is given by the formula [1]:

$$\frac{\partial}{\partial t}u(x, t) + \frac{u^2(x, t)}{2} \frac{\partial}{\partial x}u(x, t) - u(x, t)(1 - u(x, t)) = 0; \tag{12}$$

$u(x, t)$ =Degree of tempreture,
 t =Time from Zero Point, seconds,
 x =Distance from zero to 1,

Consider Eq. (12), and assume that initial condition as follow and assume that we have not any boundary condition,

$$u(x, 0) = g(x) \tag{13}$$

Where $g(x)$ =Initial condition of gas equation,

The exact solution by Adomians Decomposition method [12] has been expressed as follows:

$$v(x, t) = a(1 - e^{-x}) + at(1 - e^{-x}) - a^2t(1 - e^{-x})^2 - a^2t(e^{-x} - e^{-2x}) + \quad (14)$$

$$+ \frac{a^2t^2}{2}(1 - e^{-x}) + \frac{a^2t^2}{2}(1 + e^{-x} - e^{-2x} + ae^{-2x}) + \dots$$

5. APPLICATION OF HOMOTOPY PERTURBATION METHOD:

We consider Eq. (12) for gas dynamic equation with initial conditions as follows:
To solve Eq. (12) by means of HPM, we consider the following process after separating the linear and nonlinear parts of the equation.

A homotopy can be constructed as follows [20]:

$$H(u, p) = (1 - p)\left(\frac{\partial u(x, t)}{\partial t}\right) + p\left(\frac{\partial u(x, t)}{\partial t} + \frac{u^2(x, t)}{2} \frac{\partial u(x, t)}{\partial x} - u(x, t)(1 - u(x, t))\right). \quad (15)$$

Substituting $u = u_0 + pu_1 + \dots$ in to Eq. (15) and rearranging the resultant equation based on powers of p -terms, one has:

$$p^0 : \frac{\partial}{\partial t} u_0(x, t) = 0, \quad (16)$$

$$p^1 : \frac{1}{2} u_0^2(x, t) \frac{\partial}{\partial x} u_0(x, t) + u_0^2(x, t) + \frac{\partial}{\partial t} u_1(x, t) - u_0(x, t) = 0. \quad (17)$$

$$p^2 : 2u_0(x, t)u_1(x, t) + \frac{\partial}{\partial t} u_2(x, t) + \frac{1}{2} u_0^2(x, t) \frac{\partial}{\partial x} u_1(x, t) - \quad (18)$$

$$- u_1(x, t) + u_0(x, t)u_1(x, t) \frac{\partial}{\partial x} u_0(x, t) = 0$$

With the following initial conditions:

$$u_0(x, 0) = g(x) = a(1 - e^{-x}), \quad (19)$$

$$u_i(0, t) = 0, \quad i = 1, 2, \dots$$

$u(x, t)$ may be written as follows by solving the Eqs (16), (17) and (18):

$$u_0(x, t) = a(1 - e^{-x}) \quad (20)$$

$$u_1(x, t) = -\frac{a}{2}(a^2e^{-x} - 2a^2e^{-2x} + a^2e^{-3x} + 2a - 4ae^{-x} + 2ae^{-2x} - 2 + 2e^{-x})t \quad (21)$$

$$u_2(x, t) = -\frac{at^2}{8}(-8a^2 + 12a + 32a^2e^{-x} + a^4e^{-x} + 4e^{-x} - 24ae^{-x} - 8a^4e^{-2x} \quad (22)$$

$$- 40a^2e^{-2x} + 12a^3e^{-4x} + 16a^2e^{-3x} + 12ae^{-2x} + 18a^4e^{-3x} - 16a^4e^{-4x}$$

$$+ 5a^4e^{-5x} - 4 - 12a^3e^{-x} + 36a^3e^{-2x} - 36a^3e^{-3x})$$

In the same manner, the rest of components were obtained using the maple package.

According to the HPM, we can conclude that:

$$u(x, t) = \lim_{p \rightarrow 1} u(x, t) = u_0(x, t) + u_1(x, t) + u_2(x, t) + \dots \quad (23)$$

Therefore, substituting the values of $u_0(x, t)$, $u_1(x, t)$ and $u_2(x, t)$ from Eqs. (20), (21) and (22) in to Eq. (23) yields:

$$\begin{aligned}
 u(x, t) = & a(1 - e^{-x}) - \frac{a}{2}(a^2e^{-x} - 2a^2e^{-2x} + a^2e^{-3x} + 2a - 4ae^{-x} + 2ae^{-2x} - 2 + 2e^{-x})t \\
 & - \frac{at^2}{8}(-8a^2 + 12a + 32a^2e^{-x} + a^4e^{-x} + 4e^{-x} - 24ae^{-x} - 8a^4e^{-2x} - 40a^2e^{-2x} \\
 & + 12a^3e^{-4x} + 16a^2e^{-3x} + 12ae^{-2x} + 18a^4e^{-3x} - 16a^4e^{-4x} + 5a^4e^{-5x} - 4 \quad (24) \\
 & - 12a^3e^{-x} + 36a^3e^{-2x} - 36a^3e^{-3x})
 \end{aligned}$$

Fig. 1 shows the results of gas dynamics equation by HPM and difference with ADM result after substituting $a = .01$ and plotting for $0 < x < 1$ and $0 < t < 1$. And the variational iteration formula is obtained in the form:




Figure 1: Result of gas dynamics equation using (HPM) method [$0 < x < 1$ and $0 < t < 20$]. (A...3D, B... $x = 0.5$, C... $t = 20$)

6. APPLICATION OF VARIATIONAL ITERATION METHOD:

In this section, variational iteration method is developed for solving gas dynamic equation. Consider gas dynamics equation in Eq.12.

To solve Eq. (12) via VIM, one has to find the Langrangian multiplier, which can be identified by substituting Eq. (12) into Eq. (11), upon making it stationary leads to the following:

$$\begin{aligned}
 1 - \lambda'|_{\tau=t} &= 0 \\
 \lambda|_{\tau=t} &= 0 \\
 \lambda''|_{\tau=t} &= 0
 \end{aligned} \tag{25}$$

Solving the system of Eq. (25), Yields:

$$\lambda(\tau) = -1, \tag{26}$$

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t - \left\{ \frac{du_n(x, \tau)}{d\tau} + \frac{1}{2}u_n^2(x, \tau) \frac{du_n(x, \tau)}{d\tau} - u_n(x, \tau)(1 - u_n(x, \tau)) \right\} dt \tag{27}$$

Now, we assume that the initial approximation has the form:

$$u_0(x, t) = a(1 - e^{-x}) \quad (28)$$

Using the above variational formula (27), we have:

$$u_1(x, t) = u_0(x, t) + \int_0^t - \left\{ \frac{du_0(x, \tau)}{d\tau} + \frac{1}{2}u_0^2(x, \tau) \frac{du_0(x, \tau)}{d\tau} - u_0(x, \tau)(1 - u_0(x, \tau)) \right\} dt \quad (29)$$

Substituting Equation (28) in to Equation (29) and after simplification, we have:

$$u_1(x, t) = \frac{a}{2}(2 - 2e^{-x} - a^2te^{-x} + 2a^2te^{-2x} - a^2te^{-3x} + 2t - 2at + 4ate^{-x} - 2te^{-x} - 2ate^{-2x}) \quad (30)$$

In the same way, we obtain $u_2(x, t)$ that because it is too big we neglect to write that, Fig. 2 shows results of gas dynamics equation by HPM and difference with ADM result, after substituting $a = .01$ and plotting for $0 < x < 1$ and $t > 0$.




Figure 2: Result of gas dynamic equation using (VIM) method [$0 < x < 1$ and $t > 0$]. (A...3D, B... $x = 0.5$, C... $t = 20$)

7. CONCLUSIONS

In this paper, variational Iteration and homotopy perturbation methods have been successfully applied to find the solution of gas dynamics equation. Solution of gas dynamic equation shows that the results of proposed methods are in agreement with each other. The homotopy perturbation method which was used to solve gas dynamic equation seems to be very easy. There is less computation needed in comparison with the other methods (numerical and analytical methods). The results obtained here clearly show, that the variational iteration method, is capable of solving gas dynamic equation, with a rapid convergent successive approximation, without any restrictive assumptions or transformations, that may change the physical behavior of the problem. both of VIM and HPM solution are in agreement with decomposition method.

REFERENCES

- [1] Evans, D.J. and Bulut, H., (2002), A New approach to the gas dynamics equation: An Application of the Decomposition method. *Intern. J. Computer Math.*, vol.79(7).
- [2] He, J. H., (2004), The homotopy perturbation method for nonlinear oscillators with discontinuities, *Applied Mathematics and Computation*, 151, 287-292.
- [3] Siddiqui, A.M., Mahmood, R. and Ghori, Q.K., (2006), Thin film flow of a third grade fluid on a moving belt by Hes homotopy perturbation method, *Internat J. Nonlinear Sci. Numer. Simul*, 7(1), 714.
- [4] Abbasbandy, S., (2006), Iterated Hes homotopy perturbation method for quadratic Riccati differential equation, *Appl. Math. Comput.*, 175, 581-589.
- [5] Ganji, J. D. D. and Sadighi, A., (2006), Application of Hes Homotopy perturbation Method to Nonlinear Coupled Systems of Reaction-diffusion Equations, *International Journal of Nonlinear Science and Numerical Simulation*, 7(4), 411-418.
- [6] Rafei, M. and Ganji, D. D., (2006), Explicit Solutions of Helmholtz Equation and Fifth order Kdv Equation using Homotopy perturbation Method, *Int. J. Nonl. Sci. and Num. Simu.*, 7(3), 321-328.
- [7] Ganji, D. D. and Rajabi, A., (2006), Assessment of homotopy-perturbation and perturbation methods in heat radiation equations, *Internat. Comm. Heat Mass Transfer*, 33, 391-400.
- [8] He, J.H., (1999), Homotopy perturbation technique, *J. Comput. Math. Appl. Mech. Eng.*, 17(8), 257-262.
- [9] He, J.H., (2000), A coupling method of homotopy technique and perturbation technique for nonlinear problems, *Int. J. Non-Linear Mech.*, 35(1), 37-43.
- [10] He, J.H., (2005), Application of homotopy perturbation method to nonlinear wave equations, *Chaos, Solitons and Fractals*, 26(3), 695-700.
- [11] He, J.H., (2006), Homotopy perturbation method for solving boundary value problems, *Physics Letter A*, 350(12), 87-88.
- [12] Khaleghi, H., Ganji, D. D. and Sadighi, A., (2007), Application of variational iteration and Homotopy-perturbation methods to nonlinear heat transfer equations with variable coefficients, *Numerical Heat Transfer, Part A*, 52(1), 25-42.
- [13] He, J.H., (2006), New interpretation of Homotopy perturbation method. *Int. J. Mod. Phys. B*, 20(18), 2561-2568.
- [14] Sadighi, A. and Ganji, D. D., (2007), Solution of the generalized nonlinear Boussinesq equation using homotopy perturbation and variational iteration methods, *International Journal of Nonlinear Science and Numerical Simulation*, 8(3), 435-444.
- [15] Ganji, D. D. and A. Sadighi, (2007), Application of homotopy perturbation and variational iteration methods to nonlinear heat transfer and porous media equations, *Journal of Computational and Applied mathematics*, 207, 24-34.
- [16] He, J.H., (1999), Variational iteration method kind of non-linear analytical technique: some examples, *Internat. J. Nonlinear Mech.*, 34(4), 699-708.
- [17] He, J.H., (2000), Variational iteration method for autonomous ordinary differential systems, *Applied Mathematics and Computation*, 114, 115-123.
- [18] Abdou, M. A. and Soliman, A.A., (2005), Variational iteration method for solving Burgers and coupled Burgers equations, *J. Comput. Appl. Math.*, 181(2), 245-252.
- [19] Wazwaz, A. M., (2006), The variational iteration method for rational solutions for KdV, K(2,2) Burgers, and cubic Boussinesq equations, *J. Comput. Appl. Math.*, in press.
- [20] Sweilam, N. H. and Khader, M. M., (2007), Variational iteration method for one dimensional nonlinear thermoelasticity. *Chaos Soliton Fract.*, 32, 145-149.
- [21] Abbasbandy, S., (2006), Homotopy perturbation method for quadratic Riccati differential equation and comparison with Adomian's decomposition method, *Appl. Math. Comput.*, 172, 485-490.
- [22] Wazwaz, A.M., (2006), A comparison between the variational iteration method and Adomian decomposition method, *J. Comput. Appl. Math.*, in press.
- [23] Biazar, J., Babolian, E. and Islam, R., (2004), Solution of the system of ordinary differential equations by Adomian decomposition method, *Applied Mathematics and Computation*, 147, 713-719.



M. Kazemi-Sangsereki was born in Qaemshahr, Mazandaran, Iran, in 1984. He obtained his B.Sc. and M.Sc. degrees in Mechanical Engineering in Babol Noshiravani University of Technology, Babol, Iran. His research interests mostly include computational fluid dynamic, internal combustion engines and gas dynamics.



D. D. Ganji was born in Babol, Iran in 1975. He obtained his B.Sc. degree in Mechanical Engineering from University of Tehran in 1989. In advance, Dr. Ganji got his M.Sc. and Ph.D. degree in Mechanical Engineering from Tarbiat Modares University, Tehran in 1992 and 2004 respectively. Currently, Dr. Ganji is associate professor in department of Mechanical Engineering, Noshiravani University of Technology, Babol, Iran. He has several refereed publications in different fields of Mechanical Engineering and applied mathematics such as FDM, FVM, inverse problems, HPM, spray modeling and aerodynamics. In addition, he is editor in several international journals such as "International Journal of Nonlinear Dynamics and Engineering Science", "International Journal of Nonlinear Science and Numerical Simulation", and "International Journal of Differential Equations".



M. Gorji-Bandpy was born in Ghaemshahr, Iran in 1954. He obtained his B.Sc. and M.Sc. degree in Mechanical Engineering from University of Tehran in 1976 and 1978 respectively. In advance, Prof. Gorji-Bandpy got his Ph.D. degree in Mechanical engineering from University of Wales (UWCC) in 1991. Currently, Prof. Gorji-Bandpy is professor in department of mechanical engineering in Noshiravani University of Technology, Babol, Iran. He has several refereed publications in different fields of Mechanical Engineering such as fluid mechanics, turbomachinery, hydro engineering and renewable energies. In addition, he has editorial responsibility in several international journals.
