



Geographic distributional patterns of the genus *Bombus* (Bombini, Apidae: Hymenoptera) in northern Pakistan

Anjum SUHAİL^{*1}, Arshed Makhdoom SABİR¹, Muhammad ASGHAR¹, Muhammad Ather RAFİ², Abdul QADİR³

¹ Department of Agri. Entomology, University of Agriculture, Faisalabad, Pakistan

² National Insect Museum & Insect Pests Informatics, NARC, Islamabad, Pakistan

³ Govt. Islamia College, Sambrial, Sialkot, Pakistan

Abstract

The distributional patterns of the species of genus *Bombus* were analyzed by using cluster analysis which grouped the sites into three main geographical groups. Cluster 1 comprised of two sites i.e., Samgam and Doarian as these sites were located in the same ecological region i.e., Himalayan moist temperate forest. Cluster 2 consists of Sermik and Rama showing similar patterns of occurrence as dry temperate coniferous and steppic forest adjoin with alpine dry steppe regions while cluster 3 represented by four sites which was further divided into two sub groups viz., Naltar, Gilgit and Darkot, Hunza. Naltar and Gilgit both located in steppic forest and alpine dry steppe. Similarly sub group Darkot, Hunza belong to same ecological region i.e., alpine meadows and sub-alpine scrub. A total of 13 species of genus *Bombus* were grouped in to three clusters. Cluster 1 was represented by species (*B. (Sibricobombus) asiaticus* Morawitz, *B. (Bombus) tunicatus* Smith, *B. (Pyrobombus) subtypicus* Skorikov and *B. (Subterraneobombus) melanurus* Lepletier). These species were more or less evenly distributed in sampling sites and have wide ecological amplitude. *B. (Pyrobombus) biroi* Vogt, *B. (Bombus) lucorum* Linnaeus, *B. (Melanobombus) semenovianus* Skorikov and *B. (Mendacibombus) marussinus* Skorikov formed cluster 2. These species are less evenly distributed while cluster 3 was consisted of five species viz., *B. (Melanobombus) rufofasciatus* Smith, *B. (Alpigenobombus) kashmirensis* Friese, *B. (Orientalibombus) haemorrhoidalis* Smith, *B. (Mendacibombus) himalayanus* Skorikov and *B. (Mendacibombus) avinoviellus* Skorikov. These species are moderately distributed in the sampling areas with narrow ecological range as compared to cluster 1. These patterns reflect the complex geographic history of the fauna in the region.

Key words: Cluster analysis, Distributional patterns, Genus *Bombus*, Geography, Northern Pakistan

1. Introduction

The Northern Pakistan covers overall Northern Areas (NA), Azad Jammu & Kashmir (AJK), the upper region of North West Frontier Province (NWFP) including Chitral and some parts of the central and northern regions of Pakistan and is endowed with a great variety of flora and fauna owing to its diverse array of altitude, rainfall and climate. It is among the regions where the earth's biological wealth is most distinctive and rich (Malcolm et al., 2002). Three famous mountain ranges viz., Himalayas, the Karakorum and the Hindukush which are one of the largest mountainous regions of the world extends over an area of 132700 km² and lies between 34° 0' to 36° 50' N and 71° 12' to 75° 0' E (Hashmi and Shafiqullah, 2003) give rise to a unique blend of habitats and biological communities (Sheikh, 2000). This region represents one of the world's biologically richest ecosystems. This is due to the extreme altitudinal differences and associated flora and fauna and nowhere in the world can such diversity be seen as in this region (Anonymous, 1993).

Pollinators play a pivotal role in flowering plant reproduction and fruit set for agricultural and wild plant communities (Buchmann and Nabhan, 1996; Accorti, 2000) and bumblebees provide a vital ecological service in this regard (Sabir et al., 2007). In recent years many bumblebee (*Bombus*) species have shown serious declines in

* Corresponding author / Haberleşmeden sorumlu yazar: dranjumsuhailuaf@yahoo.com

abundance and marked contractions in many parts of the world (Rasmont, 1995; Buchmann and Nabhan, 1996; Westrich, 1996; Kevan and Phillips, 2001). There has been growing concern about the decline of natural populations of several bumblebee species in many parts of the world and many researchers think that their populations are exposed to strong unfavourable pressure because of human activities (Osborne et al., 1991). This decline may culminate in the local extinction of species which has already been recorded in various parts of the world (Rasmont, 1995; Kevan and Phillips, 2001). As the most distinctive and rich faunal and floral wealth of earth is in northern Pakistan so there is dire need to study its fauna particularly the key groups like bumblebees which play a pivotal role in conservation of ecosystems (Corbet et al., 1991; Banaszak, 1992; Barbattini, 1994; Porrini, 1999). The plight of our bumblebee fauna deserves particular attention because their loss will have negative consequences for different habitats. Owing to already scarce information, knowing bumblebees is thus extremely important to improve the pollination of our cultivated crops, to restore native plants (Sabir et al., 2008) and to evaluate the degree of deterioration of certain areas. So the main aim of this study is to investigate the geographical patterns of bumblebees in northern Pakistan.

2. Materials and methods

The studies on geographical distribution of bumblebees were conducted in the northern Pakistan, during two consecutive years 2006 and 2007. Bumblebees were searched for and caught when they landed on or right after leaving the flowers with entomological handnet despite other methods of sampling (Dafni, 1992) not only because it is an easy handling method but it also allows assessing the frequency of specific flowers visited (Quaranta et al., 2004). The data were collected on the basis of altitudinal succession of forage plants and their associated bumblebees. The bumblebees were identified up to the species level by following Williams, 1991 while the plant material (flowers) etc., was identified with the help of available literature (Stewart, 1982 and Malik and Farooq, 1984) and "Flora of Pakistan" (series 1- 204) edited by S. I. Ali and E. Nasir.

Cluster analysis (CA) was used for grouping of data set on the basis of spatial similarities (Kent and Coker, 1992; Angeler et al., 2007; Sánchez-Carrillo et al., 2007 and Qadir et al., 2008). The distributional data set was subjected to CA to identify clusters of the sampling sites indicating their similarity based on geographical similarities in the study area. CA was performed on the presence/ absence data of bumblebees studied at different sampling sites over the whole period. Euclidean distances were chosen as a measure of similarity that uses analysis of variance to evaluate the distances between clusters, attempting to minimize the sum squares of any two clusters that can be formed at each step (Kent and Coker, 1992 and Qadir et al., 2008).




Figure 1. Sampling sites in the northern Pakistan.

3. Results

It is evident (Table 1) that bumblebees of genus *Bombus* in the northern Pakistan consist of 13 species represented by 9 subgenera which are confined in the area ranging from 2291 to 5344m above sea level (a.s.l).

Table 1. Binary (presence/ absence) data of genus *Bombus* in northern Pakistan.

Bumblebees (Genus, Sub genus, Species)	Geographical Distribution							
	Hunza	Rama	Darkot	Gilgit	Naltar	Doarian	Samgam	Sermik
<i>B. (Mendacibombus) avinoviellus</i> Skorikov	-	-	-	+	+	+	+	-
<i>B. (Mendacibombus) himalayanus</i> Skorikov	-	-	+	-	+	+	+	-
<i>B. (Mendacibombus) marussinus</i> Skorikov	+	-	+	-	+	-	-	-
<i>B. (Orientalibombus) haemorrhoidalis</i> Smith	-	-	-	-	-	+	+	-
<i>B. (Subterraneobombus) melanurus</i> Lepletier	+	+	+	+	+	-	-	+
<i>B. (Alpigenobombus) kashmirensis</i> Friese	+	-	+	-	-	+	+	-
<i>B. (Pyrobombus) subtypicus</i> Skorikov	-	+	+	-	+	-	-	+
<i>B. (Pyrobombus) biroi</i> Vogt	-	-	+	-	-	-	-	-
<i>B. (Bombus) tunicatus</i> Smith	+	+	+	+	+	+	+	+
<i>B. (Bombus) lucorum</i> Linnaeus	-	-	+	+	+	-	-	-
<i>B. (Sibricobombus) asiaticus</i> Morawitz	+	-	+	+	+	+	+	+
<i>B. (Melanobombus) semenovianus</i> Skorikov	+	-	+	+	+	-	-	-
<i>B. (Melanobombus) rufofasciatus</i> Smith	+	+	+	-	-	+	+	-

3.1 Distribution of Bumblebees

Cluster analysis (CA) performed on the bumblebees' data set to evaluate geographical variations among different sampling sites. CA grouped the sites into three main geographical groups. Cluster 1 comprised of two sites i.e., Samgam and Doarian. This is because of the fact that these sites were located in the same ecological region i.e., Himalayan moist temperate forest. Cluster 2 consists of Sermik and Rama showing similar patterns of occurrence of genus *Bombus* as both ecological regions (Dry temperate coniferous and steppic forest and alpine dry steppe, respectively) adjoin there. Cluster 3 represented by four sites which was further divided into two sub groups viz., Naltar, Gilgit and Darkot, Hunza. Naltar and Gilgit both located in steppic forest and alpine dry steppe. Similarly sub group Darkot, Hunza belong to same ecological region i.e., permanent snow, alpine meadows and sub-alpine scrub.




Figure 2. Dendrogram showing different clusters of sampling sites on the basis of abundance data of genus *Bombus* in northern Pakistan.

A total of 13 species of genus *Bombus* (Table 1) were grouped in to three clusters (Fig. 2) on the basis of presence/ absence data. Cluster 1 was represented by species (*B. (Sibricobombus) asiatics* Morawitz, *B. (Bombus) tunicatus* Smith, *B. (Pyrobombus) subtypicus* Skorikov and *B. (Subterraneobombus) melanurus* Lepletier). These species were more or less evenly distributed in sampling sites and have wide ecological amplitude. *B. (Pyrobombus) biroi* Vogt, *B. (Bombus) lucorum* Linnaeus, *B. (Melanobombus) semenovianus* Skorikov and *B. (Mendacibombus) marussinus* Skorikov formed cluster 2. These species are less evenly distributed while cluster 3 was consisted of five species viz., *B. (Melanobombus) rufofasciatus* Smith, *B. (Alpigenobombus) kashmirensis* Friese, *B. (Orientalibombus) haemorrhoidalis* Smith, *B. (Mendacibombus) himalayanus* Skorikov and *B. (Mendacibombus) avinoviellus* Skorikov. These species are moderately distributed in the sampling areas with narrow ecological range as compared to cluster 1.




Figure 3. Cluster analysis showing association of genus *Bombus* in northern Pakistan.

B. asi (*B. asiatics* Morawitz), *B. tun* (*B. tunicatus* Smith), *B. sub* (*B. subtypicus* Skorikov), *B. mel* (*B. melanurus* Lepletier), *B. bir* (*B. biroi* Vogt), *B. luc* (*B. lucorum* Linnaeus), *B. sem* (*B. semenovianus* Skorikov), *B. mar* (*B. marussinus* Skorikov), *B. ruf* (*B. rufofasciatus* Smith), *B. kas* (*B. kashmirensis* Friese), *B. hae* (*B. haemorrhoidalis* Smith), *B. him* (*B. himalayanus* Skorikov) and *B. avi* (*B. avinoviellus* Skorikov)

4. Conclusions

The genus *Bombus* Latreille belongs to the monotypic tribe Bombini (Hymenoptera: Apidae), comprising 239 known species in the world over (Williams, 1985, 1998). *B. (Mendacibombus) avinoviellus* Skorikov and *B. (Mendacibombus) himalayanus* Skorikov are west Himalayan species (Williams, 1991). *B. (Mendacibombus) marussinus* Skorikov is a central Asian species also recorded from Pakistan and Kashmir (Williams, 1991)- the Hindu Kush (Reinig, 1940; Tkalcu, 1969) and the Pamir (Skorikov, 1910; Reinig, 1930; Skorikov, 1931).

B. (Orientalibombus) haemorrhoidalis Smith is a Himalayan and South East Asian species (Williams, 1991). It is also known from Kashmir and Pakistan (Richards, 1929; Frison, 1933), Assam (Bingham, 1897; Richards, 1929), south eastern Tibet (Wang, 1988), Sikkim (Friese, 1918; Richards, 1929, 1930; Frison 1935), Darjiling Bengal (Friese, 1918; Richards, 1929; Frison, 1933, 1935), Nepal (Richards, 1929; Frison, 1935; Tkalcu, 1974b), Uttar Pradesh (Richards, 1929; Frison, 1935), Himachal Pradesh (Smith, 1879; Richards, 1929; Frison, 1933, 1935), Vietnam (Tkalcu, 1968), Laos (Richards, 1929; Tkalcu, 1968), Thailand (Sakagami & Yoshikawa, 1961; Tkalcu, 1968); Burma (Frison, 1933, 1935; Skorikov, 1938; Tkalcu, 1968, 1989), Yunnan (Wang, 1987).

B. (Subterraneobombus) melanurus Lepletier is a widespread, but primarily central Asian species (Williams, 1991) also known to occur from Pakistan (Frison, 1935), the Hindu Kush (Reinig, 1940; Richards, 1951; Tkalcu, 1969), Kashmir, Xinjiang [Kunlun Shan] (Morawitz, 1886), the Hindu Raj, Karakoram, Ladakh, Zanskar, Great Himalaya and Pir Panjal ranges, in montane coniferous forest, alpine scrub and steppe and in high, subtropical semi-dscrpt (Williams, 1985), Tibet (Richards, 1928b, 1930; Wang, 1982), Mongolia (Pittioni, 1939; Skorikov, 1933a; Bischoff, 1936; Tkalcu,

1974a), the Tien Shan (Skorikov, 1931; Bischoff, 1936; Panfilov, 1957), the Pamir (Skorikov, 1912, 1931; Reinig, 1930, 1934; Bischoff, 1931), Nepal (Tkalcū, 1974b), Sikkim (Richards, 1928b, 1930), Qinghai (Morawitz, 1886; Skorikov, 1912; Panfilov, 1957; Tkalcū, 1961; Wang, 1982), Gansu (Bischoff, 1936), Inner Mongolia (Skorikov, 1933a), Shanxi (Yasumatsu, 1951), Caucasus (Skorikov, 1931) and Turkey (Skorikov, 1931; Reinig, 1971).

B. (Alpigenobombus) kashmirensis is Tibetan species known from Kashmir- the Hindu Raj, Ladakh, Zanskar, Great Himalaya and Pir Panjal ranges (Williams, 1991), Tibet (Richards, 1930; Wang, 1982), Nepal (Tkalcū, 1974b; Gansu (Morawitz, 1880; Bischoff, 1936), Qinghai (Morawitz, 1886; Bischoff, 1936; Wang, 1982), Sichuan (Morawitz, 1890; Skorikov, 1933b; Wang, 1982), Guangxi (Wang, 1982).

B. (Pyrobombus) subtypicus Skorikov is a central Asian species (Williams, 1991). It is known from Kashmir (Williams, 1985), the Hindu Kush (Reinig, 1934, 1940; Richards, 1951; Tkalcū, 1969), Pakistan (Tkalcū, 1989), the Tien Shan (Morawitz, 1880; Skorikov, 1914, 1931; Krüger, 1943) and the Pamir (Reinig, 1930, 1934; Skorikov, 1931).

B. (Pyrobombus) biroi Vogt is a central Asian species (Williams, 1991) also known from the Hindu Kush (Reinig, 1934, 1940; Tkalcū, 1969), Kashmir- Hindu Raj, Ladakh, Zanskar, Great Himalaya, Pir Panjal ranges, Mt. Apharwat. Rumbak (Zanskar ranges), Panichar near Suru and Tungri (Williams, 1985), the Tien Shan (Vogt, 1911; Skorikov, 1931; Krüger, 1943 and the Pamir (Cockerell, 1922; Reinig, 1930, 1934; Skorikov, 1931).

B. (Bombus) tunicatus Smith is a Himalayan species (Williams, 1991) also known from Pakistan (Frison, 1933, 1935), Kashmir- southern side of the Great Himalayan Range, the Pir Panjal range, Jhelum valley watershed (Vale of Kashmir and the Kishanganga valley), Hindu Kush (Tkalcū, 1969), Indian plains and Calcutta (Dover, 1922), Sikkim (Bingham, 1897; Friese, 1918; Skorikov, 1933b), Uttar Pradesh (Bingham, 1897; Frison, 1935), Himachal Pradesh (Friese, 1909; Cockerell, 1917; Frison, 1935) and Nepal (Tkalcū, 1974b).

B. (Bombus) lucorum Linnaeus is most widespread of all bumblebee species, but it is not quite circumpolar (Williams, 1991). It is recorded from Pakistan, Kashmir- the Hindu Kush (Reinig, 1940), Hindu Raj, Karakoram ranges, southern side of the Great Himalaya range, the Pir Panjal range, in the upper montane coniferous forest and in subalpine scrub (Williams, 1985), Nepal (Tkalcū, 1974b), Sikkim (Friese, 1918), Tibet (Krüger, 1951, 1958; Panfilov, 1957; Wang, 1982, 1988), Yunnan (Wang, 1987), Sichuan (Wang, 1982), Gansu (Bischoff, 1936; Krüger, 1951, 1958), in central Asia its distribution reaches southwards to the Tien Shan (Morawitz, 1880; Skorikov, 1931; Krüger, 1954; Panfilov, 1957, 1981), the Pamir (Reinig, 1930, 1934; Skorikov, 1931; Krüger, 1951, 1958; Panfilov, 1981), Turkey and the Caucasus (Krüger, 1951; Panfilov, 1981; Rasmont, 1984) to the Elburz (Rasmont, 1984) and from across the northern U.S.S.R. (Krüger, 1951; Panfilov, 1981) to Mongolia (Morawitz, 1880; Vogt, 1909; Krüger, 1954, 1958; Tkalcū, 1974a; Panfilov, 1981). Inner Mongolia (Reinig, 1936; Panfilov, 1981; Wang, 1982), Heilongjiang (Krüger, 1954, 1958), North Korea (Kim & Ito, 1987), Hokkaido in Japan (Tkalcū, 1962; Sakagami & Ishikawa, 1969) and Kamchatka (Bischoff, 1930; Krüger, 1951; Panfilov, 1981), Europe (Krüger, 1951; LÖken, 1973; Alford, 1975; Rasmont, 1984) and from Alaska almost to Hudson Bay (Milliron, 1971).

B. (Sibricobombus) asiaticus Morawitz is a central Asian and Tibetan species (Williams, 1991) also found in the Hindu Kush (Reinig, 1940; Richards, 1951; Tkalcū, 1969), Kashmir- Hindu Raj, Karakoram, Ladakh and Zanskar ranges and from across the Great Himalayan range from the northern side to at least as far as Gumri (Williams, 1985), Nepal (Tkalcū, 1974a), Tibet (Richards, 1930; Wang, 1982), Mongolia (Vogt, 1909), the Altai (Skorikov, 1931), the Tien Shan (Morawitz, 1875, 1880; Vogt, 1911; Skorikov, 1931; Panfilov, 1957), the Pamir (Friese, 1931; Reinig, 1930, 1934; Skorikov, 1931), Qinghai (Bischoff, 1936; Tkalcū, 1961; Wang, 1982) and Gansu (Bischoff, 1936).

B. (Melanobombus) semenovianus Skorikov is a central Asian species (Williams, 1991). It is known from Pakistan (Frison, 1935) and Kashmir (Hindu Raj, Karakoram, Ladakh, Zanskar, Great Himalaya ranges, Dras (Williams, 1991) and the Hindu Kush (Reinig, 1940; Richards, 1951; Tkalcū, 1969).

B. (Melanobombus) rufofasciatus Smith is a peri- Tibetan and also known from Kashmir- Hindu Raj, Great Himalaya, Pir Panjal ranges, Wangal valley, Sangisaid valley, Mt. Apharwat, Khilanmarg, Mt. Apharwat, Kishanganga valley, Great Himalaya range and Batakush (Williams, 1991), Arunachat Pradesh (Friese, 1918), Uttar Pradesh

(Richards, 1928a, 1930) Sikkim (Bingham, 1897; Friese, 1918), Gansu (Morawitz, 1880; Bischoff, 1936), Qinghai (Morawitz, 1886; Tkalcù, 1961; Wang, 1982), Tibet (Friese, 1918; Richards, 1928a, 1930; Wang, 1982, 1988), Sichuan (Morawitz, 1890; Wang, 1982) and Nepal (Tkalcù, 1974b).

As the economy of the northern Pakistan, despite tourism is mainly based on subsistence production of agricultural and horticultural crops in the form of a mountain agricultural landscape. Farmers always create their fields at the expense of forest cover and use a variety of methods to do so and burning is the most probable ancient way to obtain pastures and land for crops (Sheikh, 2000). Such anthropogenic activities are continuously deteriorating natural resources such as reduction in vegetation cover, deforestation, intensive cropping and habitat fragmentation which not only reduce the biodiversity but also a permanent threat for bumblebees' fauna and their conservation. Present study was the first effort to document the recent distribution of bumblebees' fauna keeping in view their conservation.

References

- Accorti, M. 2000. Impollinatori, economia e gestione delle risorse, In (Ed.) Pinzauti, M., Api e impollinazione, Regione Toscana, Dipartimento Sviluppo economico, Firenze, Italy. 219- 231.
- Alford, D. V. 1975. Bumblebees. Davis- Poynter, London, UK.
- Angeler, D. G., Sánchez-Carrillo, S., Rodrigo, M. A., Alvarez- Cobelas, M., Rojo, C. 2007. Does size structure reflect fish-mediated effects on water quality in a degraded semiarid wetland? *Environmental Monitoring and Assessment*. 125/1–3. 9- 17.
- Anonymous, 1993. International Center for Integrated Mountain Environment Development. Proceedings of International Symposium. Mountain Environment. Development: Constraints and Opportunities. Kathmandu, Nepal.
- Banaszak, J. 1992. Natural resources of wild bees in Poland. Pedagogical University, Bydgoszcz, Poland.
- Barbattini, R. 1994. Il ruolo delle api negli ecosistemi naturali ed agrari. L'Ape Nostra Amica. 27. 7- 12.
- Bingham, C. T. 1897. The fauna of British India, including Ceylon and Burma. Hymenoptera. Vol. I. Wasps and bees. London, UK.
- Bischoff, H. 1930. Entomologische Ergebnisse der schwedischen Kamtchatka-Expedition 1920- 1922. 29. Bombinae. (Hymen.) *Arkiv fÖr Zoology*. 21. 1- 6.
- Bischoff, H. 1931. Entomologische Ergebnisse der deutsch-russischen Alai-Pamir-Expedition 1928 (II). 4. Hymenoptera VII. Zur Kenntnis einiger Hummelnester aus dem Pamir. *Mitteilungen aus dem Zoologischen Museum in Berlin*. 16. 860- 864.
- Bischoff, H. 1936. Schwedisch-chinesische wissenschaftliche Expedition nach den nordwestlichen Provinzen Chinas, unter Leitung von Dr. Sven Hedin und Prof. Sü Ping-chang. Insekten gesammelt vom schwedischen Arzt der Expedition Dr. David Hummel 1927-1930. 56. Hymenoptera, 10, Bombinae. *Arkiv fÖr Zoology*. 27. 1- 27.
- Buchmann, S. L. Nabhan, G. P. 1996. The Forgotten Pollinators. Island Press, Washington, DC, USA.
- Cockerell, T. D. A. 1917. New social bees. *Psyche*. 24. 120-128.
- Cockerell, T. D. A. 1922. Bees in the collection of the U.S. National Museum. " Proceedings of the United States National Museum. 60. 20pp.
- Corbet, S. A., Williams, I. H. Osborne, J. L. 1991. Bees and the pollination of crops and wild flowers in the European Community. *Bee World*. 72. 47- 59.
- Dafni, A. 1992. Pollination ecology, a practical approach. Irl Press at Oxford University Press, New York.
- Dover, C. 1922. A note on bees of the genera *Xylocopa* and *Bombus* in the Indian Museum. *Record of the Indian Museum*. 24. 85- 89.
- Friese, H. 1909. Neue Varietäten von *Bombus* (Hym.). *Deutsche Entomologische Zeitchrift*. 1909. 673- 676.
- Friese, H. 1918. Über Hummelformen aus dem Himalaja, *Deutsche Entomologische Zeitchrift*. 1918. 81- 86.
- Friese, H. 1931. Über *Bombus* und *Psithyrus*. *Konowia*. 10. 300- 304.
- Frison, T. H. 1933. Records and descriptions of *Bremus* and *Psithyrus* from India (Bremidae: Hymenoptera). *Record of the Indian Museum*. 35. 331- 342.
- Frison, T. H. 1935. Records, notes and descriptions of *Bremus* from Asia (Bremidae: Hymenoptera). *Record of the Indian Museums*. 37. 339- 363.
- Hashmi, A. A. Shafiullah. 2003. Northern Areas Strategy for Sustainable Development. IUCN Pakistan, Northern Areas Programme, Gilgit. 111.
- Kent, M. Coker, P. 1992. Vegetation description and analysis: A practical approach. London: Belhaven.
- Kevan P. G. Phillips, T. P. 2001. The economic impacts of pollinator declines: an approach to assessing the consequences. *Conservation Ecology*. 5. 8.
- Kim, C. -W. & Ito, M. 1987. On the bumblebees from the Korean Peninsula (Hymenoptera: Bombidae). *Entomological Research Bulletin*. 13. 1- 42.
- Krüger, E. 1943. Über die Genitalanhänge einiger Männchen der Untergattung *Pratobombus* O. Vogt *Zeitschrift für Morphologie und Ökologie der Tiere*. 39. 527- 545.

- Krüger, E. 1951. Phänoanalytische Studien an einigen Arten der Untergattung *Terrestribombus* O. Vogt (Hymen. Bomb.). I. Teil. Tijdschrift voor Entomologie. 93 (1950). 141-197.
- Krüger, E. 1954. Phaenoanalytische Studien an einigen Arten der Untergattung *Terrrestribombus* O. Vogt. (Hymenoptera, Bombidae). II. Teil. Tijdschrift voor Entomologie. 97. 263- 298.
- Krüger, E. 1958. Phaenoanalytische Studien an einigen Arten der Untergattung *Terrestribombus* O. Vogt (Hymenoptera, Bombidae), III. Teil. Tijdschrift voor Entomologie. 101. 283- 344.
- LØken, A. 1973. Studies on Scandinavian bumble bees (Hymenoptera, Apidae). Norsk Entomologisk Tidsskrift. 20. 1- 218.
- Malcolm, J. R. Liu, C. Miller, L. B. Allnutt T. Hansen, L. 2002. Habitats at risk: global warming and species loss in terrestrial ecosystems. WWF Fund for Nature, Gland, Switzerland. 40.
- Malik, S. Farooq, S. 1984. Cultivated trees, Shrubs and Climbers of gardens of Pakistan: Taxonomic studies. PCSIR and Peshawar University.
- Milliron, H. E. 1971. A monograph of the western hemisphere bumblebees (Hymenoptera: Apidae, Bembiriae). I. The genera *Bombus* and *Megabombus* subgenus *Bombias*. Memoirs of the Entomological Society of Canada. 82. 81 pp.
- Morawilz, F. 1875. [Bees.] (Mellifera), In Fedtschenko, A.: Reise in Turkestan. II Zoologischer Teil. 303pp. Berlin.
- Morawilz, F. 1880. Ein Beitrag zur Bienen-Fauna Mittel-Asiens. Izvēstiya Imperatorskoi akademii nauk. 26. 337-379.
- Morawilz, F. 1886. Insecta in itinere cl. N. Przewalskii in Asia centrali novissime lecta. I: Apidae. Trudy Russkago entomologicheskago obshchestva. 20. 195- 229.
- Morawilz, F. 1890. Insecta a cl. G. N. Potanin in China et in Mongolia novissime lecta. XIV. Hymenoptera Aculeata. III. Apidae. Trudy Russkago entomologicheskago obshchestva. 24. 349- 385.
- Osborne, J. L. Williams, I. H. Corbet, S. A. 1991. Bees, pollination and habitat change in the European Community. Bee World. 72. 99- 116.
- Panfilov, D. V. 1957. [On the geographical distribution of bumble bees (*Bombus*) in China.] Acta geographica sinica. 23. 221- 239.
- Panfilov, D.V. 1981. [Maps 91-97.] In Gorodkov, K. B.: [Provisional atlas of the insects of the European part of the U.S.S.R. Maps 73-125] Leningrad, U.S.S.R.
- Pittioni, B. 1939. Neue und wenig bekannte Hummeln der Paläarktis (Hymenopt., Apidae). Konowia. 17(1938). 244- 263.
- Qadir, A. Malik, R. N. Husain, S. Z. 2008. Spatio-temporal variations in water quality of nullah Aik-tributary of the river Chenab, Pakistan. Environmental Monitoring and Assessment. 140. 43- 59.
- Quaranta, M., Ambroselli, S., Barro, P., Bella, S., Carini, A., Celli, G., Cogoi, P., Comba, L., Comoli, R., Felicioli, A., Floris, I., Intoppa, F., Longo, S., Maini, S., Manino, A., Mazzeo, G., Medrzycki, P., Nardi, E., Niccolini, L., Palmieri, N., Patetta, A., Piatti, C., Piazza, M. G., Pinzauti, M., Porporato, M., Porrini, C., D'Albore, G. R., Romagnoli, F., Ruiu, L., Satta A., Zandigiacomo, P. 2004. Wild bees in agroecosystems and seminatural landscapes. 1997- 2000 collection period in Italy. Bulletin of Insectology. 57. 11- 61.
- Rasmont, P. 1984. Les bourdons du genre *Bombus* Latreille sensu stricto en Europe occidentale et centrale (Hymenoptera, Apidae). Spixiana. 7. 135- 160.
- Rasmont, P. Ebmer, P. A., Banaszak J., Van Der Zanden, G. 1995. Hymenoptera Apoidea Gallica. Bulletin de la Société Entomologique de France, 100 (hors série). 10- 98.
- Reinig, W. F. 1930. Untersuchungen zur Kenntnis der Hummel fauna des Pamir-Hochlandes. Zoologische Ergebnisse der deutsch-russischen Alai-Pamir-Expedition der Notgemeinschaft der Deutschen Wissenschaft und der Akademie der Wissenschaften der U.D.S.S.R. Zeitschrift für Morphologie und Ökologie der Tiere. 17. 68- 123.
- Reinig, W. F. 1934. Entomologische Ergebnisse der deutsch-russischen Alai-Pamir-Expedition. 1928.(III). 7. Hymenoptera VIII (Gen, *Bombus* Fabr.)- Nachtrag. Deutsche entomologische Zeitschrift. 1933. 163- 174.
- Reinig, W. F. 1936. Beiträge zur Kenntnis der Hummelfauna von Mandschukuo (Hym., Apid.). Mitteilungen der Deutschen entomologischen Gesellschaft. 7. 2- 10.
- Reinig, W. F. 1940. Beiträge zur Kenntnis der Hummelfauna von Afghanistan. (Hym., Apid.). (Ergebnisse der Reise von H. und E. Kotzsch in den Hindukusch im Jahre 1936.) Deutsche entomologische Zeitschrift. 1940. 224- 235.
- Reinig, W. F. 1971. Zur Faunistik und Zoogeographie des Vorderen Orients. 3, Beitrag zur Kenntnis der Hummelen und Schmarotzerhummeln Anatoliens (Hym.. Apidae), Veröffentlichungen der Zoologischen Staatsammlung. 15. 139-165.
- Richards, O. W. 1928a. *Bombus* and *Volucella* in the Himalayas, Entomologist's Monthly Magazine. 64. 107- 108.
- Richards, O. W. 1929. A revision of the humble-bees allied to *Bombus orientalis* Smith, with the description of a new subgenus. Annals and Magazine of Natural History. 3. 378- 386.
- Richards, O. W. 1930. The humble-bees captured on the expeditions to Mt. Everest (Hymenoptera, Bombidae), Annals and Magazine of Natural History. 5. 633- 658.
- Richards, O. W. 1928b. On a collection of humble-bees (Hymenoptera, Bombidae) made in Ladakh by Col. R. Meinertzhagen. Annals and Magazine of Natural History. 2. 333- 336.

- Richards, O. W. 1951. The 3rd Danish expedition to Central Asia. Zoological results 5. Bombidae (Insccta) from Afghanistan. *Videnskabelige Meddelelser fra Dansk naturhistorisk Forening i KjØbenhavn*. 113: 191-199.
- Sabir, A. M., Suhail, A., Rafi, A., Ahmad, S., Saleem, M., Mahmood, K. 2008. Bumblebees belonging to Genus *Bombus* (Bombini: Apidae: Hymenoptera) of Northern Pakistan. 28th Pak. Congr. Zool. (Intern.), Govt. College Univ., Faisalabad, Pak. 74.
- Sabir, A. M., Suhail, A., Rafi, M. A., Mahmood, K., Ahmed, S. 2007. Foraging activity of bumblebees (*Bombus* Latr.) in relation to floral resources in agricultural and semi natural landscape. International Conference on Biological Resources of Pakistan: Problems, success and future perspectives" at University of Arid Agriculture, Rawalpindi., Pak. 25.
- Sakagami, S. F. & Ishikawa R. 1969. Note préliminaire sur la répartition géographique des bourdons japonais, avec descriptions et remarques sur quelques formes nouvelles ou peu connues. *Journal of the Faculty of Science, Hokkaido University*. 17. 152- 196.
- Sakagami, S. F. & Yoshikawa, K. 1961. Bees of Xylocopinae and Apinae collected by the Osaka City University Biological Expedition to Southeast Asia 1957-58, with some biological notes. *Nature and Life in Southeast Asia*. 1: 409- 444.
- Sánchez-Carrillo, S., Alatorre, L. C., Sánchez-Andrés R., Garatuza-Payán, J. 2007. Eutrophication and sedimentation patterns in complete exploitation of water resources scenarios: An example from northwestern semi-arid. *Environmental Monitoring and Assessment* doi:10.2007/s10661-006-9541-x.
- Sheikh, K. M. 2000. Ecological Studiesof Avifauna in the Naltar Valley, Northern Pakistan with a Conservation Perspective, Ph. D. dissertation. Department of Biological Sciences, Quaid-i- Azam University, Islamabad, Pakistan.
- Skorikov, A. S. 1910. [*Bombus mendax* Gerst. and its variations (Hymenoptera, Bombidae).] Russkoe éntomologicheskoe Obozrenie. 9. 328- 330.
- Skorikov, A. S. 1912. Neue Hummelformen (Hymenoptera, Bombidae). IV. Russkoe éntomologicheskoe Obozrenie. 12. 606- 610.
- Skorikov, A. S. 1914. *Pratobombus leuccopygos* [sic] (F.Mor.) et ses ariations (Hymenoptera, Bombidae) Russkoe entomologicheskoe Obozrenie. 14. 293- 294.
- Skorikov, A. S. 1931. Die Hummelfauna Turkestans und ihre Beziehungen zur zentralasiatischen Fauna (Hymenoptera, Bombidae). *Abhandlungen der Pamir-Expedition*. 8: 175- 247.
- Skorikov, A. S. 1933a. Zur Hummelfauna Japans und seiner Nachbarländer. *Mushi*. 6. 53- 65.
- Skorikov, A. S. 1933b. Zur Fauna und Zoogeographie der Hummeln des Himalaya. *Doklady Akademii nauk SSSR*. 5. 243- 248.
- Skorikov, A. S. 1938. Zoogeographische Gesetzmässigkeiten der Hummelfauna im Kaukasus, Iran und Anatolien (Hymenoptera, Bombinae). *Entomologicheskoe Obozrenie*. 27. 145- 151.
- Smith, F. 1979. Descriptions of new species of Hymenoptera in the collection of the British Museum. xxi+240 pp. London, UK.
- Stewart, R. R. 1982, In (Eds) Nasir, E. Ali., I. Flora of Pakistan. History exploration of plants in Pakistan and adjoining areas. University of Karachi, Pak.
- Tkalcù, B. 1961. Zur Hummelfauna der Umgebung Kuku-Nors (Hymenoptera, Bombinae), Casopis Ceskoslovenské společnosti entomologické. 58. 344- 379.
- Tkalcù, B. 1968. Revision der vier sympatrischen, homochrome geographische Rassen bildenden Hummelarten SO-Asiens (Hymenoptera, Apoidea, Bombinae). *Annotaciones Zoologicae et Botanicae*. 52. 1- 31.
- Tkalcù, B. 1969. Beiträge zur Kenntnis der Fauna Afghanistans (Sammelergebnisse von O. Jakes 1963- 64, D. Povolny 1965, D. Povolny & Fr. Tenora 1966, J. Simek 1965- 66, D. Povolny. J. Geisler, Z. Sebek & Fr, Tenura 1967). Bombinae, Apoidea, Hym. *Casopis Moravského musea v Brne*. 53. 189- 210.
- Tkalcù, B. 1974a. Ergebnisse der 1. und 2. mongolisch-tschechoslowakischen entomologisch-botanischen Expedition in der Mongolei. Nr. 29: Hymenoptera, Apoidca, Bombinae. *Sborník faunistických prací Entomologického oddělení Národního muzea v Praze*. 15. 25- 57.
- Tkalcù, B. 1974b. Eine Hummel-Ausbeute aus dem Nepal-Himalaya (Insccla, Hymenoptera, Apoidea, Bombinae). *Senckenbergiana biologica*. 55. 311- 349.
- Tkalcù, B. 1989. Neue Taxa asiatischer Hummeln (Hymenoptera, Apoidea). *Acta entomologica bohemoslovaca*. 86. 39- 60.
- Vogt, O. 1909. Studien über das Artproblem. 1, Mitteilung, über das Variieren der Hummeln. 1. Teil. *Sitzungsberichte der Gesellschaft naturforschender Freunde zu Berlin*. 1909. 28- 84.
- Vogt, O. 1911, Studien über das Artproblem. 2. Mittcilung. über das Variieren dcr Hummeln. 2. Teil. (Schluss). *Sitzungsberichte der (Gesellschaft naturforschender Freunde zu Berlin*. 1911. 31- 74.
- Wang, S.-f. 1982. Hymenoptera: Apidae - Bombus. *Insects of Xizang*. 2. 427- 447.
- Wang, S.-f. 1987. Bombus. In Huang, F.-s.: Forest insects of Yunnan. Yunnan, China.
- Wang, S.-f. 1988. Habitat use by bumble bees (*Bombus spp.*). *Ecological Entomology*. 13. 223- 237.

- Westrich, P. 1996. The problems of partial habitats. In: Matheson, A., S. L. Buchmann, C. Toole, P. Westrich, I. H. Williams, (Eds.). *The Conservation of Bees*, Linnean Society Symposium Series, vol. 18. Academic Press, London.
- Williams, P. H. 1985. A preliminary cladistic investigation of relationships among the bumble bees (Hymenoptera: Apidae). *Systemtic Entomology*. 10. 239-255.
- Williams, P. H. 1991. The bumble bees of the Kashmir Himalaya (Hymenoptera: Apidae: Bombini). *Bulletin of Natural History Museum, London (Entomology)*. 60/1. 1- 204.
- Williams, P. H. 1998. An annotated checklist of bumblebees with an analysis of patterns of description. *Bulletin of Natural History Museum, London (Entomology)*. 67. 79-152.
- Yasumalsu, K. 1951. *Bombus and Psithyrus* of Shansi, N. China (Hymenoptera, Apidae). *Mushi*. 22. 59- 62.

(Received for publication 28 December 2008)