



# On Some Difference Equations of Exponential Form

## *Üstel Biçimdeki Bazı Fark Denklemleri Üzerine*

Melih Göcen\*

Bülent Ecevit University, Art and Science Faculty, Department of Mathematics, Zonguldak, Turkey

### Abstract

In this paper, the local asymptotic behavior of positive solutions of some exponential difference equations

$$x_{n+1} = \frac{x_n + x_{n-k}}{1 + x_{n-k}e^{x_{n-k}}}, k \in \mathbb{N}, n = 0, 1, 2, \dots$$

are investigated where the initial conditions are arbitrary positive real numbers. Furthermore, some numerical examples are presented to verify our results.

2010 AMS-Mathematical Subject Classification Number: 39A10, 39A20

**Keywords:** Difference equation, Local stability, Positive solutions, Equilibrium point

### Öz

Bu makalede, başlangıç koşulları keyfi pozitif reel sayılar olmak üzere bazı

$$x_{n+1} = \frac{x_n + x_{n-k}}{1 + x_{n-k}e^{x_{n-k}}}, k \in \mathbb{N}, n = 0, 1, 2, \dots$$

üstel rasyonel fark denklemlerinin pozitif çözümlerinin lokal asimptotik davranışı araştırılmıştır. Ayrıca, sonuçlarımızı doğrulamak için bazı nümerik örnekler verilmiştir.

**Anahtar Kelimeler:** Fark denklem, Lokal kararlılık, Pozitif çözümler, Denge noktası

## 1. Introduction

In recent years, there has been a great interest in studying nonlinear difference equations and systems. Also, it is very interesting to investigate the behavior of solutions of nonlinear difference equations and to discuss the local asymptotic stability of their equilibrium points, see (5,6,7,10,12). We can see many papers and books concerning theory and applications of difference equations, see (2,3,8,9). Difference equations especially exponential type of difference equations have many applications in biology, biomathematics, bioengineering, population dynamics, genetics, etc. Therefore, the behavior of positive solutions of the difference equations of exponential form has been a great role in the theory of difference equations.

Firstly, El-Metwally et al. (4) studied the boundedness character, asymptotic behavior, periodic character of the positive solutions and stability of the positive equilibrium of the following population model:

$$x_{n+1} = a + bx_{n-1}e^{-x_n}, n = 0, 1, 2, \dots \quad (1.1)$$

where  $a, b$  are positive constants and the initial values  $x_{-1}, x_0$  are arbitrary non-negative numbers. Furthermore, authors considered  $a$  is the immigration rate and  $b$  the population growth rate. This equation may be viewed as a model in Mathematical Biology which was suggested by the people from the Harvard School of Public Health; studying the population dynamics of one species  $x_n$ .

In (11), Ozturk et al. investigated the global attractivity, the boundedness and the periodic nature of the exponential rational difference equation

$$y_{n+1} = \frac{\alpha + \beta e^{-y_n}}{\gamma + y_{n-1}}, n = 0, 1, 2, \dots \quad (1.2)$$

\*Corresponding author: [melih.gocen@beun.edu.tr](mailto:melih.gocen@beun.edu.tr)

Melih Göcen [orcid.org/0000-0001-8669-6122](https://orcid.org/0000-0001-8669-6122)

where the parameters  $\alpha, \beta$  and  $\gamma$  are positive real numbers and the initial conditions  $y_{-1}, y_0$  are arbitrary non-negative numbers.

Bozkurt (1) obtained results concerning the local and global behavior of positive solutions of the following difference equation:

$$y_{n+1} = \frac{\alpha e^{-y_n} + \beta e^{-y_{n-1}}}{\gamma + \alpha y_n + \beta y_{n-1}}, n = 0, 1, 2, \dots \tag{1.3}$$

where the parameters  $\alpha, \beta, \gamma$  and the initial conditions are arbitrary positive numbers.

Motivated by the above studies, our aim in this paper is to investigate the local asymptotic behavior of positive solutions of the following exponential rational difference equation:

$$x_{n+1} = \frac{x_n + x_{n-k}}{1 + x_{n-k} e^{x_{n-k}}}, k \in \mathbb{N}, n = 0, 1, 2, \dots \tag{1.4}$$

## 2. Preliminaries and Theorems

**Definition 2.1** Let  $I$  be an interval of real numbers  $\mathbb{R}$  and let  $f: I \times I \rightarrow I$  be a continuous function. Consider the difference equation

$$x_{n+1} = f(x_n, x_{n-1}), n = 0, 1, 2, \dots \tag{2.1}$$

where the initial conditions  $x_{-1}, x_0 \in I$ . We say that  $\bar{x}$  is an equilibrium point of Eq.(2.1) if  $\bar{x} = f(\bar{x}, \bar{x})$ .

### Definition 2.2

(i) The equilibrium  $\bar{x}$  of Eq.(2.1) is called locally stable if for every  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $x_{-1}, x_0 \in I$  with  $|x_0 - \bar{x}| + |x_{-1} - \bar{x}| < \delta$ , then  $|x_n - \bar{x}| < \epsilon$  for all  $n \geq -1$ .

(ii) The equilibrium  $\bar{x}$  of Eq.(2.1) is called locally asymptotically stable if it is locally stable, and if there exists  $\gamma > 0$  such that  $x_{-1}, x_0 \in I$  with  $|x_0 - \bar{x}| + |x_{-1} - \bar{x}| < \gamma$ , then  $\lim_{n \rightarrow \infty} x_n = \bar{x}$ .

(iii) The equilibrium  $\bar{x}$  of Eq.(2.1) is called a global attractor if for every  $x_{-1}, x_0 \in I$  we have  $\lim_{n \rightarrow \infty} x_n = \bar{x}$ .

(iv) The equilibrium  $\bar{x}$  of Eq.(2.1) is called globally asymptotically stable if it is locally stable and a global attractor.

(v) The equilibrium  $\bar{x}$  of Eq.(2.1) is called unstable if it is not stable.

**Definition 2.3** The linearized equation of (2.1) about the equilibrium point  $\bar{x}$  is

$$y_{n+1} = py_n + qy_{n-1} \tag{2.2}$$

where

$p = \frac{\partial f}{\partial x_n}(\bar{x}, \bar{x})$  and  $q = \frac{\partial f}{\partial x_{n-1}}(\bar{x}, \bar{x})$ . Then the characteristic equation of (2.2) is

$$\lambda^2 - p\lambda - q = 0. \tag{2.3}$$

The following result, known as the Linearized Stability Theorem, is very useful in determining the local stability character of the equilibrium point  $\bar{x}$  of Eq.(2.1).

### Theorem 2.1 (Linearized Stability)

Suppose that the function  $F$  is a continuously differentiable function defined on some open neighborhood of an equilibrium point  $\bar{x}$ . Then, the following statements are true:

(i) If all roots of (2.3) have absolute value less than one, then the equilibrium point of (2.1) is locally asymptotically stable.

(ii) If at least one of the roots of (2.3) has absolute value greater than one, then the equilibrium point  $\bar{x}$  of (2.1) is unstable.

## 3. Local Stability of Eq.(1.4)

Firstly, we show that Eq.(1.4) has a unique positive equilibrium point  $\bar{x}$ . The equilibrium points of Eq. (1.4) are the solutions of the equation

$$\bar{x} = \frac{\bar{x} + \bar{x}}{1 + \bar{x}e^{\bar{x}}}.$$

Set

$$f(x) = \frac{x+x}{1+x e^x} - x$$

then we have  $f\left(\frac{1}{2}\right) > 0$  and  $\lim_{x \rightarrow \infty} f(x) = -\infty$ .

Moreover, it follows from the

$$f'(x) = \frac{2 - 2x^2 e^x}{(1 + x e^x)^2} - 1 < 0$$

that Eq.(1.4) has a unique positive equilibrium point  $\bar{x}$ .

**Theorem 3.1** The positive equilibrium point  $\bar{x}$  of Eq.(1.4) is locally asymptotically stable.

**Proof.** The positive equilibrium point  $\bar{x}$  of Eq.(1.4) is the solution of the equation

$$\bar{x} = \frac{\bar{x} + \bar{x}}{1 + \bar{x}e^{\bar{x}}} \Leftrightarrow 1 + \bar{x}e^{\bar{x}} = 2 \Leftrightarrow \bar{x}e^{\bar{x}} = 1$$

and it is evaluated numerically as 0.56714. The linearized equation associated with Eq.(1.4) about the

equilibrium point  $\bar{x}$  is

$$x_{n+1} - \frac{1}{1 + \bar{x}e^{\bar{x}}}x_n - \frac{(1 - \bar{x}e^{\bar{x}} - 2\bar{x}^2 e^{\bar{x}})}{(1 + \bar{x}e^{\bar{x}})^2}x_{n-k} = 0. \tag{2.4}$$

**Case  $k = 1$**

The characteristic equation of Eq.(2.4) with  $k = 1$  about the equilibrium point  $\bar{x}$  is

$$\lambda^2 - (0.5)\lambda + 0.28357 = 0. \tag{2.5}$$

The roots of Eq.(2.5) are  $\lambda_{1,2} = 0.25 \pm 0.47018i$  and the absolute value of each root is less than one. It follows from the Theorem 2.1 that the positive equilibrium point  $\bar{x}$  of Eq.(1.4) with  $k = 1$ , is locally asymptotically stable.

**Case  $k = 2$**

The characteristic equation of Eq.(2.4) with  $k = 2$  about the equilibrium point  $\bar{x}$  is

$$\lambda^3 - (0.5)\lambda^2 + 0.28357 = 0. \tag{2.6}$$

The roots of Eq.(2.6) are  $\lambda_{1,2} = 0.51289 + 0.52562i, \lambda_3 = 0.52578$  and the absolute value of each root is less than one. It follows from the Theorem 2.1 that the positive equilibrium point  $\bar{x}$  of Eq.(1.4) with  $k = 2$ , is locally asymptotically stable.

**Case  $k = 3$**

The characteristic equation of Eq.(2.4) with  $k = 3$  about the equilibrium point  $\bar{x}$  is

$$\lambda^4 - (0.5)\lambda^3 + 0.28357 = 0. \tag{2.7}$$

The roots of Eq.(2.7) are  $\lambda_{1,2} = 0.66294 \pm 0.48501i$  and  $\lambda_{3,4} = -0.41294 \pm 0.49976i$ .

It is easy to see that absolute value of each root is less than one. This implies that the positive equilibrium point  $\bar{x}$  of Eq.(1.4) with  $k = 3$ , is locally asymptotically stable.

Thus, for all  $k \in \mathbb{N}$  values, we obtain the characteristic equation of Eq.(2.4) about the

equilibrium point  $\bar{x}$  is

$$\lambda^n - (0.5)\lambda^{n-1} + 0.28357 = 0. \tag{2.8}$$

Set  $f(\lambda) = \lambda^n$  and  $g(\lambda) = -(0.5)\lambda^{n-1} + 0.28357$ .

Then, by Rouché's theorem,  $f(\lambda)$  and  $f(\lambda) + g(\lambda)$  have same number of zeroes in an open unit disc

$|\lambda| < 1$ . Hence, all the roots of (2.8) satisfies  $|\lambda| < 1$ , and it follows from Theorem 2.1 that the unique positive equilibrium point  $\bar{x}$  of Eq.(1.4) is locally asymptotically stable.

Hence, one can obtained the desired results and the proof is complete.

**4. Numerical Examples**

In this section, some numerical examples are presented to show the positive equilibrium point  $\bar{x}$  of Eq.(1.4) is locally asymptotically stable.

**Example 4.1** Consider the difference equation (1.4) with  $k = 1$  and the initial conditions

$$x_{-1} = 0.7, x_0 = 10.$$




Figure 1. The positive solutions of Eq.(1.4) with  $k = 1$ .

**Example 4.2** Consider the difference equation (1.4) with  $k = 2$  and the initial conditions

$$x_{-2} = 0.5, x_{-1} = 0.2, x_0 = 0.4.$$




Figure 2. The positive solutions of Eq. (1.4) with  $k = 2$ .

**Example 4.3** Consider the difference equation (1.4) with  $k = 3$  and the initial conditions

$$x_{-3} = 0.1, x_{-2} = 0.5, x_{-1} = 0.2, x_0 = 0.4.$$




Figure 3. The positive solutions of Eq.(1.4) with  $k = 3$ .

## 5. References

1. **Bozkurt, F., 2013.** Stability analysis of a nonlinear difference equation. *Int. J. Mod. Nonlinear Theory Appl.*, 2: 1-6.
2. **Camouzis, E., Ladas, G. 2007.** *Dynamics of third-order rational difference equations with open problems and conjectures*, CRC Press.
3. **Elaydi, S., 1996.** *An introduction to difference equation*, New York, Springer.
4. **El-Metwally, H., Grove, E. A., Ladas, G., Levins, R., Radin, M. 2011.** On the difference equation, *Nonlinear Anal.*, 47, 4623-4634.
5. **Gocen, M., Guneyesu, M., 2018.** The global attractivity of some rational difference equations. *J. Comput. An. App.*, 25(7): 1233-1243.
6. **Gumus, M., Abo-Zeid R. 2018.** On the solutions of a  $(2k+2)$ th order difference equation. *Dyn. Contin., Disc. Imp. Sys. Ser. B*, 25: 129-143.
7. **Gumus, M. 2018.** The global asymptotic stability of a system of difference equations. *J. Diff. Eq. App.*, 24(6):976-991. doi.org/10.1080/10236198.2018.1443445.
8. **Kocić, V., Ladas, G. 1993.** *Global behavior of nonlinear difference equations of higher order with applications*, Kluwer Academic Publishers, Dordrecht.
9. **Kulenović, M. R. S., Ladas, G. 2001.** *Dynamics of second order rational difference equations*, Chapman & Hall/CRC.
10. **Okumus, I., Soykan, Y., 2018.** Some Technique To Show The Boundedness Of Rational Difference Equations, *Journal of Progressive Research in Mathematics*, 13(2), 2246-2258.
11. **Ozturk, I., Bozkurt, F., Ozen, S. 2006.** On the difference equation, *Appl. Math. Comput.*, 181, 1387-1393.
12. **Tasdemir, E., Soykan, Y., 2017.** Long-term behavior of solutions of the non-linear difference equation, 1. *Gen. Math. Not.*, 38(1): 13-31.