

Rhomboid muscle variations: notes on their naming and classification principles

Albert Gradev , Lina Malinova , Julide Kasaboglu , Lazar Jelev 

Department of Anatomy, Histology and Embryology, Medical University of Sofia, Sofia, Bulgaria

Abstract

In this report we present two cases of rhomboid muscle variations observed during routine anatomical dissections. In the first case, on the left side of an adult male cadaver, a long and slender aberrant muscle was identified starting from the lateral part of the superior nuchal line and inserting to the scapula between the rhomboid minor and levator scapulae. The muscle was identified as the rare rhomboid capitis. In the second case, in an adult female cadaver, a bilateral variation in the origin of the rhomboid major fibers was described. On the left side, the rhomboid major fibers started from spinous processes of C1–C6, while on the right side it was narrower and originating from spinous processes of C1–C3. Reviewing the literature about the rhomboid muscles variations, we conclude that one and the same aberrant structure might be named differently. We also discuss the presentation of the known variations of the rhomboids in a common scheme instead of classification.

Keywords: human; rhomboid muscle; variation

Anatomy 2020;14(1):68–71 ©2020 Turkish Society of Anatomy and Clinical Anatomy (TSACA)

Introduction

The superficial muscles of the back connect the upper limb to the axial skeleton. In the upper back region, the rhomboids (minor and major) and levator scapulae muscles can be identified inserting on the medial scapular border under the trapezius.^[1,2] The rhomboids fix the medial border of the scapula to the thoracic wall and also retract this bone superiorly and medially and rotate it to depress the glenoid cavity.^[1,3] Variations of the rhomboids have been described in different names depending on the authors' understanding of the nature of variant muscles.^[4–12]

In this report, we present two cases of variations of the rhomboids and also provide a critical review of the present literature on this topic. Additionally, we aimed to discuss the controversies with the naming of these muscle variations and their proper grouping.

Case Report

In the first case (**Figure 1**), during routine anatomical dissection of a 70-year-old Caucasian male cadaver, after cut-

ting and retracting the trapezius muscle, a small aberrant muscle bundle was identified between the rhomboid minor and levator scapulae on the left side. The complete dissection revealed that the variant muscle bundle originated from the lateral part of the superior nuchal line next to the base of the mastoid process. As it passed downwards, it crossed the fibers of the splenius capitis near its cranial insertion. Finally, the aberrant slip inserted to the superior scapular angle, between the scapular attachments of rhomboid minor and levator scapulae. Based on its origin and insertion, the variant muscle slip was identified as rhomboid capitis muscle.^[13,14] It had a length of 21 cm and a width of 6–7 mm.

In the second case (**Figure 2**), routine anatomical dissection of a 67 year-old Caucasian female cadaver revealed quite asymmetrical rhomboids. On the left side, rhomboid major had an extended origin from T1 to T6 spinous processes, as the lowest part of the muscle was mostly aponeurotic. On the right side, the rhomboid major seemed much narrower and originated from T1 to T3 spinous processes.




Figure 1. Photograph of the rhomboid capitis muscle (arrows), observed on the left side and described in Case 1. LSc: levator scapulae; RMa: rhomboid major; RMi: rhomboid minor; SCa: splenius capitis.




Figure 2. Photograph of the asymmetrical rhomboid muscles layer described in Case 2. RMa: rhomboid major; RMi: rhomboid minor.

Discussion

The rhomboids are composed usually of six flat slips that originate from the spinous processes of either C7–T5 vertebrae^[1,3] or C6–T4 vertebrae.^[2,15] The fibers of the muscles run downwards and laterally to insert on to the medial border of scapula. The upper two slips of the rhomboids belong to the rhomboid minor muscle, while the lower four slips to the rhomboid major muscle. The rhomboids are mainly supplied by the dorsal scapular nerve of the brachial plexus made up of C4 and C5 spinal nerve fibers.^[3] Additional fibers contributing from C3 and C6 spinal nerves were also reported in some studies.^[16]

Variations of the rhomboids were previously reported as case reports,^[9,14,17] or in large series.^[4–8,10] One and the same aberrant muscle is named and grouped differently in majority of these reports. A common problem is the description of this variant muscle with different names, depending on its complete separation or fusion with the usual muscle (**Figures 3a and b**). Such examples might be the rhomboid minimus (or minus)^[10,12,18] and rhomboid

tertius muscles.^[11,19] When these aberrant muscles are fused with or being a part of the rhomboids they are described as extended attachments or increased number of the slips; however, when well separated from the usual muscles, they are called by their own names (**Figures 3a and b**). Rhomboideus minimus is a small, nearly horizontal variant muscle below the rhomboid major, which was described by von Haffner.^[18] A muscle with the same morphology was named as rhomboideus minus by Mori,^[10] who also mentioned that this muscle is common in Japanese. Interestingly, a muscle also called rhomboid minimus was described in quite a different location; just superior to the rhomboid minor.^[12] Another interesting variation reported is the rhomboid capitis (rhomboid occipitalis),^[13,14] also named as occipitoscapular muscle.^[4,5] Basically, the rhomboid capitis is a common neck muscle in many lower mammals.^[14] But it is quite a rare finding in humans and great apes.^[13] Some authors still named the same muscle based on its origin and insertion simply as occipitoscapular muscle.^[17,20]




Figure 3. Diagrams presenting the normal anatomy (a) and variations (b) of the rhomboids. (b) Variations of extended origin and aberrant attachment are presented on the left side, while on the right side the red arrows show similar muscular structures, separated from the main muscles and named differently.

Some interesting descriptions of non-vertebral origins of a muscle originating from the skull and closely related to rhomboids can be noted in previous reports reviewing the variations of the levator scapulae (Figure 3b).^[12,21] This might be another example of a variation, same as rhomboid capitis, which is fused completely with the usual levator scapulae. In summary, after all these notes on the rhomboid muscles variations, it seems quite difficult to present a classification. The classification principles are simply not clear. To present one and the same structure in different groups with different names simply because it might be well-separated or non-separated is not reasonable. Instead, we propose a scheme that demonstrates the range of rhomboid muscles variations (Figure 3b).

Conclusion

All of these variant muscles can manifest, despite rare, with some clinical symptoms. The rhomboids, rarely presenting aberrant attachments which might influence the proximal myofascial pain of the upper limb.^[22] On ultrasound, CT scan and MRI, variant and non-expected mus-

cles like rhomboid capitis, can mimic a tumor.^[23] So, it is important to have knowledge about the variations of these muscles for avoiding misinterpretations of diagnostic imaging and approaching the patients with myofascial pain of the upper limb.

Acknowledgement

The authors declare no conflict of interest. The authors received no financial support for this study. Human cadavers in the authors' laboratory are provided by a donation program for teaching purposes and scientific research. The authors wish to sincerely thank those who donated their bodies to science so that anatomical research could be performed.

Author Contributions

All authors equally contributed.

References

1. Moore KL. Clinically oriented anatomy. 3rd ed. Baltimore (MD) Williams & Wilkins; 1992. p. 351, 530–33.
2. Sinelnikov RD. Atlas of human anatomy. Vol. I. Musculoskeletal system. Moscow: Mir Publishers; 1989. p. 268–72.

3. Standring S (ed). Gray's anatomy: the anatomical basis of clinical practice. 41st ed. London: Elsevier; 2016. p. 818.
4. Wood J. Variations in human myology observed during winter session of 1866–67 at King's College, London. Proc Roy Soc London 1867;15:518–46.
5. Wood J. On a group of varieties of the muscles of the human neck, shoulder, and chest, and their transitional forms and homologies in the mammalia. Phil Trans Roy Soc London 1870;160:83–116.
6. Humphrey GM. Lectures on the varieties in the muscles of man. Lecture II: The muscles of the upper limb. Br Med J 1873;2:33–7.
7. Macalister A. Additional observations on muscular anomalies in human anatomy (third series) with a catalogue of the principal muscular variations hitherto published. Proc Roy Irish Acad 1875;25:1–134.
8. Knott JF. Abnormalities in human myology. Proc Roy Irish Acad 1883;3:407–27.
9. Selden BR. Congenital absence of trapezius and rhomboideus major muscles. J Bone Joint Surg 1935;17:1058–59.
10. Mori M. Statistics on the musculature of Japanese. Okajimas Fol Anat Jap 1964;40:195–300.
11. Lee J, Jung W. A pair of atypical rhomboid muscles. Korean J Phys Anthropol 2015;28:247–51.
12. Tubbs RS, Shoja MM, Loukas M (eds). Bergman's comprehensive encyclopedia of human anatomic variation. Hoboken (NJ): John Wiley & Sons, Inc; 2016. p. 269–75.
13. Kajiyama H. The superficial dorsal muscle group in Formosan monkey. II. Second layer of the superficial muscle group (mm. atlantoscapulares anterior et posterior and m. rhomboideus). Okajimas Folia Anat Jpn 1970;47:101–20.
14. Rogawski KM. The rhomboideus capitis in man – correctly named rare muscular variation. Okajimas Folia Anat Jpn 1990;67:161–3.
15. Schuenke M, Schulte E, Schumacher U. Thieme atlas of anatomy. General anatomy and musculoskeletal System. Stuttgart: Thieme; 2010. p. 260.
16. Malesy MJ, Thomeer RT, Marani E. The dorsoscapular nerve in traumatic brachial plexus lesions. Clin Neurol Neurosurg 1993;95 Suppl:S17–23.
17. Zagyapan R, Pelin C, Mas N. A rare muscular variation: the occipito-scapularis muscle: case report. Türkiye Klinikleri Journal of Medical Sciences 2008;28:87–90.
18. von Haffner H. Eine seltene doppelseitige Anomalie des Trapezius. Internationale Monatsschrift für Anatomie und Physiologie 1903;20:313–8.
19. Jeleu L, Landzhov B. A rare muscular variation: the third of the rhomboids. Anatomy 2013;7:63–4.
20. Stanchev S, Iliev A, Malinova L, Landzhov B. A rare case of bilateral occipitoscapular muscle. Acta Morphol Anthropol 2017; 24:1–2.
21. Chotai PN, Loukas M, Tubbs RS. Unusual origin of the levator scapulae muscle from mastoid process. Surg Radiol Anat 2015;37:1277–81.
22. Dor A, Vatine JJ, Kalichman L. Proximal myofascial pain in patients with distal complex regional pain syndrome of the upper limb. Journal of Bodywork & Movement Therapies 2019;23:547–54.
23. Kim SY, Park JS, Ryu KN, Jin W, Park SY. Various tumor-mimicking lesions in the musculoskeletal system: causes and diagnostic approach. Korean J Radiol 2011;12:220–31.

ORCID ID:

A. Gradev 0000-0003-0112-1333;
 L. Malinova 0000-0002-6928-1483;
 J. Kasaboglu 0000-0001-5475-6936;
 L. Jeleu 0000-0001-8596-7867

**Correspondence to:** Lazar Jeleu, MD, PhD

Department of Anatomy, Histology and Embryology, Medical University of Sofia, blvd. Sv. Georgi Sofiyskiy, BG-1431 Sofia, Bulgaria
 Phone: +359-897-87-27-51
 e-mail: ljeleu@abv.bg

Conflict of interest statement: No conflicts declared.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported (CC BY-NC-ND3.0) Licence (<http://creativecommons.org/licenses/by-nc-nd/3.0/>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited. *Please cite this article as:* Gradev A, Malinova L, Kasaboglu J, Jeleu L. Rhomboid muscle variation: notes on their naming and classification principles. *Anatomy* 2020;14(1):68–71.