

Protecting Doctors Aged Greater than 50 Years from COVID-19

Chu-Li Tu¹   · Chien-Ming Chao²  

¹Department of Surgery, Chi Mei Medical Center, Tainan, Taiwan

²Department of Intensive Care Medicine, Chi Mei Medical Center, Liouying, Taiwan

Dear Editor,

Health, in one recent study(1), Glynn raised the concern about the need of protecting worker aged 60-69 year from COVID-19. In addition to general population, we think this issue is important within healthcare system. However, we wonder whether this estimation can be applied in the healthcare system. To answer this question, we used the same model (2, 3), to estimate the distribution of death according to the doctors' age group. First, we found that 40% and 30% of deaths are in the 70+ and 60-69 year group. Second, after excluding the doctors aged more than 70 years, it estimated that 48% of deaths developed in the 60-69-year age group (Table-1). Both of these findings

were consistent with Glynn's findings and suggest that doctors aged over 60 years should be protected. Moreover, we also noted that 31.4% of deaths was estimated to occur in the 50-59-year groups in the group aged less than 70 years. This ratio was higher in the analysis of general population by Glynn, in which 25% of death developed in the group aged 50-59 years. Overall, it suggested that doctors aged greater than 50 years should need protection within the hospital in this COVID-19 pandemic. To avoid overload of healthcare system, which was associated the poor outcome of critical COVID-19 patients and prevent the death of aged physicians, appropriate allocation of

Corresponding: Chien-Ming Chao; Department of Intensive Care Medicine, Chi Mei Medical Center, Liouying, Taiwan

ORCID ID: 0000-0003-0235-9060

E-mail: ccm870958@yahoo.com.tw

Received: Apr 8, 2020 **Accepted:** June 4, 2020

Published: June 19, 2020

This is an Open Access article distributed under the terms of Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, or reproduction in any area, original work is properly cited.

The Ulutas Medical Journal © 2020



Table 1. Estimated deaths by age-group if doctors in UK are infected with SSARS-CoV-2

Age	No. of Doctors	Portion of Doctors (%)	Infection Fatality Rate (%)	No. of Death of Patients Infected	Portion of Death (%) If Over 70s Successfully Cocooned
20-29	37967	12.6	0.031	12	1.1
30-39	90256	30	0.084	76	7.1
40-49	80399	26.7	0.161	129	12.1
50-59	56233	18.7	0.595	335	31.4
60-69	26551	8.8	1.93	512	48
70+	9652	3.2	>4.28	>676	-

physicians' works according the specificity and age (<50 vs ≥50 years) should be conducted to enhance effectiveness of healthcare resource.

Conflict of Interest

The authors declare no competing interests.

References

1. Glynn JR. Protecting workers aged 60-69 years from COVID-19. *Lancet Infect Dis.* 2020 Apr 16. pii: S1473-3099(20)30311-X. doi: 10.1016/S1473-3099(20)30311-X
2. R Verity, LC Okell, I Dorigatti, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis* 2020 March 30
3. <https://www.statista.com/statistics/698278/registered-doctors-united-kingdom-uk-by-age/> accessed on April 28, 2020

Contact Details

Chu-Li Tu

Department of Surgery, Chi Mei Medical Center, Tainan, Taiwan
Email: chule_tu@yahoo.com
ORCID: 0000-0002-3930-2925

Chien-Ming Chao

Department of Intensive Care Medicine, Chi Mei Medical Center, Liouying, Taiwan
Email: ccm870958@yahoo.com
ORCID: 0000-0003-0235-9060

How to cite?

Tu CL, Chao CM. Protecting Doctors Aged Greater than 50 Years from COVID-19. *Ulutas Med J.* 2020;6(2): 101-102

DOI: 10.5455/umj.20200513022911