



## Additions to the Knowledge of the Molluscs of the Aegean Sea with Three Species: *Crepidula fornicata* (Linnaeus, 1758), *Anadara polii* (Mayer, 1868) and *Arcuatula senhousia* (Benson in Cantor, 1842)

Alper Doğan<sup>1,\*</sup>, Bilal Öztürk<sup>1</sup>, Banu Bitlis Bakır<sup>2</sup>, Mesut Önen<sup>1</sup>

<sup>1</sup> Ege University, Faculty of Fisheries, Department of Hydrobiology, Bornova, Izmir, Turkey.

<sup>2</sup> Dokuz Eylul University, Institute of Marine Sciences and Technology, Inciralti, Izmir, Turkey

\* Corresponding Author: Tel.: +90 232 311 52 02 ; Fax: +90 232 388 36 85 ;  
E-mail: alper.dogan@ege.edu.tr

Received 26 July 2013  
Accepted 10 January 2014

### Abstract

Within the frame of surveys conducted in Izmir Bay (Aegean Sea) in November 2008, April 2009 and April-May 2012, three mollusc species (*Crepidula fornicata*, *Anadara polii* and *Arcuatula senhousia*) have been recorded. Of the identified species, *A. senhousia* is new record for the molluscan fauna of the Aegean Sea and the calyptraeid gastropod *C. fornicata* is new report for the Turkish mollusc fauna. Besides, diagnostic features of *A. polii*, a rare arcid species, is given based on specimens from Izmir Bay.

**Keywords:** New records, Gastropoda, Bivalvia, Turkey, Aegean Sea, alien mollusc.

***Crepidula fornicata* (Linnaeus, 1758), *Anadara polii* (Mayer, 1868) ve *Arcuatula senhousia* (Benson in Cantor, 1842): Üç Tür ile Ege Denizi Mollusk Bilgilerine Katkılar**

### Özet

İzmir Körfezi'nde Kasım 2008, Nisan 2009 ve Nisan-Mayıs 2012 tarihlerinde yapılan incelemelerde, üç Mollusca türü (*Crepidula fornicata*, *Anadara polii* ve *Arcuatula senhousia*) saptanmıştır. Bu türlerden, *A. senhousia* Ege Denizi Mollusca faunası için yeni kayıt, bir calyptraeid gastropod olan *C. fornicata* ise Türkiye'nin Mollusca faunası için yeni kayıttır. Ayrıca, nadir dağılımlı Arcidae türü olan *A. polii*'nin, İzmir Körfezi'nden elde edilen bireylere dayanılarak diyagnostik özellikleri verilmiştir

**Anahtar Kelimeler:** Yeni kayıt, Gastropoda, Bivalvia, Türkiye, Ege Denizi, yabancı tür

### Introduction

Izmir Bay which is located in the Aegean Sea is divided into three parts (inner part, middle part and outer part) regarding its faunistic and hydrographic features (Kocataş, 1978; 1980). The inner part, where Alsancak Harbor is located, has been suffered from intense pollution of different origins since the 1960s and it is considered to be one of the most polluted environments of the Mediterranean Sea, until the establishment of a wastewater treatment station within the frame of 'The Grand Canal Project' in February 2000. Intense shipping activities in Alsancak Harbour are another factor contributing to the pollution of the inner part of Izmir Bay. However, introduction of some alien species into the area could be accepted as another effect of marine traffic of the bay. In the studies carried out before (Demir, 1977; Doğan *et al.*,

2005; Kazak and Cavas, 2007; Öztürk and Poutiers 2005), four alien mollusc species, i.e., *Bursatella leachii* de Blainville, 1817; *Anadara transversa* (Say, 1822), *Pinctada radiata* (Leach, 1814) and *Fulvia fragilis* (Forsskål in Niebuhr, 1775) were reported from Izmir Bay.

The present study deals with other three mollusc species sampled recently from the area: *Crepidula fornicata* (Linnaeus, 1758), *Anadara polii* (Mayer, 1868) and *Arcuatula senhousia* (Benson in Cantor, 1842).

The genus *Crepidula* (Calyptraeidae) is represented with three species (*Crepidula fornicata*, *Crepidula moulinsii* and *Crepidula unguiformis*) in the Mediterranean (Gofas, 2013). The slipper limpet *Crepidula fornicata* (Linnaeus, 1758) is native to the Atlantic coast of America, from Canada to Texas (Zenetos *et al.*, 2003) was accidentally introduced in Europe at the end of nineteenth century, currently

distributed along the Atlantic coasts of Europe from Irish Sea and Norway in the north to the Spain in the south, and with small populations in the Mediterranean Sea (Richard *et al.*, 2006; Mc Neill *et al.*, 2010).

*Anadara polii*, belonging to the family Arcidae, has a strong, rectangular and gibbous shell with a straight taxodont hinge line, and mostly with well-developed byssus. The family is represented by six genera and 14 species in the Mediterranean Sea (Gofas, 2013). *Anadara granosa*, *A. inflata*, *A. kagoshimensis*, *A. natalensis* and *A. transversa* are alien species of the genus in the Mediterranean Sea (Zenetos *et al.*, 2010). *Anadara polii* is known to occur in the Mediterranean, in the Sea of Marmara as well as on the Atlantic coasts of Spain, Morocco, Portugal and Maderia (Cachia *et al.*, 2004).

The Asian date mussel *Arcuatula senhousia* is a small thin shelled and filter feeding species of Mytilidae, living in the intertidal and subtidal soft substrates of the bays and estuaries where the species builds a nest or cocoon by holding sediment with its byssal threads, which may stabilise and protect the organism within the sediment (Morton, 1973, Crooks, 1996). *Arcuatula senhousia*, which is considered as an opportunistic species with a high reproductive capacity and a fast growth rate, may constitutes large populations with over than 10,000 ind/m<sup>2</sup>, and this situation can alter the nature of the sediment where it occurs (Morton, 1973; Mistri, 2003; Munari, 2008). According to Zenetos *et al.* (2012), *A. senhousia* is one of the most successful invader which was introduced into the Mediterranean through aquaculture.

The present study includes some descriptive and distributional information on three mollusc species mentioned above, which were sampled during the surveys conducted in Izmir Bay between November 2008 and May 2012.

## Materials and Methods

The benthic samples were taken from the infralittoral and circalittoral zones of various localities in Izmir Bay with a Van Veen Grab and beam trawl by the *R/V K. Piri Reis* and *Egesif* from November 2008 to May 2012 (Figure 1).

The coordinates and characteristics of the stations are given in Table 1. The sampling material was sieved through 0.5 mm mesh, then fixed in 4% formalin, and after the sorting process in the laboratory, the specimens were preserved in 70% ethanol. The investigated specimens were deposited in the museum collections of the Faculty of Fisheries, Ege University (EFSM) (Izmir, Turkey).

## Results

The examination of the benthic material taken from different localities in Izmir Bay revealed the presence of three mollusc species: *Crepidula fornicata* (Gastropoda), *Anadara polii* (Bivalvia) and *Arcuatula senhousia* (Bivalvia). Of these, *A. senhousia* is new record for the Aegean Sea fauna, and the calyptraeid gastropod *C. fornicata* is new report for the Turkish mollusc fauna.




Figure 1. Map of the studied area with the location of the sampling sites.

***Crepidula fornicata* (Linnaeus, 1758) (Figure 2)**

**Material examined:** Sta. 2, mud with detritus, 15 m, 1 sp.; sta. 3, mud, 8 m, 8 sp.; sta. 5, mud, 11 m, 38 sp.; sta. 7, mud, 10 m, 1 sp.

**Description:** Shell oval and taller compared to the other European *Crepidula* species. No sculpture on the shell surface, except for irregular growth lines. Shell colour exteriorly cream or yellowish with brownish spots; interiorly dirty white with spots of yellowish to brown in colour. The aperture has a white septum extending to the middle part of the shell. Minimum length: 13 mm. Maximum length: 42 mm.

***Anadara polii* (Mayer, 1868) (Figure 3)**

**Material examined:** Sta. 1, sand, 76 m, 2 sp.

**Description:** Shell solid, inequivalve, almost equilateral, strongly inflated, obliquely ovate with a slightly extended posterior-ventral part, beaks just in front of midline. Umbones inflated, sculpture of 27 wide radial ribs with interspaces evidently narrower than ribs. Periostracum light brown getting dark towards margin, bearing hair-like extensions. Hinge line taxodont with numerous teeth. Length=12.8 and 9.1 mm.

***Arcuatula senhousia* (Benson in Cantor, 1842) (Figure 4)**


**Material examined:** Sta. 4, mud, 4 sp.; sta. 5, mud, 63 sp.; sta. 6, mud, 2 sp.; sta. 8, mud, 5 sp.

**Description:** Shell thin and fragile, elongate, roughly trigonal ovate in outline, umbones subterminal, very small crenulations in the hinge line,

**Table.** Coordinates and ecological characteristics of the stations where the studied species were sampled

Stations	Coordinates		Date	Depth (m)	Biotope	Sampling gear
	Bts	Btf				
1	38°40'51"N 26°30'17"E		23.11.2008	76	Sand	Grab
2	38°23'40"N 26°57'17"E	38°23'44"N 26°57'32"E	03.05.2012	15	Detritic mud	Beam trawl
3	38°25'46"N 27°02'11"E	38°25'43"N 27°02'25"E	03.05.2012	8	Mud	Beam trawl
4	38°25'11"N 27°02'51"E		10.04.2012	10	Mud	Grab
5	38°24'47"N 27°03'57"E	38°24'49"N 27°04'11"E	03.05.2012	11	Mud	Beam trawl
6	38°24'33"N 27°04'30"E		23.04.2009	12	Mud	Grab
7	38°26'44"N 27°08'18"E		03.05.2012	10	Mud	Grab
8	38°27'16"N 27°08'55"E		10.04.2012	8	Mud	Grab

(Bts: Beam trawl start, Btf: Beam trawl finish).



**Figure 2.** *Crepidula fornicata*: dorsal (A) and ventral views (B, C).

outer surface of the valves smooth with weak sculpture of numerous concentric lines, several small radial grooves in front of the umbones making the anterior part slightly crenulated; periostracum shiny and smooth, yellowish green to olive brown in colour; stripes in different tones of brown colour more evident in larger specimens, and irregular transverse bands more common in smaller specimens. Minimum length: 5.2 mm. Maximum length: 19.4 mm.

### Discussion

As a result of evaluating the benthic material taken from Izmir Bay, three mollusc species were identified. Of these species, *Crepidula fornicata* is new record for the Turkish mollusc fauna and

*Arcuatula senhousia* is new report for the Aegean Sea mollusc fauna.

*Anadara polii* is a rare species (Poppe and Goto, 1993) and was first recorded in Turkey by Sturany (1895) from the Sea of Marmara. Demir (2003) first reported this species in a species list from the Aegean coasts of Turkey without any specific data. Therefore, we provided here some ecological and distributional features of this species from Izmir Bay.

The common slipper snail *C. fornicata* can rapidly spread out of the region where it introduces, and can create huge populations effecting native communities (Mc Neill et al., 2010). The amazing success of the species can be attributed to its ecological properties. *Crepidula fornicata* is an eurytopic, eurythermal and euryhaline species and can be found in different biotopes having various




Figure 3. *Anadara polii*: right (A) and left (B) valves, and dorsal view (C).




Figure 4. *Arcuatula senhousia*: left (A) and right (B, C) valves.

ecological characteristics (Hamon and Blanchard, 1994). Unlike the grazing gastropods, *C. fornicata* is a filter-feeding species, which allows specimens to form clusters by settling on top of each other and thereby enables it to form large populations in a given area (Blanchard, 1997). Thirty eight living specimens of *C. fornicata* were found in total in the present study as they were attached to the top of each other and the clusters were free, not attached to any substrates. The other characteristic of this species is that it is a protandrous hermaphrodite and the youngest specimens being male are able to inseminate several females lying below, resulting in a high survival rate (Blanchard, 1997; Dupont et al., 2006). Although this species currently formed a scarce population (10 ind./m<sup>2</sup>) in Izmir Bay, it has a great potential to become one of the dominant components of the benthic environments in the future due to its unique biology.

*Crepidula fornicata* was introduced to the Mediterranean mainly by bivalve farming and hull fouling (Blanchard, 1997). The occurrence of this species in Izmir Bay could be attributed to shipping because of the absence of bivalve farming in the area.

*Arcuatula senhousia* was reported for the first time from the Levantine coasts of Turkey by Uysal et al. (2008) but there was no specific information about the species. Eleftheriou et al. (2011) also reported many juvenile specimens of *A. senhousia* from Iskenderun Bay. In the present study, *A. senhousia* was encountered in the polluted inner part of Izmir Bay in April 2009 (20 ind./m<sup>2</sup>), then in April 2012 (40 and 50 ind./m<sup>2</sup>). The beam-trawl samples taken in May 2012 in the area had also 63 specimens of this species. Therefore, *A. senhousia* seems to have been established in the bay and enlarged its population density during a 3 year-period. It is known that alien species can expand its populations rapidly in the polluted areas and become dominant components of benthic communities (Çınar et al., 2012).

The way of introduction of *A. senhousia* into Izmir Bay might be the same with *C. fornicata*. Thereby, the shipping transport seems to be one of the main factors for the introduction of alien species in Izmir Bay.

As a result of the present study, the number of the alien species found in Izmir Bay increased to six. Further investigations should be implemented in order to evaluate the potential impacts of the alien species on the native communities of the bay.

## References

- Blanchard, M. 1997. Spread of the slipper limpet *Crepidula fornicata* (L. 1758) in Europe. Current state and consequences. *Scientia Marina*, 61: 109–118.
- Cachia, C. Misfud, C. and Sammut, P. M. 2004. The Marine Mollusca of the Maltese Island. Part four. The classes: Caudofoveata, Solenogastres, Bivslvia, Scaphopoda & Cephalopoda. Backhuys Publishers, Leiden, 270 pp.
- Cınar, M.E., Katagan, T., Öztürk, B., Bakır, K., Dağlı, E., Açık, Ş., Doğan, A. and Bitlis, B. 2012. Spatio-temporal distributions of zoobenthos in soft substratum of Izmir Bay (Aegean Sea, eastern Mediterranean), with special emphasis on alien species and ecological quality status. *Journal of the Marine Biological Association of the United Kingdom*, 92(7), 1457–1477. doi:10.1017/S0025315412000264
- Crooks, J.A. 1996. The population ecology of an exotic mussel, *Musculista senhousia*, in a southern California Bay. *Estuaries*, 19: 42-50. doi: 10.2307/1352650
- Demir M. 1977. On the presence of *Arca (Scapharca) amygdalum* Philippi, 1847 in the harbour of Izmir, Turkey. *Istanbul University, Journal of Faculty of Science*, 42: 197-202.
- Demir, M. 2003. Shells of Mollusca Collected from the Seas of Turkey. *Turkish Journal of Zoology*, 27: 101-140.
- Doğan, A., Çınar, M.E., Önen, M., Ergen, Z. and Katağan, T. 2005. Seasonal analysis of soft-bottom zoobenthic communities in polluted and unpolluted areas of Izmir Bay (Aegean Sea). *Senckenbergiana Maritima*, 35: 133–145. doi: 10.1007/BF03043182
- Dupont, L., Richard, J., Paulet, Y.M., Thouzeau, G. and Viard, F. 2006. Gregariousness and protandry promote reproductive insurance in the invasive gastropod *Crepidula fornicata*: evidence from assignment of larval paternity. *Molecular Ecology*, 15: 3009–3021. doi: 10.1111/j.1365-294X.2006.02988.x
- Eleftheriou, A. Anagnostopoulou-Visilia, E. Anastasopoulou, E. Ates, S.A. Bachari, N. El I. Cavas, L. Cevik, C. Çulha, M. Cevik, F., Delos A.L., Derici, O.B., Ergüden, D., Fragopoulou, N., Giangrande, A., Goksan, T., Gravili, C., Gurlek, M., Hattour, A., Kapiris, K., Kouraklis, P., Lamouti, S., Prato, E., Papa, L., Papantoniou, G., Parlapiano, I., Poursanidis, D., Turan, C. and Yaghoğlu, D. 2011. New Mediterranean Biodiversity Records (December 2011). *Mediterranean Marine Science*, 12(2): 491-508.
- Gofas, S. 2013. Arcidae. in: M.J.; Costello, P.; Bouchet, G.; Boxshall, C.; Arvanitidis, W., Appeltans Eds., *European Register of Marine Species* at <http://www.marbef.org/data/aphia.php?p=taxdetails&id=208> on 2013-12-13.
- Hamon, D. and Blanchard, M. 1994. État de la prolifération de la crépidule (*Crepidula fornicata*) en baie de Saint-Brieuc. *Rapport Ifremer, Del 94-14*, 29 p.
- Kazak, M. and Cavas, L. 2007. On the occurrence of *Bursatella leachii* de Blainville, 1817 in Izmir Bay, Turkey, *Mediterranean Marine Science*, 8(2): 87-90.
- Kocatas, A. 1978. Distribution et evolution des peuplements benthiques du Golfe d'Izmir (partie interieure) soumis a des multiples pollutions. *Journees Etude Pollutions, C.I.E.S.M., Antalya*, 4: 417–421.
- Kocatas, A. 1980. Evolution cyclique du benthos dans les zones de pollution du Golfe d'Izmir (Turquie). *Journees Etude Pollutions, C.I.E.S.M.*, 5: 643–648.
- Mc Neill, G. Nunn, J. Minchin, D. 2010. The slipper limpet *Crepidula fornicata* Linnaeus, 1758 becomes established in Ireland. *Aquatic Invasions*, 5 (1):21–25. doi: 10.3391/ai.2010.5.S1.006
- Mistri, M. 2003. The non-indigenous mussel *Musculista senhousia* in an Adriatic lagoon: effects on benthic community over a ten year period. *Journal of the*

- Marine Biological Association of the UK, 83:1277–1278. doi:10.1017/S0025315403008658
- Morton, B. 1973. Some aspects of the biology, population dynamics and functional morphology of *Musculista senhousia* Benson (Bivalvia: Mytilacea). Pacific Science, 28: 19-33.
- Munari, C. 2008. Effects of the exotic invader *Musculista senhousia* on benthic communities of two Mediterranean lagoons. Hydrobiologia, 611: 29-43. doi: 10.1007/s10750-008-9459-0
- Öztürk, B. and Poutiers, J.M. 2005. *Fulvia fragilis* (Bivalvia: Cardiidae): a lessepsian mollusc species from Izmir Bay (Aegean Sea). Journal of the Marine Biological Associations of the United Kingdom, 85: 351–356. doi: 10.1017/S0025315405011252h
- Poppe, G.T. and Goto, Y. 1993. European Seashells. Vol. 2: Scaphopoda, Bivalvia, Cephalopoda. Christa Hemmen, Wiesbaden, 222 p.
- Richard, J., Huet, M., Thouzeau, G. and Paulet, Y-M. 2006. Reproduction of the invasive slipper limpet, *Crepidula fornicata*, in the Bay of Brest, France. Marine Biology, 149(4): 789-801. doi:10.1007/s00227-005-0157-4
- Sturany, R. 1895. Bestimmungsliste der von Herrn Dr Konrad Natterer auf S.M.Schiff "Taurus" im Marmara-Meere Gedredschten mollusken. Denkschriften der mathem-naturw CI 62: 119-121 (in German).
- Uysal, Z. Latif, M.A. Özsoy, E. Tuğrul, S. Kubilay, N. Beşiktepe, Ş.T. Yemenicioğlu, S. Mutlu, E. Ediger, D., Beşiktepe, Ş., Ediger, V., Ak Örek, Y., Örek, H., Demirel, M., Tunç, Ş.Ç. and Terbıyık, T. 2008. Kilikya Baseni ve Kıyusal Ekosisteminde Dolaşım, Taşınım ve Ötrofikasyon Araştırmaları. The Scientific and Technical Research Council of Turkey (TÜBİTAK) 104Y277 p. 523 (in Turkish).
- Zenetos, A. Gofas, S. Ruso, G. and Templado, J. 2003. CIESM Atlas of Exotic Species in the Mediterranean, Volume 3: Molluscs. F. Briand (Ed.), CIESM Publishers, Monaco, 376 pp.
- Zenetos, A. Gofas, S. Verlaque, M. Çınar, M.E. and Garcia Raso, J.E. 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. Mediterranean Marine Science, 11(2): 318-493.
- Zenetos, A. Gofas, S. Morri, C. Rosso, A. Violanti, D. Garcia Raso, J.E. Çınar, M.E. Almogi-Labin, A. Ates, A.S. Azzurro, E. et al. 2012. Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. Mediterranean Marine Science, 13: 328-352.