

# COMMUNICATIONS

DE LA FACULTÉ DES SCIENCES  
DE L'UNIVERSITÉ D'ANKARA

Série A<sub>1</sub>: Mathématiques

---

TOME 29

ANNÉE 1980

---

**Chebyshev Approximation With Two Null Points**

by

**Ş. YÜKSEL**

4

Faculté des Sciences de l'Université d'Ankara  
Ankara, Turquie

# Communications de la Faculté des Sciences de l'Université d'Ankara

Comité de Rédaction de la Série A<sub>1</sub>

B. Yurtsever H. Hacısalihođlu M. Oruç

Secrétaire de publication

Ö. Çakar

---

La Revue "Communications de la Faculté des Sciences de l'Université d'Ankara" est un organe de publication englobant toutes les disciplines scientifiques représentées à la Faculté.

La Revue, jusqu'à 1975 à l'exception des tomes I, II, III, était composée de trois séries:

Série A: Mathématiques, Physique et Astronomie.

Série B: Chimie.

Série C: Sciences naturelles.

A partir de 1975 la Revue comprend sept séries:

Série A<sub>1</sub>: Mathématiques

Série A<sub>2</sub>: Physique

Série A<sub>3</sub>: Astronomie

Série B: Chimie

Série C<sub>1</sub>: Géologie

Série C<sub>2</sub>: Botanique

Série C<sub>3</sub>: Zoologie

En principe, la Revue est réservée aux mémoires originaux des membres de la Faculté. Elle accepte cependant, dans la mesure de la place disponible, les communications des auteurs étrangers. Les langues allemande, anglaise et française sont admises indifféremment. Les articles devront être accompagnés d'un bref sommaire en langue turque.

Adres: Fen Fakültesi Tebliğler Dergisi Fen Fakültesi, Ankara, Turquie.

## Chebyshev Approximation With Two Null Points

Ş. YÜKSEL\*

(Received on 18 March 1980, Accepted on 16 May, 1980)

### ABSTRACT

Chebyshev approximation problem with two null points, its best approximation and uniqueness are proved.

### INDRODUCTION

Let  $CZ([0, \alpha])$  be a space of functions which are continuous in  $[0, \alpha]$  and vanish at a null point. And let this space is made up by norm

$$\|g\| = \{ \sup |g(x)| : 0 \leq x \leq \alpha \}$$

If parameter space  $P$  is selected such as an open subspace of space  $R^n$ , then an element of  $P$ , has the form  $A = (a_1, \dots, a_n)$ .

Whenever an element  $f$  of  $CZ([0, \alpha])$  is handled then the existence of approximation function  $F$  which is an element of  $CZ([0, \alpha])$  is accepted such that  $F(A, \cdot) = F$  where  $A$  is an element of parameter space  $P$ .

The essential of Chebyshev approximation problem is to search an element  $A^*$  of parameter space  $P$ , such that  $e(A) = \|f - F(A, \cdot)\|$  be minimum. Such a parameter  $A^*$  is called the best parameter and  $F(A^*, \cdot)$  the best approximation to  $f$ .

Chebyshev approximation with a null point is studied by C.B. Dunham [1] in 1972. Dunham presented "DE LA VALLE POUSSIN" type result without proof, which is useful in characterizing "near best approximation". In addition, Dunham said that the existence of Chebyshev approximation with a null point

\* Department of Physics, Science Faculty, University of Ankara.

This study is a part of Ph. D. Thesis of Ş. Yüksel.

and its uniqueness without proof, because conception of local Haar property is not sufficient for the theorem which shows uniqueness of best approximation.

In this study, the cases in which functions be zero at 0 and  $\alpha$  are investigated by new concepts. The theorems which are not proved by Dunham are presented here with proofs and "Chebyshev approximation problem with two null points" is solved.

### CHEBYSHEV APPROXIMATION WITH TWO NULL POINTS

Let  $C([0, \alpha])$  be space of continuous functions with real value on  $[0, \alpha]$ , define norm.

$$\|g\| = \{\sup |g(x)| : 0 \leq x \leq \alpha\}$$

for the space.

It is easily seen that linear space

$$CZ_0([0, \alpha]) = \{f : f \in C(0, \alpha) : f(0) = 0\}$$

and

$$CZ([0, \alpha]) = \{g : g \in CZ_0(0, \alpha) : g(\alpha) = 0\}$$

are closed in  $C([0, \alpha])$ .

Let  $R^n$  a non-empty subset of  $P$  which is a parameter space. Suppose the existence of approximation function  $F$  which is element of  $CZ([0, \alpha])$  and obey the relation  $F(A, \cdot) = F[A] = F$  for element  $f$  of  $P$ .

If there exist any element  $f$  of space  $CZ([0, \alpha])$  then, finding an element  $A$  of  $P$ , which makes

$$e(A) = \|F(A, \cdot) - f\|$$

minimum will be the essential of Chebyshev approximation problem. More clearly, we can write it such as the following

$$\begin{aligned} \rho(f) &= \inf \{ \|F(A, \cdot) - f\| : A \in P \} \\ &= \|F[A^*] - f\| \end{aligned}$$

Such a parameter  $A$  is called the "best parameter" and function  $F(A^*, \cdot)$  "best approximation to  $f$ "

*Definition 1* Suppose a subset,  $\{1, \varnothing_1, \dots, \varnothing_n, \dots\}$  of  $C([0, \alpha])$ . Let be  $c_0, c_1, \dots, c_k \in \mathbb{R}$  and call

$$p_k = \sum_{i=0}^k c_i \varnothing_i$$

If for each element  $f$  of  $C([0, \alpha])$  and each number  $\epsilon > 0$  correspond to a natural number  $k$  such that

$$\|f - p_k\| \leq \epsilon$$

then set  $\{1, \varnothing_1, \varnothing_2, \dots\}$  is called "spanner" in  $C([0, \alpha])$ .

**Theorem 1** If the set  $\{1, \varnothing_1, \dots, \varnothing_n, \dots\}$  is a spanner in  $C([0, \alpha])$  and

$$0 = \varnothing_1(0) = \varnothing_2(0) = \dots \quad (1)$$

$$0 = \varnothing_1(\alpha) = \varnothing_2(\alpha) = \dots \quad (2)$$

are fulfilled, then the following properties are exist:

$$1) CZ_o([0, \alpha]) = CZ([0, \alpha])$$

$$2) \text{ Set } \{\varnothing_1, \varnothing_2, \dots, \varnothing_n, \dots\} \text{ is a spanner in } CZ([0, \alpha]).$$

**Proof 1** Let the set  $\{1, \varnothing_1, \varnothing_2, \dots, \varnothing_n, \dots\}$  be a spanner in  $C([0, \alpha])$  and conditions be fulfilled.

Suppose  $f \in CZ_o([0, \alpha])$ . So  $f \in C([0, \alpha])$ , then for each  $\epsilon > 0$  there are real numbers  $c_0, c_1, \dots, c_k$  such that

$$\|f - p_k\| \leq \epsilon \quad (3)$$

Equation (3) gives us property

$$|f(x) - (c_0 + c_1\varnothing_1(x) + c_2\varnothing_2(x) + \dots + c_k\varnothing_k(x))| \leq \epsilon$$

for each  $x \in [0, \alpha]$ . The fact that  $f(0) = 0$  gives  $|c_0| \leq \epsilon$  and the inequality gives rise to  $c_0 = 0$ . As a result, we get relation

$$|f(x) - (c_1\varnothing_1(x) + c_2\varnothing_2(x) + \dots + c_k\varnothing_k(x))| \leq \epsilon$$

for each  $x \in [0, \alpha]$ . If we put  $\alpha$  instead of  $x$ , for each  $x > 0$  in the above relation we get  $f(\alpha) = 0$  from the fact  $|f(\alpha)| \leq \epsilon$ . Therefore

$$CZ_o([0, \alpha]) = CZ([0, \alpha])$$

**Proof 2** On the other hand, if we use the proof technique above mentioned, can be easily seen that the set  $\{\varnothing_1, \varnothing_2, \dots, \varnothing_n, \dots\}$  be a spanner in  $CZ([0, \alpha])$ .

**Definition 2** If there exist point set  $\{x_1, x_2, \dots, x_m\}$  such that  $0 < x_1 < x_2 < \dots < x_m < \alpha$  and

$\|g\| = |g(x_i)|$ ,  $g(x_i) = (-1)^i g(x_1)$ ;  $i = 1, 2, \dots, m$   
 for an element  $g$  of  $CZ([0, \alpha])$  then  $g$  alternates  $m$  times in  $[0, \alpha]$ .  
 If function  $g$  alternates  $m$  times but not  $m+1$  times then we will  
 say that  $g$  alternates definitely  $m$  times.

*Definition 3* Let  $A = (a_1, a_2, \dots, a_n)$  and  $F(A, \cdot)$  be an approximation function of  $CZ([0, \alpha])$ . Let  $i = 1, 2, \dots, n$  and vector space  $D(A, \dots)$  with dimension  $m$  which is made up of  $\partial F(A, \cdot) / \partial a_i$ . Evidently  $m \leq n$ .

If the condition  $F(A, \cdot) = F(B, \cdot)$  requires that  $F(A, \cdot) - F(B, \cdot)$  has  $m$  null points at  $[0, \alpha]$  then approximation function  $F$  has property (CZ).

Now let us try to have result DE LA VALLE-POUSSIN type.

*Theorem 2* If an approximation function  $F(A, \cdot)$  of space  $CZ([0, \alpha])$  has property (CZ) and function  $f$  is an element of  $CZ([0, \alpha])$  such that function  $F(A, \cdot) - f$  alternates definitely  $m+1$  times then each element  $B$  of parameter space  $P$  has the following property

$$\begin{aligned} \text{Max } \{ |F(B, x_i) - f(x_i)| : i = 0, \dots, m \} &\geq \\ &\geq \text{Min } \{ |F(A, x_i) - f(x_i)| : i = 0, \dots, m \} \end{aligned} \quad (4)$$

*Proof*

1) Relation

$$(F(A, x_i) - f(x_i)) (F(A, x_i) - F(B, x_i)) \leq 0 \quad (5)$$

is exist, otherwise

$$(F(A, x_i) - f(x_i)) (F(A, x_i) - F(B, x_i)) > 0 \quad (6)$$

would be true. Function  $F(A, \cdot) - f$  alternates on space  $\{x_0, x_1, \dots, x_m\}$  According to Equation (6) it is clear that function  $F(A, x_i) - F(B, x_i)$  will have the same alternant. Let us assume  $F(A, x_0) - f(x_0) > 0$  without violating the generality, so that the relations

$$F(A, x_0) - F(B, x_0) > 0$$

$$F(A, x_1) - F(B, x_1) < 0$$

.....

prove that  $F(A,.) - F(B,.)$  has  $m$  zeros. This requires  $F(A,.) = F(B,.)$  in accordance with the property (CZ). This result contrasts the assumption (6).

$$2) \text{ Max } \{ |F(B, x_i) - f(x_i)| ; i=0,1,\dots,m \} = |F(B, x_p) - f(x_p)| \geq$$

$$0 \leq p \leq m$$

$$\text{Min } \{ |F(A, x_i) - f(x_i)| ; i=0,1,\dots,m \} = |F(A, x_q) - f(x_q)|$$

$$0 \leq q \leq m$$

If the relation (4) was correct then we would have found

$$|F(B, x_p) - f(x_p)| < |F(A, x_q) - f(x_q)| \quad (7)$$

It is clear that the inequality (7) is true for each  $x_i: i = 0,1, \dots, m$  and then we have

$$|F(B, x_i) - f(x_i)| < |F(A, x_i) - f(x_i)| \quad (8)$$

If we add  $f(x_i)$  to the second factor of relation (5) we find,

$$(F(A, x_i) - f(x_i)) ( (F(A, x_i) - f(x_i)) - (F(B, x_i) - f(x_i)) )$$

$$\leq 0 \quad (5')$$

Function  $F(A,.) - f$  alternates on set  $\{x_0, x_1, \dots, x_m\}$

so we can take  $F(A, x_0) - f(x_0) > 0$  without violating the generality.

Under these circumstances from the inequality (5') we can find

$$(F(A, x_0) - f(x_0)) \leq (F(B, x_0) - f(x_0))$$

We know that  $F(A, x_0) - f(x_0) > 0$ , so  $F(B, x_0) - f(x_0)$  is also positive. Hence we have

$$|F(A, x_0) - f(x_0)| \leq |F(B, x_0) - f(x_0)|$$

However, this relation contrasts to the equation (8).

*Definition 4* Let an element  $f$  of space  $CZ([0, \alpha])$  and  $F(A,.)$  be best to  $f$ . For the best approximation to  $f$  necessary and sufficient condition is that  $F(A,.)$  alternates definitely  $m+1$  times then it is said that  $F(A,.)$  has property (CS) with degree  $m+1$

*Definition 5* Let  $A = (a_1, a_2, \dots, a_n)$  and  $F(A,.)$  be an approximation function at  $CZ([0, \alpha])$  and if the following conditions exist then approximation  $F(A,.)$  has local Haar property with null points  $m$  degree at  $A$ .

(i)  $\partial F(A,.) / \partial a_i$  are exist and continious for each  $i$ ;  
 $i = 1, 2, \dots, n$

(ii) Let

$$B = (b_1, b_2, \dots, b_n) \text{ and } D(A, B, x) = \sum_{i=1}^n b_i \frac{\partial F(A, x)}{\partial a_i}$$

When  $\| B \|$  is sufficientl small and relation

$$F(A-B, x) - F(A, x) = D(A, B, x) + R(A, B, x)$$

is correct, then

$$R(A, B, x) = O \| B \|^2$$

(iii) Element  $A$  has a neighbourhood which is contained by  $P$ .

(iv) Linear space  $D(A,.)$  is an Haar space with dimension  $m$  at  $[0, \alpha]$ .

*Theorem 3* If approximation function  $F$  of space  $CZ([0, \alpha])$  has local Haar property at  $A$  and  $F(A,.)$  be best to element  $f$  of  $CZ([0, \alpha])$  then in that case  $F$  has property (CS) with degree  $m+1$ .

*Proof* Suppose  $F$  has not property (CS) with degree  $m+1$ . Now, function  $F(A,.) - f$  alternates at  $[0, \alpha]$  less than  $m+1$ . There is relation

$$F(A, 0) - f(0) = 0$$

So there is a number  $\gamma$  such that

$$| F(A, x) - f(x) | < e(A) / 2 : 0 \leq x \leq \gamma \leq \alpha$$

where  $e(A) = \| F(A,.) - f \|$

At the point  $\alpha$ , we can write

$$F(A, \alpha) - f(\alpha) = 0$$

So there exist  $\delta > 0$  such that

$$| F(A, x) - f(x) | < e(A) / 2 ; \delta \leq x \leq \alpha$$

If approximation function  $F(A,.)$  is best to  $f$  at  $[0, \alpha]$  then  $F(A,.)$  will be best to  $f$  at the interval  $[\gamma, \delta]$  ( [2], Theorem 3)

Under these circumstances of the function  $F(A,.) - f$  alternates  $m+1$  times at  $[0, \alpha]$  then  $F(A,.) - f$  also alternates  $m+1$  times at the interval  $[\gamma, \delta]$  ( [3], Theorem 5).



If the function  $F(A, \cdot) - f$  alternates less than  $m+1$  times we find an element  $B$  of parameter space  $P$  with the help of Theorem 2, under the condition  $\gamma \leq x \leq \delta$ , such that

$$\begin{aligned} \| F(B, \cdot) - F(A, \cdot) \| &< e(A)/2 \\ | F(B, x) - f(x) | &< e(A) \end{aligned}$$

Then we have

$$\begin{aligned} | F(B, x) - f(x) | &\leq | F(B, x) - F(A, x) | + | F(A, x) - f(x) | \\ &< e(A)/2 + e(A)/2 \\ &< e(A) \end{aligned}$$

From this and the fact that  $e(B) = \| F(B, \cdot) - f \|$  we have  $e(B) < e(A)$ . However this contrasts to the best approximation. As a result, function  $F(A, \cdot)$ ,  $f$  alternates definitely  $m+1$  times.

Combining Theorem 2 and 3 we get the following result:

*Theorem 4* If approximation function  $F(A, \cdot)$  has the property (CZ) and local Haar property with null points degree  $m$ , then, necessary and sufficient condition for approximation function  $F(A, \cdot)$  being best to  $f$  is that  $F(A, \cdot)$  has property (CS) with degree  $m+1$ .

*Theorem 5* Let approximation function  $F(A, \cdot)$  has the condition of Theorem 4. If  $F(A, \cdot)$  is best it will be unique.

*Proof* Let us choose  $\gamma$  and  $\delta$ ,  $\gamma \leq x \leq \delta$ , such that

$$| F(A, x) - f(x) | < e(A)/2$$

If approximation function  $F(A, \cdot)$  be best to  $f$  at the interval  $[0, \alpha]$ ,  $F(A, \cdot)$  is also best to  $f$  at  $[\gamma, \delta]$  ([2]; Theorem 3)

In that case, if function  $F(A, \cdot) - f$  alternates  $m+1$  times at interval  $[0, \alpha]$  and also alternates  $m+1$  times at  $[\gamma, \delta]$  ([3]; Theorem 5)

So it is sufficient to prove the theorem at interval  $[\gamma, \delta]$ .

Let us suppose  $F(A, \cdot)$  is not the best and unique approximation to  $f$  at the interval  $[\gamma, \delta]$ , take two approximation functions as  $F(A, \cdot)$  and  $F(B, \cdot)$ .

Function  $F(A, \cdot) - f$  alternates  $m+1$  times at the interval  $[\gamma, \delta]$  so there is the property:

$$F(A, x_i) - f(x_i) = - (F(A, x_{i-1}) - f(x_{i-1})) \quad ; \quad i = 0, 1, \dots, m$$

In accordance with the Theorem 2 we get

$$F(A, x_0) - F(B, x_0) \leq 0$$

$$F(A, x_1) - F(B, x_1) \geq 0$$

.....

If there exist definite inequality, then function  $F(A, \cdot) - F(B, \cdot)$  will have definite  $m+1$  null points at  $[\gamma, \delta]$  and from Haar condition one can get equation

$$F(A, \cdot) = F(B, \cdot)$$

$D(B, \cdot, \cdot)$  fulfills Haar condition so  $F(B, \cdot)$  has local Haar property in the case of equality.

Let  $0 \leq t \leq 1$  and  $C$  be an element of parameter space  $P$ , if we apply property (ii) of local Haar condition to function

$$F(B, x) - F(B-tC, x)$$

we will get

$$F(A, x) - F(B-tC, x) = F(A, x) - F(B, x) + tD(B, C, x) + R(B, C, x)$$

If we choose  $t$  sufficiently small we will have system

$$F(A, x_0) - F(B-tC, x_0) < 0$$

$$F(A, x_1) - F(B-tC, x_1) > 0$$

.....

Function  $F(A, \cdot) - F(B-tC, \cdot)$  has  $m$  null points at the interval, and while  $t$  approaching zero,  $F(A, \cdot) = F(B, \cdot)$  equality will be exist.

Here, function  $F(A, \cdot)$  is the best unique approximation to  $f$  at the interval  $[\gamma, \delta]$ . If there had been another best approximation at  $[0, \alpha]$  it would have also been best approximation at  $[\gamma, \delta]$ , that is impossible.

#### REFERENCES

- [1] Dunham, C.B., *Chebyshev Approximation with a Null Point*, *Z. Angew. Math. Mech.* 52, 239 (1972).
- [2] Meinardus, G., and Schwedt, D., *Nicht-Lineare Approximationen*, *Arch. Rationel Mech. Anal.*, 17, 297 (1964).
- [3] Dunham, C. B., *Alternating Chebyshev Approximation*, *Transactions of the American Math. Soc.* Vol. 178, (1973).

#### ÖZET

İki noktada sıfır olan Chebyshev yaklaşım problemi, en iyi yaklaşımı ve bunun tekliği ispatlanmaktadır.

**Prix de l'abonnement annuel**

Turquie : 15 TL; Étranger: 30 TL.

Prix de ce numéro : 5 TL (pour la vente en Turquie).

Prière de s'adresser pour l'abonnement à : Fen Fakültesi  
Dekanlığı Ankara, Turquie.