

## AN APPLICATION OF THE FIBRATION THEOREM OF EHRESMANN

DORIN ANDRICA., LIVIU MARE

(CLUJ - NAPOCA)

(Received April 17, 1992; Accepted July 14, 1992)

### ABSTRACT

The main purpose of the paper is to prove that the map (7) and also its restriction to  $GL^+(n, \mathbb{R})$  is a locally trivial fibration.

From the general theory of fiber bundles we know that a bundle map between two  $C^\infty$ -differentiable manifolds is a surjective submersion. Here arise a natural problem: given  $M$  and  $N$  two  $C^\infty$ -differentiable manifolds and  $f: M \rightarrow N$  a smooth surjective submersion, find sufficient conditions in order that  $f$  be a locally trivial fibration. A such condition is given by:

**Theorem.** (Ehresmann [3, Th. 8.12, p. 84]). If  $f: M \rightarrow N$  is a proper surjective submersion then  $f$  is a locally trivial fibration.

We shall consider

$$GL(n, \mathbb{R}) = \{X \in M_n(\mathbb{R}) : \det X \neq 0\} \quad (1)$$

the real general linear group, which is a  $n^2$  dimensional  $C^\infty$ -differentiable manifold, as an open subset of  $M_n(\mathbb{R})$ . It is known that  $GL(n, \mathbb{R})$  has two connected components:

$$GL^+(n, \mathbb{R}) = \{X \in GL(n, \mathbb{R}) : \det X > 0\} \text{ and}$$

$$GL^-(n, \mathbb{R}) = \{X \in GL(n, \mathbb{R}) : \det X < 0\},$$

both open in  $GL(n, \mathbb{R})$ .

We also consider

$$S_n(\mathbb{R}) = \{X \in M_n(\mathbb{R}) : {}^tX = X\}, \quad (2)$$

the set of symmetric matrices. Clearly we can identify  $S_n(\mathbb{R})$  with the Euclidean space  $\mathbb{R}^{n(n+1)/2}$ . In the following we shall denote by  $S_n^+(\mathbb{R})$  the subset of  $S_n(\mathbb{R})$  formed of all positive definite matrices.

Finally denote by

$$O_n(\mathbb{R}) = \{X \in GL(n, \mathbb{R}) : {}^tX \cdot X = I_n\}, \quad (3)$$

the set of orthogonal matrices.

In the sequel we shall use the following two results:

(a) Diagonal form of symmetric matrices). For any  $A \in S_n(\mathbb{R})$  there exists  $T \in O_n(\mathbb{R})$  such that

$${}^tT A T = \begin{bmatrix} \lambda_1 & & & \\ & \cdot & & \\ & & \cdot & \\ & & & \lambda_n \\ 0 & & & & \end{bmatrix}$$

where  $\lambda_1, \dots, \lambda_n$  are the eigenvalues of the matrix  $A$ . (see, for example, [2, Th. 2, p. 83]).

(b) (Polar decomposition in  $GL(n, \mathbb{R})$ ). Any  $X \in GL(n, \mathbb{R})$  admits a unique decomposition in the form:

$$X = OS \quad (4)$$

with  $O \in O_n(\mathbb{R})$  and  $S \in S^+_n(\mathbb{R})$ . Moreover the application  $O_n(\mathbb{R}) \times S^+_n(\mathbb{R}) \rightarrow GL(n, \mathbb{R})$  given by

$$(O, S) \longrightarrow O.S \quad (5)$$

is a diffeomorphism.

In this paper, by using the above mentioned result of Ehresmann, we shall obtain a locally trivial fibration of  $GL(n, \mathbb{R})$  (and respectively of  $GL^+(n, \mathbb{R})$ ) and we will put in evidence an interesting connection with the trivial fibration given by  $\det: GL^+(n, \mathbb{R}) \longrightarrow \mathbb{R}^*_+$  (we denote by  $\mathbb{R}^*_+$  the set of real positive numbers)

Let begin with the proof of two helping results:

**Lemma 1.** The set  $S^+_n(\mathbb{R})$  is open in  $S_n(\mathbb{R})$ .

**Proof:** Observe that (a) supply us with the following relation:

$$S_n(\mathbb{R}) = \bigsqcup_{T \in O_n(\mathbb{R})} T \{D(\lambda_1, \dots, \lambda_n) : \lambda_i \in \mathbb{R}\} T^t$$

where

$$D(\lambda_1, \dots, \lambda_n) = \begin{bmatrix} \lambda_1 & & & 0 \\ & \cdot & & \\ & & \cdot & \\ & & & \lambda_n \\ 0 & & & & \end{bmatrix}$$

In this relation we have

$$S_n^+(\mathbb{R}) = \bigsqcup_{T \in O_n(\mathbb{R})} T \{D(\lambda_1, \dots, \lambda_n) : \lambda_i > 0 \text{ for all } i = 1, \dots, n\} t_T$$

But every  $T \{D(\lambda_1, \dots, \lambda_n) : \lambda_i > 0 \text{ for all } i = 1, \dots, n\} t_T$  is clearly open in  $T \{D(\lambda_1, \dots, \lambda_n) : \lambda_i \in \mathbb{R}\} t_T$ ; consequently  $S_n^+(\mathbb{R})$  is open in  $S_n(\mathbb{R})$ . Lemma 1 is proved.

Consider the following sets: for any  $A \in GL(n, \mathbb{R}) \subset S_n(\mathbb{R})$  put

$$O_n(A, \mathbb{R}) = \{X \in GL(n, \mathbb{R}) : {}^tXX = A\}$$

and if  $\det A > 0$

$$O_n^+(A, \mathbb{R}) = \{X \in O_n(A, \mathbb{R}) : \det X = \sqrt{\det A}\}$$

$$O_n^-(A, \mathbb{R}) = \{X \in O_n(A, \mathbb{R}) : \det X = -\sqrt{\det A}\}$$

Clearly  $O_n(I_n, \mathbb{R}) = O_n(\mathbb{R})$  and  $O_n^+(I_n, \mathbb{R}) = SO_n(\mathbb{R})$  where  $SO_n(\mathbb{R})$  represents the special orthogonal group.

**Lemma 2.** (i) The map  $\varphi: S_n^+(\mathbb{R}) \rightarrow S_n^+(\mathbb{R})$

$$X \varphi(X) = X^2 \tag{6}$$

is a proper bijection.

(ii) We have the following chain of equivalences:

$$O_n(A, \mathbb{R}) \neq \emptyset \Leftrightarrow O_n^+(A, \mathbb{R}) \neq \emptyset \Leftrightarrow A \in S_n^+(\mathbb{R}).$$

**Proof:** (i) The fact that  $\varphi$  is one-to-one is an immediate consequence of (a). Let's show that  $\varphi$  is proper: for  $K \subset S_n^+(\mathbb{R})$  compact we have to prove that  $\varphi^{-1}(K)$  is bounded.

A very useful norm on  $M_n(\mathbb{R})$ , equivalent with the Euclidean norm is

$$\|A\| = [\max\{|\lambda_i| : \lambda_i \text{ eigenvalue of } {}^tAA\}]^{1/2}.$$

But if  $A \in S_n^+(\mathbb{R})$  then

$$\|A\| = [\max\{\lambda_i^2 : \lambda_i \text{ eigenvalue of } A\}]^{1/2},$$

so that  $\|\varphi^{-1}(A)\| = \sqrt{\|A\|}$ . Because  $K$  is bounded,  $\varphi^{-1}(K)$  is bounded, too.

(ii) The first equivalence holds because  $\det(J_n A) = -\det A$ ,

where  $J_n = \begin{bmatrix} -1 & & 0 \\ & 1 & \\ & & \cdot \\ 0 & & 1 \end{bmatrix} \in O_n(\mathbb{R}).$

Assume  $O_n(A, \mathbb{R}) \neq \emptyset$ . Therefore  $A = {}^tXX$ , thus  $A$  is symmetric. If  $\lambda$  is an eigenvalue of  $A$  and  $x \in \mathbb{R}^n$  is an eigenvector corresponding to  $\lambda$ , then

$$\lambda = \frac{\|Xx\|}{\|x\|^2} > 0.$$

It follows that  $A \in S_n^+(\mathbb{R})$ .

Conversely, if  $A \in S_n^+(\mathbb{R})$ , by the surjectivity of  $\varphi$  one obtains that  $O_n(A, \mathbb{R}) \neq \emptyset$ .

Now, we are in position to state the main result of this paper.

**Theorem (i)** The map  $f: GL(n, \mathbb{R}) \rightarrow S_n^+$  given by

$$X f(\chi) = {}^t\chi\chi \quad (7)$$

is a fibration of  $GL(n, \mathbb{R})$  with the type fiber  $O_n(\mathbb{R})$ .

(ii) The restriction  $f \Big|_{GL^+(n, \mathbb{R})} : GL^+(n, \mathbb{R}) \rightarrow S_n^+(\mathbb{R})$

is also a fibration, with the type fiber  $SO_n(\mathbb{R})$ .

**Proof:** First, we will show that  $f$  is a submersion. Using the well-known result concerning the equality of the Frechet and Gateaux differentials for smooth maps, we obtain:

$$\begin{aligned} (df)_B(C) &= \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} [f(B + \lambda C) - f(B)] = \\ &= \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} [{}^t(B + \lambda C)(B + \lambda C) - {}^tBB] = {}^tBC + {}^tCB. \end{aligned}$$

It follows that, for every  $B \in GL(n, \mathbb{R})$  the differential  $(df)_B: M_n(\mathbb{R}) \rightarrow S_n(\mathbb{R})$  is surjective. Let  $D \in S_n(\mathbb{R})$  and choose  $C = {}^t(B^{-1})D/2$ . Then

$$(df)_B(C) = {}^tB {}^t(B^{-1})D/2 + {}^tDB^{-1}B/2 = D/2 + D/2 = D.$$

We prove now that  $f$  is a proper map. To this end, let's observe that, by using the polar decomposition (b) it follows that  $f(X) = S^2$ . So that  $f = \varphi \circ h$ , where  $\varphi$  is given by (6) and  $h: GL(n, \mathbb{R}) \rightarrow S_n^+(\mathbb{R})$  is given by

$$h(X) = S. \tag{8}$$

By Lemma 2, (i), it remains to show that  $h$  is proper. If  $K \subset S^+_n(\mathbb{R})$  is compact then by (5) the set  $h^{-1}(K)$  is diffeomorphic to  $K \times O_n(\mathbb{R})$ , so that is compact (don't forget that  $O_n(\mathbb{R})$  is compact).

Now, we can apply Ehresmann's theorem and deduce that  $f: GL(n, \mathbb{R}) \rightarrow S^+_n(\mathbb{R})$  is a locally trivial fibration. The fiber along the identic matrix  $I_n$  is, as we have already seen,  $O_n(\mathbb{R})$ .

The assertion (ii) follows observing that the restriction of  $f$  to the open set  $GL^+(n, \mathbb{R})$  is a proper submersion and applying then Ehresmann's theorem.

**Remarks 1.** Notice that both fibrations obtained in Theorem are in fact trivial, since the polar decomposition gives allways a diffeomorphism. In addition, we can give an explicit formula for the fiber along a matrix  $A \in S^+_n(\mathbb{R})$  for both fibrations:

if  $X_0 \in O^+_n(A, \mathbb{R}) \subset O_n(A, \mathbb{R})$  then the fiber is  $O_n(\mathbb{R}) X_0 = \{XX_0 : X \in O_n(\mathbb{R})\}$  for the first fibration, respectively  $SO_n(\mathbb{R}) X_0$  for the second one.

2. It is worth to mention the following interesting connection between the fibration (ii) and the bundle map

$$g: GL^+(n, \mathbb{R}) \rightarrow \mathbb{R}^*_+, g(X) = (\det X)^2.$$

The correspondence  $X \rightarrow ((\det X)^2, \frac{1}{n \sqrt{\det X}} X)$

gives a diffeomorphism between  $GL^+(n, \mathbb{R})$  and  $\mathbb{R}^*_+ \times SL(n, \mathbb{R})$  which shows that  $g$  is a trivial fibration of the fiber  $SL(n, \mathbb{R})$ . We denote the fiber along  $\alpha \in \mathbb{R}^*_+$  by  $SL^{(\alpha)}(n, \mathbb{R})$ ; so that

$$SL^{(\alpha)}(n, \mathbb{R}) = \{X \in GL^+(n, \mathbb{R}) : \det X = \sqrt{\alpha}\}$$

Denoting  $S_n^{(\alpha)}(\mathbb{R}) = \{A \in S^+_n(\mathbb{R}) : \det A = \alpha\}$  the

restriction  $f \Big|_{SL^{(\alpha)}(n, \mathbb{R})} : SL^{(\alpha)}(n, \mathbb{R}) \rightarrow S_n^{(\alpha)}(\mathbb{R})$

will be a proper submersion. We obtain a fibration of the fiber  $SL^{(\alpha)}(n, \mathbb{R})$  with the fiber along  $A \in S_n^{(\alpha)}(\mathbb{R})$

$$\{X \in SL^{(\alpha)}(n, \mathbb{R}) : {}^tXX = A\} = \{X \in GL^+(n, \mathbb{R}) : {}^tXX = A \text{ and } \det X = \sqrt{\det A}\} = SO_n(A, \mathbb{R}),$$

the same fiber as in (ii).

3. There are serious reasons to believe that some of the results presented here remain valid in the case when  $GL(n, \mathbb{R})$  is replaced by the automorphism group of an infinite dimensional Hilbert space.