

ON p - VALENT FUNCTIONS WITH NEGATIVE AND MISSING COEFFICIENTS II

M.K. AOUF and H.E. DARWISH

Department of Mathematics, Faculty of Science, University of Mansoura, Mansoura, Egypt.

(Received June 14, 1994; Accepted Jan. 12, 1995)

ABSTRACT

Let $P_k(p, A, B, \alpha)$ be the class of functions $f(z) = z^p + \sum_{n=k}^{\infty} |a_{p+n}| z^{p+n}$, $k \geq 2$, analytic and p -valent in the unit disc $U = \{z : |z| < 1\}$ and satisfying the condition

$$\left| \frac{\frac{zf'(z)}{f(z)} - p}{[pB + (A - B)(p - \alpha)] - B - \frac{zf'(z)}{f(z)}} \right| < 1, \quad z \in U,$$

where $-1 \leq B < A \leq 1$, $-1 \leq B < 0$, and $0 \leq \alpha < p$.

In this paper we obtain coefficient estimate, distortion and closure theorems and the radius of convexity for the class $P_k(p, A, B, \alpha)$. We also obtain class preserving integral operators of the form

$$F(z) = \frac{c+p}{z^c} \int_0^z t^{c-1} f(t) dt, \quad c > -p$$

for the class $P_k(p, A, B, \alpha)$. Conversely when $F(z) \in P_k(p, A, B, \alpha)$, radius of p -valence of $f(z)$ has been determined.

1. INTRODUCTION

Let S_p ($p \geq 1$) denote the class of functions of the form

$$f(z) = z^p + \sum_{n=1}^{\infty} a_{p+n} z^{p+n} \tag{1.1}$$

which are analytic and p -valent in the unit disc $U = \{z : |z| < 1\}$. For $-1 \leq B < A \leq 1$, $-1 \leq B < 0$, and $0 \leq \alpha < p$, let $P^*(p, A, B, \alpha)$ be the class of those functions $f(z)$ of S_p for which

$\frac{zf'(z)}{f(z)}$ is subordinate to $\frac{p + [pB + (A-B)(p-\alpha)]z}{1+Bz}$. In other words

$f(z) \in P^*(p, A, B, \alpha)$ if and only if there exists a function $w(z)$ satisfying $w(0) = 0$ and $|w(z)| < 1$ for $z \in U$ such that

$$\frac{zf'(z)}{f(z)} = \frac{p + (pB + (A-B)(p-\alpha))w(z)}{1 + Bw(z)}, \quad z \in U. \quad (1.2)$$

The condition (1.2) is equivalent to

$$\left| \frac{\frac{zf'(z)}{f(z)} - p}{[pB + (A-B)(p-\alpha)] - B \frac{zf'(z)}{f(z)}} \right| < 1, \quad z \in U. \quad (1.3)$$

Let T_p denote the subclass of S_p consisting of functions analytic and p -valent which can be expressed in the form

$$f(z) = z^p - \sum_{n=k}^{\infty} |a_{p+n}| z^{p+n} \quad (k \geq 2). \quad (1.4)$$

Let us define

$$P_k(p, A, B, \alpha) = P^*(p, A, B, \alpha) \cap T_p.$$

Aouf [1, 2], Gupta and Jain [5, 6], Goel and Sohi [4], Sarangi and Uralegaddi [29], Shukla and Dashrath [10], and Silverman [11] have studied certain subclasses of analytic functions with negative coefficients and Kumar [7] and Aouf [3] have recently studied certain subclasses of analytic functions with negative and missing coefficients.

In this paper, under the assumption $-1 \leq B < 0$, and $k \geq 2$, we obtain coefficient estimate, distortion theorem, covering theorem and radius of convexity for the class $P_k(p, A, B, \alpha)$. We also obtain the class preserving integral operators of the form

$$F(z) = \frac{c+p}{z^c} \int_0^z t^{c-1} f(t) dt, \quad c > -p \quad (1.5)$$

for the class $P_k(p, A, B, \alpha)$. Conversely, when $F(z) \in P_k(p, A, B, \alpha)$, we determine the radius of p -valence of $f(z)$ defined by (1.5). Lastly

we show that the class $P_k(p, A, B, \alpha)$ is closed under "arithmetic mean" and "convex linear combinations".

Remark: We observe that our distortion theorem (Theorem 2) improves the results of Sarangi and Patil [8, Theorem 3].

2. COEFFICIENT ESTIMATE

Theorem 1. Let the function $f(z)$ be defined by (1.4). Then $f(z) \in P_k(p, A, B, \alpha)$ if and only if

$$\sum_{n=k}^{\infty} [(1-B)n + (A-B)(p-\alpha)] |a_{p+n}| \leq (A-B)(p-\alpha). \quad (2.1)$$

The result is sharp.

Proof: Let $|z| = 1$, then

$$|zf'(z) - pf(z)| = |[pB + (A-B)(p-\alpha)] f(z) - Bzf'(z)|$$

$$= | - \sum_{n=k}^{\infty} n |a_{p+n}| z^{p+n} | - |(A-B)(p-\alpha) z^p |$$

$$+ \sum_{n=k}^{\infty} [nB + (B-A)(p-\alpha)] |a_{p+n}| z^{p+n} |$$

$$\leq \sum_{n=k}^{\infty} [(1-B)n + (A-B)(p-\alpha)] |a_{p+n}| - (A-B)(p-\alpha) \leq 0$$

(since $-1 \leq B < 0$).

Hence by the principle of maximum modulus $f(z) \in P_k(p, A, B, \alpha)$.

Conversely, suppose that

$$\begin{aligned} & \left| \frac{zf'(z)}{f(z)} - p \right| \\ & \leq \left| [pB + (A-B)(p-\alpha)] - B \frac{zf'(z)}{f(z)} \right| \\ & = \left| - \sum_{n=k}^{\infty} n |a_{p+n}| z^{p+n} \right| \\ & \leq (A-B)(p-\alpha) z^p + \sum_{n=k}^{\infty} [nB + (B-A)(p-\alpha)] |a_{p+n}| z^{p+n} < 1, \quad z \in U. \end{aligned}$$

Since $|Re(z)| \leq |z|$ for all z , we have

$$\operatorname{Re} \left\{ \frac{\sum_{n=k}^{\infty} n |a_{p+n}| z^{p+n}}{(A-B)(p-\alpha) z^p + \sum_{n=k}^{\infty} [nB + (B-A)(p-z)] |a_{p+n}| z^{p+n}} \right\} < 1. \quad (2.2)$$

Choose values of z on the real axis so that $\frac{zf'(z)}{f(z)}$ is real. Upon clearing the denominator in (2.2) and letting $z \rightarrow 1^-$ through real values, we obtain

$$\sum_{n=k}^{\infty} [(1-B)n + (A-B)(p-\alpha)] |a_{p+n}| \leq (A-B)(p-\alpha).$$

This completes the proof of the theorem.

Sharpness follows if we take

$$f(z) = z^p - \frac{(A-B)(p-\alpha)}{(1-B)n + (A-B)(p-\alpha)} z^{p+n} \quad (n \geq k, k \geq 2). \quad (2.3)$$

3. DISTORTION PROPERTIES

Theorem 2. If a function $f(z)$ defined by (1.4) is in the class $P_k(p, A, B, \alpha)$, then for $|z| = r$

$$\begin{aligned} r^p - \frac{(A-B)(p-\alpha)}{(1-B)k + (A-B)(p-\alpha)} r^{p+k} &\leq |f(z)| \\ &\leq r^p + \frac{(A-B)(p-\alpha)}{(1-B)k + (A-B)(p-\alpha)} r^{p+k} \end{aligned} \quad (3.1)$$

and

$$\begin{aligned} p r^{p-1} - \frac{(A-B)(p-\alpha)(p+k)}{(1-B)k + (A-B)(p-\alpha)} r^{p+k-1} &\leq |f'(z)| \\ &\leq p r^{p-1} + \frac{(A-B)(p-\alpha)(p+k)}{(1-B)k + (A-B)(p-\alpha)} r^{p+k-1} \end{aligned} \quad (3.2)$$

All the inequalities are sharp.

Proof: From Theorem 1, we have

$$\begin{aligned} & [(1-B) k + (A-B)(p-\alpha)] \sum_{n=k}^{\infty} |a_{p+n}| \\ & \leq \sum_{n=k}^{\infty} [(1-B) n + (A-B)(p-\alpha)] |a_{p+n}| \leq (A-B)(p-\alpha). \end{aligned}$$

This implies that

$$\sum_{n=k}^{\infty} |a_{p+n}| \leq \frac{(A-B)(p-\alpha)}{(1-B)k + (A-B)(p-\alpha)}. \quad (3.3)$$

Thus

$$\begin{aligned} |f(z)| & \leq |z|^p + \sum_{n=k}^{\infty} |a_{p+n}| |z|^{p+n} \\ & \leq r^p + r^{p+k} \sum_{n=k}^{\infty} |a_{p+n}| \\ & \leq r^p + \frac{(A-B)(p-\alpha)}{(1-B)k + (A-B)(p-\alpha)} r^{p+k}. \end{aligned}$$

Also

$$\begin{aligned} |f(z)| & \geq |z|^p - \sum_{n=k}^{\infty} |a_{p+n}| |z|^{p+n} \\ & \geq |z|^p - r^{p+k} \sum_{n=k}^{\infty} |a_{p+n}| \\ & \geq r^p - \frac{(A-B)(p-\alpha)}{(1-B)k + (A-B)(p-\alpha)} r^{p+k}. \end{aligned}$$

Further

$$\begin{aligned} |f'(z)| & \leq p r^{p-1} + \sum_{n=k}^{\infty} (p+n) |a_{p+n}| |z|^{p+n-1} \\ & \leq p r^{p-1} + r^{p+k-1} \sum_{n=k}^{\infty} (p+n) |a_{p+n}|. \end{aligned} \quad (3.4)$$

In view of Theorem 1.

$$\sum_{n=k}^{\infty} (1-B) \left[p + n - \frac{p(1-B) + (B-A)(p-\alpha)}{(1-B)} \right] |a_{p+n}| \leq (A-B)(p-\alpha)$$

or

$$\begin{aligned} \sum_{n=k}^{\infty} (1-B) (p + n) |a_{p+n}| &\leq (A-B) (p-\alpha) \\ &+ \left[p (1-B) + (B-A) (p-\alpha) \right] \sum_{n=k}^{\infty} |a_{p+n}| \end{aligned} \quad (3.5)$$

(3.5), with the help of (3.3), implies that

$$\sum_{n=k}^{\infty} (p+n) |a_{p+n}| \leq \frac{(A-B)(p-\alpha)(p+k)}{(1-B)k + (A-B)(p-\alpha)}. \quad (3.6)$$

A substitution of (3.6) into (3.4) yields the right-hand inequality of (3.2).

On the other-hand,

$$\begin{aligned} |f'(z)| &\geq p r^{p-1} - r^{p+k-1} \sum_{n=k}^{\infty} (p+n) |a_{p+n}| \\ &\geq p r^{p-1} - \frac{(A-B)(p-\alpha)(p+k)}{(1-B)k + (A-B)(p-\alpha)} r^{p+k-1} \end{aligned}$$

This completes the proof of Theorem 2.

Equality in (3.1) and (3.2) is obtained if we take

$$f(z) = z^p - \frac{(A-B)(p-\alpha)}{(1-B)k + (A-B)(p-\alpha)} z^{p+k} (z = \pm r).$$

Corollary 1. If $f(z) \in P_k(p, A, B, \alpha)$, then the disc U is mapped by $f(z)$ onto a domain that contains the disc

$$|w| < \frac{(1-B)k}{(1-B)k + (A-B)(p-\alpha)}.$$

The result is sharp with the extremal function

$$f(z) = z^p - \frac{(A-B)(p-\alpha)}{(1-B)k + (A-B)(p-\alpha)} z^{p+k}.$$

Putting $\alpha = 0$ in Theorem 2 and Corollary 1 we get:

Corollary 2. If a function $f(z)$ defined by (1.4) is in the class $P_k(p, A, B)$, then for $|z| = r$

$$r^p - \frac{(A-B)p}{(1-B)k + (A-B)p} r^{p+k} \leq |f(z)| \leq r^p + \frac{(A-B)p}{(1-B)k + (A-B)p} r^{p+k}$$

and

$$\begin{aligned} p r^{p-1} - \frac{(A-B)p(p+k)}{(1-B)k + (A-B)p} r^{p+k-1} &\leq |f'(z)| \\ &\leq p r^{p-1} + \frac{(A-B)p(p+k)}{(1-B)k + (A-B)p} r^{p+k-1} \end{aligned}$$

The result is sharp, with the external function

$$f(z) = z^p - \frac{(A-B)p}{(1-B)k + (A-B)p} z^{p+k} (z = \pm r).$$

Corollary 3. If $f(z) \in P_k(p, A, B)$, then the disc U is mapped by $f(z)$ onto a domain that contains the disc

$$|w| < \frac{(1-B)k}{(1-B)k + (A-B)p}.$$

The result is sharp with the external function

$$f(z) = z^p - \frac{(A-B)p}{(1-B)k + (A-B)p} z^{p+k}.$$

4. INTEGRAL OPERATORS

Theorem 3. Let c be a real number such that $c > -p$. If $f(z) \in P_k(p, A, B, \alpha)$, then the function $F(z)$ defined by (1.5) also belongs $P_k(p, A, B, \alpha)$.

Proof: Let $f(z) = z^p - \sum_{n=k}^{\infty} |a_{p+n}| z^{p+n}$. Then from the representation of $F(z)$, it follows that

$$F(z) = z^p - \sum_{n=k}^{\infty} |b_{p+n}| z^{p+n},$$

where

$$|b_{p+n}| = \left(\frac{c+p}{c+p+n} \right) |a_{p+n}|.$$

Therefore using Theorem 1 for the coefficients of $F(z)$ we have

$$\begin{aligned} & \sum_{n=k}^{\infty} [(1-B)n + (A-B)(P-\alpha)] |b_{p+n}| \\ &= \sum_{n=k}^{\infty} [(1-B)n + (A-B)(P-\alpha)] \left(\frac{c+p}{c+p+n} \right) |a_{p+n}| \\ &\leq (A-B)(p-\alpha) \end{aligned}$$

since $\frac{c+p}{c+p+n} < 1$ and $f(z) \in P_k(p, A, B, \alpha)$. Hence $F(z) \in P_k(p, A, B, \alpha)$

Theorem 4. Let c be a real number such that $c > -p$. If $F(z) \in P_k(p, A, B, \alpha)$, then the function $f(z)$ defined by (1.5) is p -valent in $|z| < R^*$, where

$$R^* = \inf_{n \geq k \geq 2} \left[\left(\frac{c+p}{c+p+n} \right) \left[\frac{(1-B)n + (A-B)(P-\alpha)}{(A-B)(p-\alpha)} \right] \left(\frac{p}{p+n} \right) \right]^{\frac{1}{n}}.$$

The result is sharp.

Proof: Let $F(z) = z^p - \sum_{n=k}^{\infty} |a_{p+n}| z^{p+n}$. It follows then from (1.5) that

$$f(z) = \frac{z^{1-c}}{c+p} [z^c F(z)] = z^p - \sum_{n=k}^{\infty} \left(\frac{c+p+n}{c+p} \right) |a_{p+n}| z^{p+n}.$$

In order to obtain the required result it suffices to show that

$$\left| \frac{f'(z)}{z^{p-1}} - p \right| < p \text{ in } |z| < R^*.$$

Now

$$\begin{aligned} \left| \frac{f'(z)}{z^{p-1}} - p \right| &= \left| - \sum_{n=k}^{\infty} (p+n) \left(\frac{c+p+n}{c+p} \right) |a_{p+n}| z^n \right| \\ &\leq \sum_{n=k}^{\infty} (p+n) \left(\frac{c+p+n}{c+p} \right) |a_{p+n}| |z|^n. \end{aligned}$$

Thus

$$\left| \frac{f'(z)}{z^{p-1}} - p \right| < p, \text{ if } \sum_{n=k}^{\infty} (p+n) \left(\frac{c+p+n}{c+p} \right) |a_{p+n}| |z|^n < p. \quad (4.1)$$

But Theorem 1 confirms that

$$\sum_{n=k}^{\infty} p \left[\frac{(1-B)n + (A-B)(p-\alpha)}{(A-B)(p-\alpha)} \right] |a_{p+n}| \leq p.$$

Hence (4.1) will be satisfied if

$$(p+n) \left(\frac{c+p+n}{c+p} \right) |a_{p+n}| |z|^n \leq p \left[\frac{(1-B)n + (A-B)(p-\alpha)}{(A-B)(p-\alpha)} \right] |a_{p+n}|,$$

$$n \geq k \geq 2$$

or if

$$|z| \leq \left\{ \left(\frac{c+p}{c+p+n} \right) \left[\frac{(1-B)n + (A-B)(p-\alpha)}{(A-B)(p-\alpha)} \right] \left(\frac{p}{p+n} \right) \right\}^{\frac{1}{n}}$$

$$n \geq k \geq 2.$$

Therefore $f(z)$ is p -valent in $|z| < R^*$. Sharpness follows if we take

$$F(z) = z^p - \frac{(A-B)(p-\alpha)}{(1-B)n + (A-B)(p-\alpha)} z^{p+n}, \quad n \geq k \geq 2.$$

5. RADIUS OF CONVEXITY

Theorem 5. If $f(z) \in P_k(p, A, B, \alpha)$, then $f(z)$ is p -valently convex in the disc $|z| < R_p$, where

$$R_p = \inf_{n \geq k \geq 2} \left\{ \left[\frac{(1-B)n + (A-B)(p-\alpha)}{(A-B)(p-\alpha)} \right] \left(\frac{p}{p+n} \right)^2 \right\}^{\frac{1}{n}}.$$

The result is sharp.

Proof: In order to establish the required result it suffices to show that

$$\left| \left[1 + \frac{zf''(z)}{f'(z)} \right] - p \right| \leq p \text{ for } |z| < R_p.$$

Let $f(z) = z^p - \sum_{n=k}^{\infty} |a_{p+n}| z^{p+n}$. Then we have

$$\begin{aligned} \left| \left[1 + \frac{zf''(z)}{f'(z)} \right] - p \right| &= \left| \frac{- \sum_{n=k}^{\infty} n(p+n) |a_{p+n}| z^n}{p - \sum_{n=k}^{\infty} (p+n) |a_{p+n}| z^n} \right| \\ &\leq \frac{\sum_{n=k}^{\infty} n(p+n) |a_{p+n}| |z|^n}{p - \sum_{n=k}^{\infty} (p+n) |a_{p+n}| |z|^n}. \end{aligned}$$

Therefore $\left| \left[1 + \frac{zf''(z)}{f'(z)} \right] - p \right| \leq p$ if

$$\sum_{n=k}^{\infty} n(p+n) |a_{p+n}| |z|^n \leq p^2 - \sum_{n=k}^{\infty} p(p+n) |a_{p+n}| |z|^n$$

or if

$$\sum_{n=k}^{\infty} \left(\frac{p+n}{p} \right)^2 |a_{p+n}| |z|^n \leq 1. \quad (5.1)$$

From Theorem 1, we have

$$\sum_{n=k}^{\infty} \frac{(1-B)n + (A-B)(p-\alpha)}{(A-B)(p-\alpha)} |a_{p+n}| \leq 1.$$

Hence (5.1) will be satisfied if

$$\left(\frac{p+n}{p} \right)^2 |z|^n \leq \left[\frac{(1-B)n + (A-B)(p-\alpha)}{(A-B)(p-\alpha)} \right]$$

or if

$$|z| \leq \left\{ \left[\frac{(1-B)n + (A-B)(p-\alpha)}{(A-B)(p-\alpha)} \right] \left(\frac{p}{p+n} \right)^2 \right\}^{\frac{1}{n}}, \quad n \geq k, \quad k \geq 2.$$

Therefore $f(z)$ is p -valently convex in the disc $|z| < R_p$. The result is sharp with the extremal function $f(z)$ defined by (2.3).

6. CLOSURE PROPERTIES

In this section we show that the class $P_k(p, A, B, \alpha)$ is closed under “arithmetic mean” and “convex linear combinations”.

Theorem 6. If $f_j(z) = z^p - \sum_{n=k}^{\infty} |a_{p+n}|_j z^{p+n}$, $j = 1, 2, \dots, m$.

If $f_j(z) \in P_k(p, A, B, \alpha)$ for each $j = 1, 2, \dots, m$, then the function $h(z)$

$= z^p - \sum_{n=k}^{\infty} |b_{p+n}| z^{p+n}$ also belongs to $P_k(p, A, B, \alpha)$, where $b_{p+n} =$

$$\frac{1}{m} \sum_{j=1}^m a_{p+n}.$$

Proof: Since $f_j(z) \in P_k(p, A, B, \alpha)$, it follows from Theorem 1 that

$$\sum_{n=k}^{\infty} [(1-B)n + (A-B)(p-\alpha)] |a_{p+n}|_j \leq (A-B)(p-\alpha), \quad j = 1, 2, \dots, m.$$

Therefore

$$\sum_{n=k}^{\infty} [(1-B)n + (A-B)(p-\alpha)] |b_{p+n}|$$

$$\begin{aligned} &\leq \sum_{n=k}^{\infty} [(1-B)n + (A-B)(p-\alpha)] \left\{ \frac{1}{m} \sum_{j=1}^m |a_{p+n-j}| \right\} \\ &\leq (A-B)(p-\alpha). \end{aligned}$$

Hence, by Theorem 1, $f(z) \in P_k(p, A, B, \alpha)$.

Theorem 7. Let $f_p(z) = z^p$ and

$$f_{p+n}(z) = z^p - \frac{(A-B)(p-\alpha)}{(1-B)n + (A-B)(p-\alpha)} z^{p+n} (n \geq k, k \geq 2).$$

Then $f(z) \in P_k(p, A, B, \alpha)$ if and only if it can be expressed in the form $f(z) = \lambda_p f_p(z) + \sum_{n=k}^{\infty} \lambda_n f_{p+n}(z)$, where $\lambda_n \geq 0$ and $\lambda_p + \sum_{n=k}^{\infty} \lambda_n = 1$.

Proof: Let us assume that

$$\begin{aligned} f(z) &= \lambda_p f_p(z) + \sum_{n=k}^{\infty} \lambda_n f_{p+n}(z) \\ &= [1 - \sum_{n=k}^{\infty} \lambda_n] z^p + \sum_{n=k}^{\infty} \lambda_n [z^p - \frac{(A-B)(p-\alpha)}{(1-B)n + (A-B)(p-\alpha)} z^{p+n}] \\ &= z^p - \sum_{n=k}^{\infty} \frac{(A-B)(p-\alpha)}{(1-B)n + (A-B)(p-\alpha)} \lambda_n z^{p+n}. \end{aligned}$$

Then from Theorem 1 we have

$$\begin{aligned} &\sum_{n=k}^{\infty} [(1-B)n + (A-B)(p-\alpha)] \left[\frac{(A-B)(p-\alpha)\lambda_n}{(1-B)n + (A-B)(p-\alpha)} \right] \\ &= (A-B)(p-\alpha) \sum_{n=k}^{\infty} \lambda_n \leq (A-B)(p-\alpha). \end{aligned}$$

Hence $f(z) \in P_k(p, A, B, \alpha)$.

Conversely, let $f(z) \in P_k(p, A, B, \alpha)$. It follows from Theorem 1 that

$$|a_{p+n}| \leq \frac{(A-B)(p-\alpha)}{(1-B)n + (A-B)(p-\alpha)} \quad (n = k, k+1, \dots, k \geq 2).$$

Setting

$$\lambda_n = \frac{(1-B)n + (A-B)(p-\alpha)}{(A-B)(p-\alpha)} |a_{p+n}|, \quad (n = k, k+1, \dots, k \geq 2).$$

and

$$\lambda_p = 1 - \sum_{n=k}^{\infty} \lambda_n,$$

we have

$$\begin{aligned} f(z) &= z^p + \sum_{n=k}^{\infty} |a_{p+n}| z^{p+n} \\ &= z^p + \sum_{n=k}^{\infty} \lambda_n z^p + \sum_{n=k}^{\infty} \lambda_n z^p - \sum_{n=k}^{\infty} \lambda_n \frac{(A-B)(p-\alpha)}{(1-B)n + (A-B)(p-\alpha)} z^{p+n} \\ &= [1 - \sum_{n=k}^{\infty} \lambda_n] z^p + \sum_{n=k}^{\infty} \lambda_n [z^p - \frac{(A-B)(p-\alpha)}{(1-B)n + (A-B)(p-\alpha)} z^{p+n}] \\ &= \lambda_p f_p(z) + \sum_{n=k}^{\infty} \lambda_n f_{p+n}(z). \end{aligned}$$

This completes the proof of Theorem 7.

Remarks:

(1) Putting $\alpha = 0$ in Theorems 1, 3, 5, 6 and 7, we get the corresponding results obtained by Sarangi and Patil [8].

(2) We observe that our results in Corollary 2 and Corollary 3 improves the results of Sarangi and Patil [8, Theorem 3 and its Corollary].

REFERENCES

- [1] M.K. AOUF., A generalization of the multivalent functions with negative coefficients, J. Korean Math. Soc. 25. (1988), no. 1, 53-66.

- [2] M.K. AOUF., A generalization of the multivalent functions with negative coefficients. II, Bull. Korean Math. Soc. 25 (1988), no. 2, 221-232.
- [3] M.K. AOUF On p-valent functions with negative and missing coefficients, J. Math. Res. Exposition 10 (1990), no. 2, 249-256.
- [4] R.M. GOEL, and N.S. SOHIL, Multivalent functions with negative coefficients, Indian J. Pure Appl. Math. 12 (1981), no. 7, 844-853.
- [5] V.P. GUPTA, and P.K. JAIN., Certain classes of univalent functions with negative coefficients, Bull. Austral. Math. Soc. 14 (1976), 409-416.
- [6] V.P. GUPTA, and P.K. JAIN., Certain classes of univalent functions with negative coefficients. II, Bull. Austral. Math. Soc. 15 (1976), 467-473.
- [7] VINOD, KUMAR. On univalent functions with negative and missing coefficients, J. Math. Res. Exposition 4 (1984), no. 1, 27-34.
- [8] S.M. SARANCI, and V.J. PATIL., On multivalent functions with negative and missing coefficients, J. Math. Res. Exposition 10 (1990), no. 3, 341-348.
- [9] S.M. SARANCI, and B.A. URALAGADDI., The radius of convexity and starlikeness for certain classes of analytic functions with negative coefficients. I, Acad. Naz. Lincei Rend. 65 (1978), 38-42.
- [10] S.L. SHUKLA, and DASHRATH., On certain classes of multivalent functions with negative coefficients, Pure Appl. Math. Sci. 20 (1984), 1-2, 63-72.
- [11] H. SILVERMAN., Univalent functions with negative coefficients, Proc. Amer. Math. Soc. 51 (1975) 109-116.