


ISSN-1308-416X

# Cell Membranes and Free Radical Research

Volume 1 Number 2 1 December 2008

EDITOR-IN-CHIEF  
Mustafa Nazıroğlu, Isparta, TURKEY



# Unifying Mechanism for Eye Toxicity: Electron Transfer, Reactive Oxygen Species, Antioxidant Benefits, Cell Signaling and Cell Membranes

Peter Kovacic,<sup>1</sup> Ratnasamy Somanathan<sup>1,2</sup>

<sup>1</sup>Department of Chemistry, San Diego State University, San Diego, CA 92182-1030 USA

<sup>2</sup>Centro de Graduados e Investigación del Instituto Tecnológico de Tijuana, Apdo postal 1166, Tijuana, B.C.

---

## ABSTRACT

Eye toxicity comprises an important part in the category of insults to major organs. There is considerable literature comprising industrial chemicals, atmospheric pollutants, metals, pesticides, therapeutic drugs, abused drugs and household chemicals. The list includes both endogenous and exogenous materials. The various constituents of the eye are affected. This review provides extensive evidence for involvement of oxidative stress (OS) and electron transfer (ET) as a unifying theme. Successful application of the mechanistic approach is made to all of the main classes of toxicants, in addition to large numbers of miscellaneous types. We believe it is not coincidental that the vast majority of these substances incorporate ET functions (quinone, metal complex, ArNO<sub>2</sub>, or conjugated iminium) either per se or in metabolites, potentially giving rise to reactive oxygen species (ROS) by redox cycling. Some categories, e.g., peroxide and radiation, appear to generate ROS by non-ET routes. Beneficial effects of antioxidants are documented extensively, in addition to literature on cell signaling.

**Key words:** eye toxins, electron transfer, oxidative stress, antioxidants, cell membranes.

---

Cell membranes play an important role, particularly the lipid constituents which undergo oxidation to toxic hydroperoxides. Recent literature is discussed which links pathogenesis of various eye components to toxicity resulting from oxidative injury to membrane lipids. Our framework should increase understanding and contribute to preventative measures, such as use of antioxidants. This review is the first to provide a comprehensive approach to the ET-ROS-OS mechanism. Also, it is unique in applying a multifaceted relationship to the toxicity mode of action.

## INTRODUCTION

Extensive literature exists on the subject of eye toxicants which fall into various classes, such as industrial chemicals, atmospheric pollutants, metals, pesticides, therapeutic drugs, abused drugs, and household chemicals. The list includes endogenous and exogenous substances. The various constituents of the eye are affected resulting in adverse reactions ranging from minor irritation to permanent blindness. The preponderance of bioactive substances and their metabolites incorporate electron transfer (ET) functions, which, we believe, play an important role in physiological responses. The main groups include quinones (or phenolic precursors), metal complexes (or complexors), aromatic nitro compounds (or reduced hydroxylamine and nitroso derivatives), and conjugated imines (or iminium species). In vivo redox cycling with oxygen can occur giving rise to oxidative stress (OS) through generation of reactive oxygen species (ROS), such as hydrogen peroxide, hydroperoxides, alkylperoxides, and diverse radicals (hydroxyl, alkoxy, hydroperoxyl, and superoxide). In some cases, ET results in

interference with normal electrical effects, e.g., in respiration or neurochemistry. Generally, active entities possessing ET groups display reduction potentials in the physiologically responsive range, i.e., more positive than -0.5 V. ET, ROS, and OS have been increasingly implicated in the mode of action of drugs and toxins, e.g., antiinfective agents (Kovacic and Becvar, 2000), anticancer drugs (Kovacic and Osuna, 2000), carcinogens (Kovacic and Jacintho, 2001a), and toxicants, namely reproductive (Kovacic and Jacintho, 2001b), hepatic (Kovacic et al. 2002), renal (Poli et al. 1989), cardiovascular (Kovacic and Thurn, 2005), neural (Kovacic and Somanathan, 2005), mitochondrial (Kovacic et al. 2005), abused drugs (Kovacic and Cooksy, 2005), ear (Kovacic and Somanathan, 2008), and various other categories including human illnesses (Halliwell and Gutteridge, 1999a).

There is a plethora of experimental evidence supporting the OS theoretical framework, including generation of the common ROS, lipid peroxidation, degradation products of oxidation, depletion of AOs, and DNA oxidation and cleavage products, as well as electrochemical data. This comprehensive, unifying mechanism is in keeping with the frequent observations that many ET substances display a variety of activities, e.g., multiple drug properties, as well as toxic effects. Knowledge of events at the molecular level can result in practical application in medicine.

It is instructive to examine the basic biochemistry of ET functionalities in more detail. Redox cycling occurs between hydroquinone and p-benzoquinone, and between catechol and o-benzoquinone with generation of superoxide via ET to oxygen. Semiquinones act as intermediates. Various amino acids can operate

as electron donors. Superoxide serves as precursor to a variety of other ROS. The quinones can belong in either the endogenous or exogenous category. In the case of aromatic nitro compounds, the reduced nitroso and hydroxylamine metabolites can similarly enter into redox cycling, including an oxy radical intermediate. This class is only in the exogenous group. Less known are conjugated iminium compounds, of which paraquat is a predominant member (see below). Electron uptake yields resonance contributors. The imine precursor, commonly formed by condensation of protein primary amino groups with carbonyls, is readily converted to iminium by protonation. The metal category is treated in a section below.

This review demonstrates that the ET-ROS-OS unifying theme, which has been successful for many other classes of toxicants, can also be applied to eye toxicity. The toxic groups include a wide variety of structurally diverse substances. Not surprisingly, numerous reports exist on the beneficial effects of antioxidants (AOs). Cell signaling and cell membranes are also addressed. Our summary reveals that the mechanistic framework serves as a common thread for the large majority of reported toxins. This review is the only comprehensive, unifying and integrated one based on ET-ROS-OS, AOs and cell signaling. However, it should be emphasized that physiological activity of endogenous and exogenous substances is often complex and multifaceted. Our objective does not encompass extensive treatment of other modes of action. A number of original references may be found in the reviews and articles cited. The literature covered is mainly from more recent years.

## POTENTIAL MECHANISMS OF EYE TOXICITY

### Electron Transfer

A brief summary of the basic biochemistry of ET functionalities would be helpful as foundation for ensuing sections. Quinones undergo redox cycling with the corresponding hydroquinone or catechol with intermediate involvement of semiquinones (Kovacic and Becvar, 2000). A pertinent example is naphthoquinone (see Naphthalene section). A similar situation pertains to nitroso and hydroxylamine metabolites of ArNO<sub>2</sub> involving intermediate oxy radicals (Kovacic and Becvar, 2000). A relevant example is chloramphenicol in the Therapeutic Drugs section. In some cases, the parent nitro compound is capable of electron uptake. For metal complexes, higher valence metals provide a large surface for electron delocalization and electron attraction by positive charge. Conjugated iminium species entail a similar electronic attraction and resonance stabilization via delocalization.

### Reactive Oxygen Species and Oxidative Stress

There is extensive literature dealing with the toxic effect of ROS on various constituents of the eye. Glaucoma is associated with OS which contributes to pathogenesis and might be a target for prevention and therapy (Saccá et al. 2007). The ROS adversely affect the human trabecular meshwork (TM) whose integrity is compromised. There is correlation between oxidative DNA damage in the TM and visual field defects. The OS may contribute to homeostasis between NO and endothelins, with possible involvement in ganglionic cell death. Furthermore, a genetic disposition may make for greater susceptibility to ROS damage. Evidence supports involvement of the immune system in glaucomatous neurodegeneration (Tezel et al. 2007). Findings support ROS regulation of the immune response by stimulating the antigen-presenting ability of glial cells and functioning as co-stimulatory agents for antigens. Increasing evidence points to a key role for ROS in this condition (Izzotti et al. 2006a). Oxidative DNA damage is enhanced in the optical epithelium regulating aqueous humour out flow. The pathogenic role of ROS has implications for prevention (see AO section). There is increase in 8-OH-dG (8-hydroxy-2-deoxyguanosine) from DNA oxidative degradation in these patients (Izzotti et al. 2006b). Several lines of evidence support the pathogenic role of ROS, including the following items: (a) high levels of hydrogen peroxide, (b) TM possesses AO activity, (c) increases in SOD and GSH peroxidase activities in response to the threat by ROS, (d) ROS compromise TM integrity.

Reports provide considerable evidence in support of a role for ROS in cataract formation. Aging is by far the biggest risk factor for cataract (Truscott, 2005). Oxidation is the hallmark of this age-related illness. Loss of thiol groups and methionine oxidation increase as the condition worsens. There may be no significant protein oxidation in the lens center with advancing age. The key element in oxidative prevention appears to be the level of GSH. Evidence indicates that the lens barrier is importantly involved. About 25% of the population aged >65 years and about 50% aged >80 years experience a serious loss of vision from cataracts (Babizhayev et al. 2004). There is support for involvement of free radical-induced lipid oxidation in cataract development. Initial stages are characterized by accumulation of primary lipid peroxidation products, whereas in later stages there is prevalence of fluorescent end products. Oxidation of thiol groups results from a decrease in GSH. Data show that N-acetylcarnosine is effective in the management of cataract reversal, as well as prevention. Two major processes can occur with proteins in cataractogenesis (Davies, 2001). One involves direct photo-oxidation arising from UV radiation, thereby generating excited states or radicals. The other entails indirect oxidation

of protein by UV generated ROS. Mechanistic aspects are discussed. The phacoemulsification and aspiration (PEA) technique has become the most popular cataract surgery (Takahashi, 2005). However, radical formation from ultrasound is believed to be a cause of corneal endothelium damage in PEA. In aqueous solution, ultrasound induces cavitation, resulting in generation of highly toxic hydroxyl radicals. The major ingredient in ophthalmic viscosurgical devices is hyaluronate, a free radical scavenger which is expected to provide protection against OS.

A related aspect comprises insult to the eye lens. Oxidative stress was induced in cultured lens epithelial cells and in intact mouse lenses by hydrogen peroxide (Petersen, 2007). The lenses exposed to the peroxide suffered loss of transparency, decreased proteasome activity and lowered GSH levels. Inhibition of proteasome in the cells produced increase in apoptosis and disturbed redox balance. Evidence was provided for involvement of an oxidative process by ROS that mediates apoptosis induced by transforming growth factor beta-2 in human lens epithelial cells (Yao et al. 2007). Results indicate redox regulation of lens epithelial cells (Wilhelm et al. 2007). Hydrogen peroxide and two AOs were used in the study. Both AOs completely inhibited proliferation.

Considerable work has been done on the relationships of OS to macular degeneration and retinal damage. New insights were recently provided on retinopathy which appears to be derived from OS (Hardy et al. 2005). Retinal tissue is rich in polyunsaturated fat which is sensitive to lipid peroxidation by ROS, and the tissue responds to pathogenic stimuli by activation of phospholipases that is the first step in the synthesis of important classes of lipid second messengers. There is discussion of the role of retinal oxygen metabolism in origin of macular degeneration (Nowak et al. 2005). The ROS are believed to be responsible for apoptotic cell death and development of pathological changes in the age-related macular degeneration. Many AO systems in the retina provide protection from damage by ROS. ROS appear to be involved in diabetic retinopathy (Yamato et al. 2007). Evidence from ESR was obtained for free radical species in the eyes of mice with induced diabetes. Data are reported dealing with the effect of OS on mammalian retina (Yu et al. 2007). Oxidation reversibly inhibited phosphatase activity due to disulfide formation from Cys residues. Treatment with hydrogen peroxide caused an increase in phosphatase expression, as did constant exposure to light which induced photooxidative stress. Both oxidation and overexpression of phosphatase are enhanced by inhibition of the GSH AO system.

Various other articles deal with the subject of eye damage by ROS-OS. Oxidative stress from ROS is related to many chronic disorders, as pointed out in

the Introduction, including cataracts and macular degeneration (Tsubota, 2007.). The ocular-surface is subject to OS from UV light, air oxygen changes in oxygen pressure due to blinking and inflammation. Mice deficient in SOD exhibited clinical aspects of macular degeneration. A case was reported of optic neuropathy related to inhalation of hydrogen peroxide (Domagala et al. 2007). Evidence points to susceptibility of the ocular lens to tissue damage resulting from exposure to nitric oxide (Varma, 2007). Adverse effects include cataract formation. Another investigation also deals with NO-related cellular damage in the eye (Ozdek, 2007). Macular iron levels increase with age, which has the potential of contributing to retinal degeneration (He et al. 2007; Wong et al. 2007). Post-mortem retinas from patients with macular degeneration have increased levels of iron. Compounds of this metal may play a role in a broad range of other ocular diseases, including glaucoma, cataracts and hemorrhage. Prevention by iron chelation has clinical potential. References to toxins in the Introduction discuss the action of iron as a Fenton catalyst in generation of toxic hydroxyl radicals from hydrogen peroxide. In addition to the adverse effects by ROS on the eye already mentioned, others include uveitis and retrolental fibroplasias (Ohia et al. 2005). Various peroxides exert toxic reactions of the anterior uvea, especially on sympathetic nerves and smooth muscles of the iris-ciliary bodies. Calcium and arachidonic acid metabolites appear to be involved. Effects caused by the peroxides in the retina are mediated by second messengers, such as NO. A variation in ocular perfusion may lead to an increase in ROS that could induce an apoptotic cascade in retinal ganglion cells (Haufroid and Collignon-Robe, 2004). An article puts focus on free-radical oxidation in the pathogenesis of eye diseases (Kravchuk, 2004). In addition to research already treated, there is a report on light-induced photoreceptor damage (Meyer-Rochow, 2001). A large number of variables are involved, including free radicals and lipid peroxides. The chemistry is discussed in more detail in a review on mechanism of carcinogenesis (Kovacic and Jacintho, 2001a.). A widespread endogenous source of ROS comprises the mitochondria which leak a small percent of electrons from the ET chain, resulting in formation of superoxide from reaction with oxygen (Kovacic et al. 2005). Various reports deal with the adverse effects of OS from ROS on eye constituents (Ottonello et al. 2000; Ohia et al. 2005; Moreno et al. 2004; Santosa and Jones 2005; Beatty, et al. 2000; Marsili et al 2004; Pradhan et al. 2004; Varma and Hegde, 2007; Spector, 1995; Seetharam and Sujatha, 1998)

## Reactive Nitrogen Species

The most important member is nitric oxide which is treated in a subsequent section. Peroxynitrite, formed



by reaction of NO with superoxide, can be beneficial or induce toxic effects (Jacintho and Kovacic, 2003). Aslan et al. (2006), using rat models of ocular injury, found that both elevated intraocular pressure and ocular inflammation augment inducible nitric oxide synthase (NOS2) expression, retinal protein nitration and apoptosis. Aslan et al. (2007) report that extensive formation of nitrated proteins and reactive nitrogen species in the cornea contributes to tissue destruction in uveitis. Hence, inhibition of NOS2 may have beneficial effects.

## SURVEY OF EYE TOXICANTS

### Ultraviolet (UV) Radiation

Normally, light transmission through the eye is benign. However, intense light exposure can be hazardous (Roberts, 2002, Roberts, 2001). Before middle age, there is protection by an efficient AO system. In addition, protective pigments dissipate energy. A brief review presents evidence that solar UV radiation increases the risk of several diseases of the eye, such as cataract and neoplasms (Gallagher and Lee, 2006). Various studies deal with cataract formation due to UV radiation (McCarty and Taylor, 2002; Ayala et al. 2000), involvement of free radicals, lipid peroxidation and protein modification (van Kuijk, 1991), as well as ROS formation and DNA damage (Wolf et al. 2008). Recent investigations have been carried out on the adverse effects on the eye of increased exposure to UV radiation resulting from ozone depletion in the atmosphere (West et al. 2005; de Gruijl et al. 2003). Radiation induces increase in ROS which can be countered by repair processes (Risa et al. 2004). Because of involvement by ROS in toxicity, research was carried out on AOs which were found to be beneficial (Karslioglu et al. 2005, Anwar and Moustafa, 2001; Fu et al. 2000). In other research, radiation induced proto-oncogene expression and cell death in ocular tissues (Wickert et al. 1999). In connection with other high energy radiation, laser eye therapy can generate ROS and RNS, which in turn initiate lipid peroxidation, protein damage or DNA modification (Mileva and Zlateva, 2005). Vitamin E supplementation reduces OS significantly.

### ENDOGENOUS AGENTS

Some of these are discussed in earlier sections, such as ROS.

### Nitric Oxide (NO)

This free radical serves a variety of functions in the body, including action as second messenger. The diverse effects involve vascular tone, neurotransmission, immune cytotoxicity and many others. A review deals with physiological and pathological functions, e.g., in the eye (Becquet et al. 1997). Adverse in-

fluences are inflammation and degenerative diseases (glaucoma and retinal insult). The toxicity of NO is reviewed from the standpoint of ROS-OS (Jacintho and Kovacic, 2003).

### 4-Hydroxy-2-nonenal (HNE)

This toxin has attracted much attention as a degradation product of lipid oxidation. Cataract formation is associated with increased levels of HNE (Lassen et al. 2007). More recently, the role of the more toxic 4-keto oxidation product has been addressed in relation to ET-ROS (Kovacic, 2006.).

### Advanced Glycation End-products (AGE)

Formation of AGE is a key pathological event linked to a range of human diseases (Stitt, 2001). AGE are receiving attention as modulators of important visual disorders. Reports indicate widespread AGE accumulation at sites of ocular pathology. These products also induce apoptosis of bovine retinal pericytes (Denis et al. 2002). Treatment with various AOs completely inhibited apoptosis, suggesting involvement of OS. Other investigations associate AGE with generation of OS (Halliwell and Gutteridge, 1999c). The overall process entails formation of toxic ROS, such as hydrogen peroxide, superoxide and hydroxyl radical.

### XENOBIOTICS

There are various reports dealing with large numbers or mixtures of eye toxicants, including common ones (McCaa, 1982.), environmental pollutants (Saxena et al. 2003; Jankauskienė et al. 2007), organic solvents (Gong et al. 2003), volatile chemicals (Doty et al. 2004), industrial chemicals (Gobba, 2003), and drugs and ocular toxicants (Chiou, 1999). Mode of action based on ET-ROS-OS is discussed in reviews on organ toxins in the Introduction. Individual members are addressed in the following treatment.

### OTHER ENVIRONMENTAL TOXICANTS

#### Naphthalene

Naphthalene metabolites were examined in relation to lens cell toxicity (Russell et al. 1991). The quinone product was toxic to most cells and caused depletion of GSH levels. *o*-Naphthoquinone caused cortical cataract formation (Martynkina et al. 2002). Mitochondria in lens epithelial cells are the target of the toxicity. Oxidative stress has been implicated (Pandya et al. 2000). Apoptosis also occurs, which is attenuated by the AO curcumin. Additional discussion of OS via ROS from quinone metabolites is discussed elsewhere (Kovacic and Somanathan, 2005).

#### Styrene

Exposure of the eye to this polymer monomer im-

pairs color vision even at relatively low concentrations (Gong et al. 2002). Metabolites that may generate OS are the sidechain oxide and 4-vinylphenol (Kovacic and Somanathan 2006b).

### Toluene

Authors concluded that impairment of color vision in industry can be chronic (Zavali et al. 1998). Metabolic analogy has been drawn between benzene and toluene entailing hydroxylation as precursor to quinone formation (Kovacic and Jacintho, 2001a). The origin of ROS has been addressed, entailing hydroxylated metabolites, quinone methide and epoxide (Kovacic and Somanathan, 2005).

### Ethylene Oxide

This reagent, widely used as an industrial intermediate for ethylene glycol, may produce cataracts (Fujishiro et al. 1991). It is instructive that changes in the GSH redox cycle may be involved. A prior review documents the ability of alkylating agents to generate ROS (Kovacic and Osuna, 2000).

### Sulfur Mustard

This chemical warfare agent mainly affects the eye, skin and pulmonary system (Vidan et al. 2002). Ocular injury is the most immediate and distressing. Inflammation and free radical formation are evidently involved (Morad et al. 2005). The poison causes destructive ocular lesions on the surface and cornea, leading to visual deterioration and irritation (Javadi et al. 2005). As in the case of the prior section, this chemical is an alkylating agent with the ability to generate OS (Kovacic and Osuna, 2000). DNA alkylation generates a conjugated iminium.

### Methyl Methacrylate

In a study of this industrial monomer in the polymer industry, the following insults were observed: corneal edema, neurovascularization, iris engorgement, inflammation, limbal hyperemia and cataract (Holyk and Eifrig, 1979). The monomer resulted in depletion of liver GSH (Elovaara et al. 1983). It is proposed that activation occurs by oxidation to the 2,3-epoxy derivative (Boylard and Chasseaud, 1967). Hence, the mechanism may be similar to that of the above alkylating agents.

### Poynuclear Aromatic Hydrocarbons (PAHs)

Experiments with fish demonstrate an interaction between UV radiation and PAHs as a possible contributing factor to cataract induction (Laycock et al. 2000). Much research, particularly in the cancer and anticancer drug areas, reveal metabolism to an epoxide which functions as the toxic alkylator (Kovacic and

Jacintho, 2001). Above sections address ROS formation.

### Ethanol

There is increased risk of cataract formation from daily consumption of three drinks of alcoholic beverages (Prickett et al. 2004). On the other hand, there is decreased risk from intake of one drink daily. As discussed in a review, the higher risk may be attributed to enhanced production of ROS, whereas the beneficial effect may be due to AO action (Kovacic and Somanathan, 2006a).

### Cyanide

Results with rabbits suggest that cyanide-induced cataractogenesis may be a consequence of disruption of vitreous human and lenticular Ca homeostasis (Okolie and Audu, 2004). Mitigation results from AO vitamin administration (Okolie and Asonye, 2004). Neurotoxicity is accompanied by increases in superoxide and lipid peroxidation (Kovacic and Somanathan, 2005). The adverse effects are attenuated by aspirin. Part of the pro-oxidant effect may reflect inhibition of AO enzymes, such as SOD and catalase.

### Sulfur Dioxide

Various effects on the eye from inhalation exposure were examined (Kilic, 2000). Visual-evoked potentials are altered for which lipid peroxidation could play a role. There is potential for induction of protective AO enzymes. Decrease in activities of SOD and GSH was noted in studies as a heart toxin (Kovacic and Thurn, 2005). In water, sulfite is generated which is oxidized to radical anions of SO<sub>3</sub>, SO<sub>4</sub> and SO<sub>5</sub>. Peroxidation of rat mitochondria can be induced by sulfite.

### Anesthetics

Topical ocular anesthesia has been part of ophthalmology for more than a century (McGee and Fraunfelder, 2007). The most commonly used drugs are generally well tolerated, but can also be toxic to the eye, especially when abused. The most common insults are to the ocular surface, with others including corneal infiltrates, ulceration and perforation.

### Therapeutic Drugs

There has been substantial research on this subject. The reports include various ocular toxic reactions brought about by a variety of therapeutic agents, including tamoxifen, retinoids, sulfa drugs, anticancer agents, aminoglycosides and antibiotics (Moorthy and Valluri, 1999; Fraunfelder, 2003; Santaella and Fraunfelder, 2007; Abdollahi et al. 2004; Li et al. 2008, Fraunfelder, 2006; Fraunfelder and Fraunfelder, 2004; Widmer and Helbig, 2006; al-Twei-

geri et al. 1996). Research on ocular toxicity has also centered on individual drugs, such as fluoroquinolones (Thompson, 2007), desferrioxamine (Blake et al. 1985), phenothiazine (Boet, 1970), 5-fluorouracil (Shapiro et al. 1985), chloramphenicol (Cole et al. 1957), chlorpromazine (Siddall, 1966), amphotericin B (Barza et al. 1985), radiosensitizer (Rootman et al. 1982), and carbencilline and gentamycin (Jain and Hussain, 1982). The unifying mechanistic framework for these drugs, based on ET-ROS-OS is available in reviews on anti-infective agents (Kovacic and Becvar, 2000) and anti-cancer drugs (Kovacic and Osuna, 2000).

### Abused Drugs

Abuse of drugs, both legal and illegal, can cause ocular injury and disease (Mclane and Carroll, 1986). The drugs are grouped into five categories: opiates, marijuana, stimulants, depressants and hallucinogens. Cocaine is reported to cause corneal disturbances ranging from superficial punctate keratitis to perforation (Pilon and Scheiffle, 2006). The keratopathy may involve corneal epithelial disruption and stromal ulceration. A key cocaine metabolite is the nitroxide radical. A unifying mechanism for abused drug action and toxicity is proposed based on ET-ROS-OS (Kovacic and Cooksy, 2005).

### Cigarette Smoke

Reports reveal strong association between smoking and common eye diseases, including Graves' ophthalmopathy, macular degeneration, glaucoma and cataract (Cheng et al. 2000). It causes morphological and functional changes to the lens and retina, which can occur by direct contact or via the lung. There is enhanced generation of radicals and depletion of AOs. Ocular damage may lead to permanent blindness. Various studies deal with a link between smoking and cataract development, such as epidemiological data (Sulochana, 2001). Smokeless tobacco use was rather strongly associated with cataract (Raju et al. 2006). Decreased levels of the AOs, namely SOD, GSH and GSH peroxidase in lens of smokers suggests oxidative insult involved in cataract formation (Sulochana et al. 2002). Smoking produces deteriorating effects on the lipid layer of the ocular surface (Altinors et al. 2006). Cigarette smoke condensate causes OS and negatively affects detoxifying agents, such as GSH and GSH-S-transferase (Mathur et al. 2005). Various reviews in the Introduction deal with toxic components of smoke and their relation to ET-ROS-OS.

### Pesticides and Herbicide

An association between retinal degeneration and fungicide use was observed among farmer applicators of pesticides (Kirrane et al. 2005). A recent review

addresses ocular toxicity from pesticide application (Jaga and Dharmani, 2006). Exposure is related to retinopathy, neuropathy and abnormal ocular movement. Ocular surface toxicity resulted from paraquat accidentally introduced into the eyes of a fruit farmer (McKeag et al. 2002). The mechanism of these two classes in relation to the ET-ROS-OS approach is described in organ toxicity reviews presented in the Introduction.

### Metals

Members of this class display various toxicities as presented in the references on toxins in the Introduction. Heavy metal compounds usually possess reduction potentials quite amenable to ET in vivo. Toxicity is characterized by generation of ROS, lipid peroxidation, DNA cleavage and decrease in AO concentration. The negative effects are alleviated by AOs of various types.

Some investigations on adverse effects of metals on the eye are presented herein. Exposure to soluble silver compounds may produce toxic effects on the eye (Drake and Hazelwood, 2005). Studies involving the effect of Cu, Zn and Pb on crab eyes revealed eye pigmentary abnormalities, particularly those involving the retina (Lavolpe et al. 2004). The relationship between concentrations of Cu and Fe in the aqueous humour and intraocular pressure may help explain the role of these metals in the pathogenesis of glaucoma (Iqbal et al. 2002). Potential exposure of the eye to lead is a matter of concern resulting from use of certain cosmetics throughout the world (Nnorom et al. 2005). Their continuous use could produce an increase in the trace metal levels in the ocular system with accompanying toxic consequences. The levels of toxic metals (As, Cd and Pb) were significantly higher in scalp hair samples of children exhibiting ocular problems (Kazi et al. 2006). Administration of higher concentrations of zinc to the insulted retina exacerbates the condition and also acts as a toxin (Ugarte and Osborne, 2001). Zn supplements in the diet must be taken with caution. The effect of oral zinc on macular degeneration has been addressed (Newsome et al. 1988).

## ANTIOXIDANTS AND EYE TOXICITY

A large amount of research, partly addressed in prior sections, has been carried out on the protective effect of AOs on eye injury, which lends support to involvement of ROS toxins in OS. Lutein and zeaxanthin, major components of macular pigment, are part of the retinal AOs (Chucair et al. 2007). In human retinal endothelial cells pretreated with hydrogen peroxide, clusterin protected against apoptotic cell death (Kim et al. 2007). A randomized clinical trial revealed a significantly lower incidence of age-related maculopathy in a cohort of patients treated with high doses of

AOs than in a placebo group (Algvare et al. 2006). Fruits and vegetables, a rich source of carotenoids and other AOs, are thought to provide health benefits by decreasing the risk of various diseases, including those of the eye (Krinsky, 2005, Guggenbühl, 2006). A randomized trial revealed a beneficial effect of AOs (beta-carotene, vitamin C and vitamin E) on slowing progression of age-related macular degradation (Evans and Henshaw, 2008). Supplementation by omega-3 polyunsaturated fatty acids is proposed for certain subjects at risk for age-related macular degeneration, and supplementation with an AO cocktail is recommended in other cases (Desmettre et al. 2004). Melanin can act as a photoprotector by quenching ROS (Dayhaw-Barker, 2002). In animal studies, AOs were found to inhibit retinopathy in diabetes (Kowluru and Kennedy, 2001). Increasing the diversity of AOs provides significantly more protection than using any individual AO. In an investigation of optical neuropathy, a decrease in GSH occurred with changes in thiol redox potential (Schoeler et al. 2006). Corneal damage occurs following exposure to ultrasound energy due to radical formation (Nemet et al. 2007). Addition of AOs, such as ascorbic acid, exerted a protective effect. A combination of high glucose and absence of SOD increases formation of cataracts (Olofsson et al. 2007). Apparently, NO contributes to the process. L-Carnitine may protect against damage by gamma radiation by scavenging radicals and increasing SOD activity (Kocer et al. 2007). A similar protective effect was observed in the case of peroxide damage (Shamsi et al. 2007). 2-Cys peroxiredoxins are AO enzymes that eliminate hydrogen peroxide in the ciliary body of the eye, thus exerting a protective influence (Hong et al. 2007). Epigallocatechin gallate, an AO in green tea, attenuates damage to retina caused by ischemia/reperfusion and by hydrogen peroxide (Zhang et al. 2007).

Melatonin, endogenously produced in the eye, may act as a protective agent in ocular conditions, such as photokeratitis, cataract, glaucoma, retinopathy and ischemia/reperfusion injury (Siu et al. 2006). However, production of the AO is minimal in newborns, with gradual waning with aging. Supplementation may be beneficial. Influence on cataract formation and lipid peroxidation is in keeping with free-radical scavenging properties (Li et al. 1997). Results indicate that the AO may protect against UVB-induced cataract development by directly quenching lipid peroxidation (Bardak et al. 2000). Data indicate that melatonin has a promising role in the treatment of glaucoma (Lundmark et al. 2007). There are other studies relevant to the AO properties (Abe et al. 1994; Yağci et al. 2006)

Pathophysiological mechanisms involved in nutritional toxic neuropathies entail correction of OS and quenching of free radicals (Orssaud et al. 2007). An article deals with lipid peroxidation and AO defenses

in eye diseases (Shaimova, 2002). A 2006 report addresses molecular mechanisms of neuroprotection in the eye (Barnstable, 2006). A general discussion of AO mechanisms is available in a book chapter (Kovacic and Somanathan, 2006c).

Additional, relevant reports on AOs can be found in the following areas: Graves ophthalmopathy (Bouzas et al. 2000), ocular surface of diabetics (Peponis et al. 2002), association of ocular disorders with mortality (Clemons et al. 2004), ageing macula (Gerster, 1991), diabetic retinopathy (Lin et al. 2006), photoprotectants (Stahl and Sies, 2005), radiation-induced cataracts (Karslioğlu et al. 2005), diabetes-induced cataracts (Osakabe et al. 2004; Hegde and Varma, 2004; Kyselova et al. 2005; Özmen et al. 2000), cataract and macular degeneration (Moeller et al. 2000), senile cataract (Miratashi et al. 2005), Behçet's disease (Taysi et al. 2002), eye tear fluid (Choy et al. 2000), lipid membrane oxidation (Bhosale and Bernstein, 2005), ocular diseases (Boonefoy et al. 2002), retinal albumin leakage (Rota et al. 2004), retinal neurons (Lee et al. 2001), ocular melanogenesis (Sarangarajan and Apte, 2004), retinal pigment epithelial cells (Lu et al. 2002), retinal cells (Areias et al. 2001), retinal neurons (Lee et al. 2001), ocular disease and visual dysfunction (Whatham et al. 2008) and cornea and conjunctiva (Demir et al. 2005).

## CELL SIGNALING

A study demonstrates that pro-inflammatory cytokines increase ROS through mitochondrial and NADH oxidase in cultured human retinal pigment epithelial cells (Yang et al. 2007). Because of the pathogenic role of cytokines, antagonistic cytokines have been tested with positive results in Grave's ophthalmopathy (Maccocci, 2004). Antioxidant therapy might be envisioned since *in vitro* studies have shown possible involvement of ROS (Maccocci, 2004, Bartalena, 2003). There is a report on upregulation of the thioredoxin system via the Nrf2-AO responsive element pathway in adaptive-retinal neuroprotection (Tanito et al. 2007). In a study of retinoid signaling, results revealed that retinoic acid inhibits Ca elevation and over activation of calpains, suggesting the potential of calpain-targeting therapies for cataract (Nishikiori, 2007). The NF-kappa  $\square$  signal pathway may be important in hydrogen peroxide-induced damage in human lens epithelial cells that is involved in cataractogenesis (Jin et al. 2007). Data demonstrate selective induction of activated protein kinases (MAPK) in human lens epithelial cells by UV radiation (Bomser, 2002). These kinases modulate many cellular events, such as cell growth, death, differentiation and migration. Three major MAPK cascades have been identified. Prior work demonstrates the importance of kinase in cataract development. Activation of nuclear transcription factor in the retina of diabetics is inhibited by AOs,



suggesting involvement of ROS (Kowluru et al. 2003).

Considerable research has been done on the basic mechanism of cell signaling based on radicals and electrons. More than 10 years ago, ROS attracted attention in relation to cell signaling. Since then, several books have addressed this aspect (Forman and Cadenas, 1997; Hancock, 2005), including RNS, e.g., NO. Evidence has accumulated that ROS, such as hydrogen peroxide, superoxide, and the hydroxyl radical, are important chemical mediators that regulate the transduction of signals by modulating protein activity via redox chemistry (Veriweij and Gringhuis, 2002). Authors have proposed that ROS have been conserved throughout evolution as universal second messengers (Schulze-Osthoff et. al. 1997). Nearly every step in signal transfer is sensitive to ROS, which can function as primary signals and as second messengers in the activation of transcription factors (Forman and Cadenas, 1997). The approach is elaborated in a recent review (Kovacic and Pozos, 2006). A 2004 review (Hormuzdi et al. 2004) summarizes the present status of electrophysiological effects and deserves special attention. After a burst of research dealing with electrical coupling, gap junctions became less popular among neurobiologists versus the ionic approach. Recent reports have brought gap junctions back into the spotlight, suggesting that this type of cell-cell signaling may be interrelated with, rather than an alternative to, chemical transmission. We believe the thesis is credible, since electromagnetic effects of electrons and radicals in motion should have an influence on positive and negative charges associated with the central nervous system (CNS).

Other studies have addressed the electrical framework (Kovacic and Pozos, 2006.). Results suggest that nanosecond-pulsed electric fields modulate cell signaling from the plasma membrane to intracellular structures and function (Beebe et al. 2003). This technology could provide a powerful, unique tool to recruit signaling mechanisms that can eliminate aberrant cells by apoptosis. A 1993 symposium (Frey, 1993) focused on bioelectricity of cell signaling. Living creatures can be regarded as complex electrochemical systems that evolved over billions of years. Organisms interacted with and adapted to an environment of electrical and magnetic fields. Humans are now immersed in a man-made environment of such fields whose long-term effects are unknown.

## CELL MEMBRANES

There is extensive literature on the role of lipid peroxidation in toxicity. Cell membranes play an important part since lipids are essential constituents. Unsaturated lipid hydroperoxides comprise the principal actors that supply ROS. A toxic product from lipid degradation by ROS is 4-hydroxy-2-nonenal which has been the object of much research. Herein, recent examples

are provided of application to eye membranes.

A study by Marin-Constaño et al. (2005) showed that oxidant injury to human retinal pigment epithelial cells causes cell membrane blebbing. Apparently, oxidative attack disrupts the cell-specific surface proteases. Data by Yamada et al. (2008) support the hypothesis that oxidized lipoproteins comprise a trigger for initiating early events in the pathogenesis of age-related macular degeneration. The accumulation of certain lipoproteins in Bruch membrane evidently is involved. Oxidized LDL induced alterations in genes related to lipid metabolism, OS, inflammation and apoptosis. Florence (1995) noted that ROS cause lipid peroxidation in cell membranes and inactivate membrane-bound enzymes. These radicals that play a role in various diseases can be countered by use of AOs. In a recent report on oxygen toxicity in premature infants, Weinberger et al. (2002) invoked tissue injury by oxygen through formation of ROS which leads to peroxidation of membrane lipids. Protective measures include limiting oxygen and light exposure, as well as use of AOs and antiinflammatory agents. Evans et al. (1998) claim that linoleic hydroperoxide, an important source of biomembrane damage, is implicated in the onset of a variety of illnesses. A detoxification role for GSH peroxidase was indicated. Glutamate exposure, carried out by Mawatari et al. (1996), increased membrane oxidation two-fold, as judged by lipid peroxidation. ROS accumulate in response to OS after GSH depletion resulting in glioma cell death by Glu. Debbasch et al. (2001) discovered a significant decrease of membrane intergrety after exposure to quaternary ammonium salts. An apoptotic mechanism appears to be present. In this case, superoxide may play an important role in the tissue damage induced in the ocular surface disorders. In an investigation of action mechanism during OS-induced cataract, Spector (1995) observed damage to DNA and membrane pumps systems. Extensive oxidation of lens protein and lipid is associated with human cataract in older individuals and with the presence of elevated levels of hydrogen peroxide. Carmody et al. (1999) noted the ability of AOs to ameliorate light-induced retinal degeneration, suggesting a role for OS in cell death. Increase in ROS is accompanied by depletion in GSH in a model of photoreceptor apoptosis. These changes in cellular redox state precede disruption of mitochondrial transmembrane potential and DNA nicking that are events in apoptotic cell death. The ability of AOs to inhibit apoptosis through the scavenging of ROS establishes a role for ROS in photoreceptor apoptosis. In a report on uveitis, Goto et al. (1992) detected lipid peroxidation initiated by phagocyte-derived ROS. The presence of inflammation-mediated lipid peroxidation was confirmed by the generation of TBARS. Varma et al. (1984) examined the effect of OS on the lens and on cataract formation. Light induced lipid peroxidative degradation of tissue lipids.

The adverse effect was attenuated by scavengers of ROS. Lipid degradation was prevented by vitamins C and E. The results suggest that photodynamic injury to the lens and membrane lipids is incumbent upon initial generation of superoxide, followed by conversion to other ROS. Albender et al. (2007), in a study involving synaptosomal membranes, reported protection by melatonin against Al-induced oxidative damage, indicating a neuroprotective effect based on AO

activity. In a related study (Karbownik et al. 2000), melatonin reduced the toxic effects of phenylhydrazine and iron in cell membranes, apparently due to beneficial AO action.

### Acknowledgement

Editorial assistance by Thelma Chavez is acknowledged.

### References

- Abdollahi M, Shafiee A, Bathaiee F. S, Sharifzadeh M, Nikfar S. 2004. Drug-induced toxic reactions in the eye: an overview. *J Infus Nurs* 27: 386-398.
- Abe M, Reiter RJ, Orhii PB, Hara M, Poeggeler B. 1994. Inhibitory effect of melatonin on cataract formation in newborn rats: evidence for an antioxidative role for melatonin. *J Pineal Res* 17: 94-100.
- Albendea CD, Gómez-Trullén EM, Fuentes-Broto L, Miana-Mena FJ, Millán-Plano S, Reyes-Gonzales MC, Martínez-Ballarín E, García JJ. 2007. Melatonin reduces lipid and protein oxidative damage in synaptosomes due to aluminium. *J Trace Element Med Biol* 21: 261-268.
- Algvere P. V, Marshall J, Seregard S, 2006. Age-related maculopathy and the impact of blue light hazard. *Acta Ophthalmol Scand* 84: 4-15.
- Altinors D. D, Akçaya S, Akova Y. A, Bilezikçi B, Goto E, Dogru M, Tsubota K. 2006. Smoking associated with damage to the lipid layer of the ocular surface. *Am J Ophthalmol* 141: 1016-1021.
- al-Tweigeri T, Nabholz J. M, Mackey J. R, 1996. Ocular toxicity and cancer chemotherapy. A review. *Cancer* 78: 1359-1373.
- Anwar M. M, Moustafa M. A. 2001. The effect of melatonin on eye lens of rats exposed to ultraviolet radiation. *Compar Biochem Physiol Part C* 129: 57-63.
- Areias F. M, Rego A. C, Oliveira C. R, Seabra R. M, 2001. Antioxidant effect of flavonoids after ascorbate/Fe<sup>2+</sup>-induced oxidative stress in cultured retinal cells. *Biochem Pharmacol* 62: 111-118.
- Aslan M, Yücel I, Akar Y, Yücel G, Ciftçioğlu MA, Sanlioglu S. 2006. Nitrotyrosine formation and apoptosis in rat models of ocular injury. *Free Rad Res* 40: 147-153.
- Aslan M, Yucel I, Ciftçioğlu A, Savaş B, Akar Y, Yucel G, Sanlioglu S. Corneal protein nitration in experimental uveitis. 2007. *Exp Biol Med* 232: 1308-1313.
- Ayala M. N, Michael R, Söderberg P. G. 2000. In vivo cataract after repeated exposure to ultraviolet radiation. *Exp Eye Res* 70: 451-456.
- Babizhayev M. A, Deyev A. I, Yermakova V. N, Brikman I. V, Bours J. 2004. Lipid peroxidation and cataracts: N-acetylcarnosine as therapeutic tool to manage age-related cataracts in human and in canine eyes. *Drugs in R & D* 5: 125-139.
- Bardak Y, Ozertürk Y, Ozgüner F, Durmuş M, Dekibaş N. 2000. Effect of melatonin against oxidative stress in ultraviolet-B exposed rat lens. *Curr Eye Res* 20: 225-230.
- Barnstable C. J, Tombran-Tink J, 2006. Molecular mechanisms of neuroprotection in the eye. *Adv Exp Med Biol* 572: 291-295.
- Bartalena L. 2003. Oxidative stress and Graves' ophthalmopathy: in vitro studies and therapeutic implications. *Biofactors* 19: 155-163.
- Barza M, Baum J, Szoka F, D'Amico D. J. 1985. Ocular toxicity of intravitreally injected liposomal amphotericin B in rhesus monkeys. *Am J Ophthalmol* 100: 259-263.
- Beatty S, Koh, H-H, Henson, D, Boulton M. 2000. The role of oxidative stress in the pathogenesis of age-related macular degeneration. *Surv Ophthalmol* 45: 115-134.
- Becquet F, Courtois Y, Goureau O, 1997. Nitric oxide in the eye: multifaceted roles and diverse outcomes. In: Cotlier, E, Weinreb, R, eds., *Current Research, Survey Ophthalmol* 42: 71-82.
- Beebe S. J, Fox P. M, Rec L. J. 2003. Nanosecond, high-intensity pulsed electric fields induce apoptosis in living cells. *FASEB J* 17: 1493-1495.
- Bhosale P, Bernstein P. S. 2005. Synergistic effects of zeaxanthin and its binding protein in the prevention of lipid membrane oxidation. *Biochem Biophys Acta* 1740: 116-121.
- Blake D. R, Winyard P, Lunec J, Williams A, Good P. A, Crewes S. J, Gutteridge J. M. C, Rowley D, Halliwell B, Cornish A, Hider R. C. 1985. Cerebral and ocular toxicity induced by desferrioxamine. *Quart J Med New Series*: 56, 345-355.
- Boet D. J. 1970. Toxic effects of phenothiazines on the eye. *Doc Ophthalmol* 28: 1-69.
- Bomser J. A. 2002. Selective induction of mitogen-activated protein kinases in human lens epithelial cells by ultraviolet radiation. *J Biochem Mol Toxicol* 16: 33-40.
- Boonefey M, Draai J, Kostka T. 2002, Antioxidants to slow ageing, facts and perspectives. *Presse Med* 31: 1174-1184.
- Bouzas E. A, Karadimas P, Mastorakos G, Koutras D. A. 2000. Antioxidant agents in the treatment of Grave's ophthalmopathy. *Am J Ophthalmol* 129: 618-622.
- Boyland E, Chasseaud L. F. 1967. Enzyme-catalyzed conjugations of glutathione with unsaturated compounds. *Biochem J* 104: 95-102.
- Carmody RJ, McGowan AJ, Cotter TG. 1999. Reactive oxygen species as mediators of photoreceptor apoptosis in vitro. *Exp Cell Res* 248: 520-530.
- Cheng A. C, Pang C. P, Leung A. T, Chua J. K, Fan D. S, Lam D. S. 2000. The association between cigarette smoking and ocular diseases. *Hong Kong Med J* 6: 195-200.
- Chiou, G. C. Y. Ed. 1999, *Ophthalmic Toxicology*, 2nd. Edition, Taylor & Francis, Philadelphia.
- Choy C. K. M, Benzie I. F. F, Cho P. 2000. Ascorbic acid concentration and total antioxidant activity of human tear fluid measured using the FRASC assay. *Invest Ophthalmol Vis Sci* 41: 3293-3298.
- Chucair A. J, Rotstein, N. P, Sangiovanni J. P, Doring A, Chew E. Y, Politi L. E. 2007. Lutein and zeaxanthin protect photoreceptors from apoptosis induced by oxidative stress. *Invest Ophthalmol Vis Sci* 48: 5168-5177.
- Clemons T. E, Kurinij N, Sperduto D, Bressler S. B. 2004. Associations of mortality with ocular disorders and an intervention of high-dose antioxidants and zinc in the age-related eye disease

- study – AREDS report no.13. *Arch Ophthalmol* 122: 716-726.
- Cole J. G, Cole H. G, Janoff L. A. 1957. A toxic ocular manifestation of chloramphenicol therapy; report of a case of optic neuritis. *Am J Ophthalmol* 44: 18-20.
- Davies M. J. 2001. Photo-oxidation of proteins and its role in cataractogenesis. *J Photochem Photobiol* 63: 114-125.
- Dayhaw-Barker, P. 2002. Retinal pigment epithelium melanin and ocular toxicity. *Int J Toxicol* 21: 451-454.
- Debbasch C, Brignole F, Pisella P-J, wernet J-M, rat P, Baudouin C. 2001. Quaternary ammonium and other preservative's contribution in oxidative stress and apoptosis on Chang conjunctive cells. *Invest Ophthalmol Vis Sci* 42: 642-652.
- de Gruijl F. R, Longstreth J, Norval M, Cullen A. P, Slaper H, Kripke M. L, Takizawa Y, van der Leun J. C, 2003. Health effects from stratospheric ozone depletion and interactions with climate change. *Photochem Photobiol Sci* 2: 16-28.
- Demir U, Demir T, Ilhan N. 2005. The protective effect of alpha-lipoic acid against oxidative damage in rabbit conjunctiva and cornea exposed to ultraviolet radiation. *Ophthalmologica* 219: 49-53.
- Denis U, Lecomte M, Paget C, Ruggiero D, Wiernsperger N, Lagarde M. 2002. Advanced glycation end-products induce apoptosis of bovine retinal pericytes in culture: involvement of diacylglycerol/ceramide production and oxidative stress induction. *Free Rad Biol Med* 33: 236-247.
- Desmettre T, Lecerf J. M, Souied E. H, 2004. Nutrition and age-related macular degeneration. *J Fr Ophthalmol* 27: 3538-56.
- Doma F. M, Koer A, Tanidir R. 2007. Optic neuropathy related to hydrogen peroxide inhalation. *Clin Neuropharmacol* 30: 55-57.
- Doty R. L, Cometto-Muoz J. E, Jalowayski A. A, Dalton P, Kendel-Reed M, Hodgson, M. 2004. Assessment of upper respiratory tract and ocular irritative effects of volatile chemicals in humans. *Crit Rev Toxicol* 34: 85-142.
- Drake P, Hazelwood K. J. 2005. Exposure-related health effects of silver and silver compounds: a review. *Ann Occup Hyg* 49: 575-585.
- Elovaara E, Kivistö H, Vainio H. 1983. Effects of methyl methacrylate on non-protein thiols and drug metabolizing enzymes in rat liver and kidneys. *Arch Toxicol* 52: 109-121.
- Evans J. R, Henshaw K. 2008. Antioxidant vitamin and mineral supplements for slowing the progression of age-related macular degeneration. *Cochrane Database Syts Rev* (2). *Cochrane AN*: CD000254.
- Evans MV, Turton HE, Grant CM, Dawes IW. 1998. Toxicity of linoleic acid hydroperoxide to *Saccharomyces cerevisiae*; involvement of a respiration-related process for maximal sensitivity and adaptive response. *J Bacteriol* 483-490.
- Florence TM. 1995. The role of free radicals in disease. *Aust NZ J Ophthalmol* 23: 3-7.
- Forman H.G, Cadenas, E. Eds., 1997. *Oxidative Stress and Signal Transduction*. New York; Chapman and Hall. 475 pp.
- Fraunfelder F. W, 2006. Corneal toxicity from topical ocular and systemic medications. *Cornea* 25, 1133-1138.
- Fraunfelder F. W. 2003. Ocular adverse drug relations. *Expert Opin Drug Saf* 2: 411-420.
- Fraunfelder F. W, Fraunfelder F. T. 2004. Adverse ocular drug reactions recently identified by the national registry of drug-induced ocular side effects. *Ophthalmology* 111: 1275-1279.
- Frey A.H. 1993. Electromagnetic field interaction with biological systems. *FASEB J* 7: 272-281.
- Fu Y-C, Jin X-P, Wei S-M, Lin H-F, Kacew S. 2000. Ultraviolet radiation and reactive oxygen generation as inducers of keratinocyte apoptosis: protective role of tea polyphenols. *J Toxicol Environ Health Part A* 61: 177-188.
- Fujishiro K, Mori, K, Inoue N. 1991. Effects of inhaled ethylene oxide on the lens glutathione redox cycle in rats. *Arch Toxicol* 65: 606-607.
- Gallagher R. P, Lee T. K. 2006. Adverse effects of ultraviolet radiation: a brief review. *Progr Biophys Mol Biol* 92: 119-131.
- Garg TK, Chang JY. 2004. 15-deoxy-delta 12, 14-prostaglandin J2 prevents reactive oxygen species generation and mitochondrial membrane depolarization induced by oxidative stress. *BMC Pharmacol* 4: 6.
- Gerster H.1991. Review: antioxidant protection of the ageing macula. *Age Ageing* 20: 60-69.
- Gobba F. 2003. Occupational exposure to chemicals and sensory organs: a neglected research field. *Neuro Toxicol* 24: 675-691.
- Gong Y, Kishi R, Kasai S, Katakura Y, Fujiwara K, Umemura T, Kondo T, Sato T, Sata, F, Tsukishima E, Tozaki S, Kawai T, Miyama Y. 2003. Visual dysfunction in workers exposed to a mixture of organic solvents. *Neuro Toxicol* 24: 703-710.
- Gong Y. Y, Kishi R, Katakura Y, Tsukishima E, Fujiwara K, Kasai S, Satoh T, Sata F, Kawai T. 2002. Relation between color vision loss and occupational styrene exposure level. *Occup Environ Med* 59: 824-829.
- Goto H, Wu G-S, Chen F, Kristeva M, Sevanian A, Rao NA. 1992. Lipid peroxidation in experimental uveitis: sequential studies. *Curr Eye Res* 11: 489-499.
- Guggenbühl N. 2006. The source of antioxidants. *Bull Soc Belge Ophthalmol* 301: 41-45.
- Halliwell B, Gutteridge J. M. C. 1999. *Free Radicals in Biology and Medicine*. Oxford, New York, (a) pp. 1-859, (b) 548-553, (c) 643.
- Hancock G. T. 2005. *Cell Signaling*. Oxford University Press, New York. 296 pp.
- Hardy P, Beauchamp M, Sennlaub F, Gobeil F Jr, Tremblay L, Mwaikambo B, Lachapelle P, Chemtob S. 2005. New insights into the retinal circulation: inflammatory lipid mediators in ischemic retinopathy. *Prostaglandins Leukot Essent Fatty Acids* 72: 301-325.
- Haufroid Y, Collignon-Robe N. J. 2004. Ocular vasospastic syndrome. *Rev Med Liege* 59 : 227-231.
- He X, Hahn P, Iacovelli J, Wong R, King C, Bhisitkul R, Massaro-Giordano M, Dunaief J. L. 2007. Iron homeostasis and toxicity in retinal degeneration. *Prog Retin Eye Res* 26: 649-673.
- Hegde K. R, Varma S. D. 2004. Morphogenetic and apoptic changes in diabetic cataract: prevention by pyruvate. *Mol Cell Biochem* 262: 233-237.
- Holyk P. R, Eifrig D. E. 1979. Effects on monomeric methylmethacrylate on ocular tissues. *Am J Ophthalmol* 88: 385-395.
- Hong S, Kim C. Y, Lee J. H, Seong G. J. 2007. Immunohistochemical localization of 2-Cys peroxiredoxins in human ciliary body. *Tissue Cell* 39 : 365-368.
- Hormuzdi S. G, Filippov M. A, Mitropoulou G, Monyer H, Bruzzone R. 2004. Electrical synapses: a dynamic signaling system that shapes the activity of neuronal networks. *Biochem Biophys Acta* 1662: 113-137.
- Iqbal Z, Mohammad Z, Shah M. T, Bashir S, Khan T, Khan, M. D. 2002. Relationship between the concentration of copper and iron in the aqueous humour and intraocular pressure in rabbits treated with topical steroids. *Clin Exp Ophthalmol* 30: 28-35.
- Izzotti A, Bagnis A, Saccá S. C. 2006a. The role of oxidative stress in glaucoma. *Mutat Res* 612: 105-114.
- Izzotti A, Di Marco B, De Flora S, Saccá S, 2006b. Open angle

- glaucoma: epidemiology, pathogenesis and prevention. *Recent Prog Med* 97: 37-45.
- Jain M. R, Hussain S. 1982. Bioestimation of carbenicillin and gentamycin after their intravitreal injection and evaluation of their toxic effect on ocular tissues. *Indian J Ophthalmol* 30: 195-200.
- Jacinto J. D, Kovacic P. 2003. Neurotransmission and neurotoxicity by nitric oxide, catecholamines and glutamate: unifying themes of reactive oxygen species and electron transfer. *Curr Med Chem* 10: 2693-2703.
- Jaga K, Dharmani C. 2006. Ocular toxicity from pesticide exposure: a recent review. *Environ Health Preven Med* 11: 102-107.
- Jankauskienė J, Imbrasienė D, Stanislovaitytė D, Jarušaitienė D, Narkevičiūtė R. 2007. Influence of environmental pollution on the status of patients with endocrine ophthalmopathy in Kaunas. *Ekologia* 53: 10-14.
- Javadi M-A, Yazdani S, Sajjadi H, Jadidi K, Karimian F, Einollahi B, Ja'farinasab M-R, Zare M., 2005. Chronic and delayed-onset mustard gas keratitis. *Ophthalmology* 112: 617-625.
- Jin X. H, Ohgami K, Shiratori K, Koyana Y, Yoshida K, Kase S, Ohno S. 2007. Inhibition of nuclear factor-kappa B activation attenuates hydrogen peroxide-induced cytotoxicity in human lens epithelial cells. *Br J Ophthalmol* 91: 369-371.
- Karbownik M, Reiter RJ, Garcia JJ, Tan D. 2000. Melatonin reduces phenylhydrazine-induced oxidative damage to cellular membranes: evidence for the involvement of iron. *Int J Biochem Cell Biol* 32: 1045-1054.
- Karslioglu I, Ertekin M. V, Taysi S, Kocer I, Sezen O, Gepdiremen A, Kocer M, Bakan N. 2005. Radioprotective effects of melatonin on radiation-induced cataract. *J Radiat Res* 46: 277-282.
- Kazi T, Afridi H. I, Kazi G. H, Jamali M. K, Arain M. B, Jalbani N. 2006. Evaluation of essential and toxic metals by ultrasound-assisted acid leaching from scalp hair samples of children with macular degeneration patients. *Clin Chim Acta* 369: 52-60.
- Kilic D. 2003. The effects of ageing and sulfur dioxide inhalation exposure on visual-evoked potentials, antioxidant enzyme systems, and lipid-peroxidation levels of the brain. *Neurotoxicol Teratol* 25: 587-598.
- Kim J. H, Kim J. H, Yu Y. S, Min B. H, Kim K. W. 2007. The role of clusterin in retinal development and free radical damage. *Br J Ophthalmol* 91: 1541-1546.
- Kirrane E. F, Hoppin J. A, Kamel F, Umbach D. M, Boyes W. K, Deroos A. J, Alavanja M, Sandler D. P. 2005. Retinal degeneration and other eye disorders in wives of farmer pesticide applicators enrolled in the agricultural health study. *Am J Epidemiol* 161: 1020-1029.
- Kocer I, Tayasi S, Ertekin M. V, Karslioglu I, Gepdiremen A, Sezen O, Serifoglu K. 2007. The effect of L-carnitine in the prevention of ionizing radiation-induced cataracts: a rat model. *Graefes Arch Clin Exp Ophthalmol* 245: 588-594.
- Kovacic P. 2006. Novel electrochemical approach to enhanced toxicity of 4-oxo-2-nonenal vs. 4-hydroxy-2-nonenal (role of imine): oxidative stress and therapeutic modalities. *Med Hypotheses* 67: 151-156.
- Kovacic P, Becvar L.E. 2000. Mode of action of anti-infective agents: emphasis on oxidative stress and electron transfer. *Curr Pharmaceut Des* 6: 143-167.
- Kovacic P, Cooks A.L. 2005. Unifying mechanism for toxicity and addiction by abused drugs: electron transfer and reactive oxygen species. *Med Hypotheses* 64: 367-366.
- Kovacic P, Jacinto J.D. 2001a. Mechanisms of carcinogenesis: focus on oxidative stress and electron transfer. *Curr Med Chem* 8: 773-796.
- Kovacic P, Jacinto J.D. 2001b. Reproductive toxins: pervasive theme of oxidative stress and electron transfer. *Curr Med Chem* 8: 863-892.
- Kovacic P, Osuna J.A. 2000. Mechanisms of anticancer agents: emphasis on oxidative stress and electron transfer. *Curr Pharmaceut Des* 6: 277-309.
- Kovacic P, Pozos R. S. 2006. Cell signaling (mechanism and reproductive toxicity): redox chains, radicals, electrons, relays, conduit, electrochemistry, and other medical implications. *Birth Defects Res Part C* 78: 333-344.
- Kovacic P, Pozos R.S, Somanathan R, Shangari R, O'Brien P.J. 2005. Mechanism of mitochondrial uncouplers, inhibitors, and toxins: focus on electron transfer, free radicals, and structure-activity relationships. *Curr Med Chem* 5: 2601-2623.
- Kovacic P, Sacman A, Wu-Weis M. 2002. Nephrotoxins: Widespread role of oxidative stress and electron transfer. *Curr Med Chem* 9: 823-847.
- Kovacic P, Somanathan R. 2008. Ototoxicity and noise trauma; electron transfer, reactive oxygen species, cell signaling, electrical effects, and protection by antioxidants; practical medical aspects. *Med Hypotheses* 70: 914-923.
- Kovacic, P, Somanathan, R. 2006a. Alcohol mechanisms: cell signaling, toxicity, addiction, prevention, therapy and beneficial effects, in: Brozner, E. Y.(ed.), *New Research on Alcohol Abuse and Alcoholism*, Nova Science Publishers, New York. pp. 40-102.
- Kovacic, P, Somanathan R. 2006b. Mechanism of teratogenesis: electron transfer, reactive oxygen species and antioxidants. *Birth Defects Res Part C* 78: 308-325.
- Kovacic P, Somanathan R. 2005. Neurotoxicity: The broad framework of electron transfer, oxidative stress and protection by antioxidants. *Curr Med Chem-CNS Agents* 5: 249-258.
- Kovacic P, Thurn L.A. 2005. Cardiovascular toxins from the perspective of oxidative stress and electron transfer. *Curr Vasc Pharmacol* 3: 107-117.
- Kovacic P, Somanathan R (2006c) Beneficial effects of antioxidants in relation to carcinogens, toxins and various illnesses. In: Panglossi, H. V, editor. *Frontiers in Antioxidants Research*, Nova Science Publishers, Hauppauge, NY. pp. 1-38.
- Kovacic P. 2006. Novel electrochemical approach to enhanced toxicity of 4-oxo-2-nonenal vs. 4-hydroxy-2-nonenal (role of imine): oxidative stress and therapeutic modalities. *Med Hypotheses* 67: 151-156.
- Kowluru R, Kennedy A. 2001. Therapeutic potential of antioxidants and diabetic retinopathy. *Expert Opin Investig Drugs* 10: 1665-1675.
- Kowluru R. A, Koppolu P, Chakrabarti S, Chen S. 2003. Diabetes-induced activation of nuclear transcriptional factor in the retina, and its inhibition by antioxidants. *Free Rad Res* 37, 1169-1180.
- Kravchuk E. A. 2004. Free-radical oxidation in the pathogenesis of eye diseases. *Vestn Oftalmol* 120: 48-51.
- Krinsky N. I. 2005. Carotenoid action and their relation to health and disease. *Mol. Aspects Med* 26: 459-516.
- Kyselova Z, Gajdosik A, Gajdosikova A, Ulicna O, Mihalova, D, Karasu C, Stefek M. 2005. Effect of the pyridindole antioxidant stobadine on development of experimental diabetic cataract and on lens protein oxidation in rats: comparison with vitamin E and BHT. *Mol Vis* 11: 56-65.
- Lassen N, Bateman J. B, Estey T, Kuszak J. R, Nees D. W, Pitigorsky J, Duester G, Day B. J., Huang J, Hines L. M, Vasilioiu V. 2007. Multiple and additive functions of ALDH3A1 and ALDH1A1 cataract phenotype and ocular oxidative damage in Aldh1a1 (-1-)/Aldh1a1 (-1-) knock-out mice. *J Biol Chem* 282:



25668-25676.

- Lavolpe M, Greco L. L, Kesselman D, Rodriguez E. 2004. Differential toxicity of copper, zinc and lead during the embryonic development of *Chasmagnathus Granulatus* (Brachyura, Varunidae). *Environ Toxicol Chem* 23: 960-967.
- Laycock N. L. C, Schirmer K, Bols N. C, Sivak J. G. 2000. Optical properties of rainbow trout lenses after in vitro exposure to polycyclic aromatic hydrocarbons in the presence or absence of ultraviolet radiation. *Exp Eye Res* 70: 205-214.
- Lee M-C, Chung Y-T, Lee J-H, Jung J-J, Kim H-S, Kim S-U. 2001. Antioxidant effect of melatonin in human retinal neuron cultures. *Exp Neurol* 172: 407-415.
- Li J, Tripathi R. C, Tripathi B. J. 2008. Drug-induced ocular disorders. *Drug Saf* 31: 127-141.
- Li ZR, Reiter RJ, Fujimori O, Oh CS, Duan YP. 1997. Cataractogenesis and lipidoxidation in newborn rats treated with buthionine sulfoximine: preventive actions of melatonin. *J Pineal Res* 22: 117-123.
- Lin J, Bierhaus, A, Bugert P, Dietrich, N, Feng Y, vom Hagen F, Nawroth P, Brownlee M, Hammes H-P. 2006. Effect of R-(+)- $\alpha$ -lipoic acid on experimental diabetic retinopathy. *Diabetologia* 49: 1089-1096.
- Lin D, Lee H-G, Liu Q, Perry G, Smith MA, Sayre LM. 2005. 4-Oxo-2-nonenal is both more neurotoxic and more protein reactive than 4-hydroxy-2-nonenal. *Chem Res Toxicol* 18: 1219-1231.
- Lu H, Hunt D. M, Ganti R, Davis A, Dutt K, Alam J, Hunt R. C. 2002. Metallothionein protects retinal pigment epithelial cells against apoptosis and oxidative stress. *Exp Eye Res* 74: 83-92.
- Lundmark P. O, Pandi-Perumal S. R, Srinivasan V, Cardinali D. P. Rosenstein R. E. 2007. Melatonin in the eye: implications for glaucoma. *Exp Eye Res* 84: 1021-1030.
- Marcocci C. 2004. Novel aspects of immunosuppressive and radiotherapy management of Graves' ophthalmopathy. *J Endocrinol Invest* 27: 272-280.
- Marin-Castaño ME, Csaky KG, Cousins SW. 2005. Nonlethal oxidant injury to human retinal pigment epithelium cells causes cell membrane blebbing but decreased MMP-2 activity. *Invest Ophthalmol Vis Sci* 46: 3331-3340.
- Martynkina L. P, Qian W, Shichi H. 2002. Naphthoquinone cataract in mice: mitochondrial change and protection by superoxide dismutase. *J Ocular Pharmacol Therap* 18: 231-239.
- Mathur R, Gupta K. S, Joshi, S, Velpandian T. 2005. Cigarette smoke condensate reduces the detoxifying capabilities of rat lens. *Indian J Pharmacol* 37: 37-45.
- Marsili S, Salganik R. I, Albright C. D, Freely C. D, Johnsen S, Peiffer R. L, Costello M. J. 2004. Cataract formation in a strain of rats selected for high oxidative stress. *Exp Eye Res* 79: 595-612.
- Mawatari K, Yau Y, Sugitani K, Takadera T, Kato S. 1996. Reactive oxygen species involved in the glutamate toxicity of C6 glioma cells via XC antiporter system. *Neuroscience* 73: 201-208.
- McCaa C. S. 1982. The eye and visual nervous system: anatomy, physiology and toxicology. *Environ Health Perspect* 44: 1-8.
- McCarty C. A, Taylor H. R. 2002. A review of the epidemiologic evidence linking ultraviolet radiation and cataracts. In: Hockwin O, Kojima M, Takahashi N, Sliney D. H. Basel, Karger, eds., *Progress in Lens and Cataract Research Dev Ophthalmol* 35: 21-31.
- McGee H. T, Fraunfelder F. W. 2007. Toxicities of topical ophthalmic anesthetics. *Expert Opin Drug Saf* 6: 637-640.
- McKeag D, Maini R, Taylor H. R. 2002. The ocular surface toxicity of paraquat. *Br J Ophthalmol* 86: 350-351.
- Mclane N. J, Carroll D. M. 1986. Ocular manifestation of drug abuse. *Surv Ophthalmol* 30: 298-313.
- Meyer-Rochow V. B. 2001. The crustacean eye: dark/light adaptation, polarization sensitivity, flicker fusion frequency, and photoreceptor damage. *Zool Sci* 18: 1175-1197.
- Mileva M, Zlateva G. 2005. Comparing effects of low-intensity laser irradiation and vitamin E on oxidative damages in experimental model of diquat-induced cataract in rabbits. *Trakia J Sci* 3: 6-9.
- Miratashi A. M, Besharati M. R, Shoja M. R. 2005. Vitamin C concentration of aqueous humour in senile cataract. *J Res Med Sci* 10: 269-273.
- Moeller S. M, Jacques P. F, Blumberg J. B. 2000. The potential role of dietary xanthophylls in cataract and age-related macular degeneration. *J Am Coll Nutr* 19: 522S-527.
- Moorthy R. S, Valluri S. 1999. Ocular toxicity associated with systemic drug therapy. *Curr Opin Ophthalmol* 10: 438-446.
- Morad Y, Banin E, Averbukh E, Berenshtein E, Obolensky A, Chevion M. 2005. Treatment of ocular tissues exposed to nitrogen mustard: beneficial effects of zinc desferrioxamine combined with steroids. *Invest Ophthalmol Vis. Sci.* 46, 1640-1646.
- Moreno M. C, Campanelli J, Sande P, Sáenz D. A, Sarmiento M. I. K, Rosenstein R. E. 2004. Retinal oxidative stress induced by high intraocular pressure. *Free Rad Biol Med* 37: 803-812.
- Nemet A. Y, Assia E. I, Meyerstein D, Gedanken A, Topaz M. 2007. Protective effect of free-radical scavengers on corneal endothelial damage in phacoemulsification. *J Cataract Refract Surg* 33: 310-315.
- Nnorom I. C, Igwe J. C, Oji-Nnorom C. G. 2005. Trace metal contents of facial (make-up) cosmetics commonly used in Nigeria. *J Biotechnol* 4: 1133-1138.
- Newsome D. A, Swartz M, Leone N. C, Elston R. C, Miller E. 1988. Oral zinc in macular degeneration. *Arch Ophthalmol* 106: 192-198.
- Nishikiori N, Osanai M, Chiba H, Kojima T, Ohguro H, Sawada N. 2007. Inhibitory effects of retinoic acid receptor alpha stimulant on murine cataractogenesis through suppression of deregulated calpains. *Invest Ophthalmol Vis Sci* 48: 2224-2249.
- Nowak M, Gnitecki W, Jurowski P. 2005. The role of retinal oxygen metabolism in origin of age-related degeneration (AMD). *Klin Oczna* 107: 715-718.
- Ohia S. E, Opere C. A, Leday A. M. 2005. Pharmacological consequences of oxidative stress in ocular tissues. *Mutat Res* 579: 22-36.
- Okolie N. P, Asonye C. C, 2004. Mitigation of cataractogenic potential of cyanide by antioxidant vitamin administration. *J Biomed Res* 3: 48-52.
- Okolie N. P, Audu K. 2004. Correlation between cyanide-induced decreases in ocular Ca<sup>2+</sup>-ATPase and lenticular opacification. *J Biomed Sci* 3: 37-41.
- Olofsson E. M, Marklund S. L, Behndig A. 2007. Glucose-induced cataract in CuZn-SOD null lenses: an effect of nitric oxide? *Free Rad Biol Med* 42: 1098-1105.
- Orssaud C, Roche O, Dufier J. L. 2007. Nutritional optic neuropathies. *J Neurol Sci* 262: 158-164.
- Osakabe N, Yamagishi M, Natsume M, Yasuda A, Osawa T. 2004. Ingestion of proanthocyanidins derived from cacao inhibits diabetes-induced cataract formation in rats. *Exp Biol Med* 229: 33-39.
- Ottone S, Foroni C, Carta A, Petrucco S, Maraini G. 2000. Oxidative stress and age-related cataract. In *Aging and Age-Related Ocular Disease*. Ed., Lütjen-Drecoll, E.; Erlangen-Nürnberg. *Ophthalmologica* 214: 78-85.

- Ozdek S, Urgancioglu B, Turku U. O, Bilgihan A. 2007. A possible mechanism for transpupillary thermotherapy: nitric-oxide-related cellular damage. *Can J Ophthalmol* 42: 609-612.
- Özmen D, Özmen B, Erkin E, Güner I, Habif S, Bayindir. 2000. Lens superoxide dismutase and catalase activities in diabetic cataract. *Turk J Endocrinol Metabol* 1: 1-4.
- Pandya U, Saini M. K, Jin G. F, Awasthi S, Godley B. F, Awasthi Y. C. 2000. Dietary curcumin prevents ocular toxicity of naphthalene in rats. *Toxicol Lett* 115: 195-204.
- Peponis V, Papathanasiou M, Kapranou A, Magkou C, Tylogada A, Melidonis A, Drosos T, Sitaras N. M. 2002. Protective role of oral antioxidant supplementation in ocular surface of diabetic patients. *Br J Ophthalmol* 86: 1369-1373.
- Petersen A, Carlsson T, Karlsson J. O, Zetterberg M. 2007. The proteasome and intracellular redox status: implications for apoptotic regulation in lens epithelial cells. *Curr Eye Res* 32: 871-882.
- Pilon A. F, Scheiffle J. 2006. Ulcerative keratitis associated with crack-cocaine abuse. *Cont Lens Anterior Eye* 29: 263-267.
- Poli G, Cheeseman K.H, Dianzani M.U, Slater T.F. Ed.; 1989. *Free Radicals in the Pathogenesis of Liver Injury*, Pergamon, New York.
- Pradhan A. K, Shukla A. K, Reddy M. V. R, Garg, N. 2004. Assessment of oxidative stress and antioxidant status in age related cataract in a rural population. *Indian J Clin Biochem* 19: 83-87.
- Prickett C. D, Lister E, Collins M, Trevithick-Sutton C. C, Hirst M, Vinson J. A, Noble E, Trevithick J. R. 2004. Nonlinearity in biology, toxicology, and medicine. *J. Dose-Response* 2: 353-370.
- Raju P, George R, Ramesh S. V, Arvind H, Baskaran M, Vijaya L. 2006. Influence of tobacco use on cataract development. *Br J Ophthalmol* 90: 1374-1377.
- Risa O, Soether O, Löfgren S, Söderberg P. G, Krane J, Midelfart A. 2004. Metabolic changes in rat lens after in vivo exposure to ultraviolet irradiation; measurements by high resolution MAS 1H NMR spectroscopy. *Invest Ophthalmol Vis Sc.* 45: 1916-1921.
- Roberts J. E. 2001. Ocular phototoxicity. *J Photochem Photobiol B: Biol* 64: 136-143.
- Roberts J. E. 2002. Screening for ocular phototoxicity. *Int J Toxicol* 21: 491-500.
- Rootman J, Joseph P. D, Adomat H, Palcic B. 1982. Ocular absorption and toxicity of a radiosensitizer and its effect on hypoxic cells. *Arch Ophthalmol* 100: 468-471.
- Rota R, Chiavaroli C, Garay R. P, Hannaert P. 2004. Reduction of retinal albumin leakage by the antioxidant calcium dobesilate in streptozotocin-diabetic rats. *Eur J Pharmacol* 495: 217-224.
- Russell P, Yamada T, Xu G. T, Garland D, Zigler J. S. Jr. 1991. Effects of naphthalene metabolites on cultured cells from eye lens. *Free Rad Biol Med* 10: 255-261.
- Saccá S. C, Izzotti A, Rossi P, Traverso C. 2007. Glaucomatous outflow pathway and oxidative stress. *Exp Eye Res* 84: 389-399.
- Santaella R. M, Fraunfelder F. W. 2007. Ocular adverse effects associated with systemic medications: recognition and management. *Drugs* 67: 75-93.
- Santosa S, Jones P. J. H. 2005. Oxidative stress in ocular disease: does lutein play a protective role? *CMAJ* 173: 861-862.
- Sarangarajan R, Apte S. P. 2004. Ocular melanogenesis: the role of antioxidants. *Ophthalm Res* 36: 303-311.
- Saxena R, Srivastava S, Trivedi D, Anand E, Joshi S, Gupta S. K. 2003. Impact of environmental pollution on the eye. *Acta Ophthalmol Scand* 81: 491-494.
- Schoeler S, Winkler-Stuck K, Szibor R, Haroon M. F, Gellerich F. N. 2006. Glutathione depletion in antioxidant defense of differentiated NT2-LHON cybrids. *Neurol Dis* 25: 536-544.
- Schulze-Osthoff K, Bauer M. K, Vogt M, Wesselborg S. 1997. Oxidative stress and signal transduction. *Int J Vitam Nutr Res* 67: 336-342.
- Seetharam B. K, Sujatha N. 1998. Flavin nucleotide in human lens: regional distribution in Brunescent cataracts. *Indian J Ophthalmol* 46: 233-237.
- Shaimova V. A. 2002. Lipid peroxidation and antioxidant defense in eye diseases. *Vestn Oftalmol* 118: 56-57.
- Shamsi F. A, Chaudhry I. A, Boulton M. E, Al-Rajhi A. A. 2007. L-Carnitine protects human retinal pigment epithelial cells from oxidative damage. *Curr Eye Res* 32: 574-584.
- Shapiro M. S, Thoft R. A, Friend J, Parrish R. K, Gressel M. G. 1985. 5-Fluorouracil toxicity to the ocular surface epithelium. *Invest Ophthalmol Vis Sci* 26: 580-583.
- Siddall J. R. 1966. Ocular toxic changes with chlorpromazine and thioridazine. *Can J Ophthalmol* 1: 190-198.
- Siu AW, Maldonado M, Sanchez-Hidalgo M, tan DX, Reiter RJ. 2006. Protective effects of melatonin in experimental free radical-related ocular diseases. *J Pineal Res* 40: 101-109.
- Spector A. 1995. Oxidative stress-induced cataract: mechanism of action. *FASEB J* 9: 1173-1182.
- Stahl W, Sies H. 2005. Bioactivity and protective effects of natural carotenoids. *Biochem Biophys Acta* 1740: 101-107.
- Stitt A. W. 2001. Advanced glycation: an important pathological event in diabetic and age related ocular disease. *Br J Ophthalmol* 85: 746-753.
- Sulochana K. N. 2001. Smoking and cataract. *Curr Sci* 81: 1151.
- Sulochana K. N, Punitham R, Ramakrishnan S. 2002. Effect of cigarette smoking on cataract: antioxidant enzymes and constituent minerals in the lens and blood of humans. *Indian J Pharmacol* 34: 428-431.
- Takahashi H. 2005. Free radical development in phacoemulsification cataract surgery. *J Nippon Med Sch* 72: 4-12.
- Tanito M, Agbaga M. P, Anderson R. E. 2007. Upregulation of thioredoxin system via Nrf2-antioxidant responsive element pathway in adaptive-retinal neuroprotection in vivo and vitro. *Free Rad Biol Med* 42: 1838-1850.
- Taysi S, Kocer I, Memisogullari R, Kiziltunc A. 2002. Serum oxidant/antioxidant status in patients with Behçet's disease. *Ann Clin Lab Sci* 32: 377-382.
- Tezel G, Yang X, Luo C, Peng Y, Sun S. L, Sun D. 2007. Mechanisms of immune system activation in glaucoma: oxidative stress-stimulated antigen presentation by the retina and optic nerve head glia. *Invest Ophthalmol Vis Sci* 48: 705-714.
- Thompson A. M. 2007. Ocular toxicity of fluoroquinolones. *Clin Exp Ophthalmol* 35: 566-577.
- Truscott R. J. 2005. Age-related nuclear cataract-oxidation is the key. *Exp Eye Res* 80: 709-725.
- Tsubota K. 2007. Oxidative stress and inflammation:hypothesis for the mechanism of aging. *Nippon Ganka Gakkai Zasshi* 111: 193-205.
- Ugarte M, Osborne N. N. 2001. Zinc in the retina. *Progress Neurobiol* 64: 219-249.
- van Kuijk F. J. 1991. Effects of ultraviolet light on the eye: role of protective glasses. *Environ Health Perspect* 96: 177-184.
- Varma S. D. 2007. Susceptibility of the ocular lens to nitric oxide: implications in cataractogenesis. *J Ocul Pharmacol Therap* 23: 188-195.
- Varma S. D, Hegde K. R. 2007. Oxidative stress and cataract formation: horizons on its medical prevention. *Expert Rev Ophthalmol*

thalamol 2: 779-801.

Varma SD, Chand D, Sharma YR, Kuck JF, Richards RD. 1984. Oxidative stress on lens and cataract formation: role of light and oxygen. *Curr Eye Res* 3: 35-58.

Veriweij C. L, Gringhuis S. J. 2002. Oxidants and tyrosine phosphorylation; role of acute and chronic oxidative stress in T- and B-lymphocyte signaling. *Antioxid Redox Signal* 5: 543-551.

Vidan A, Luria S, Eisenkraft A, Hourvitz A. 2002. Ocular injuries following sulfur mustard exposure: clinical characteristics and treatment. *IMAJ* 4: 577-578.

Weinberger B, Iaskin DL, Heck DE, Iaskin JD. 2002. Oxygen toxicity in premature infants. *Toxicol Appl Pharmacol* 181: 60-67.

West S. K, Longstreth J. D, Munoz B. E, Pitcher H. M, Duncan D. D. 2005. Model of risk of cortical cataract in the US population with exposure to increased ultraviolet radiation due to stratosphere ozone depletion. *Am J Epidemiol* 162: 1080-1088.

Whatham A, Bartlett H, Eperjesi F, Blumenthal C, Allen J, Suttle C, Gaskin K. 2008. Vitamin and mineral deficiencies in the developed world and their effect on the eye and vision. *Ophthalm Physiol Opt* 28: 1-12.

Wickert H, Zaar K, Grauer A, John M, Zimmermann M, Gillardon F. 1999. Differential induction of proto-oncogene expression and cell death in ocular tissues following ultraviolet irradiation of the rat eye. *J Ophthalmol* 83: 225-230.

Widmer S, Helbig, H. 2006. Presumed macular toxicity of intravitreal antibiotics. *Klin Monatsbl Augenheilkd.* 223: 456-458.

Wilhelm J, Smistík Z, Mahelková G, Vytásek R. 2007. Redox regulation of proliferation of lens epithelial cells in culture. *Cell Biochem. Funct* 25: 317-321.

Wolf N, Pendergrass W, Singh N, Swisshelm K, Schwartz J. 2008. Radiation cataracts: mechanisms involved in their long delayed occurrence but then rapid progression. *Mol Vis* 14: 274-285.

Wong R. W, Richa D. C, Hahn P, Green W. R, Dunaief J. L. 2007. Iron toxicity as a potential factor in AMD. *Retina* 27: 997-1003.

Yağci R, Aydın B, Erdurmuş M, Karadağ R, Gürel A, Durmuş M, Yiğitoğlu R. 2006. Use of melatonin to prevent selenite-induced cataract formation in rat eyes. *Curr Eye Res* 31: 845-850.

Yamada Y, Tian J, Yang Y, Cutler RG, Wu T, Telljohann RS, Mattson MP, Handa JT. 2008. Oxidized low density lipoprotein induce a pathologic response by retinal pigmented epithelial cells. *J Neurochem* 105: 1187-1197.

Yamato M, Matsumoto S, Ura K, Yamada K, Naganuma T, Inoguchi, T, Watanabe T, Utsumi H. 2007. Are free radical reactions increased in the diabetic eye? *Antioxid Redox Signal* 9: 367-373.

Yang D, Elnor S. G, Bian Z. M, Till G. O, Petty H. R, Elnor V. M. 2007. Pro-inflammatory cytokines increase reactive oxygen species through mitochondria and NADPH oxidase in cultured RPE cells. *Exp Eye Res* 85: 462-472.

Yao K, Tan J, Gu W. Z, Ye P. P, Wang K. J. 2007. Reactive oxygen species mediates the apoptosis induced by transforming growth factor beta(2) in human lens epithelial cells. *Biochem Biophys Res Commun* 354: 278-283.

Yu L, Kelly U, Ebricht J. N, Malek G, Saloupis, P, Rickman D. W, McKay B. S, Arshavsky V. Y, Bowes Rickman C. 2007. Oxidative stress-induced expression and modulation of phosphatase of regenerating liver-1 (PRL-1) in mammalian retina. *Biochem Biophys Acta* 1773: 1473-1482.

Zavalic M, Mandic Z, Turk R, Bogadi-Sare A, Plavec D, Gomzi M, Skender L. J. 1998. Assessment of color vision impairment in male workers exposed to toluene generally above occupational

exposure limits. *Occup Med* 48: 175-180.

Zhang B, Safa R, Rusciano D, Osborne N. N. 2007. Epigallocatechin gallate, an active ingredient from green tea, attenuates damaging influences to the retina caused by ischemia/reperfusion. *Brain Res* 1159: 40-53.