

# New host findings of two parasitic isopods (Flabellifera: Cymothoidae) found on some fish caught in the Dardanelles Strait (Çanakkale, Turkey)

Ruhay ALDIK<sup>1\*</sup>, Ahmet ÖKTENER<sup>2</sup>, Fikret ÇAKIR<sup>3</sup>, Yusuf ŞEN<sup>3</sup>,  
Gençtan Erman UĞUR<sup>3</sup>, Sezginer TUNCER<sup>3</sup>

<sup>1</sup>Department of Fisheries Technology, Faculty of Applied Sciences, Canakkale Onsekiz Mart University,  
TR, 17100, Canakkale, Turkey

<sup>2</sup>Department of Fisheries, Sheep Research Institute, Çanakkale Road, 7.km, Bandırma, Balıkesir

<sup>3</sup>Faculty of Marine Sciences and Technology, Canakkale Onsekiz Mart University, TR, 17100, Çanakkale,  
Turkey

Geliş Tarihi (Received Date): 07.09.2021

Kabul Tarihi (Accepted Date): 14.12.2021

## Abstract

Seven cymothoid species are identified in parasitological studies conducted in the Dardanelles Strait between May 2018- May 2020. The Cymothoids identified are as follows: *Mothocya epimerica* Costa, 1851, *Nerocila bivittata* (Risso, 1816), *Anilocra frontalis* H. Milne Edwards, 1840, *Ceratothoa oestroides* (Risso, 1826), *Ceratothoa parallela* (Otto, 1828), *Ceratothoa italica* Schiödte & Meinert, 1883, *Emetha audouini* (H. Milne Edwards, 1840). *A. frontalis* and *C. italica* are found for the first time from *Sciaena umbra* Linnaeus, 1758 and *Boops boops* (Linnaeus, 1758) respectively. Although several reports of these parasites have been given from Turkey, the morphological characters of three species have not been given. The drawings, descriptions of the mouthparts, pereopods, and pleopods of *N. bivittata*, *A. frontalis*, *C. italica* are given in this study.

**Keywords:** Turkey, cymothoid, Dardanelles Strait, *Anilocra*, *Ceratothoa*, *Nerocila*

\*Ruhay ALDIK, ruhayaldik@gmail.com, <http://orcid.org/0000-0001-5791-7491>

Ahmet ÖKTENER, ahmetoktener@yahoo.com, <http://orcid.org/0000-0003-0858-0532>

Fikret ÇAKIR, fikretcakir17@yahoo.com, <http://orcid.org/0000-0001-5261-2365>

Yusuf ŞEN, yusuf.sen@comu.edu.tr, <http://orcid.org/0000-0002-0595-4618>

Gençtan Erman UĞUR, ermanugur@hotmail.com, <http://orcid.org/0000-0002-3131-9239>

Sezginer TUNCER, stuncer@comu.edu.tr, <http://orcid.org/0000-0002-6634-7109>

## Çanakkale Boğazı' nda (Çanakkale, Türkiye) yakalanan bazı balıklarda bulunan iki parazitik izopod türünün (Flabellifera: Cymothoidae) yeni konak bulguları

### Öz

Çanakkale Boğazı' nda yedi cymothoid türü Mayıs 2018- Mayıs 2020 tarihleri arasında teşhis edilmiştir. Teşhis edilen parazit türleri şunlardır: *Mothocya epimerica* Costa, 1851, *Nerocila bivittata* (Risso, 1816), *Anilocra frontalis* H. Milne Edwards, 1840, *Ceratothoa italica* Schiöde & Meinert, 1883, *Ceratothoa parallela* (Otto, 1828), *Ceratothoa oestroides* (Risso, 1826), *Emetha audouini* (H. Milne Edwards, 1840). *A. frontalis* ve *C. italica* sırasıyla ilk kez eşkina balığından, *Sciaena umbra* Linnaeus, 1758 ve kupez balığından, *Boops boops* (Linnaeus, 1758) bulunmuştur. Türkiye' de her ne kadar bu parazitlerin çeşitli raporları verilmekle birlikte, üç türün morfolojik karakterleri verilmemiştir. Bu çalışmada *N. bivittata*, *A. frontalis*, *C. italica*' nın ağız parçaları, pereopod ve pleopodlarının tanımlamaları ve çizimleri verilmiştir.

**Anahtar kelimeler:** Türkiye, cymothoid, çanakkale boğazı, *Anilocra*, *Ceratothoa*, *Nerocila*

### 1. Introduction

The members of the family Cymothoidae (Crustacea, Isopoda) are ectoparasites living on marine fishes. They are mostly reported on the body surface, fins of fishes (*Anilocra* Leach, 1818, *Nerocila* Leach, 1818); in the buccal cavity of fishes (*Ceratothoa* Dana, 1852, *Emetha* Schiöedte & Meinert, 1883), and in the gill cavity of fishes (*Mothocya* Costa in Hope, 1851, *Elthusa* Schiöedte & Meinert, 1884). They are rarely reported in a pouch of fishes [1, 2].

Cymothoids are mostly known as tongue-biters. There are various comparative taxonomic studies on the presence of cymothoids on wild fish [3-6]. In addition, there are also records of parasitic isopods from fish farms and several studies on the effects of parasites on the host fish. Bragoni et al. [7] examined the effects of *Nerocila orbignyi* (Guérin-Méneville, 1832) on farmed sea bass in Corsica. They identified that the decrease in the body condition of fishes and increase in blood urea on sea bass infested with the parasite. They found significant differences in the blood parameters of the host fish. Çolak et al. [8] searched the effects of *Ceratothoa oestroides* on cultured meagre in Croatian. They found that fish infested with cymothoid is smaller than non-infested fish. Horton and Okamura [2] examined the hematology parameters of farmed sea bass infested by *C. oestroides* in Turkey. They mentioned that infested fish had lowered erythrocyte counts, hematocrit, and hemoglobin values and significantly increased leucocyte counts. Papapanagiotou and Trilles [9] observed serious lesions on farmed sea bream infested with *Ceratothoa parallela* (Otto, 1828) in Greece.

Although there are various reports about cymothoids that are presented only with photographs in Turkey, the studies on their morphology are quite a few. It can sometimes be difficult to reach studies on the morphology of Cymothoids in the

Mediterranean and the Black Sea. Cymothoids are not only found in parasitological studies. For example, researchers studying fish systematics and population dynamics can find these parasites in the field or the laboratory. As a result, more research into cymothoid biology is needed to better understand parasite relationships. The aim of this study is to update the morphological features and make diagnosis easier. New hosts for *C. italica*, *A. frontalis*, are also described in the study.

## 2. Material and method

Fish samples were obtained from the commercial fisherman at the landing, Çanakkale Strait, Sea of Marmara during May 2018 – May 2020. Cymothoid samples were preserved in 70% alcohol and dissected using a stereomicroscope. The drawings of extremities of parasites were done with a microscope drawing tube. Measurements were given in millimeters (mm). Species names and synonyms were controlled with an online database [10]. The ecological characters (feeding, habitat, etc) of the host were examined according to Fishbase [11]. Identifications of cymothoids were carried out according to Trilles [3], Bruce [4], Horton [5], Hadfield [6], Aneesh et al. [12].

## 3. Results

In this parasitological study, seven species of cymothoid isopods from Çanakkale Strait were identified (Table 1).

Table 1. Cymothoid species found and their hosts

<b>Cymothoid Species</b>	<b>Host</b>	<b>Examined Fish</b>	<b>Number of fish samples examined</b>	<b>Site of Infestation</b>	<b>Prevalence (%)</b>
<i>Mothocya epimerica</i> (Figure 1a)	Big-scale sand smelt, <i>Atherina boyeri</i> Risso 1810	132	4	Gill cavity	3
<i>Nerocila bivittata</i> (Figure 1b)	Red scorpionfish, <i>Scorpaena scrofa</i> Linnaeus 1758	42	3	Body surface	7
<i>Anilocra frontalis</i> (Figure 1c,1d,1e)	Brown meagre, <i>Sciaena umbra</i> Linnaeus 1758	24	2	Body surface	8.3
<i>Ceratothoa oestroides</i>	Bogue, <i>Boops boops</i> (Linnaeus 1758)	171	23	Buccal cavity	13
<i>Ceratothoa oestroides</i>	Blotched picarel, <i>Spicara maena</i> (Linnaeus 1758)	90	18	Buccal cavity	20
<i>Ceratothoa oestroides</i>	Atlantic horse mackerel, <i>Trachurus trachurus</i>	325	1	Buccal cavity	0.3
<i>Ceratothoa parallela</i>	Bogue, <i>Boops boops</i> (Linnaeus 1758)	171	14	Buccal cavity	8
<i>Ceratothoa parallela</i>	Blotched picarel, <i>Spicara maena</i> (Linnaeus 1758)	90	24	Buccal cavity	26
<i>Ceratothoa italica</i> (Figure.1f, 1g)	Bogue, <i>Boops boops</i> (Linnaeus 1758)	171	2	Buccal cavity	1.1
<i>Emetha audouini</i>	Blotched picarel, <i>Spicara maena</i> (Linnaeus 1758)	90	5	Buccal cavity	5.5




Figure. 1. a. *Mothocya epimerica*, b. *Nerocila bivittata*, c. *Anilocra frontalis*, d. *A. frontalis* on *Sciæna umbra*, e. Attachment of *A. frontalis* on the body surface, f. *Ceratothoa italica*, g. fifth pleonite of *Ceratothoa italica*

***Cymothoidae Leach, 1818 (Isopoda)******Nerocila Leach, 1818****Nerocila bivittata* (Risso, 1816) (Figures 1b, 2, 3)

Female: Length 10.9 mm, width 6.3 mm. Body length is 1.7 times its width. The widest and narrowest of pereonites are 5 and 1, respectively. Cephalon dome-shaped and visible, cephalon length is 0.5 times the width. Eyes well developed; one eye 0.2 times width of the cephalon, 0.4 times length of the cephalon.

Pereonites increased gradually and slightly from 1 to 5; decreasing progressively from 5 to 7 in width. Pereonites progressively increased from 1 to 5 except pereonite 2, and later decreased from 5 to 7 in length. Pereon widest at pereonite 4, the narrowest at pereonite 7. Pereonites 5 longest, pereonite 2 shortest. Coxae 1-4 partly visible in dorsal view; posteroventral angles of coxae 1-4 extending beyond posterior pereonites. Coxae 5-7 are slightly visible. Pleonite 1 slightly visible and partly concealed by pereonite 7, other pleonites visible; pleonites lengths increased progressively; pleonite 5 longest; pleonite 1 shortest. Pleonite on widest at pereonite 5, the narrowest at pereonite 1 in width. Pleotelson length is 0.65 times the width, lateral and posterior margins slightly rounded.

Antennula and antennula consist of eight and eleven articles, respectively. Mandible palp article 1 largest, article 2 and 3 same length; article 3 with 9 setae on distolateral margin decreasing in size from longest distal seta, article 2 with 2 setae. Maxillula with four terminal robust setae. Maxilliped distal palp with five apical recurved setae. Lateral and medial lobes of maxilla with 2 setae.

Pereopods 1-5 without seta. Propodus with three setae, carpus with two setae of pereopod 6. Propodus with 10 setae; carpus with five setae and merus with three setae of pereopod 7. Pereopods 5-7 longer 1-4 similar in size. Pereopod 1 basis length 1.6 times the widest; ischium length 0.4 times basis length; propodus length 1.5 times width; dactylus length 1.9 times propodus length and 4 times basal width. Pereopod 7 basis length 2.8 times widest; ischium length 0.3 times basis length; merus length 0.75 times width and 0.3 times ischium; carpus 0.8 times width and 0.8 times ischium; propodus 1.8 times width and 1.1 times ischium; dactylus 1.1 times propodus and 3 times basal width. Pleopods gradually decreases from 1 to 5; pleopods 1-2 and 3-5 are similar in size. Endopod of all pleopods shorter than exopod. Pleopod 1 exopod 1.3 times width, lateral and medial margins rounded; endopod 1.5 times width, medial and lateral margin straight, distally rounded; peduncle 3 times length.

Protopod medial margin of pleopods 1-5 with four coupling hooks. Appendix masculina o pleopod 2 about 0.4 times endopod length. Proximomedial lobes of pleopods 3-5 well developed. Pleopods 3-5 with folds.

Pleotelson length 0.7 times anterior width; dorsal surface of pleotelson smooth; length 0.7 times pleotelson anterior width; margins of lateral and posterior slightly rounded. Endopod shorter than exopod of uropod; posterior margin of pleotelson extends beyond endopod. Endopod 1.8 times the greatest width; median and lateral margins slightly rounded; apice without seta. Exopod extends beyond to apice of endopod; 4 times greatest width, medial and lateral margins straight; apices with two setae. Apices of exopod and endopod rounded. Uropod peduncle with one seta apically.




Figure 2. *Nerocila bivittata*, female a. antennula, b. antenna, c. maxilliped, d. maxilla, e. mandible, f. maxillule




Figure 3. *Nerocila bivittata*, female a pereopods, b. pleopods, c. Uropod

**Remarks:** *Nerocila bivittata* was described by Risso [13]. The redescription of this species was made only by Trilles [14] and Öktener et al. [15]. The mouthparts (maxilliped, maxilla, maxillule, mandible), pereopods, and pleopods found in this study are compatible with Trilles [14] and Öktener et al. [15]'s findings.



**Genus *Anilocra* Leach, 1818**

*Anilocra frontalis* H. Milne Edwards, 1840 (Figure 1c, 1d, 1e, 4, 5)

Male: Length 12 mm, width 4 mm. Body elongate, length is 3 times its width. The widest and narrowest of pereonites are 4 and 7, respectively. Cephalon triangle-shaped and visible, length 0.57 times width. Eyes very developed; one eye 0.3 times cephalon width, 0.5 times cephalon length.

Pereonites increased gradually and slightly from 1 to 4; decreasing progressively from 5 to 7 in width. Pereonites progressively increased from 1 to 5 and later decreased from 5 to 7 in length. Pereon widest at pereonite 4, the narrowest at pereonite 7. The longest and shortest of pereonites are 5 and 7, respectively. Coxae 2-7 partially visible in dorsal, posteroventral angles of coxae 2-7 not extending beyond posterior pereonites.

Pleonite 1 is slightly visible in the dorsal. Pleonite 1 partly concealed by pereonite 7, other pleonites visible; pleonites lengths increased progressively; pleonite 5 longest; pleonite 1 shortest, pleonites 2-3 subequal in length. Pleonite 1 is slightly visible and pereonite 7 partially covers it. Pleonite on widest at pereonite 5, the narrowest at pereonite 1 in width. Pleotelson length 0.7 times width, dorsal surface smooth; posterior and lateral margins slightly rounded.

Antenna longer than antennula. Antennula and antenna include 8 and 9 articles, respectively. Maxilliped distal palp with three apical recurved setae. Mandible palp article 1 largest, article 2 and 3 same in length; article 3 with 14 setae on distolateral margin decreasing in size from longest distal seta; article 2 with 1 seta. Maxillula with four terminal robust setae. Lateral and medial lobes of maxilla with 2 recurved setae.

Pereopods with seta except pereopod 1. Pereopod 2-6 with one seta on the anterior margin of merus. Pereopods 5-6 with two setae on the posterior margin of propodus. Pereopod 7 with 27 setae on posterior margin of propodus; carpus with 17 setae and merus with seven setae. Pereopod 1 basis length 2 times greatest width; ischium 0.5 times basis; propodus 2.1 times width; dactylus slender, 1.5 times propodus, 2.1 times basal width. Pereopod 7 basis 2 times greatest width; ischium 0.7 times basis; merus 2 times width, 0.7 times ischium; carpus 1.5 times width, 0.6 times ischium length; propodus length 2.9 times width, as long as ischium; dactylus slender 1.1 times propodus, 3.6 times basal width.

Pleopods gradually decrease from 3 to 5; pleopods 1-3 and 4-5 are similar in size. Pleopod exopods are longer than endopods. Pleopod 1 exopod 1.4 times width, medial margin convex; endopod 1.7 times width, medial margin straight; peduncle 3 times as wide as long. Distal and lateral margins of exopod and endopod are slightly rounded. Protopod medial margin of pleopods with four coupling hooks. Appendix masculina of second pleopod 0.6 times endopod length. Proximomedial lobes of pleopods 3-5 well developed.

Pleotelson 0.7 times anterior width, posterior and lateral margins slightly rounded. Endopod shorter uropod exopod, endopod extending beyond posterior margin of pleotelson. Endopod 2.5 times width, lateral and medial margin slightly rounded, all of the margins without seta. Exopod extending to end of endopod, 5.2 times width, lateral



and medial margin straight, apices without seta. Distal of endopod and exopod rounded. Uropod peduncle margins with a seta.




Figure 4. *Anilocra frontalis* male, a. antenna, b. antennula, c. maxilliped, d. maxilla, e. mandible, f. maxillule




Figure 5. *Anilocra frontalis* male, a. pereopods, b. pleopods, c. Uropod

**Remarks:** *Anilocra frontalis* was described by Milne Edwards [16]. The redescription of this species was made only by Trilles [14]. The number of seta on maxilliped, maxilla, maxillule, mandible found in this study is compatible with Trilles [14] 's findings. The number of articles on the antenna in this study is compatible with Trilles [14], but the number of articles on antennule is different which was found by Trilles [14]. The dispersion and sequence of spines on pereopods articles and coupling hooks on pleopods found in this study are compatible with Trilles [14]. The male of *A. frontalis* resembles *A. physodes*. *A. frontalis* can be separated from *A. physodes* redescribed by Trilles [14], by the lateral lobe of maxilla with 2 setae; medial lobe with 2 large setae (two robust setae on lateral lobe and one robust seta on medial lobe in *A. physodes*), mandible palp second article with 3 setae and third article with 11-13 setae (9 setae on the distolateral margin of articles 3 respectively in *A. physodes*); pleopods and pleotelson without setae (posterior side of pleopods and pleotelson with setae in *A. physodes*).

### **Genus *Ceratothoa* Dana, 1852**

*Ceratothoa italica* Schiödte & Meinert, 1883 (Figure 1f, 1g, 6, 7, 8, 9)

Female: Length 22.14 mm, width 8.2 mm. Body elongate, 2.7 times width. The widest and narrowest of pereonites are 4 and respectively. Cephalon triangle-shaped, and visible, cephalon length 0.4 times the width. Eyes well developed; one eye 0.23 times cephalon width, 0.5 times cephalon length.

Pereonites increased gradually and slightly from 1 to 4; decreasing progressively from 5 to 7 in width. Pereonites progressively increased from 2 to 4 and later decreased from 5 to 7 in length. Pereon widest at pereonite 4, the narrowest at pereonite 7. Pereonite 1 longest, pereonite 7 shortest. Coxae 4-7 partly visible in dorsal view; posteroventral angles of coxae 4-7 extending beyond posterior pereonites.

Pleonites are visible in the dorsal. Pleonites lengths increased progressively (except 4); pleonite 5 longest; pleonite 1 shortest. Pleonite on widest at pereonite 3, the narrowest at pereonite 1 in width. Pleotelson 0.47 times anterior width; posterior and lateral margins slightly rounded.

Antenna slightly longer than antennula, consisting of 7 articles. Antennula consists of 6 articles. Maxilliped distal palp with three apical recurved setae on article 3. Mandible palp article 1 largest, article 1 shortest, article 3 with 3 setae on distolateral margin. Maxillula with four terminal robust setae. Lateral and medial of maxilla with 9 and 5 robust setae.

Pereopods without seta. Pereopod 1 basis 1.7 times greatest width; ischium 0.6 times basis; propodus 1.1 times width; dactylus slender, 1.2 times propodus, 1.1 times basal width. Pereopod 7 basis 1.5 times greatest width; ischium 0.6 times basis; merus 0.5 times width, 0.4 times ischium; carpus 0.4 times width, 0.3 times ischium; propodus 1.1 width, 0.5 times ischium; dactylus slender 1.1 times propodus, 1.8 times basal width.

Pleopods gradually decrease from 3 to 5; pleopod 1 biggest and pleopod 5 smallest. Endopod of all pleopods shorter than exopod. Pleopod 1 exopod 1.1 times width, endopod 1.2 times width. Distal, lateral margin of exopod and endopod slightly

rounded, medial margin strongly convex. Peduncle 2.3 times as width as length. Pleopods without coupling hooks on protopod medial margin. Pleopods without folds.

Pleotelson 0.5 times anterior width; lateral margins straight; posterior margin slightly rounded. Uropod exopod slightly shorter than endopod; endopod not extending beyond posterior margin of pleotelson. Endopod, apically rounded, 5 times as long as greatest width. Exopod extending to end of endopod, 4.4.5 times width. Medial, lateral margins of endopod and exopod slightly straight, margins and apices without seta. Uropod peduncle margins without seta.




Figure 6. *Ceratothoa italica* female, a. antenna, b. antennule, c. maxilliped, d. maxilla, e. mandible, f. maxillule




Figure 7. *Ceratothoa italica* female, a pereopods, b. pleopods, c. uropod

Male: Length 8.5 mm, width 3.28 mm. Body elongate, 2.6 times width. The widest and narrowest of pereonite are 5 and 7 respectively. Cephalon triangle-shaped, visible, 0.45 times longer than width. Eyes developed; one eye 0.36 times cephalon width and 0.8 times length of the cephalon.

Pereonites increased gradually and slightly from 1 to 5; decreasing progressively from 5 to 7 in width. Pereonites progressively increased from 2 to 4 and later decreased from 5 to 7 in length. The widest and narrowest of pereonite are 5 and 1, respectively. The longest and shortest of pereonites are 1 and 7 respectively. Coxae 4-7 partly visible in dorsal view; posteroventral angles of coxae 4-7 slightly extending beyond posterior pereonites.

Pleonite 1 is slightly visible in the dorsal. Pereonite 7 partially covers pleonite 1. Pleonites lengths increased progressively; pleonite 5 longest; pleonite 2 shortest in

length. Pleonite on widest at pereonite 3, the narrowest at pereonite 1 in width. Pleotelson 0.47 times anterior width, posterior and lateral margins slightly rounded.

Antennula consists of 7 articles. Antenna consists of nine articles. Maxilliped distal palp with three apical recurved setae on article 3. Mandible palp article 1 largest, article 2 shortest, article 3 with 12 setae on distolateral margin; article 2 with 6 setae. Maxillula with four terminal robust setae. Lateral and medial of maxilla with 3 recurved robust setae.

Pereopods with one seta on the anterior margin of merus. Pereopod 1 basis 1.5 greatest width; ischium 0.5 times basis; propodus 1.8 times width; dactylus 0.9 times propodus, 2.2 times basal width. Pereopod 7 basis 1.6 times greatest width; ischium 0.7 times basis; merus 0.6 times width, 0.3 times ischium; carpus 0.6 times width, 0.3 times ischium; propodus 1.6 times width, 0.7 times ischium; dactylus slender 0.8 times propodus, 1.6 times basal width.

Pleopods gradually decrease from 3 to 5; pleopods 1-2 and 3-5 are similar in size. Endopod of all pleopods shorter than exopod. Pleopod 1 exopod 1.1 times width, distal and medial margins slightly rounded, lateral margin strongly convex; endopod 1.6 times width, distal and medial, lateral margin slightly rounded and medial margin straight; peduncle 2.3 times as wide as long. Protopod medial margin of pleopods 1-5 with four coupling hooks. Pleopods 1 with well-developed proximomedial lobe. Pleopods without folds.

Pleotelson 0.5 times anterior width; lateral slightly convex; posterior slightly rounded. Uropod endopod slightly shorter than exopod; endopod not extending beyond posterior margin of pleotelson. Endopod 4.5 times the greatest width; exopod extending to end of endopod, 4.4 times the greatest width. Medial, lateral, distal margins of exopod and endopod slightly rounded, medial margins straight. Margins and apices of exopod and endopod without seta. Uropod peduncle margins with two setae.7




Figure 8. *Ceratothoa italica* male, a. antenna, b. antennule, c. maxilliped, d. maxilla, e. mandible




Figure 9. *Ceratothoa italica* male, a. pereopods, b. pleopods, c. Uropod

**Remarks:** The female and second pullus stage of *Ceratothoa italica* was described by Schiödte and Meinert [17], except males. Horton [5] redescribed seven species of *Ceratothoa* genus. Schiödte and Meinert [17] and Horton [5]’s descriptions do not include all of the mouthparts of females and males. The antennule with six articles and antenna with nine articles of females found in the present study is different from the antennule with eight articles and antenna with nine articles found by Schiödte and Meinert [17]. Pereopod morphological characters of *Ceratothoa italica* were given in a key to the *Ceratothoa* genus prepared by Horton [5] as “Pleotelson much wider than long and as wide or wider than pereonite VII. Prominent merus expansions on pereopods I-III, less prominent on pereopods V-VII”. Pereopod characters found in this study are compatible with Horton [5]. Neither the mouthparts of the female nor the morphologies of the male of *C. italica* are given in Schiödte and Meinert [17] and Horton [5]. Therefore, no comparison has been made in this study.

#### 4. Discussion

Species belonging to the genera *Nerocila* (*Nerocila bivittata* (Risso, 1816), *Nerocila orbigny* (Guérin-Méneville, 1832), *Nerocila milesensis* Öktener, Trilles, Tuncer, 2020), *Anilocra* (*Anilocra physodes* (Linnaeus, 1758), *Anilocra frontalis* H. Milne Edwards, 1840), *Mothocya* (*Mothocya epimerica* Costa, 1851, *Mothocya belonae* Bruce, 1986, *Mothocya taurica* (Czerniavsky, 1868)), *Elthusa* (*Elthusa poutassouiensis* (Penso, 1939), *Elthusa sinuata* (Koelbel, 1879)), *Ceratothoa* (*Ceratothoa parallela* (Otto, 1828), *Ceratothoa capri* (Trilles, 1964), *Ceratothoa italica* Schiödte & Meinert, 1883, *Ceratothoa oestroides* (Risso, 1826), *Ceratothoa oxyrrhynchaena* Koelbel, 1878), *Emetha* (*Emetha audouini* (H. Milne Edwards, 1840)) have been reported from marine fish in Turkey [15, 18].

*Mothocya epimerica* is a typical parasite of Atherinidae family. This species was only reported from *Atherina boyeri* Risso 1810 (syn. *Atherina mochon*, *Atherina rissoi*), and *Atherina hepsetus* Linnaeus 1758 [1, 19]. It is known from the Mediterranean, Black Sea, Adriatic Sea, and Atlantic Ocean [1, 19, 20]. In this study, this parasite is found only on *Atherina boyeri*.

*Ceratothoa italica* is a parasite that typically settles in the buccal cavity of fish. It prefers especially fish belonging to the family Sparidae such as *Lithognathus mormyrus* (Linnaeus 1758), *Pagellus erythrinus* (Linnaeus 1758), *Oblada melanura* (Linnaeus 1758), *Sargus* sp, *Diplodus sargus* (Linnaeus 1758), *Diplodus annularis* (Linnaeus 1758) [1, 21, 22]; rarely reported from *Spicara maena* (Linnaeus 1758), *Dicentrarchus labrax* (Linnaeus 1758) [18, 23]. This species has already been recorded in the Mediterranean, Adriatic Sea, and Sea of Marmara [1]. *Boops boops* (Linnaeus 1758) belonging to the family Sparidae has been found as a new host for *C. italica*. The fact that this parasite was previously reported from sparids confirms our finding.

*Anilocra frontalis* is a typical external parasite, especially prefers body surfaces and fins of fish. This species was especially reported on the Labridae (*Labrus bergylta* Ascanius 1767 (syn. *Labrus maculatus*), *Labrus mixtus* Linnaeus 1758 (syn. *Labrus vetula*), *Labrus merula* Linnaeus 1758, *Symphodus melops* (Linnaeus 1758) (syn. *Labrus melops*), *Symphodus cinereus* (Bonnaterre 1788) (syn. *Crenilabrus cinereus*), *Symphodus ocellatus* (Linnaeus 1758) (syn. *Crenilabrus ocellatus*), *Symphodus rostratus* (Bloch 1791) [1].

*Anilocra frontalis* has also been recorded from the other fish families such as *Pollachius pollachius* (Linnaeus 1758) (syn. *Merlangus pollachius*), *Lipophrys pholis* (Linnaeus 1758) (syn. *Blennius pholis*), *Taurulus bubalis* (Euphrasen 1786) (syn. *Cottus bubalis*), *Redigobius bikolanus* (Herre 1927) (syn. *Gobius flavescens*), *Gobius paganellus* Linnaeus 1758, *Pomatoschistus minutus* (Pallas 1770) (syn. *Gobius minutus*), *Gaidropsarus mediterraneus* (Linnaeus 1758) (syn. *Onos mustella*), *Spinachia spinachia* (Linnaeus 1758) (syn. *Spinachia vulgaris*), *Boops boops* (Linnaeus 1758), *Spondyliosoma cantharus* (Linnaeus 1758), *Sarpa salpa* (Linnaeus 1758) [1], *Mullus barbatus* Linnaeus 1758, *Umbrina canariensis* Valenciennes 1843, *Pagellus acarne* (Risso 1827), *Diplodus annularis* (Linnaeus 1758), *Lithognathus mormyrus* (Linnaeus 1758), *Solea solea* (Linnaeus 1758) (syn. *Solea vulgaris*) [22], *Umbrina cirrosa* (Linnaeus 1758), *Argyrosomus regius* (Asso, 1801) (syn. *Sciena aquila*) [24].



It is known in the Mediterranean, Atlantic Ocean, and Adriatic [1]. *Sciaena umbra* Linnaeus 1758 is a new host for *A. frontalis*. *Sciaena umbra* belongs to Sciaenidae. This parasite was reported from *Umbrina canariensis* Valenciennes 1843 (Sciaenidae) by Ramdane et al. [22] and from *Umbrina cirrosa*, *Argyrosomus regius* by Ramdane and Trilles [24].

*Nerocila bivittata* is recorded from Labridae, Sparidae, and Scorpaenidae in the Mediterranean [1]. Host selectivity confirms the finding in this study. It is reported for the first time from in the Dardanelles Strait.

*Ceratothoa oestroides* is known from North Atlantic Ocean, the Mediterranean Sea. Hosts of this parasite constitute mainly Sparidae and Centranchidae families [1]. In the present study, the parasite was frequently found in blotched picarel, bogue.

*Ceratothoa parallela* is mostly distributed in the Mediterranean and Atlantic Ocean. It like *C. oestroides* has been reported from many host species of Sparidae [1].

## References

- [1] Trilles, J.P., Les Cymothoidae (Crustacea, Isopoda) du monde (Prodrome pour une faune), **Studia Marina**, 21/22,1-2, 1-288, (1994).
- [2] Horton, T. and Okamura, B., Post-haemorrhagic anaemia in sea bass, *Dicentrarchus labrax* L., caused by blood-feeding of *Ceratothoa oestroides* (Isopoda: Cymothoidae), **Journal of Fish Diseases**, 26, 401-406, (2003).
- [3] Trilles, J.P., Recherches sur les Isopodes Cymothoidae des côtes françaises. Vol. I: Bionomie et parasitisme, Vol. II: Biologie Générale et Sexualité. Thèse Doctorat ès Sciences, Montpellier, 793 pp, (1968).
- [4] Bruce, N.L., Australian species of *Nerocila* Leach, 1818, and *Creniola* n. gen. (Isopoda: Cymothoidae), crustacean parasites of marine fishes, **Records of the Australian Museum**, 39, 355-412, (1987).
- [5] Horton, T., *Ceratothoa steindachneri* (Isopoda: Cymothoidae) new to British waters with a key to north-east Atlantic and Mediterranean *Ceratothoa*, **Journal of the Marine Biological Association of the United Kingdom**, 80, 1041-1052, (2000).
- [6] Hadfield, K.A., The biodiversity and systematics of marine fish parasitic isopods of the family Cymothoidae from southern Africa, **Philosophiae Doctor, University of Johannesburg, Faculty of Science**, Johannesburg, 445 pp, (2012).
- [7] Bragoni, G., Romestand, B. and Trilles, J.P., Cymothoadian parasitosis of the sea-dace (*Dicentrarchus labrax* Linnaeus 1758) during breeding. II. Parasitic ecophysiology in the Diana pond (upper Corsica), **Annales de Parasitologie Humaine et Comparee**, 58, 593-609, (1983).
- [8] Čolak, S., Kolega, M., Mejdandžić, D., Župan, I., Šarić, T., Piplović, E. and Mustačić, B., Prevalence and effects of the cymothoid isopod (*Ceratothoa oestroides*, Risso 1816) on cultured meagre (*Argyrosomus regius*, Asso 1801) in the Eastern Adriatic Sea, **Aquaculture Research**, 49, 1001-1007, (2018).
- [9] Papapanagiotou, E.P. and Trilles, J.P., Cymothoid parasite *Ceratothoa parallela* inflicts great losses on cultured gilthead sea bream *Sparus aurata* in Greece, **Diseases of Aquatic Organisms**, 45, 237-239, (2001).

- [10] WoRMS Editorial Board, World Register of Marine Species, 2021. <https://www.marinespecies.org> at VLIZ, (Accessed 2021-05-23).
- [11] Froese, R. and Pauly D., FishBase.World Wide Web electronic publication, (2021). [www.fishbase.org](http://www.fishbase.org), version 02/2021, (Accessed 2021-05-21).
- [12] Aneesh, P.T., Hadfield, K.A., Smit, N.J. and Kumar, A.B., A new genus and species of fish parasitic cymothoid isopod (Crustacea) from Indian waters, with a key to the branchial-attaching cymothoid genera, **Marine Biology Research**, 16, 8-9, 565-584, (2020).
- [13] Risso, A., **Histoire Naturelle des Crustacés des Environs de Nice**, Librairie Grecque-Latine-Allemande, Paris. 175 p, (1816).
- [14] Trilles, J.P., Les Cymothoidae (Isopoda, Flabellifera) des cotes françaises. II. Les Anilocridae Schioedte et Meinert, 1881. Genres *Anilocra* Leach, 1818, et *Nerocila* Leach, 1818, **Bulletin du Muséum d'histoire naturelle**, 3e sér. 290 (Zool 200), 347-378, (1975).
- [15] Ökter, A., Tuncer, S. and Trilles, J.P., *Nerocila milesensis* n.sp. (Isopoda: Cymothoidae) parasitic on Devil Firefish *Pterois miles* (Bennett, 1828) from the South Aegean Sea, Turkey, **Vie et milieu - Life and environment**, 70, 1, 7-17, (2020).
- [16] Milne Edwards, H., **Histoire Naturelle des Crustacés, Comprenant l'Anatomie, la Physiologie et la Classification de ces Animaux**, Encyclopédique Roret, Paris. Vol. III, 638 p. (1840).
- [17] Schioedte, J.C. and Meinert, F.W., Symbolae ad Monographiam Cymothoarum Crustaceorum Isopodum Familiae 3. Saophridae. 4. Ceratothoinae, **Naturhistorisk Tidsskrift**, 3, 13, 281-378, (1883).
- [18] Ökter, A. and Trilles, J.P., Report on Cymothoids (Crustacea, Isopoda) collected from marine fishes in Turkey, **Acta Adriatica**, 45, 2, 145-154, (2004).
- [19] Ökter, A. and Sezgin, M., *Mothocya epimerica* Costa, 1851 Flabellifera: Cymothoidae), an Isopod Parasite in the Branchial Cavities of the Black Sea Silverfish *Atherina boyeri* Risso, 1810 (Perciformes, Atherinidae), **Journal of the Black Sea / Mediterranean Environment**, 6, 23-29, (2000).
- [20] Charfi-Cheikhrouha, F., Zghidi, W., Ould Yarba, L. and Trilles, J.P., Les Cymothoidae (Isopodes parasites de poissons) des côtes tunisiennes: écologie et indices parasitologiques, **Systematic Parasitology**, 46, 146-150, (2000).
- [21] Rokicki, J., Biology of adult Isopoda (Crustacea) parasitizing fishes of North-West Africa shelf, **Acta Ichthyologica et Piscatoria**, 15, 1, 95-122, (1985).
- [22] Ramdane, Z., Bensouilah, M.A. and Trilles, J.P., The Cymothoidae (Crustacea, Isopoda), parasites on marine fishes, from the Algerian fauna, **Belgian Journal of Zoology**, 137, 1, 67-74, (2007).
- [23] Bariche, M. and Trilles, J.P., Preliminary check-list of Cymothoids (Crustacea, Isopoda) from Lebanon, parasiting on marine fishes, **Zoology in the Middle East**, 34, 5-12, (2005).
- [24] Ramdane, Z. and Trilles, J.P., Cymothoidae and Aegidae (Crustacea, Isopoda) from Algeria, **Acta Parasitologica**, 53, 2, 173-178, (2008).