

COMMUNICATIONS

DE LA FACULTÉ DES SCIENCES
DE L'UNIVERSITÉ D'ANKARA

Série B: Chimie

TOME: 28

ANNÉE: 1982

**Effect Of Substituents On The Polarographic And Spectrophotometric
Behaviour Of 2- [N-(phenyl)- acetimidoyl] -1,3-Indandione**

by

M.A. MORSI, A.M.A. HELMY

16

Faculté des Sciences de l'Université d'Ankara
Ankara, Turquie

Communications de la Faculté des Sciences de l'Université d'Ankara

Comité de Redaction de la Série B

E. Alper – S. Gümüř – T. Gündüz – Y. Sankaya – C. Tüzün
Secrétaire de Publication
Ö. Çakar.

La Revue "Communications de la Faculté des Sciences de l'Université d'Ankara" est un organe de publication englobant toutes les diciplines scientifique représentées à la Faculté des Sciences de l'Université d'Ankara.

La Revue, jusqu'à 1975 à l'exception des tomes I, II, III etait composé de trois séries

Série A: Mathématiques, Physique et Astronomie,

Série B: Chimie,

Série C: Sciences Naturelles.

A partir de 1975 la Revue comprend sept séries:

Série A₁: Mathématiques,

Série A₂: Physique,

Série A₃: Astronomie,

Série B: Chimie,

Série C₁: Géologie,

Série C₂: Botanique,

Série C₃: Zoologie.

En principe, la Revue est réservée aux mémoires originaux des membres de la Faculté des Sciences de l'Université d'Ankara. Elle accepte cependant, dans la mesure de la place disponible les communications des auteurs étrangers. Les langues Allemande, Anglaise et Française seront acceptées indifféremment. Tout article doit être accompagnés d'un resume.

Les articles soumis pour publications doivent être remis en trois exemplaires dactylographiés et ne pas dépasser 25 pages des Communications, les dessins et figures portes sur les feuilles séparées devant pouvoir être reproduits sans modifications.

Les auteurs reçoivent 25 extraits sans couverture.

l'Adresse : Dergi Yayın Sekreteri,
Ankara Üniversitesi,
Fen Fakültesi,
Beşevler-Ankara

Effect Of Substituents On The Polarographic And Spectrophotometric Behaviour Of 2- [N-(phenyl)- acetamidoyl] -1,3-Indandione

M.A. MORSI, A.M.A. HELMY

Chemistry Department, Faculty of Science Elmansoura University, Elmansoura, Egypt


(Received November 3, 1982; accepted November 29, 1982)

ABSTRACT:

The effect of a variety of substituents on the polarographic behaviour and acid-base equilibrium (revealed spectrophotometrically) of 2-[N-(phenyl) - acetamidoyl]-1,3-indandione has been studied in wide pH range buffer solutions (pH~3-12). The application of the linear free energy relationships using $E_{1/2}$ values with different σ sets showed a weak dependence while pK_a - σ sets showed a good correlation.


Introduction

1,3-indandione and some of its substituted derivatives showed wide applications in medicinal chemistry^{1,2,3}, as antiinflammatory agent⁴ and miticides⁵. Relevant studies to the electrochemistry of this class of compounds were recently reviewed by Stradins⁶. Due to its wide applicability our group was involved with the polarographic and spectometric studies for different groups of substituted 1,3-indandione^{7,8}. The polarographic reduction of 2-[N-(phenyl) acetamidoyl] -1,3-indandione (I_a) was found to proceed along two-electronic, $E_{1/2}$ -pH dependent wave (A), which is assigned to reduction of the $>C=N$ - linkage according to:




Wave A

At $\text{pH} > 8$ wave (A) started to decrease in height with appearance of one electronic $E_{1/2}$ -pH independent wave (B) assigned to reduction of the carbonyl group resulting from the alkaline hydrolysis of the azomethine bond:



In continuation to this direction we report here the effect of electron-withdrawing groups (p-Cl, m-NO₂, p-NO₂) and electron-donating group (p-CH₃) on the polarographic reduction waves (A) and (B) and the degree of ionization of H in position 2 of 2-[N-(phenyl) acetimidoyl]-1,3-indandione:



	I _a	I _b	I _c	I _d	I _e
R =	H	CH ₃ -p	Cl-p	NO ₂ -m	NO ₂ -p

Meanwhile the fidelity of existence of linear free energy relationship (LFER) was tested through correlation of the experimentally determined $E_{1/2}$ and pK_a values with different σ sets.

Experimental

Organic Syntheses:

Compounds I_{a-e} were prepared following procedures previously reported by Amer et al.⁹ A mixture of 2-acetyl-1, 3-indandione (0.01

mole) and aniline, or the respective substituted aniline, in 30 ml acetic acid was heated at 80°C for 30 minutes. The solid that obtained after cooling or dilution with water, was recrystallized from ethanol to give I_{a-c} . Credit of purity was checked by m.p. and elemental analysis which agreed well with the theoretical values.

Polarography:

(a) Apparatus. Polarographic curves were recorded with an E 506 pen-type recorder polarograph (Metrohm, Switzerland). The capillary possessed the following characteristics in H_2O open circuit: $t = 5.1$ s/drop, $m = 1.49$ mg s^{-1} for $h = 40$ cm.

(b) Solution. $10^{-3}M$ -Stock solutions were prepared by dissolving an accurately weighed quantity of material in the appropriate volume of absolute ethanol. Britton-Robinson modified universal buffers¹⁰ were used as supporting electrolyte.

(c) Measurements. All experiments were carried at $25 \pm 2^\circ C$. The half-wave potentials were measured graphically and expressed versus the saturated calomel electrode (SCE) with an accuracy ± 0.005 V.

(d) Procedure. Ethanol (7 ml) and the appropriate buffer solution (12 ml) were introduced in the polarographic cell. The mixture was then deaerated by bubbling a stream of hydrogen for 12 min. 1 ml of $10^{-3}M$ depolarizer was then introduced into the cell so that the final concentration was $5 \times 10^{-5}M$ in 20 ml of 40 % (V/V) ethanolic buffer. Controlled Potential Electrolysis:

Mercury pool electrolysis was carried out 40 % V/V ethanol $10^{-2}M$ -HCl solution (200 ml) and substance (I_a) (200 mg) taken as a typical example. The electrolysis cell was a 250 ml conical flask in which the reference, auxiliary electrodes, and the gas inlet were added by means of a cork. The potential was controlled by a tutorial T⁶ transistorized potentiostat at -1.0 V vs. SCE (i.e. on the limiting current plateau of wave A). The progress of the electrolysis was followed by recording the decrease in current with time and the number of electrons was computed from $i-t$ curves following the procedure outlined by Lingane¹¹ and found to be 2e. After disconnecting the electrolysis cell, 1 ml of the resulting solution was withdrawn and the presence of aniline

in this solution was revealed by a standard spot test¹². Trials to separate and crystalize a solid product were unsuccessful where almost resinous material was obtained.

Spectrophotometric Measurements and Determination of the Apparent Dissociation Constant:

These were carried out using PYE Unicam 1800 spectrophotometer supplemented with Unicam SP 1805 programme controller automatic recording unit. The runs in the visible and UV range were carried out for $5 \times 10^{-5}M$ I_{a-c} , 40 % alcoholic Britton - Robinson buffer solutions at roughly increasing 0.5 pH unit. The pH of the solution was measured and the spectrophotometric run was recorded within few minutes. The pK_a was then calculated using the equation¹³.

$$pH = pK_a + \log (A - A_{min}) / (A_{max} - A)$$


where A_{max} is the absorbance of the undissociated acid, A is the absorbance of the organic reagent at a given pH value and A_{min} is the absorbance of the completely dissociated acid at high pH medium. The intersection of the straight line of pH vs. $\log (A - A_{min}) / (A_{max} - A)$ with the pH axis, at $\log (A - A_{min}) / (A_{max} - A) = 0$ gave directly the pK_a .

Results And Discussion


Polarographic Reduction:

The polarographic behaviour of the p-CH₃ and p-Cl substituents (compounds I_b & I_c) were found identical with that reported for the parent compound I_a ⁸. On the other hand the polarograms obtained with the m-NO₂ (I_d) and p-NO₂ (I_e) derivatives (schematically given in Figs. 1 and 2 respectively) are somewhat complicated due to the interference of the reduction waves of the nitro group with that of the $>C=N-$ linkage. Thus at pH < 4 the nitro group of the m-NO₂ substituent (Fig. 1) showed 4-electronic and 2-electronic waves successively; these were followed by a two-electronic wave at more negative potentials assigned to reduction of the $>C=N-$ linkage (wave A). At $6.5 < pH < 4$ the 2-electronic wave of the nitro group amalgamates with wave (A) and at pH 7.38 the amalgamated wave splits again into two two-electronic waves: the more negative wave (assigned to reduction of $>C=N-$


Fig 1: Schematic representation of the polarograms of $5 \times 10^{-5}M$ 2-N-(m-NO₂-phenyl) aceto-
midoyl-1,3-indandione (I_D) in 40 % V/V ethanolic Britton-Robinson buffers.



group) decreases in height with pH increase⁸ and the more positive wave to reduction of the NO₂ group. At pH > 9 the reduction waves of the nitro decreases in height to a four electronic process and remains constant up to the highest pH. In acidic, neutral and fairly alkaline media (pH < 9) the m-NO₂ group reduces to the respective amine along six electronic process as follows:



At pH > 9 the m-NO₂ is reduced via four-electronic process to the corresponding hydroxylamine:



The p-NO₂ derivative (I_e, Fig. 2) showed similar behaviour except at pH > 9 where the wave height associated with the nitro group reduction still corresponded to 6-electronic process. This is well known behaviour accompanying the polarographic reduction of the aromatic nitro group in the p-position where usually a quinoid^{14,15,16} form is involved in the electrodic process which can be schemed as follows:





Fig 2: Schematic representation of the polarograms of $5 \times 10^{-5} \text{M}$ 2- N-(p-NO₂-phenyl) aceto-
midoyl 1,3-indandione (I₆) in 40 % V/V ethanolic Britton - Robinson buffers.







Fig 3: (a) Electronic absorption spectra of $5 \times 10^{-5} \text{ M}$ (I_a) in 40 % V/V ethanolic Britton-Roninson buffers. (b) $\log (A - A_{\min}) / (A_{\max} - A)$ - pH plot.

pK_a values, calculated from plots of pH vs. $\log (A - A_{\min}) / (A_{\max} - A)$ are compiled in table 1.

Table 1
Spectrophotometric pK_a for compounds $I_a\beta_e$.

Compound	I_a , H	I_b ,	I_c , p-Cl	I_d , m-NO ₂	I_e , p-NO ₂
pK_a	10.50	10.80	10.30	9.60	9.40

Structure-Reactivity Relationship:

The influence of substituents of the reduction wave of the $>C=N$ -group (wave A) and the acid-base equilibrium constant pK_a for the series under investigation were examined. The correlation of half-wave potentials for I_{a-e} at pH 4.0, 6.0 and 8.0 with sigma constants σ as taken from the tabulation of Retchie and Sagar¹⁷ is shown in Fig. 4. The coordinates for the nitro derivatives were not included in the correlation at pH > 4 due to the interference of the reduction waves of the nitro group with that of the azomethine linkage; also a better correlation could be observed when using values for NH_2 which results from reduction of the NO_2 group. The data were treated statistically using the least square method according to Jaffe¹⁸. Detailed calculations were summarized in Table 2. It is obvious that a weak correlation exists between $E_{1/2}-\sigma$ since the reactivity parameter ρ oscillates between $+ 0.03$ and $+ 0.07$. This is to be expected since water molecule is involved in the electrodic process (cf. introduction). For wave (B) the correlation showed complete independency of $E_{1/2}$ on σ values. This confirms the electroreduction scheme previously suggested for these compounds⁸ since the substituents is separated in the aniline portion and thus has absolutely no effect.

On the other hand the correlation of $pK_a - \sigma$ sets (σ , σ^o , σ^+) shown in Fig. 5 reflects a quite fair correlation which is also in accordance with expectation, i.e. p-CH₃(I_b) exhibits the higher pK_a whereas the p-NO₂ (I_e) exhibits the lower one. It is obvious from Table 3 that the values of ρ are positive, signifying that the substrate with an electron withdrawing substituents facilitates the ionization process, hence lower pK_a values compared to the standard substrate. From the other hand, electron donating substituents increases the negative cloud at the ionization center leading to higher pK_a values than that of the parent compound.




Fig 4: Dependence of $E_{1/2}$ on σ constants at pH 4.6 and 8.




Fig 5: Dependence of the dissociation constant pK_a on σ constants.

Table 2

Values of reaction constant q from $E_{1/2} - \sigma$ plots for compounds I_{a-e} at different pH's (wave A).

pH	$+ q$	r^a	\pm s.d.
4.0	0.03	0.892	0.002
6.0	0.07	0.901	0.002
8.0	0.06	0.852	0.005

a) Correlation coefficient

b) Standard deviation

Table 3

Values of reaction constant q from plots of $pK_a - \sigma$ sets.

$+ q$			r^a	\pm s.d. ^b
σ	σ^o	σ^+		
+ 1.4	+ 1.4	+ 1.4	0.997	0.007

References

- 1) L.B. Kilgore, J.H. Ford, W.C. Wolfe, Ind. Eng. Chem. 34, 494 (1942).
- 2) D. Molho, Proc. Intern. Conf. Thrombosis and Embolism, Ist, Basel, 193 (1954).
- 3) M. Weiner, B.B. Brodie, J.J. Burns, Proc. Intern. Conf. Thrombosis and Embolism, Ist Basel 181 (1954).
- 4) S. Rosini, L. Trallori, S. Silverstri, Farmaco, Ed. Sci., 31, 315 (1976); C.A., 85, 13972 y (1976).
- 5) J.A. Durden, A.A. Sousa, U.S. Pat. 3, 954, 998 (1976); C.A., 85, 73467 x (1976).
- 6) J. Stradins, "Structure and tautomeric transformations of β -dicarbonyl compounds" ch. 4 "Electrochemical behaviour of cyclic β -diketones" pp. 236-312, Zinatnu Akad. Vestis, Riga, USSR (1977).
- 7) M.A. Morsi, A.H.M. Helmy, H.M. Fahmy, Polarographic and Spectrophotometric Behaviour of Substituted 1,3-Indandione: I. 2-phenylazo-1,3-indandione J. Electroanal. Chem. (In Press).
- 8) M.A. Morsi, A.H.M. Helmy, H.M. Fahmy, Polarographic and Spectrophotometric Behaviour of Substituted 1,3-Indandione: II. 2-(phenyl) acetomidoyl]-1,3-indandione; 1,3-indandione-mono-anil and di-anil J. Electroanal. Chem. (In Press).
- 9) F.A. Amer, E. Afsah, H. Etman, Z. Naturforsch., 34b, 867 (1979).
- 10) H.T.S. Britton, "Hydrogen Ions" Vol. 1, Chapman and Hall, London, p. 365 (1955).
- 11) J.J. Lingane, J. Amer. Chem. Soc. 67, 1916 (1945).
- 12) F. Feigl, Spot Tests, 4th ed., Vol. 2, Elsevier, p. 109(1954).
- 13) R.M. Issa, A.H. Zwait, J. Chem. U.A.R. 14, 161 (1971).
- 14) D. Stocesova, Collect. Czech. Chem. Commun., 14, 615 (1950).

- 15) **J. Pearson**, *Trans. Faraday Soc.*, 44, 683 (1943) & 44, 692 (1948).
- 16) **H. Lund** "Cathodic Reduction of Nitro Compounds" in *Organic Electrochemistry: an Introduction and a Guide*. Ed. by M.M. Baizer, Marcel Dekker Inc., N.Y. Ch. VII, P. 325 (1973).
- 17) **C.P. Retchie, W.F. Sager**, "Progress in Physical Organic Chemistry" Vol. II, Interscience. New York (1964).
- 18) **H.H. Jaffe**, *Chem. Rev.* 53, 1916 (1953).