

PENILE FRACTURE WITH URETHRAL INJURY

(Received 21 December, 1993)

B. Çetinel, M.D.^{} / E. Akkuş M.D.^{**} / O. Arar M.D.^{***} / V. Solok M.D.^{*}**

^{*} Professor, Department of Urology, Cerrahpaşa Faculty of Medicine, İstanbul University, İstanbul, Turkey.

^{**} Instructor, Department of Urology, Cerrahpaşa Faculty of Medicine, İstanbul University, İstanbul, Turkey.

^{***} Research Assistant, Department of Urology, Cerrahpaşa Faculty of Medicine, İstanbul University, İstanbul, Turkey.

SUMMARY

An unusual case of penile fracture (rupture of corpus cavernosum) that accidentally occurred during sexual intercourse is presented. The injury also involved the rupture of the urethra. The need for immediate surgery is emphasized when compared to the complications of conservative treatment.

Key Words: Penile fracture, Urethra, Trauma

INTRODUCTION

Associated rupture of both corpora cavernosa and urethra during coitus is a urologic emergency. When recognized early and surgically repaired, a good functional result regarding micturition and erection can be expected.

CASE REPORT

A 37-year-old man presented with penile swelling, hematoma, pain and deformity 4 hours after sexual intercourse. He also complained of urethral bleeding. 14F Foley catheter was tried to be inserted, but the procedure was unsuccessful with the feeling of "false route". The patient underwent immediate exploration with circumcision incision, and it was recognized that

both right corpus cavernosum and urethra (3 cm and 1,5 cm respectively) were injured (Figure). Evacuation of hematoma, antibiotic irrigation of corpus cavernosum, and repair of corpus cavernosum and the urethra were performed during this emergency operation. The follow-up controls revealed excellent functional results including normal erection and micturition.

DISCUSSION

Penile fracture with urethral injury, is one of the rare urological experiences. It is usually the result of blunt trauma during intercourse or masturbation. Either conservative treatment or immediate surgical exploration are the therapeutic options.(2) Conservative treatment usually results with penile plaque, fibrosis or curvature and perhaps impotence (2,3). Urethral injury accompanying penile fracture cases were treated successfully if immediate surgery was performed.(1,3,4)

In our case with immediate surgical exploration and repair of the injuries, excellent functional results including erection and micturition were achieved. We therefore recommend immediate surgical treatment with circumcision incision which provides excellent exposure in cases of penile fracture with or without urethral injury.



Fig 1. Associated rupture of both corpora cavernosa and urethra were exposed via circumcised approach.


REFERENCES

1. Bitker MO, Berlin J, Jardin A, Chatelain C. 'Faux pas du coit' with associated rupture of both corpora cavernosa and urethra. *Urology* 1988;32:447-448.
2. Tan LB, Chiang CP, Huang CH. Traumatic rupture of the corpus cavernosum. *Br J Urol* 1991;68:626-628.
3. Anselmo G, Fandella A, Faggiano L. Fractures of the penis: Therapeutic approach and long term results. *Br J Urol* 1991;67:509-511.
4. Tsang T, Demby AM. Penile fracture with urethral injury. *J Urol* 1992;147:466-468.

Ürün Bilgisi:

Bileşimi: Trental® 400: Her drajede yavaş salınımlı 400 mg Pentoksifilin Trental® ampul: Her ampulde [5 ml] 100 mg Pentoksifilin. **Eddikasyonları:** Periferik, serebral, göz ve iç kulagın dolaşım bozuklukları. **Kontrendikasyonlar:** Pentoksifilin'e aşırı duyarlılık, masif kanamalar, akut miyokard enfarktüsü, gebelik. **Yan etkileri:** Gastrointestinal yanıklıklar [bulantı, kusma vb], başağrısı, duyarlılık reaksiyonları [kusma, ürtiker vb], ender hipotansiyon. **İlaç etkileşimi:** Hipotansifler, insülin ve glibenklamid ile birlikte kullanımında doz ayarlanması gerekebilir. **Pozoloji:** Trental® 400: Günde üç kez birer draje. Trental ampul: bkz prospektüs. **Ticari şekilleri ve fiyatı:** 20 drajelik kutu: 122.700 TL.. 5 ampullük kutu: 55.400 TL.. [Mart 1994] Reçete ile satılır. Ayrintılı bilgi için firmamızla başvurunuz. Türk Hoechst San. ve Tic. A.Ş. Davutpaşa Cad. No.145 34020 Topkapı - İstanbul.

Hoechst 



etken-madden: pentoksitilin

Trental

®

400

dolaşım bozukluklarında
hemoreolojik ve antitrombositer
tedavi yaklaşımı

Acknowledgement to the Reviewers

We wish to express our deep appreciation to the following reviewers for the time they have spent reviewing manuscripts of Volume VII.

Editorial Board

Akdaş, F. (Ph.D.)	Durmuşoğlu, F. (M.D.)	Sivas, A. (M.D.)
Akoğlu, E. (M.D.)	Esemenli, T. (M.D.)	Söyletir, G. (M.D.)
Akoğlu, T. (M.D.)	Erer, M. (M.D.)	Şehitoğlu, M.A. (M.D.)
Aktan, S. (M.D.)	Erez, S. (M.D.)	Şenvar, A. (M.D.)
Aktan, Ö. (M.D.)	Gökyigit, A. (M.D.)	Şimşek, F. (M.D.)
Ander, H. (M.D.)	Gürmen, N. (M.D.)	Taga, Y. (M.D.)
Asena, U. (M.D.)	Hatemi, H. (M.D.)	Taşyürekli, M. (M.D.)
Aytekin, Y. (Ph.D.)	İlker, Y. (M.D.)	Taner, D. (M.D.)
Babacan, F. (M.D.)	İnce, U. (M.D.)	Tekelioğlu, M. (M.D.)
Barış, İ. (M.D.)	Johansson, C. (Ph.D.)	Tözün, N. (M.D.)
Başaran, M. (M.D.)	Kırkali, Z. (M.D.)	Turgan, Ç. (M.D.)
Cengiz, C. (M.D.)	Külli, S. (M.D.)	Turoğlu, T. (M.D.)
Ceryan, K. (M.D.)	Numanoğlu, A. (M.D.)	Ulusoy, NB. (M.D.)
Ceyhan, N. (M.D.)	Okar, İ. (M.D.)	Uysal, M. (M.D.)
Çamlıbel, T. (M.D.)	Oktay, Ş. (M.D., Ph.D.)	Üneri, C. (M.D.)
Çavdar, S. (Ph.D.)	Özer, F. (M.D.)	Vural, F. (M.D.)
Çavdar, T. (M.D.)	Pekin, S. (M.D.)	Yalaz, K. (M.D.)
Çelebiler, Ö. (M.D.)	Renda, Y. (M.D.)	Yalçın, S. (Ph.D.)
Çırakoğlu, B. (M.D.)	Siva, A. (M.D.)	Yurtkuran, M. (M.D.)
Çolgar, U. (M.D.)		

Author Index for Volume VII

Issue, Page	Issue, Page		
Acan, NL	3: 115	Karaca, C	1: 40
Adalet, I	4: 160	Karakoç, F	3: 111
Afşar, N	1: 22	Kars, A	1: 14
Akan, H	4: 192	Kayhan, B	1: 14
Akkuş, E	4: 196	Kılıç, T	3: 111
Akman, MA	1: 33	Kır, M	1: 10; 2: 86
Akoğlu, E	2: 54; 3: 126	Kışlaoglu, E	4: 155
Aktan, S	1: 22	Korkmaz, H	1: 18
Alican, İ	4: 178	Kutlay, LC	1: 10; 2: 59
Alpdoğan, Ö	3: 126	Lawrence, RA	2: 54
Arar, O	4: 196	Leylek, O	1: 33; 2: 86
Arbak, S	4: 166, 171, 178	Macoska, JA	4: 187
Arda, S	3: 102	Mamikoğlu, B	1: 18; 2: 74
Arsan, B	4: 150	Menteş, BB	2: 81
Artvinli, M	1: 6	Micale, MA	4: 187
Aslan, N	4: 178	Mülazimoğlu, L	2: 78
Atabey, A	1: 40	Nurlu, H	2: 81
Avkan, V	2: 78	Oktay, Ş	4: 178
Aykut, C	1: 22	Ozan, H	2: 88
Baltalı, E	1: 14	Öbek, C	3: 130
Barutçu, A	1: 40	Özdemir, A	4: 155
Başaran, M	3: 111	Özdemir, TK	1: 6
Batman, AÇ	1: 18; 2: 74	Özek, E	3: 111
Belgin, E	3: 106	Özener, Ç	2: 54; 3: 126
Berkman, K	4: 178	Özgür, Ş	3: 126
Bilgehan, A	2: 81	Özyıldan, E	1: 14
Bilgen, H	3: 111	Özyıldan, Ö	1: 14
Bilgin, T	2: 88	Pamir, N	3: 111
Bilsel, S	1: 36	Peker, F	4: 155
Canatay, H	1: 33	Pekin, S	1: 10; 2: 59, 86.
Canbulat, EC	2: 69	Pekin, T	2: 86
Cantez, S	4: 160	Pontes, JE	4: 187
Cantürk, NZ	3: 102	Powell, IJ	4: 187
Cengiz, C	2: 88	Rakunt, C	4: 192
Ceyhan, B	3: 126	Sakr, W	4: 187
Ceyhan, N	2: 59, 86	Sarioğlu, T	3: 106
Çavdar, S	4: 171	Sav, A	2: 74
Çayıcı, B	2: 81	Sıralı, M	4: 160
Çelikel, T	3: 126	Solak, N	2: 64; 4: 150
Çetinel, B	3: 130; 4: 196	Solok, V	3: 130; 4: 196
Dizdar, H	2: 74	Söyletiř, G	1: 27
Durak, N	4: 155	Şehitoğlu, MA	2: 74
Dülger, M	3: 102	Şen, F	4: 178
Eker, S	4: 160	Şenel, A	4: 192
Emerk, K	1: 36	Tanık, A	4: 192
Ener, B	3: 133	Taşer, B	2: 78
Erbengi, T	4: 166	Telatar, H	1: 14
Ercan, F	4: 178	Tellaloğlu, S	3: 122
Erdiš, E	2: 64; 4: 150	Tekuzman, G	1: 14
Erdoğru, T	3: 122	Tezcan, EF	3: 115
Erenus, M	1: 10, 33; 2: 59, 90	Tolunay, Ş	2: 88
Erkilinci, Ş	2: 64; 4: 150	Topeli, A	1: 14
Ersoy, A	4: 160	Tutkun, A	1: 18; 2: 74
Ersöz, Ö	2: 54	Türkalp, I	2: 69; 4: 146
Erşahin, Ç	1: 36	Türkeri, LN	4: 187
Ertem, D	3: 111	Türközkan, N	2: 81
Eskitürk, A	1: 27	Ütkah, NZ	3: 102
Ferahkőse, Z	2: 81	Üneri, C	1: 18; 2: 74
Fırat, D	1: 14	Üstün, M	2: 64; 4: 150
Gençoğlu, A	1: 6	Vahaboğlu, MH	2: 78
Göğüş, Y	1: 33	Yalçın, O	4: 160
Gözükara, İ	3: 119	Yalçiner, A	1: 36
Grignon, DJ	4: 187	Yalın, A	4: 171
Gülkilik, A	2: 64; 4: 150	Yeğen, BC	4: 178
Gürsel, B	3: 106	Yenilmez, E	4: 166
Hergenç, G	3: 119	Yıldırım, I	2: 78
Kadioğlu, A	3: 122	Yüksel, F	4: 155
Kaptanağası, A	4: 146	Wolman, SR	4: 187

Contents of Volume VII

Is Hypocapnia an Important Factor in Exercise - Induced Asthma? T.K. Özdemir / M. Artvinli / A. Gençoğlu	6
Comparison of the Efficacy of Lisuride Versus Bromocriptine in the Treatment of Hyperprolactinemia M. Erenus / M. Kir / L.C. Kutlay / S. Pekin	10
Prevalence of Hepatitis C Virus Antibody in Cancer Patients from Turkey A. Topeli / Ö. Özyikan / E. Özyikan / A. Kars / G. Tekuzman E. Baltalı / B. Kayhan / H. Telatar / D. Fırat	14
The Effect of Antibiotic Therapy after Tonsillectomy A.Ç. Batman / A. Tutkun / B. Mamikoğlu / H. Korkmaz / C. Üneri	18
Risk Factors and Specific Stroke Types of Cerebrovascular Accident Cases at Marmara Medical School Neurology Department N. Afşar / S. Aktan / C. Aykut	22
Isolation Rate of Unusual Bacterial Pathogens from Stool Cultures A. Eskitürk / G. Söyletir	27
Single Dose Intravenous Ondansetron in the Prophylaxis of Postoperative Nausea and Vomiting M. Erenus / Ö. Leylek / H. Canatay / M.A. Akman / Y. Göğüş	33
Effect of Defibrotide on Cultured Endothelial Cell Cycle Progression Analysed by Flow Cytometry Ç. Erşahin / S. Bilsel / A. Yalçiner / K. Emerk	36
Facial Cleft and Maxillary Duplication A. Barutçu / C. Karaca / A. Atabay	40
Erythropoietin in the treatment of anaemia of chronic renal failure I.Ç. Özener / R.A. Lawrence / Ö. Ersöz / E. Akoğlu	54
The value of nonstress test in the assessment of fetal well being before 34 weeks of gestation - A prospective analysis M. Erenus / L. Kutlay / N. Ceyhan / S. Pekin	59
Color Doppler sonography in the differential diagnosis of ovarian masses M. Üstün / Ş. Erkilinç / A. Gülkılık / N. Solak / E. Erdiş	64
Acute myocardial infarction risk predicted by serum levels of lipoprotein (a) in Turkish population E.C. Canbulat / I. Türkålpa	69
The distribution of the nucleolar organiser regions in the dysplastic epithelial lesions and in the epidermoid carcinomas of the vocal cords Ç. Batman / A. Tutkun / A. Sav / B. Mamikoğlu / H. Dizdar C. Üneri / M.A. Şehitoğlu	74
Nasopharyngeal carriage rate and antimicrobial resistance of Haemophilus influenzae in İstanbul - Türkiye M.H. Vahaboglu / L. Mülazimoğlu / I. Yıldırım / V. Avkan / B. Taşer	78
Protective effects of verapamil on experimental ischemia - reperfusion - induced liver injury in cirrhotic rats Z. Ferahköse / H. Nurlu / B.B. Menteş / N. Türközkan / A. Bilgehan / B. Çaycı	81
An intraabdominal troublemaker: Endometrioma Ö. Leylek / N. Ceyhan / T. Pekin / M. Kir / S. Pekin	86
Mature cystic teratoma with malignant degeneration T. Bilgin / C. Cengiz / Ş. Tolunay / H. Ozan	88
Ovarian cancer and reproduction M. Erenus	90

Alteration of complement levels following thermal injury and surgery S. Arda / N.Z. Cantürk / N.Z. Utkan / M. Dülger	102
The effect of glycerol on static acoustic impedance in Meniere's disease T. Saroğlu / B. Gürsel / E. Belgin	106
Transcranial Doppler sonography: Physiological influences on cerebral hemodynamics E. Özak / D. Ertem / F. Karakoç / T. Kılıç / H. Bilgen N. Pamır / M. Başaran	111
Inhibition of brain glutathione reductase by sodium arsenite N.L. Acan / E.F. Tezcan	115
Osteocalcin levels in Graves' disease and euthyroid controls G. Hergenç / İ. Gözükara	119
Intrapenile arterial anatomic details examined by color Doppler ultrasonography A. Kadıoğlu / T. Erdoğru / S. Tellaloğlu	122
Goodpasture's syndrome mimicking idiopathic pulmonary hemosiderosis B. Ceyhan / T. Çelikel / E. Akoğlu / Ç. Özener / Ö. Alpdoğan / Ş. Özgün	126
Bulbous urethral cyst in a man B. Çetinel / C. Öbek / V. Solok	130
Candida infections in ENT B. Ener	133
Plasma lipid peroxide levels in type II diabetics : Relationship with long term diabetic complications I. Türkalp / A. Kaptanagasi	146
Color Doppler sonography in the differential diagnosis of uterine masses M.Üstün / Ş. Erkilinç / A. Qulkilik / N. Solak / E. Erdiş / B. Arsan	150
Gunshot wounds to the face N. Durak / E. Kişlaoglu / F. Yüksel / F. Peker / A. Özdemir	155
Radiouclide hysterosalpingography : Does it have a role in the management of female infertility? I. Adalet / O. Yalçın / A. Ersoy / S. Eker / M. Sıralı / S. Cantez	160
Effects of nickel sulphate on rat testes seminiferous tubules : An ultrastructural investigation E. Yenilmez / S. Arbak / T. Erbengi	166
Vincula tendineum and vascularisation of the flexor pollicis longus tendon A. Yalın / S. Çavdar / S. Arbak	171
Stress ulcer : A morphological approach S. Arbak / F. Şen / F. Ercan / İ. Alican / N. Aslan B.Ç. Yeğen / S. Oktay / K. Berkman	178
Combined polymerase chain reaction (PCR) and fluorescence in situ hybridization (FISH) analysis of Y chromosome loss related to age and disease status in prostate cancer L.N. Türkeri / W. Sakr / S.R. Wolman / M.A. Micale / D.J. Grignon I.J. Powell / J.E. Pontes / J.A. Macoska	187
Pycnodysostosis : Report of two cases A. Şenel / C. Rakunt / H. Akan / A. Tanık	192
Penile fracture with urethral injury B. Çetinel / E. Akkuş / O. Arar / V. Solok	196
Acknowledgement to the Reviewers	
Author index to Volume VII	
Contents of Volume VII	

MARMARA MEDICAL JOURNAL INSTRUCTIONS TO AUTHORS

1. Manuscripts, letters and editorial correspondence should be sent to 'Editor, Marmara Medical Journal, Marmara University, Faculty of Medicine, Istanbul-Turkey' by first class mail (airmail for overseas).
2. Submissions considered for publication are received with the understanding that no part of the submission has previously appeared elsewhere in any but abstract form.
3. Manuscripts should be typed double-spaced on standard size typewriter paper with margins of at least 2.5 cm. This includes references, tables and figure-legends. The original typescript and two high-quality copies of the manuscripts should be submitted.
4. Number pages consecutively in order and place author(s) name, highest degree, institutional affiliations and address below the title.
5. Marmara Medical Journal invites papers on original research, case reports, reviews, short communications for practical applications, letters, editorials, book reviews and announcements. The number of typewritten pages should not exceed 10 for original articles, 12 for reviews, 4 for case reports and 1 for letters.
6. Original articles and research papers should normally be divided into following sections:
A. (1) An informative summary for not more than 200 words must be included and should appear at the beginning of the paper. (2) Key words. (3) Introduction. (4) Materials and Methods. (5) Results. (6) Discussion and (7) References.
B. References must be typed in double spacing and numbered consecutively as they are cited. The style of references is that of the Index Medicus. List all authors when there are six or fewer, when there are seven or more, list the first three, then 'et al'. Sample references follow:
 1. Steward JH, Castaldi PA. Uremic bleeding: a reversible platelet defect corrected by dialysis. QJ Med. 1967; 36 : 409 - 23.
 2. Bearn AG. Wilson's Disease. In: Stanbury JB, Wyngaarden JB, Fredrickson DS, eds. The metabolic basis of inherited disease. New York : McGraw - Hill, 1972: 103-50.
7. Tables should be as few as possible and should include only essential data. Tables should be typed in double spacing on separate sheets and have a legend for each. Diagrams or illustrations should be drawn with black Indian ink on white paper and should be given Roman numerals. Each illustration should be accompanied by a legend clearly describing it. All legends should be grouped and typewritten (double spaced) on a separate sheet of paper. Photographs and photomicrographs should be unmounted high-contrast glossy black-on-white prints and should not be retouched. Each photograph or illustration should be marked on the back with the name(s) of the author(s), should bear an indication of sequence number and the top should be marked with an arrow. All measurements should be given in metric units.
8. Manuscripts are examined by the editorial board usually sent to outside referees. The editor reserves the right to reject or to return the manuscript to the author(s) for additional changes if all the guidelines and requirements are not uniformly completed. Only two copies of the rejected papers are returned to the author(s).
9. Proofs will be submitted to the author responsible for proof correction and should be returned to the editor within 5 days. Major alterations from the text cannot be accepted.
10. Correspondence and communications regarding manuscripts and editorial material subscriptions and payments should be sent to:

The Editor
Marmara Medical Journal
Marmara University, Faculty of Medicine
Haydarpaşa - İstanbul.