



Analysis of trace elements accumulation in some landscape plants as an indicator of pollution in an urban environment: Case of Ankara

Elnaji Abdulaziz Ahmida Saleha^b, Öznur Işınkaralar^{b,*}

^aDepartment of Geography College of Education, Al-Qubba Branch, University of Derna, Libya

^bDepartment of Landscape architecture, Faculty of Engineering Architecture, Kastamonu University, Kastamonu, Türkiye

*Corresponding Author: obulan@kastamonu.edu.tr

Received: March 16, 2022 ◆ Accepted: June 10, 2022 ◆ Published Online: June 27, 2022

Abstract: As a result of the increase in structure and population densities in urban areas, many problems arise in cities that need to be solved. With the effect of keeping the planned approaches to air pollution in the city's background, urban air pollution has started to feel serious with the adverse effects it has created on living things. Heavy metals, which are one of the components of air pollution, seriously threaten living things because they cannot be eliminated from metabolism. For this reason, heavy metal pollution should be monitored in order to make the necessary interventions. The research aims to interpret the trace element level by using *Cupressus sempervirens* L. *Thuja occidentalis* L. *Platanus orientalis* L. *Prunus cerasifera* Ehrh. and *Photinia serrulata* Lindl. plants in the city center of Ankara as biomonitors. In this context, variance analysis was performed on lead and cadmium concentration levels with the SPSS package program. The mean values were subjected to the Duncan test for multiple comparison purposes. The difference in the accumulation levels of plant species and organs was revealed due to the research.

Keywords: Urban Air Quality, Trace Elements, Accumulation, Biomonitoring, Ankara

Öz: Kentsel alanlarda yapı ve nüfus yoğunluklarının zamanla artmasının bir sonucu olarak kentlerde pek çok çözülmesi gereken sorun ortaya çıkmaktadır. Kentlerde hava kirliliğine yönelik plan yaklaşımlarının geri planda tutulmasının etkisiyle kentsel hava kirliliği, kendini canlıların üzerinde yarattığı olumsuzluklar ile ciddi bir şekilde hissettirmeye başlamıştır. Hava kirliliği bileşenlerinden biri olan ağır metaller metabolizmadan tamamen atılmadığından canlı yaşamını ciddi bir şekilde tehdit eder. Bu nedenle gerekli müdahalelerin yapılabilmesi için ağır metal kirliliğinin izlenmesi gerekmektedir. Araştırmada Ankara kent merkezinde yer alan *Cupressus sempervirens* L. *Thuja occidentalis* L. *Platanus orientalis* L. *Prunus cerasifera* Ehrh. *Photinia serrulata* Lindl. Bitkilerinin biyomonitör olarak kullanılmasıyla trace element düzeyinin yorumlanması amaçlanmaktadır. Bu bağlamda kurşun ve kadmiyum konsantrasyon düzeylerine SPSS paket programıyla varyans analizi yapılmıştır. Ortalama değerler çoklu karşılaştırma yapılması amacıyla Duncan teste tabi tutulmuştur. Araştırma sonucunda, bitki türü ve organların birikim düzeylerinin farklılığı ortaya konmuştur.

Anahtar Kelimeler: Kentsel Hava Kalitesi, Eser elementler, Birikim, Biyoizleme, Ankara

1. Introduction

Urban researches are studies that need to be handled multidimensionally, such as transportation [1-3], urban design [4-8], urban economy and development [9-11], landscape planning [12, 13], urban policy [14, 15] environment and infrastructure [16-18]. Increasing the quality of life in cities, which constitutes the main stage of daily activities in today's conditions, depends on determining the sources of the problems and interventions [19]. The main reasons for reducing the quality of urban life can be counted as population density, various industrial activities, use of poor quality fuel, the density of traffic, and the use of chemical products harmful to nature in many areas [20-23]. Problems such as climate change, global warming, and air pollution that arise from these problems affect all ecological systems [24, 25]. Air pollution, which is one of the problems that occur, is highly affected by the release from sources, the behavior of pollutants in the atmosphere, their transport by wind, and the silhouette and morphology of the city that cause their accumulation [26, 27]. In this context, air pollution research is of vital importance in terms of urban planning.

The accumulation of air pollutants on various living things can represent air quality on an urban scale. To determine the effect of air quality on living things, it has become very important to determine how far they have been moved and their accumulation levels, starting from their source [28]. However, the detection of pollutants in metropolitan cities is a complex research topic [29]. Among the main reasons, many externalities, such as the fact that it cannot be followed instantly, cannot be detected at the source, and the effect of weather events (such as meteorological movements) on the accumulation affects the atmospheric accumulation. Therefore, various studies are being carried out to use some species as biomonitors since it is difficult to monitor and the cost is high. Furthermore, heavy metals from these pollutants are

hazardous in high concentrations as they can have a toxic effect [30]. However, the direct measurement of heavy metals in the air does not provide enough information in terms of environmental impact and monitoring [31, 32].

Some materials such as mosses, indoor plant species, and the rings and organs of perennial trees used in landscape afforestation were used in some studies. High concentrations of metals that can accumulate in these species have been found. However, they have found that various heavy metals can get in multiple areas at a higher rate than others. For example, the differences between the element types found at the industrial sites and the metals found in the areas with high traffic indicate that passive sampling is appropriate in the search for a method. These papers have shown the absorption capacity of accumulation varies between species [33, 34]. Therefore, it is essential to investigate the suitability of the species that can be preferred as biomonitors for the selection of target pollutants. This study was aimed to determine the accumulation levels of Lead (Pb) and Cadmium (Cd) in the organs of 5 different landscape plants in areas with heavy traffic in the city center of Ankara.

2. Material and Method

This paper is conducted on the species grown in the urban center of Ankara, where the traffic is dense. In selecting the study area, the traffic's multi-port and low flow rate were taken into account. Samples were collected from *Cupressus sempervirens* L. (SCs), *Thuja occidentalis* L. (STo), *Platanus orientalis* L. (SPo), *Prunus cerasifera* Ehrh. (SPc) and *Photinia serrulata* Lindl. (SPs) plant, which is frequently used in landscape studies. Samples are from the last year's exile. They were collected in late October towards the end of the vegetation season of 2021, bagged and labeled, and brought to the laboratory. It was then dissected into organs without the use of metal tools. Some of the bark and leaves have been washed. After rinsing with distilled water, all samples were labeled.

In the next stages of the study, The US EPA 3052 was used to apply the method. Organs, leaf, bark and wood samples were also shown with their codes. Labeled samples were kept at room conditions for two weeks after pretreatment without being exposed to direct sunlight. It was then dried in an oven at 45°C for two weeks.

The dried plant samples were ground into powder and weighed 0.5 g, and placed in tubes designed for microwaving. 10 mL of 65% HNO₃ was added to the samples. The prepared samples were then burned in a microwave device at 280 PSI and 180 °C for 20 minutes. After the processes were completed, the tubes were removed from the microwave and left to cool. Deionized water was added to the cooled samples to make up to 50 ml. After the prepared samples were filtered through filter paper, they were read at appropriate wavelengths in the ICP-OES device. The obtained values were multiplied by the dilution factor, and the element concentrations of the study were calculated.

SPSS program was used to compare the mean values of the analyses performed with three repetitions. Analysis of variance was applied to the data. Homogeneous groups were obtained by applying the Duncan test to statistically different groups, at least at the 95% confidence level. The analysis results are expressed with F values in terms of the storage capacity of the plants.

3. Results

When the high concentration Pb values obtained within the scope of the study were examined among the species, high values were obtained in SPc in Table 1. Although the lowest accumulation among the organs was in the wood, the accumulation level in SPc was relatively high with 6726 ppb. While the lowest value in wood was obtained with 877.66 ppb in SPo, it was found at 6995.2 ppb in outer bark and 6255.53 ppb in leaf. When ranking from high to low in terms of the values of Pb among the organs, they are listed as outer bark, leaf, and wood.

Table 1. Deposition of Pb concentrations (ppb) based on organs

Species	Organ			F value
	Leaf	Bark	Wood	
SCs	8738.46 Cde	8881.60 De	2448.66 Ag	32.9***
STo	8702.00 Bcd	8808.40 Dd	1097.86 Ac	51.5***
SPo	6255.53 Cb	6995.20 Ea	877.66 Aa	1156.5***
SPc	9176.46 Ei	8806.33 Dd	6726.00 Ak	
SPs	8843.66 Eg	8688.33 Bc	2092.13 Af	
F value	9169.43***	2338.07***	16905.84***	

*** expresses $p < 0.001$, lower case letters represent vertical directions, whereas capital letters represent horizontal directions.

When the concentration values in the leaf were compared, the highest value was obtained with SPs 28221.93 ppb, while the lowest value was obtained with STo 7953.20 ppb. Regarding values in the crust, the highest accumulation level was observed in SPo at 15712.86 ppb, while the minor accumulation was observed in SPc at 2940.06 ppb. When the accumulation levels in wood were compared, the accumulation in STo was analyzed, with 1842 ppb being the lowest.

Table 2. Deposition of Cd concentrations (ppb) based on organs

Species	Organ			F value
	Leaf	Bark	Wood	
SCs	9447.86 Ee	6525.33 Bd	2612.73 Ae	18034.62***
STo	7953.20 Dd	8805.86 Ei	1842.00 Ab	23105.08***
SPo	17015.13 EI	15712.86 DI	8109.20 An	8552.04***
SPc	19624.06 Dm	2940.06 Ba	2196.66 Ad	35244.1***
SPs	28221.93 Dn	7308.60 Bf	3185.46 Af	33703.61***
F value	44078.52***	13783.27***	30418.68***	

*** expresses $p < 0.001$, lower case letters represent vertical directions, whereas capital letters represent horizontal directions.

4. Discussion and Conclusion

Heavy metal accumulation can cause serious health problems, a multidimensionality researched issue. For example, Conti et al. [35] determined the element levels in bees and edible hive products and interpreted the health risk status with reference doses. Another study analyzed the variation in Zn and Cd levels by exposing algae to cigarette smoke in the living area and investigated its effect on indoor air quality [36]. Therefore, the accumulation of toxic substances in living things can be handled in many contexts.

Heavy metal accumulation was investigated in 5 different plant species selected in the study. Işınkaralar [37] investigated the Cd concentration in the range of 120-250 ppb in *Ailanthus altissima* in his research conducted in another location in Ankara. In addition, the outer bark values of Cd are about twice that of wood. Similarly, a higher concentration was obtained in the bark than in the wood in all species in this study. However, *Thuja occidentalis* L. bark accumulation was approximately four times higher than in wood. Another study [38] compared the heavy metal uptake capacity of 14 species, which also followed the accumulation of Cd. *Chamaecyparis lawsoniana* was found to be quite suitable for monitoring. According to Turkyılmaz et al. [34], the change in Cd amounts according to traffic density was statistically significant at the 99.9% confidence only in the usage of *S. babylonica* according to their analysis of 4 different species. The highest Pb values were obtained in *A. hippocastanum*. Various studies reveal that traffic density and industrial emissions affect Pb and Cd concentrations [39]. Zinikovskaia et al. [40] used the algae biomonitoring technique in air pollution monitoring. Possible sources of air pollution in the case area, including Pb and Cd and other nine metals, have been identified as resuspension of soil particles, agricultural applications, vehicles, industry, and thermal power plants.

5. Suggestions

The most obvious result of the decrease in air quality in cities every day is seen in living things. Growth disorders, stress conditions, and early deaths, especially seen in landscape plants in urban areas, are the leading causes of toxic metals at very high levels. Although the ability of trees used in urban areas to absorb heavy metals in their bodies differs between species, it has been observed that all of them have the ability to absorb. The presence of Pb and Cd-containing metals, which are among the species' traffic-induced releases, is increasing due to the formation of both fuel-oriented and motor vehicles caused by wear and corrosion. More comprehensive information about the sources can be obtained by diversifying the locations in future studies.

Competing Interest / Conflict of Interest

The authors declare that they have no competing interests.

Funding

Not applicable

Author Contribution

We declare that all Authors equally contribute.

6. References

- [1] Cleophas, C., Cottrill, C., Ehmke, J. F., & Tierney, K. (2019). Collaborative urban transportation: Recent advances in theory and practice. *European Journal of Operational Research*, 273(3), 801-816.
- [2] Glazener, A., Sanchez, K., Ramani, T., Zietsman, J., Nieuwenhuijsen, M. J., Mindell, J. S., ... & Khreis, H. (2021). Fourteen pathways between urban transportation and health: A conceptual model and literature review. *Journal of Transport & Health*, 21, 101070.

- [3] Taylor, M. A. (2008). Critical transport infrastructure in Urban areas: impacts of traffic incidents assessed using accessibility-based network vulnerability analysis. *Growth and Change*, 39(4), 593-616.
- [4] Buttazzoni, A., Dean, J., & Minaker, L. (2022). Urban design and adolescent mental health: A qualitative examination of adolescent emotional responses to pedestrian-and transit-oriented design and cognitive architecture concepts. *Health & Place*, 76, 102825.
- [5] Naboni, E., Natanian, J., Brizzi, G., Florio, P., Chokhachian, A., Galanos, T., & Rastogi, P. (2019). A digital workflow to quantify regenerative urban design in the context of a changing climate. *Renewable and Sustainable Energy Reviews*, 113, 109255.
- [6] Şahin, F., Öztürk, S., & Işınkaralar, Ö. (2019). An Assessment about Barrier Free Design in Urban Green Areas. In *Academic Studies* p. 289.
- [7] Tian, Y., Gu, K., & Wei, T. (2022). Preserving the cultural identity of Chinese cities in urban design through a typomorphological approach. In *Urban Morphology, Architectural Typology and Cities in Transition* (pp. 172-198). EDP Sciences.
- [8] Öztürk, S., Işınkaralar, Ö., & Ayan, E. (2018). Visibility Analysis in Historical Environments: The case of Kastamonu Castle and its Surrounding. *Journal of Current Researches on Social Sciences*, 8(4), 405-412.
- [9] Işınkaralar, Ö., and Varol, C. (2021). Kent Merkezlerinde Ticaret Birimlerin Mekansal Örüntüsü Üzerine Bir Değerlendirme: Kastamonu Örneği. *Mimarlık Bilimleri ve Uygulamaları Dergisi*, 6(2), 396-403.
- [10] Öztürk, S., Işınkaralar, Ö., & Yılmaz, D. (2019). Cittaslow Hareketi ve Toplumsal Farkındalık: Gerze (Sinop) Örneği. *Academic Studies on Social and Education Sciences*.
- [11] Öztürk, S., Işınkaralar, Ö., & Kesimoğlu, F. (2021). An Assessment on Shopping Centers as Consumption Place. *Kastamonu University Journal of Engineering and Sciences*, 7(1), 65-73.
- [12] Öztürk, S., İsmail, T. S. Y., & Işınkaralar, Ö. (2020). İnanç Turizmi Odağında Benli Sultan Külliyesi Peyzaj Tasarım Önerileri. *Türk Turizm Araştırmaları Dergisi*, 4(1), 349-360.
- [13] Hüsam, A., Öztürk, S., & Dönmez, Y. (2021). Parkların Peyzaj Mimarlığı Açısından İncelenmesi: Karabük Kent Merkezi Örneği. *Journal of Humanities and Tourism Research*, 11(2), 339-346.
- [14] Yılmaz, D., & Işınkaralar, Ö. Climate Action Plans Under Climate-Resilient Urban Policies. *Kastamonu University Journal of Engineering and Sciences*, 7(2), 140-147.
- [15] Li, L., Ma, S., Zheng, Y., & Xiao, X. (2022). Integrated regional development: Comparison of urban agglomeration policies in China. *Land Use Policy*, 114, 105939.
- [16] Kanai, J. M., & Schindler, S. (2022). Infrastructure-led development and the peri-urban question: Furthering crossover comparisons. *Urban Studies*, 00420980211064158.
- [17] Kandpal, V., Tyagi, V., & Kaur, H. (2022). Smart City: Sustainable City for Tackling Urban Challenges. In *Infrastructure Planning and Management in India* (pp. 147-157). Springer, Singapore.
- [18] Loures, L. (2015). Post-industrial landscapes as drivers for urban redevelopment: Public versus expert perspectives towards the benefits and barriers of the reuse of post-industrial sites in urban areas. *Habitat International*, 45, 72-81.
- [19] Bennedjai, R., & Bencherif, M. (2022). Local Urban Management for Improving the Quality of Life in Algerian Collective Housing Estates. *Journal of Urban Planning and Development*, 148(2), 05022003.
- [20] Majewska, A., Denis, M., Jarecka-Bidzińska, E., Jaroszewicz, J., & Krupowicz, W. (2022). Pandemic resilient cities: Possibilities of repairing Polish towns and cities during COVID-19 pandemic. *Land Use Policy*, 113, 105904.
- [21] Jha, R., Dwivedi, S., & Modhera, B. (2022). Measurement and practices for hazardous waste management. In *Hazardous Waste Management* (pp. 89-115). Elsevier.
- [22] Aydar, E., & Akkaş, E. (2022). The emission of natural harmful particulate matters by wind erosion and possible impact areas, Cappadocia province, Central Anatolia, Turkey. *Bulletin of Engineering Geology and the Environment*, 81(1), 1-22.
- [23] Forouhar, N., Forouhar, A., & Hasankhani, M. (2022). Commercial gentrification and neighbourhood change: A dynamic view on local residents' quality of life in Tehran. *Land Use Policy*, 112, 105858.
- [24] Fuller, R., Landrigan, P. J., Balakrishnan, K., Bathan, G., Bose-O'Reilly, S., Brauer, M., ... & Yan, C. (2022). Pollution and health: a progress update. *The Lancet Planetary Health*.
- [25] Grifoni, M., Franchi, E., Fusini, D., Voccianta, M., Barbafieri, M., Pedron, F., ... & Petruzzelli, G. (2022). Soil Remediation: Towards a Resilient and Adaptive Approach to Deal with the Ever-Changing Environmental Challenges. *Environments*, 9(2), 18.
- [26] Ukaogo, P. O., Ewuzie, U., & Onwuka, C. V. (2020). Environmental pollution: causes, effects, and the remedies. In *Microorganisms for sustainable environment and health* (pp. 419-429). Elsevier.
- [27] Tian, Y., Yao, X., & Chen, L. (2019). Analysis of spatial and seasonal distributions of air pollutants by incorporating urban morphological characteristics. *Computers, Environment and Urban Systems*, 75, 35-48.

- [28] Matthaios, V. N., Kramer, L. J., Sommariva, R., Pope, F. D., & Bloss, W. J. (2019). Investigation of vehicle cold start primary NO₂ emissions inferred from ambient monitoring data in the UK and their implications for urban air quality. *Atmospheric Environment*, 199, 402-414.
- [29] Peng, Y., Fang, W., Krauss, M., Brack, W., Wang, Z., Li, F., & Zhang, X. (2018). Screening hundreds of emerging organic pollutants (EOPs) in surface water from the Yangtze River Delta (YRD): occurrence, distribution, ecological risk. *Environmental Pollution*, 241, 484-493.
- [30] Vardhan, K. H., Kumar, P. S., & Panda, R. C. (2019). A review on heavy metal pollution, toxicity and remedial measures: Current trends and future perspectives. *Journal of Molecular Liquids*, 290, 111197.
- [31] Isinkaralar, K. (2022). Temporal Variability of Trace Metal Evidence in *Cupressus arizonica*, *Platanus orientalis*, and *Robinia pseudoacacia* as Pollution-Resistant Species at an Industrial Site. *Water Air Soil Pollut* 233, 250.
- [32] Kang, S., Zhao, K., Yu, D. G., Zheng, X., & Huang, C. (2022). Advances in biosensing and environmental monitoring based on electrospun nanofibers. *Advanced Fiber Materials*, 1-32.
- [33] Chen, D., Liu, X., Bian, R., Cheng, K., Zhang, X., Zheng, J., ... & Li, L. (2018). Effects of biochar on availability and plant uptake of heavy metals—A meta-analysis. *Journal of Environmental Management*, 222, 76-85.
- [34] Turkyilmaz, A., Cetin, M., Sevik, H., Isinkaralar, K., & Saleh, E. A. A. (2020). Variation of heavy metal accumulation in certain landscaping plants due to traffic density. *Environment, Development and Sustainability*, 22(3), 2385-2398.
- [35] Conti, M. E., Astolfi, M. L., Finoia, M. G., Massimi, L., & Canepari, S. (2022). Biomonitoring of element contamination in bees and beehive products in the Rome province (Italy). *Environmental Science and Pollution Research*, 1-18.
- [36] Świsłowski, P., Śmiechowicz, B., & Rajfur, M. (2022). Effects of tobacco smoke on indoor air quality: the use of mosses in biomonitoring. *Journal of Environmental Health Science and Engineering*, 1-9.
- [37] Isinkaralar, K. (2022). The large-scale period of atmospheric trace metal deposition to urban landscape trees as a biomonitor. *Biomass Conversion and Biorefinery*, 1-10.
- [38] Isinkaralar, K., Koc, I., Erdem, R., & Sevik, H. (2022). Atmospheric Cd, Cr, and Zn deposition in several landscape plants in Mersin, Türkiye. *Water, Air, & Soil Pollution*, 233(4), 1-10.
- [39] Isinkaralar, K. (2022). Atmospheric deposition of Pb and Cd in the *Cedrus atlantica* for environmental biomonitoring. *Landscape and Ecological Engineering*, 1-10.
- [40] Zinicovscaia, I., Chaligava, O., Yushin, N., Grozdov, D., Vergel, K., & Hramco, C. (2022). Moss Biomonitoring of Atmospheric Trace Element Pollution in the Republic of Moldova. *Archives of Environmental Contamination and Toxicology*, 82(3), 355-366.