

INTEGRALS AND A MASCHKE-TYPE THEOREM FOR WEAK HOPF π -COALGEBRAS

Chen Quan-guo and Wang Shuan-hong

Received: 3 October 2010; Revised: 14 February 2011

Communicated by Stefaan Caenepeel

ABSTRACT. Let π be a discrete group. We shall introduce the more general concept of an integral of a weak Doi-Hopf π -datum (H, A, C) , where H is a weak Hopf π -coalgebra coacting on an algebra A and acting on a π -coalgebra $C = \{C_\alpha\}_{\alpha \in \pi}$. We prove that there exists a total integral $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha-1}, A)\}_{\alpha \in \pi}$, then any representation of (H, A, C) is injective in a functorial way, as a corepresentation of C and vice versa. As the application of the existence of a total integral, we prove the Maschke-type Theorem for weak Doi-Hopf π -modules.

Mathematics Subject Classification(2010): 16T05

Keywords: weak Hopf π -coalgebra, integral, weak Doi-Hopf π -datums

1. Introduction

The category ${}^C\mathcal{U}(H)_A$ of Doi-Hopf modules over the bialgebra H was introduced in [9]. It is the category of the modules over the algebra A which are also comodules over the coalgebra C and satisfy certain compatibility condition involving H . The study of ${}^C\mathcal{U}(H)_A$ turned out to be very useful: it was shown in [3, 9] that many categories such as the module and comodule categories over bialgebras, the Hopf modules category [13], and the Yetter-Drinfeld category [11, 25] are special cases of ${}^C\mathcal{U}(H)_A$. Many results known for module categories over bialgebras or Hopf algebras were generalized to Hopf group-coalgebra [21].

Hopf group-algebras appeared in the work of Turaev [14] on homotopy quantum field theories as a generalization of ordinary Hopf algebras. Let us note that there exists a symmetric monoidal category, the so-called Turaev category, constructed by Caenepeel and De Lombaerde [4] the Hopf algebras which are the same as Hopf group-coalgebras. A purely algebraic study of Hopf group-coalgebras can be found in the references Virelizier ([19, 20]), Wang ([21-24]) and Zunino ([26-27]).

The work was partially supported by the NNSF of China (No.10871042) and the NNSF of Jiangsu Province (No.BK2009258).

As a generalization of ordinary Hopf group-coalgebras and weak Hopf algebras [1], weak Hopf group-coalgebras were studied in the work of Van Daele and Wang [18]. Weak Hopf group-coalgebras not only provide examples of weak multiplier Hopf algebras ([17, 18]), but also provide an approach to construct a class of new braided crossed categories in the sense of Turaev. A purely algebraic study of weak Hopf group-coalgebras can be found in [17]. Many results in Hopf group-coalgebra had been generalized to weak Hopf group-coalgebra. For example, the Fundamental Theorem of weak Hopf group-comodules was given in [17]. In the sequel, the fundamental theorem had been generalized to weak relative Hopf group-comodules in [12].

In this paper, we generalize the definition of Doi-Hopf π -modules to the case when H is a weak Hopf group-coalgebra and develop the theory of weak Doi-Hopf π -modules.

This article is organized as follows:

In Section 2, we recall definitions and basic results related to separable functors and (weak) Hopf group-coalgebras and in Section 3, we recall some important adjoint functors and introduce some weak Doi-Hopf π -modules.

In Section 4, inspired by the idea adopted by Menini and Militaru ([10]) or Caenepeel et al.([5]), we introduce the notion of integral for weak Doi-Hopf π -datums and prove that if there exists $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ a total integral, then the natural transformation $\rho : F_A \circ 1_{\pi - c_{\mathcal{U}(H)_A}} \rightarrow F_A \circ G \circ F^C$ splits. Conversely, if the natural transformation $\rho : F_A \circ 1_{\pi - c_{\mathcal{U}(H)_A}} \rightarrow F_A \circ G \circ F^C$ splits, then there exists $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ a total integral(see Theorem 4.4). The application of the integral is also considered (see Theorem 4.7).

In Section 5, we prove the Maschke-type theorem for weak Doi-Hopf π -modules (see Theorem 5.2).

2. Preliminaries

Throughout this paper, we always let π be a discrete group with a neutral element e and k a field. If U and V are k -spaces, $T_{U,V} : U \otimes V \rightarrow V \otimes U$ will denote the flip map defined by $T_{U,V}(u \otimes v) \rightarrow v \otimes u$, for all $u \in U$ and $v \in V$.

2.1. Separable Functors. Separable functors were introduced in [8]: futher applications and properties have been discussed in [28]and [6]. Separable functors of the second kind were introduced in [7]. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ and $\mathcal{H} : \mathcal{C} \rightarrow \mathcal{E}$ be covariant functors. We then have functors

$$\text{Hom}_{\mathcal{C}}(\bullet, \bullet), \text{Hom}_{\mathcal{D}}(F(\bullet), F(\bullet)), \text{Hom}_{\mathcal{E}}(\mathcal{H}(\bullet), \mathcal{H}(\bullet)) : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \underline{\text{Sets}}$$

and natural transformations

$$\mathcal{F} : \text{Hom}_{\mathcal{C}}(\bullet, \bullet) \rightarrow \text{Hom}_{\mathcal{D}}(F(\bullet), F(\bullet)), \mathfrak{H} : \text{Hom}_{\mathcal{C}}(\bullet, \bullet) \rightarrow \text{Hom}_{\mathcal{E}}(H(\bullet), H(\bullet))$$

given by

$$\mathcal{F}_{C,C'}(f) = F(f), \mathfrak{H}_{C,C'}(f) = H(f)$$

for $f : C \rightarrow C'$ in \mathcal{C} . The functor F is called \mathcal{H} -separable if there exists a natural transformation

$$\mathcal{P} : \text{Hom}_{\mathcal{D}}(F(\bullet), F(\bullet)) \rightarrow \text{Hom}_{\mathcal{E}}(H(\bullet), H(\bullet))$$

such that $\mathcal{P} \circ \mathcal{F} = \mathfrak{H}$.

We recall two properties: The first is Maschke's Theorem for \mathcal{H} -separable functors (see [7, Prop. 2.4]): Let F be an \mathcal{H} -separable functor. If $f : C \rightarrow C'$ in \mathcal{C} is such that $F(f)$ has a left, right, or two-sided inverse in \mathcal{D} , then $H(f)$ has a left, right, or two-sided inverse in \mathcal{E} . The second one is called Rafael's Theorem for \mathcal{H} -separability (see [7, Theorem 2.7]). If F has a right adjoint G , then F is \mathcal{H} -separable if and only if there exists a natural transformation $\xi : \mathcal{H}GF \rightarrow \mathcal{H}$ such that $\xi_C \circ \mathcal{H}(\eta_C) = I_{\mathcal{H}(C)}$ for any $C \in \mathcal{C}$.

2.2. The π -Coalgebras. Recall from Turaev ([14]) and Virelizier ([19]) that a π -coalgebra is a family of k -spaces $C = \{C_\alpha\}_{\alpha \in \pi}$ together with a family of k -linear maps $\Delta = \{\Delta_{\alpha,\beta} : \Delta_{\alpha\beta} \rightarrow \Delta_\alpha \otimes \Delta_\beta\}_{\alpha,\beta \in \pi}$ (called a *comultiplication*) and a k -linear map $\varepsilon : C_e \rightarrow k$ (called a *counit*) such that Δ is coassociative in the sense that

$$(\Delta_{\alpha,\beta} \otimes id_{C_\gamma}) \circ \Delta_{\alpha\beta,\gamma} = (id_{C_\alpha} \otimes \Delta_{\beta,\gamma}) \circ \Delta_{\alpha,\beta\gamma}, \quad (2.1)$$

$$(id_{C_\alpha} \otimes \varepsilon) \circ \Delta_{\alpha,e} = id_{C_\alpha} = (\varepsilon \otimes id_{C_\alpha}) \circ \Delta_{e,\alpha}, \quad (2.2)$$

for any $\alpha, \beta, \gamma \in \pi$.

Remark 2.1. $(C_e, \Delta_{e,e}, \varepsilon)$ is an ordinary coalgebra in the sense of Sweedler. Following the Sweedler's notation for π -coalgebras, for any $\alpha, \beta \in \pi$ and $c \in C_{\alpha\beta}$, one writes

$$\Delta_{\alpha,\beta}(c) = c_{(1,\alpha)} \otimes c_{(2,\beta)}. \quad (2.3)$$

The coassociativity axiom (2.1) gives that, for any $\alpha, \beta, \gamma \in \pi$ and $c \in C_{\alpha\beta\gamma}$,

$$c_{(1,\alpha\beta)(1,\alpha)} \otimes c_{(1,\alpha\beta)(2,\beta)} \otimes c_{(2,\gamma)} = c_{(1,\alpha)} \otimes c_{(2,\beta\gamma)(1,\beta)} \otimes c_{(1,\beta\gamma)(2,\gamma)}, \quad (2.4)$$

which is written as $c_{(1,\alpha)} \otimes c_{(2,\beta)} \otimes c_{(3,\gamma)}$. Inductively, we can define $c_{(1,\alpha_1)} \otimes c_{(2,\alpha_2)} \otimes \cdots c_{(n,\alpha_n)}$, for any $c \in C_{\alpha_1\alpha_2\cdots\alpha_n}$. The axiom (2.2) gives that, for any $\alpha \in \pi$ and $c \in C_\alpha$,

$$\varepsilon(c_{(1,e)})c_{(2,\alpha)} = c = c_{(1,\alpha)}\varepsilon(c_{(2,e)}). \quad (2.5)$$

2.3. The π -C-Comodules. Let C be a π -coalgebra. A left π - C -comodule is a k -vector space V endowed with a family of k -linear maps $\rho^V = \{\rho_\alpha^V : V \rightarrow C_\alpha \otimes V\}_{\alpha \in \pi}$ such that for all $\alpha, \beta \in \pi$ and $v \in V$,

$$v_{<-1,\alpha>} \otimes v_{<0,0>} <-1,\beta> \otimes v_{<0,0>} <0,0> = v_{<-1,\alpha\beta>(1,\alpha)} \otimes v_{<-1,\alpha\beta>(2,\beta)} \otimes v_{<0,0>}, \quad (2.6)$$

$$\varepsilon(v_{<-1,e>})v_{<0,0>} = v, \quad (2.7)$$

where we use the standard notation $\rho_\alpha^V(v) = v_{<-1,\alpha>} \otimes v_{<0,0>}.$

2.4. Weak Hopf π -Coalgebras. We recall from Van Daele and Wang ([18]) that a weak semi-Hopf π -coalgebra $H = \{H_\alpha, m_\alpha, 1_\alpha, \Delta, \varepsilon\}_{\alpha \in \pi}$ is a family of algebras $\{H_\alpha, m_\alpha, 1_\alpha\}_{\alpha \in \pi}$ and at the same time a π -coalgebra $\{H_\alpha, \Delta = \{\Delta_{\alpha,\beta}\}, \varepsilon\}_{\alpha, \beta \in \pi}$ such that

- (i) The comultiplication $\Delta_{\alpha,\beta} : H_{\alpha\beta} \rightarrow H_\alpha \otimes H_\beta$ is a homomorphism of algebras (not necessary unit-preserving) such that

$$(\Delta_{\alpha,\beta} \otimes id_{H_\alpha})\Delta_{\alpha\beta,\gamma}(1_{\alpha\beta\gamma}) = (\Delta_{\alpha,\beta} \otimes 1_\gamma)(1_\alpha \otimes \Delta_{\beta,\gamma}(1_{\beta\gamma})), \quad (2.8)$$

$$(\Delta_{\alpha,\beta} \otimes id_{H_\alpha})\Delta_{\alpha\beta,\gamma}(1_{\alpha\beta\gamma}) = (1_\alpha \otimes \Delta_{\beta,\gamma}(1_{\beta\gamma}))(\Delta_{\alpha,\beta}(1_{\alpha,\beta}) \otimes 1_\gamma), \quad (2.9)$$

for all $\alpha, \beta, \gamma \in \pi$.

- (ii) The counit $\varepsilon : H_e \rightarrow k$ is a k -linear map satisfying the identity

$$\varepsilon(gxh) = \varepsilon(gx_{(2,e)})\varepsilon(x_{(1,e)}h) = \varepsilon(gx_{(1,e)})\varepsilon(x_{(2,e)}h), \quad (2.10)$$

for all $g, h, x \in H_e$.

A weak Hopf π -coalgebra is a weak semi-Hopf π -coalgebra $H = \{H_\alpha, m_\alpha, 1_\alpha, \Delta, \varepsilon\}_{\alpha \in \pi}$ endowed with a family of k -linear maps $S = \{S_\alpha : H_\alpha \rightarrow H_{\alpha^{-1}}\}$ (called *antipode*) such that the following data hold:

$$m_\alpha(S_{\alpha^{-1}} \otimes id_{H_\alpha})\Delta_{\alpha^{-1},\alpha}(h) = 1_{(1,\alpha)}\varepsilon(h1_{(2,e)}), \quad (2.11)$$

$$m_\alpha(id_{H_\alpha} \otimes S_{\alpha^{-1}})\Delta_{\alpha,\alpha^{-1}}(h) = \varepsilon(1_{(1,e)}h)1_{(2,\alpha)}, \quad (2.12)$$

$$S_\alpha(g_{(1,\alpha)})g_{(2,\alpha^{-1})}S_\alpha(g_{(3,\alpha)}) = S_\alpha(g), \quad (2.13)$$

for all $h \in H_e$, $g \in H_\alpha$ and $\alpha \in \pi$.

Remark 2.2. $H = \{H_\alpha, m_\alpha, 1_\alpha, \Delta, \varepsilon\}_{\alpha \in \pi}$ is an ordinary weak Hopf algebra. The set of axioms of the above definition is not self-dual. A weak Hopf π -coalgebra H is said to be of finite type if, for all $\alpha \in \pi$, H_α is finite-dimensional as a k -vector space. Note that it does not mean that $\bigoplus_{\alpha \in \pi} H_\alpha$ is finite-dimensional (unless $H_\alpha = 0$ for

all but a finite number of $\alpha \in \pi$). The antipode $S = \{S_\alpha\}_{\alpha \in \pi}$ of the weak Hopf π -coalgebra H is said to be bijective if each S_α is bijective.

Let H be a weak hopf π -coalgebra. Define the family of linear maps $\varepsilon^t = \{\varepsilon_\alpha^t : H_e \rightarrow H_\alpha\}_{\alpha \in \pi}$ and $\varepsilon^s = \{\varepsilon_\alpha^s : H_e \rightarrow H_\alpha\}_{\alpha \in \pi}$ by the formulars

$$\varepsilon_\alpha^s(h) = m_\alpha(S_{\alpha^{-1}} \otimes id_{H_\alpha})\Delta_{\alpha^{-1}, \alpha}(h) = 1_{(1, \alpha)}\varepsilon(h1_{(2, e)}), \quad (2. 14)$$

$$\varepsilon_\alpha^t(h) = m_\alpha(id_{H_\alpha} \otimes S_{\alpha^{-1}})\Delta_{\alpha, \alpha^{-1}}(h) = \varepsilon(1_{(1, e)}h)1_{(2, \alpha)}, \quad (2. 15)$$

for any $h \in H_e$ and $\alpha \in \pi$, where $\varepsilon^t, \varepsilon^s$ are called the π -target and π -source counital maps. Introduce the notations $H^t = \varepsilon^t(H) = \{\varepsilon_\alpha^t(H_e)\}_{\alpha \in \pi}$ and $H^s = \varepsilon^s(H) = \{\varepsilon_\alpha^s(H_e)\}_{\alpha \in \pi}$ for their images.

Let H be a weak hopf π -coalgebra. Then we have the following properties, we refer to [18] for full detail.

- (W1) $\Delta_{\alpha, \beta}(1_{\alpha\beta}) \in H_\alpha^s \otimes H_\beta^t$, for all $\alpha, \beta \in \pi$;
- (W2) $\varepsilon_\alpha^t(gh) = \varepsilon_\alpha^t(g\varepsilon_e^t(h)), \varepsilon_\alpha^t(\varepsilon_e^t(g)h) = \varepsilon_\alpha^t(g)\varepsilon_\alpha^t(h)$, for all $g, h \in H_e$;
- (W3) $\Delta_{\alpha, \beta}(H_{\alpha\beta}^t) \subseteq H_\alpha \otimes H_\beta^t, \Delta_{\alpha, \beta}(H_{\alpha\beta}^s) \subseteq H_\alpha^s \otimes H_\beta$;
- (W4) $x_{(1, \alpha)} \otimes \varepsilon_\beta^t(x_{(2, e)}) = 1_{(1, \alpha)}x \otimes 1_{(2, \beta)}$, for all $\alpha, \beta \in \pi$ and $x \in H_e$;
- (W5) $\varepsilon_\beta^s(x_{(1, e)}) \otimes x_{(2, \alpha)} = 1_{(1, \beta)} \otimes x1_{(2, \alpha)}$, for all $\alpha, \beta \in \pi$ and $x \in H_e$;
- (W6) $\varepsilon_\alpha^t(h)\varepsilon_\alpha^s(g) = \varepsilon_\alpha^s(g)\varepsilon_\alpha^t(h)$, for all $g, h \in H_e$;
- (W7) $S_\alpha(xy) = S_\alpha(y)S_\alpha(x)$, for all $\alpha \in \pi$ and $x, y \in H_\alpha$;
- (W8) $S_\alpha(1_\alpha) = 1_{\alpha^{-1}}$, for all $\alpha \in \pi$;
- (W9) $\Delta_{\beta^{-1}, \alpha^{-1}} \circ S_{\alpha\beta} = T_{H_\alpha^{-1}, H_\beta^{-1}} \circ (S_\alpha \otimes S_\beta) \circ \Delta_{\alpha, \beta}$, for all $\alpha, \beta \in \pi$;
- (W10) $\varepsilon_\alpha^t \circ S_e = \varepsilon_\alpha^t \circ \varepsilon_e^s = S_{\alpha^{-1}} \circ \varepsilon_{\alpha^{-1}}^s$;
- (W11) $x_{(1, \alpha)} \otimes \varepsilon_\beta^s(x_{(2, e)}) = x1_{(1, \alpha)} \otimes S_{\beta^{-1}}(1_{(2, \beta^{-1})})$, for all $\alpha, \beta \in \pi$ and $x \in H_\alpha$;
- (W12) $\varepsilon_\beta^t(x_{(1, e)}) \otimes x_{(2, \alpha)} = S_{\beta^{-1}}(1_{(1, \beta^{-1})}) \otimes 1_{(2, \alpha)}x$, for all $\alpha, \beta \in \pi$ and $x \in H_\alpha$;
- (W13) If H is of finite type, then the antipode S is bijective.

Definition 2.3. Let H be a weak Hopf π -coalgebra over the field k . A k -algebra A is called a *weak left π -H-comodule algebra* if there exists a family of maps $\rho^A = \{\rho_\alpha^A : A \rightarrow H_\alpha \otimes A\}$ such that

$$(id_{H_\alpha} \otimes \rho_\beta^A) \circ \rho_\alpha^A = (\Delta_{\alpha, \beta} \otimes id_A) \circ \rho_{\alpha\beta}^A, \quad (2. 16)$$

$$(\varepsilon \circ id_A) \circ \rho_e^A = id_A, \quad (2. 17)$$

$$\rho_\alpha^A(1_A) = (\varepsilon_\alpha^s \circ id_A) \circ \rho_e^A(1_A), \quad (2. 18)$$

$$\rho_\alpha^A(ab) = \rho_\alpha^A(a)\rho_\alpha^A(b), \quad (2. 19)$$

for all $\alpha, \beta \in \pi$ and $a, b \in A$. We use the standard notation $\rho_\alpha^A(a) = a_{<-1, \alpha>} \otimes a_{<0, 0>}$.

Lemma 2.4. *Let A be a weak left π - H -comodule algebra. Then, for all $\alpha \in \pi$ and $a \in A$,*

$$1_{<-1,\alpha>} \otimes 1_{<0,0><-1,e>} \otimes 1_{<0,0><0,0>} = 1_{\alpha(1,\alpha)} \otimes 1_{\alpha(2,e)} 1'_{<-1,e>} \otimes 1'_{<0,0>} , \quad (2.20)$$

$$\varepsilon_\alpha^s(a_{<-1,e>}) \otimes a_{<0,0>} = 1_{<-1,\alpha>} \otimes a 1_{<0,0>} , \quad (2.21)$$

$$\varepsilon_\alpha^t(a_{<-1,e>}) \otimes a_{<0,0>} = S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes 1_{<0,0>} a . \quad (2.22)$$

Proof. The proof is straightforward. \square

Definition 2.5. A π -coalgebra C is called a *weak right π - H -module coalgebra* if there exists a family of maps $\cdot : C_\alpha \otimes H_\alpha \rightarrow C_\alpha$ such that

$$(c \cdot h) \cdot h' = c \cdot hh' , \quad (2.23)$$

$$c \cdot 1_\alpha = c , \quad (2.24)$$

$$c \cdot \varepsilon_\alpha^t(g) = \varepsilon_C(c_{(1,e)} \cdot g)c_{(2,\alpha)} , \quad (2.25)$$

$$\Delta_{\alpha,\beta}(c' \cdot g') = c'_{(1,\alpha)} \cdot g'_{(1,\alpha)} \otimes c'_{(2,\beta)} \cdot g'_{(2,\beta)} , \quad (2.26)$$

for all $c \in C_\alpha$, $c' \in C_{\alpha\beta}$, $h, h' \in H_\alpha$, $g' \in H_{\alpha\beta}$, $g \in H_e$ and $\alpha, \beta \in \pi$.

Lemma 2.6. E.q (2. 25) is equivalent to

$$\varepsilon_C(c \cdot h) = \varepsilon_C(c \cdot \varepsilon_e^t(h)) , \quad (2.27)$$

for all $c \in C_e$ and $h \in H_e$.

Proof. Assume E.q (2.27) holds. For all $c \in C_\alpha$ and $g \in H_e$, we have

$$\begin{aligned} \varepsilon_C(c_{(1,e)} \cdot g)c_{(2,\alpha)} &\stackrel{(2.27)}{=} \varepsilon_C(c_{(1,e)} \cdot \varepsilon_e^t(g))c_{(2,\alpha)} \\ &\stackrel{(2.26)}{=} \varepsilon_C(c_{(1,e)} \cdot 1_{(1,e)} \varepsilon_e^t(g))c_{(2,\alpha)} \cdot 1_{(2,\alpha)} \\ &\stackrel{(W3)}{=} \varepsilon_C(c_{(1,e)} \cdot \varepsilon_\alpha^t(g)_{(1,e)})c_{(2,\alpha)} \cdot \varepsilon_\alpha^t(g)_{(2,\alpha)} \\ &\stackrel{(2.26,2.5)}{=} c \cdot \varepsilon_\alpha^t(g). \end{aligned}$$

Thus E.q (2. 25) holds.

Conversely, assume E.q (2. 25) holds. Taking $\alpha = e$, $c \in C_e$ and $g \in H_e$, we have

$$c \cdot \varepsilon_e^t(g) = \varepsilon_C(c_{(1,e)} \cdot g)c_{(2,e)}.$$

So

$$\varepsilon_C(c \cdot \varepsilon_e^t(g)) \stackrel{(2.25)}{=} \varepsilon_C(c_{(1,e)} \cdot g)\varepsilon_C(c_{(2,e)}) \stackrel{(2.5)}{=} \varepsilon_C(c \cdot g).$$

The proof of the Lemma is completed. \square

3. Weak Doi-Hopf π -Module: Functors And Structure

Let H be a weak Hopf π -coalgebra. A *weak Doi-Hopf π -datum* is a triple (H, A, C) , where A is a weak left π - H -comodule algebra and C a weak right π - H -module coalgebra.

A *weak Doi-Hopf π -module* M is a right A -module which is also a left π - C -comodule with the coaction structure $\rho^M = \{\rho_\lambda^M : M \rightarrow C_\lambda \otimes M\}_{\lambda \in \pi}$ such that the following compatible condition holds:

$$\rho_\alpha^M(m \cdot a) = m_{<-1,\alpha>} \cdot a_{<-1,\alpha>} \otimes m_{<0,0>} \cdot a_{<0,0>}, \quad (3. 1)$$

for all $\alpha \in \pi$ and $m \in M, a \in A$.

The set of weak Doi-Hopf π -modules together with both an A -module maps and a π - C -comodule maps will form a category of weak Doi-Hopf π -modules and will be denoted by ${}^{\pi-C}\mathcal{U}(H)_A$ (called a *weak Doi-Hopf π -modules category*).

Let $F^C : {}^{\pi-C}\mathcal{U}(H)_A \rightarrow \mathcal{U}_A$ be the forgetful functor which forgets the π - C -coaction and

$$G : \mathcal{U}_A \rightarrow {}^{\pi-C}\mathcal{U}(H)_A, \quad M \mapsto G(M) = \bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes M}$$

its right adjoint, where $\overline{C_\alpha \otimes M} = \{c \cdot 1_{<-1,\alpha>} \otimes m \cdot 1_{<0,0>} | m \in M, c \in C_\alpha\}$ and the weak Doi-Hopf π -module structures on $\overline{C_\alpha \otimes M}$ are given by

$$(c \cdot 1_{<-1,\alpha>} \otimes m \cdot 1_{<0,0>}) \cdot a = c \cdot a_{<-1,\alpha>} \otimes m \cdot a_{<0,0>}, \quad (3. 2)$$

$${}^l\rho_\beta^{G(M)}(c \cdot 1_{<-1,\alpha>} \otimes m \cdot 1_{<0,0>}) = c_{(1,\beta)} \otimes c_{(2,\beta^{-1}\alpha)} \cdot 1_{<-1,\beta^{-1}\alpha>} \otimes m \cdot 1_{<0,0>}, \quad (3. 3)$$

for all $c \in C_\alpha, a \in A, m \in M$ and $\alpha, \beta \in \pi$. The unit of the adjoint pair (F^C, G) is

$$\rho : 1_{\pi-C}\mathcal{U}(H)_A \rightarrow G \circ F^C$$

defined by $\rho_M : M \rightarrow G(M), \rho_M(m) = \bigoplus_{\alpha \in \pi} m_{<-1,\alpha>} \otimes m_{<0,0>}$, for all $m \in M$. Since A is a right A -module, one has that $\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A}$ is a weak Doi-Hopf π -module via

$$(c \cdot 1_{<-1,\alpha>} \otimes a \cdot 1_{<0,0>}) \cdot b = c \cdot b_{<-1,\alpha>} \otimes a \cdot b_{<0,0>}, \quad (3. 4)$$

$${}^l\rho_\beta^{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a \cdot 1_{<0,0>}) = c_{(1,\beta)} \otimes c_{(2,\beta^{-1}\alpha)} \cdot 1_{<-1,\beta^{-1}\alpha>} \otimes a \cdot 1_{<0,0>} \quad (3. 5)$$

for all $a \in A, c \in C_\alpha$ and $\beta \in \pi$.

Lemma 3.1. *The vector space $\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A}$ is a right π - C -comodule via*

$$\begin{aligned} & {}^r \rho_\beta^{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \\ = & c_{(1,\alpha\beta^{-1})} \cdot 1_{<-1,\alpha\beta^{-1}>} \otimes a_{<0,0>} 1_{<0,0>} \otimes c_{(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}), \end{aligned}$$

for any $\beta, \alpha \in \pi$, $c \in C_\alpha$ and $a \in A$.

Proof. First, we can prove that $(id_{G(A)} \otimes \varepsilon_C) \circ {}^r \rho_e^{G(A)} = id_{G(A)}$. In fact, for all $c \in C_\alpha$ and $a \in A$, we have

$$\begin{aligned} & (id_{G(A)} \otimes \varepsilon_C) \circ {}^r \rho_e^{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \\ = & \varepsilon(c_{(2,e)} S_e(a_{<-1,e>})) c_{(1,\alpha)} \cdot 1_{<-1,\alpha>} \otimes a_{<0,0>} 1_{<0,0>} \\ \stackrel{(2.27)}{=} & \varepsilon(c_{(2,e)} \varepsilon_e^t(S_e(a_{<-1,e>}))) c_{(1,\alpha)} \cdot 1_{<-1,\alpha>} \otimes a_{<0,0>} 1_{<0,0>} \\ \stackrel{(W10)}{=} & \varepsilon(c_{(2,e)} \varepsilon_e^t \varepsilon_e^s(a_{<-1,e>})) c_{(1,\alpha)} \cdot 1_{<-1,\alpha>} \otimes a_{<0,0>} 1_{<0,0>} \\ \stackrel{(2.21)}{=} & \varepsilon(c_{(2,e)} \varepsilon_e^t(1'_{<-1,e>})) c_{(1,\alpha)} \cdot 1_{<-1,\alpha>} \otimes a 1'_{<0,0>} 1_{<0,0>} \\ \stackrel{(2.27,2.18)}{=} & \varepsilon(c_{(2,e)} \varepsilon_e^s(1'_{<-1,e>})) c_{(1,\alpha)} \cdot 1_{<-1,\alpha>} \otimes a 1'_{<0,0>} 1_{<0,0>} \\ \stackrel{(W3,2.26)}{=} & \varepsilon((c \cdot \varepsilon_\alpha^s(1'_{<-1,e>}))_{(2,e)}) (c \cdot \varepsilon_\alpha^s(1'_{<-1,e>}))_{(1,\alpha)} \cdot 1_{<-1,\alpha>} \otimes a 1'_{<0,0>} 1_{<0,0>} \\ \stackrel{(2.5)}{=} & c \cdot \varepsilon_\alpha^s(1'_{<-1,e>}) 1_{<-1,\alpha>} \otimes a 1'_{<0,0>} 1_{<0,0>} \\ \stackrel{(2.18)}{=} & c \cdot 1'_{<-1,\alpha>} 1_{<-1,\alpha>} \otimes a 1'_{<0,0>} 1_{<0,0>} \\ \stackrel{(2.19)}{=} & c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}. \end{aligned}$$

For all $c \in C_\alpha$ and $a \in A$, we have

$$\begin{aligned} & ({}^r \rho_\gamma^{G(A)} \otimes id_{C_\beta}) \circ {}^r \rho_\beta^{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \\ = & c_{(1,\alpha\beta^{-1})(1,\alpha\beta^{-1}\gamma^{-1})} \cdot 1_{<-1,\alpha\beta^{-1}\gamma^{-1}>} \otimes a_{<0,0><0,0>} 1_{<0,0>} \\ & \otimes c_{(1,\alpha\beta^{-1})(2,\gamma)} S_{\gamma^{-1}}(a_{<0,0><-1,\gamma^{-1}>}) \otimes c_{(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}) \\ \stackrel{(2.4)}{=} & c_{(1,\alpha\beta^{-1}\gamma^{-1})} \cdot 1_{<-1,\alpha\beta^{-1}\gamma^{-1}>} \otimes a_{<0,0><0,0>} 1_{<0,0>} \\ & \otimes c_{(2,\gamma\beta)(1,\gamma)} S_{\gamma^{-1}}(a_{<0,0><-1,\gamma^{-1}>}) \otimes c_{(2,\gamma\beta)(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}) \\ \stackrel{(2.16)}{=} & c_{(1,\alpha\beta^{-1}\gamma^{-1})} \cdot 1_{<-1,\alpha\beta^{-1}\gamma^{-1}>} \otimes a_{<0,0>} 1_{<0,0>} \\ & \otimes c_{(2,\gamma\beta)(1,\gamma)} S_{\gamma^{-1}}(a_{<-1,\beta^{-1}\gamma^{-1}>(2,\gamma^{-1})}) \otimes c_{(2,\gamma\beta)(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}\gamma^{-1}>(1,\beta^{-1})}) \\ \stackrel{(W9)}{=} & c_{(1,\alpha\beta^{-1}\gamma^{-1})} \cdot 1_{<-1,\alpha\beta^{-1}\gamma^{-1}>} \otimes a_{<0,0>} 1_{<0,0>} \otimes \\ & c_{(2,\gamma\beta)(1,\gamma)} S_{\beta^{-1}\gamma^{-1}}(a_{<-1,\beta^{-1}\gamma^{-1}>})_{(1,\gamma)} \otimes c_{(2,\gamma\beta)(2,\beta)} S_{\beta^{-1}\gamma^{-1}}(a_{<-1,\beta^{-1}\gamma^{-1}>})_{(2,\beta)} \end{aligned}$$

and also

$$\begin{aligned}
& (id_{G(A)} \otimes \Delta_{\gamma, \beta}) \circ^r \rho_{\gamma\beta}^{G(A)}(c \cdot 1_{<-1, \alpha>} \otimes a 1_{<0, 0>}) \\
&= (id_{G(A)} \otimes \Delta_{\gamma, \beta})(c_{(1, \alpha\beta^{-1}\gamma^{-1})} \cdot 1_{<-1, \alpha\beta^{-1}\gamma^{-1}>} \\
&\quad \otimes a_{<0, 0>} 1_{<0, 0>} \otimes c_{(2, \gamma\beta)} S_{(\gamma\beta)^{-1}}(a_{<-1, \beta^{-1}\gamma^{-1}>})) \\
&= c_{(1, \alpha\beta^{-1}\gamma^{-1})} \cdot 1_{<-1, \alpha\beta^{-1}\gamma^{-1}>} \otimes a_{<0, 0>} 1_{<0, 0>} \\
&\quad \otimes c_{(2, \gamma\beta)(1, \gamma)} S_{(\gamma\beta)^{-1}}(a_{<-1, \beta^{-1}\gamma^{-1}>})(1, \gamma) \otimes c_{(2, \gamma\beta)(2, \beta)} S_{(\gamma\beta)^{-1}}(a_{<-1, \beta^{-1}\gamma^{-1}>})(2, \beta).
\end{aligned}$$

Thus we prove that $({}^r\rho_{\gamma}^{G(A)} \otimes id_{C_{\beta}}) \circ^r \rho_{\beta}^{G(A)} = (id_{G(A)} \otimes \Delta_{\gamma, \beta}) \circ^r \rho_{\gamma\beta}^{G(A)}$.

The proof is completed. \square

The vector space

$$\bigoplus_{\beta \in \pi} \bigoplus_{\gamma \in \pi} \overline{C_{\beta} \otimes C_{\gamma} \otimes A} = G \circ F^C \left(\bigoplus_{\gamma \in \pi} \overline{C_{\gamma} \otimes A} \right) = GF^C G(A) = G^2(A)$$

is also an object in ${}^{\pi-C}\mathcal{U}(H)_A$, i.e.,

$$\begin{aligned}
& G \circ F^C \left(\bigoplus_{\gamma \in \pi} \overline{C_{\gamma} \otimes A} \right) \\
&= \left\{ \bigoplus_{\beta \in \pi} (c_{\beta} \cdot 1'_{<-1, \beta>} \otimes (\bigoplus_{\gamma \in \pi} d_{\gamma} \cdot 1_{<-1, \gamma>} \otimes a 1_{<0, 0>})) \cdot 1'_{<0, 0>} \mid \forall a \in A \right\} \\
&= \left\{ \bigoplus_{\beta \in \pi} \bigoplus_{\gamma \in \pi} (c_{\beta} \cdot 1_{<-1, \beta>} \otimes d_{\gamma} \cdot 1_{<0, 0><-1, \gamma>} \otimes a 1_{<0, 0><0, 0>}) \mid \forall a \in A \right\}
\end{aligned}$$

and also have

$$\left\{
\begin{aligned}
& (c \cdot 1_{<-1, \alpha>} \otimes d \cdot 1_{<0, 0><-1, \gamma>} \otimes a 1_{<0, 0><0, 0>}) \cdot b \\
&= c \cdot b_{<-1, \alpha>} \otimes d \cdot b_{<0, 0><-1, \gamma>} \otimes ab_{<0, 0><0, 0>} , \\
& {}^l\rho_{\beta}^{G^2(A)}(c \cdot 1_{<-1, \alpha>} \otimes d \cdot 1_{<0, 0><-1, \gamma>} \otimes a 1_{<0, 0><0, 0>}) \\
&= c_{(1, \beta)} \otimes c_{(2, \beta^{-1}\alpha)} \cdot 1_{<-1, \beta^{-1}\alpha>} \otimes d \cdot 1_{<0, 0><-1, \gamma>} \otimes a 1_{<0, 0><0, 0>} , \\
& {}^r\rho_{\beta}^{G^2(A)}(c \cdot 1_{<-1, \alpha>} \otimes d \cdot 1_{<0, 0><-1, \gamma>} \otimes a 1_{<0, 0><0, 0>}) \\
&= c \cdot 1_{<-1, \alpha>} \otimes d_{(1, \gamma\beta^{-1})} \cdot 1_{<0, 0><-1, \gamma\beta^{-1}>} \\
&\quad \otimes a_{<0, 0>} 1_{<0, 0><0, 0>} \otimes d_{(2, \beta)} S_{\beta^{-1}}(a_{<-1, \beta^{-1}>}),
\end{aligned}
\right.$$

for all $c \in C_{\alpha}$, $d \in C_{\gamma}$, $a, b \in A$ and $\beta, \alpha, \gamma \in \pi$.

Now, let $F_A : {}^{\pi-C}\mathcal{U}(H)_A \rightarrow {}^{\pi-C}\mathcal{U}$ be the other forgetful functor, which forgets the A -action and

$$\hat{G} : {}^{\pi-C}\mathcal{U} \rightarrow {}^{\pi-C}\mathcal{U}(H)_A, \quad \hat{G}(N) = \{n \hat{\otimes} a = \varepsilon_C(n_{<-1, e>} \cdot 1_{<-1, e>}) n_{<0, 0>} \otimes 1_{<0, 0>} a\}$$

its left adjoint, where for $N \in {}^{\pi-C} \mathcal{U}$, $\hat{G}(N) \in {}^{\pi-C} \mathcal{U}(H)_A$ via the structures: for all $\beta \in \pi$, $n \in N$ and $a, b \in A$,

$$(n \hat{\otimes} a) \cdot b = n \hat{\otimes} ab, \quad (3.6)$$

$$\rho_{\beta}^{\hat{G}(N)}(n \hat{\otimes} a) = n_{<-1,\beta>} \cdot a_{<-1,\beta>} \otimes n_{<0,0>} \otimes a_{<0,0>}. \quad (3.7)$$

Since $\mathcal{C} = \bigoplus_{\alpha \in \pi} C_{\alpha}$ is a left π - C -comodule via ${}^l\rho_{\beta}^{\mathcal{C}}(c) = c_{(1,\beta)} \otimes c_{(2,\beta^{-1}\alpha)}$, for all $c \in C_{\alpha}$ and $\beta \in \pi$. We have $\hat{G}(\mathcal{C}) = \bigoplus_{\alpha \in \pi} \hat{G}(C_{\alpha})$, where

$$\begin{aligned} \hat{G}(C_{\alpha}) &= \{ \varepsilon_C(c_{(1,e)} \cdot 1_{<-1,e>}) c_{(2,\alpha)} \otimes 1_{<0,0>} a \} \\ &\stackrel{(2.25)}{=} \{ c \cdot \varepsilon_{\alpha}^t(1_{<-1,e>}) \otimes 1_{<0,0>} \} \\ &\stackrel{(2.18,W10)}{=} \{ c \cdot S_{\alpha^{-1}}(\varepsilon_{\alpha^{-1}}^s(1_{<-1,e>})) \otimes 1_{<0,0>} a \} \\ &\stackrel{(2.18)}{=} \{ c \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes 1_{<0,0>} a \}. \end{aligned}$$

We can view $\hat{G}(\mathcal{C})$ as a weak Doi-Hopf π -module via

$$(c \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes 1_{<0,0>} a) \cdot b = c \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes 1_{<0,0>} ab, \quad (3.8)$$

$$\begin{aligned} &{}^l\rho_{\beta}^{\hat{G}(\mathcal{C})}(c \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes 1_{<0,0>} a) \\ &= c_{(1,\beta)} \cdot a_{<-1,\beta>} \otimes c_{(2,\beta^{-1}\alpha)} \cdot S_{\beta^{-1}\alpha}(1_{<-1,\beta^{-1}\alpha>}) \otimes 1_{<0,0>} a_{<0,0>}, \end{aligned} \quad (3.9)$$

for all $c \in C_{\alpha}$, $a, b \in A$ and $\alpha, \beta \in \pi$.

From the discussion above, we have two types of weak Doi-Hopf π -modules, i.e., weak Doi-Hopf π -module $\bigoplus_{\alpha \in \pi} \overline{C_{\alpha} \otimes A}$ via (3.4) and (3.5) and weak Doi-Hopf π -module $\hat{G}(\mathcal{C})$ via (3.8) and (3.9). The following proposition will reveal the relation between them.

Proposition 3.2. *two types of weak Doi-Hopf π -modules $\bigoplus_{\alpha \in \pi} \overline{C_{\alpha} \otimes A}$ and $\hat{G}(\mathcal{C})$ are isomorphic in the category ${}^{\pi-C} \mathcal{U}(H)_A$.*

Proof. We construct the maps as follows: for all $c \in C_{\alpha}$, $a \in A$ and $\alpha \in \pi$,

$$u : \bigoplus_{\alpha \in \pi} \overline{C_{\alpha} \otimes A} \rightarrow \bigoplus_{\alpha \in \pi} \hat{G}(C_{\alpha}),$$

$$u(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) = c \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes a_{<0,0>}, \quad (3.10)$$

$$v : \bigoplus_{\alpha \in \pi} \hat{G}(C_{\alpha}) \rightarrow \bigoplus_{\alpha \in \pi} \overline{C_{\alpha} \otimes A},$$

$$v(c \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes 1_{<0,0>} a) = c \cdot a_{<-1,\alpha>} \otimes a_{<0,0>}. \quad (3.11)$$

For all $c \in C_\alpha$, $a \in A$ and $\alpha \in \pi$, since

$$\begin{aligned}
& v \circ u(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \\
&= v(c \cdot S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>}) \otimes a_{<0,0>}) \\
&= c \cdot S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>}) a_{<0,0><-1,\alpha>} \otimes a_{<0,0><0,0>} \\
&\stackrel{(2.16)}{=} c \cdot S_{\alpha^{-1}}(a_{<-1,e>(1,\alpha^{-1})}) a_{<-1,e>(2,\alpha)} \otimes a_{<0,0>} \\
&\stackrel{(2.14)}{=} c \varepsilon_\alpha^s(a_{<-1,e>}) \otimes a_{<0,0>} \stackrel{(2.21)}{=} c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}.
\end{aligned}$$

So we have $v \circ u = id_{G(A)}$. On the other hand, for all $c \in C_\alpha$, $a \in A$ and $\alpha \in \pi$, since

$$\begin{aligned}
& u \circ v(c \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes 1_{<0,0>} a) \\
&= c \cdot a_{<-1,\alpha>} S_{\alpha^{-1}}(a_{<0,0><-1,\alpha^{-1}>}) \otimes a_{<0,0><0,0>} \\
&\stackrel{(2.16)}{=} c \cdot a_{<-1,e>(1,\alpha)} S_{\alpha^{-1}}(a_{<-1,e>(2,\alpha^{-1})}) \otimes a_{<0,0>} \\
&\stackrel{(2.15)}{=} c \cdot \varepsilon_\alpha^t(a_{<-1,e>}) \otimes a_{<0,0>} \stackrel{(2.22)}{=} c \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) \otimes 1_{<0,0>} a.
\end{aligned}$$

Thus we have $u \circ v = id_{\hat{G}(\mathcal{C})}$. The proof that u is both an A -module map and a π - C -comodule map is straightforward. \square

4. Integral of a weak Doi-Hopf π -Datum

Definition 4.1. Let (H, A, C) be a weak Doi-Hopf π -datum. A family of k -linear maps $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ are called an integral of (H, A, C) if

$$c_{(1,\alpha)} \otimes \theta_\beta(c_{(2,\beta)})(d) = d_{(2,\alpha)}(\theta_{\alpha\beta}(c)(d_{(1,(\alpha\beta)^{-1})}))_{<-1,\alpha>} \otimes (\theta_{\alpha\beta}(c)(d_{(1,(\alpha\beta)^{-1})}))_{<0,0>}, \quad (4.1)$$

for all $\alpha, \beta \in \pi$ and $c \in C_{\alpha\beta}$, $d \in C_{\beta^{-1}}$. An integral $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ is called total if

$$\sum_{\alpha \in \pi} \theta_\alpha(c_{(1,\alpha)})(c_{(2,\alpha^{-1})}) = \varepsilon_C(c \cdot 1_{<-1,e>}) 1_{<0,0>}, \quad (4.2)$$

for all $\alpha \in \pi$, $c \in C_e$.

We shall now prove that the existence of an integral $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}$ permits the deformation of a k -linear map between two weak Doi-Hopf π -modules until it becomes a π - C -colinear map.

Proposition 4.2. *Let (H, A, C) be a weak Doi-Hopf π -datum, $N \in {}^{\pi-C} \mathcal{U}$, $M \in {}^{\pi-C} \mathcal{U}(H)_A$ and $u : N \rightarrow M$ a k -linear map. Suppose that there exists $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ an integral. Then*

(1) *The map*

$$\tilde{u} : N \rightarrow M, \quad \tilde{u}(n) = \sum_{\alpha \in \pi} u(n_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>})(u(n_{<0,0>})_{<-1,\alpha^{-1}>}),$$

for all $n \in N$, is left π - C -colinear,

(2) *If θ is a total integral and $f : M \rightarrow N$ is a morphism in ${}^{\pi-C} \mathcal{U}(H)_A$ which is a k -split injection (resp. a k -split surjection), then f has a π - C -colinear retraction (resp. a section).*

Proof. (1) For $n \in N, \beta \in \pi$, we have

$$\begin{aligned} & \rho_\beta^M \circ \tilde{u}(n) \\ = & \sum_{\alpha \in \pi} u(n_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>})(u(n_{<0,0>})_{<-1,\alpha^{-1}>})_{<-1,\beta>}' \\ & \otimes u(n_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>})(u(n_{<0,0>})_{<-1,\alpha^{-1}>})_{<0,0>}' \\ \stackrel{(2.6)}{=} & \sum_{\alpha \in \pi} u(n_{<0,0>})_{<-1,\alpha^{-1}\beta>} \theta_\alpha(n_{<-1,\alpha>}) \\ & (u(n_{<0,0>})_{<-1,\alpha^{-1}\beta>})_{(1,\alpha^{-1})} \\ & \otimes u(n_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>})(u(n_{<0,0>})_{<-1,\alpha^{-1}\beta>})_{<-1,\alpha^{-1}>} \\ \stackrel{\nu=\alpha^{-1}\beta}{=} & \sum_{\nu \in \pi} u(n_{<0,0>})_{<-1,\nu>} \theta_{\beta\nu^{-1}}(n_{<-1,\beta\nu^{-1}>}) \\ & (u(n_{<0,0>})_{<-1,\nu>})_{(1,\nu\beta^{-1})} \\ & \otimes u(n_{<0,0>})_{<0,0>} \theta_{\beta\nu^{-1}}(n_{<-1,\beta\nu^{-1}>})(u(n_{<0,0>})_{<-1,\nu>})_{<-1,\beta\nu^{-1}>} \\ \stackrel{(4.1)}{=} & \sum_{\nu \in \pi} n_{<-1,\beta\nu^{-1}>} (1,\beta) \otimes u(n_{<0,0>})_{<0,0>} \\ & \theta_{\nu^{-1}}(n_{<-1,\beta\nu^{-1}>})_{(2,\nu^{-1})} (u(n_{<0,0>})_{<-1,\nu>}) \\ \stackrel{\alpha=\nu^{-1}}{=} & \sum_{\alpha \in \pi} n_{<-1,\beta\alpha>} (1,\beta) \otimes u(n_{<0,0>})_{<0,0>} \\ & \theta_\alpha(n_{<-1,\beta\alpha>})_{(2,\alpha)} (u(n_{<0,0>})_{<-1,\alpha^{-1}>}) \\ \stackrel{(2.6)}{=} & \sum_{\alpha \in \pi} n_{<-1,\beta>} \otimes u(n_{<0,0>})_{<0,0>} \\ & \theta_\alpha(n_{<0,0>})_{<-1,\alpha>} (u(n_{<0,0>})_{<-1,\alpha^{-1}>}) \\ = & (id \otimes \tilde{u}) \circ \rho_\beta^N(n). \end{aligned}$$

Hence \tilde{u} is a left π - C -colinear.

(2) Let $u : N \rightarrow M$ be a k -linear retraction (resp. section) of f . Then $\tilde{u} : N \rightarrow M$, is a left π - C -colinear retraction (resp. section) of f . Assume first that u is a retraction of f . Then, for $m \in M$, one has

$$\begin{aligned} (\tilde{u} \circ f)(m) &= \sum_{\alpha \in \pi} u(f(m)_{<0,0>})_{<0,0>} \theta_\alpha(f(m)_{<-1,\alpha>})(u(f(m)_{<0,0>})_{<-1,\alpha^{-1}>}) \\ &= \sum_{\alpha \in \pi} u(f(m_{<0,0>}))_{<0,0>} \theta_\alpha(m_{<-1,\alpha>})(u(f(m_{<0,0>}))_{<-1,\alpha^{-1}>}) \\ &= \sum_{\alpha \in \pi} m_{<0,0>} <0,0> \theta_\alpha(m_{<-1,\alpha>})(m_{<0,0>} <-1,\alpha^{-1}>) \\ &\stackrel{(2.6)}{=} \sum_{\alpha \in \pi} m_{<0,0>} \theta_\alpha(m_{<-1,e>(1,\alpha)})(m_{<-1,e>(2,\alpha^{-1})}) \\ &\stackrel{(4.2)}{=} m_{<0,0>} \cdot 1_{<0,0>} \varepsilon(m_{<-1,e>} \cdot 1_{<-1,e>}) = m. \end{aligned}$$

Hence $\tilde{u} : N \rightarrow M$ is a left π - C -colinear retraction of f .

On the other hand, if u is a section of f , then, for $n \in N$, we have

$$\begin{aligned} (f \circ \tilde{u})(n) &= \sum_{\alpha \in \pi} f(u(n_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>})(u(n_{<0,0>})_{<-1,\alpha^{-1}>})) \\ &= \sum_{\alpha \in \pi} f(u(n_{<0,0>})_{<0,0>}) \theta_\alpha(n_{<-1,\alpha>})(u(n_{<0,0>})_{<-1,\alpha^{-1}>}) \\ &= \sum_{\alpha \in \pi} f(u(n_{<0,0>}))_{<0,0>} \theta_\alpha(n_{<-1,\alpha>})(f(u(n_{<0,0>}))_{<-1,\alpha^{-1}>}) \\ &= \sum_{\alpha \in \pi} n_{<0,0>} <0,0> \theta_\alpha(n_{<-1,\alpha>})(n_{<0,0>} <-1,\alpha^{-1}>) \\ &\stackrel{(2.6)}{=} \sum_{\alpha \in \pi} n_{<0,0>} \theta_\alpha(n_{<-1,e>(1,\alpha)})(n_{<-1,e>(2,\alpha^{-1})}) \\ &\stackrel{(4.2)}{=} n_{<0,0>} \cdot 1_{<0,0>} \varepsilon(n_{<-1,e>} \cdot 1_{<-1,e>}) = n, \end{aligned}$$

i.e., $\tilde{u} : N \rightarrow M$ is a left π - C -colinear section of f . Thus the proof is completed. \square

For weak Doi-Hopf π -modules $\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A}$ and $\bigoplus_{\beta \in \pi} \bigoplus_{\gamma \in \pi} \overline{C_\beta \otimes C_\gamma \otimes A}$, we define a map as follows:

$$\begin{aligned} \rho_{G(A)} : \bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A} &\rightarrow \bigoplus_{\beta \in \pi} \bigoplus_{\gamma \in \pi} \overline{C_\beta \otimes C_\gamma \otimes A}, \\ \rho_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) &= \bigoplus_{\gamma \in \pi} c_{(1,\gamma)} \cdot 1_{<-1,\gamma>} \otimes c_{(2,\gamma^{-1}\alpha)} \cdot 1_{<0,0>} <-1,\gamma^{-1}\alpha> \otimes a 1_{<0,0>} <0,0>, \quad (4.3) \end{aligned}$$

for all $c \in C_\alpha$, $a \in A$ and $\alpha \in \pi$.

Lemma 4.3. $\rho_{G(A)}$ is a morphism in the category ${}^{\pi-C}\mathcal{U}^{\pi-C}$.

Proof. It is sufficient to prove that for all $\lambda \in \pi$, the following identities

$${}^l \rho_{\lambda}^{G^2(A)} \circ \rho_{G(A)} = (id_{C_{\lambda}} \otimes \rho_{G(A)}) \circ {}^l \rho_{\lambda}^{G(A)}$$

and

$${}^r \rho_{\lambda}^{G^2(A)} \circ \rho_{G(A)} = (\rho_{G(A)} \otimes id_{C_{\lambda}}) \circ {}^r \rho_{\lambda}^{G(A)}$$

hold. Now we shall check the first. In fact, for all $\lambda, \alpha \in \pi$ and $c \in C_{\alpha}$, we have

$$\begin{aligned} & {}^l \rho_{\lambda}^{G^2(A)} \circ \rho_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a1_{<0,0>}) \\ = & \bigoplus_{\gamma \in \pi} c_{(1,\gamma)(1,\lambda)} \otimes c_{(1,\gamma)(2,\lambda^{-1}\gamma)} \cdot 1_{<-1,\lambda^{-1}\gamma>} \\ & \quad \otimes c_{(2,\gamma^{-1}\alpha)} \cdot 1_{<0,0><-1,\gamma^{-1}\alpha>} \otimes a1_{<0,0><0,0>} \\ \stackrel{(2.1)}{=} & \bigoplus_{\gamma \in \pi} c_{(1,\lambda)} \otimes c_{(2,\lambda^{-1}\alpha)(1,\lambda^{-1}\gamma)} \cdot 1_{<-1,\lambda^{-1}\gamma>} \\ & \quad \otimes c_{(2,\lambda^{-1}\alpha)(2,\gamma^{-1}\alpha)} \cdot 1_{<0,0><-1,\gamma^{-1}\alpha>} \otimes a1_{<0,0><0,0>} \\ \stackrel{\gamma=\lambda\omega}{=} & \bigoplus_{\omega \in \pi} c_{(1,\lambda)} \otimes c_{(2,\lambda^{-1}\alpha)(1,\omega)} \cdot 1_{<-1,\omega>} \\ & \quad \otimes c_{(2,\lambda^{-1}\alpha)(2,\omega^{-1}\lambda^{-1}\alpha)} \cdot 1_{<0,0><-1,\omega^{-1}\lambda^{-1}\alpha>} \otimes a1_{<0,0><0,0>} \\ = & (id_{C_{\lambda}} \otimes \rho_{G(A)}) \circ {}^l \rho_{\lambda}^{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a1_{<0,0>}). \end{aligned}$$

Similarly, we can check the other. Thus the proof of the Lemma is completed. \square

Theorem 4.4. Let (H, A, C) be a weak Doi-Hopf π -datum. The following conditions are equivalent:

- (1) There exists $\theta = \{\theta_{\alpha} : C_{\alpha} \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ a total integral,
- (2) the natural transformation

$$\rho : F_A \circ 1_{\pi-C} \mathcal{U}(H)_A \rightarrow F_A \circ G \circ F^C$$

splits,

- (3) The maps

$$\begin{aligned} \rho_{G(A)} : \bigoplus_{\alpha \in \pi} \overline{C_{\alpha} \otimes A} & \rightarrow \bigoplus_{\beta \in \pi} \bigoplus_{\alpha \in \pi} \overline{C_{\beta} \otimes C_{\alpha} \otimes A}, \\ \rho_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a1_{<0,0>}) & \\ = & \bigoplus_{\beta \in \pi} c_{(1,\beta)} \cdot 1_{<-1,\beta>} \otimes c_{(2,\beta^{-1}\alpha)} \cdot 1_{<0,0><-1,\beta^{-1}\alpha>} \otimes a1_{<0,0><0,0>}, \end{aligned} \quad (4.4)$$

for all $c \in C_{\alpha}$, $a \in A$ and $\alpha \in \pi$, splits in $\pi-C \mathcal{U}^{\pi-C}$.

Proof. (1) \Rightarrow (2). Let $\theta = \{\theta_{\alpha} : C_{\alpha} \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ be a total integral. We have to construct a natural transformation ξ that splits ρ . Let $M \in \pi-C \mathcal{U}(H)_A$

and $u_M : G(M) \rightarrow M$ be the k -linear retraction of $\rho_M : M \rightarrow G(M)$ given by

$$u_M\left(\bigoplus_{\alpha \in \pi} c_\alpha \cdot 1_{<-1,\alpha>} \otimes m_\alpha \cdot 1_{<0,0>}\right) = \varepsilon_C(c_e \cdot 1_{<-1,e>})m_e \cdot 1_{<0,0>}.$$

In fact, for all $m \in M$, we have

$$\begin{aligned} u_M \circ \rho_M(m) &= u_M\left(\bigoplus_{\alpha \in \pi} m_{<-1,\alpha>} \otimes m_{<0,0>}\right) \\ &= \varepsilon_C(m_{<-1,e>})m_{<0,0>} = m. \end{aligned}$$

We define

$$\xi_M = \tilde{u}_M : G(M) \rightarrow M,$$

$$\begin{aligned} \xi_M\left(\bigoplus_{\alpha \in \pi} c_\alpha \cdot 1_{<-1,\alpha>} \otimes m_\alpha \cdot 1_{<0,0>}\right) \\ = \sum_{\alpha \in \pi} m_{\alpha<0,0>} \cdot 1_{<0,0><0,0>} \theta_\alpha(c_\alpha \cdot 1_{<-1,\alpha>})(m_{\alpha<-1,\alpha^{-1}>} \cdot 1_{<0,0><-1,\alpha^{-1}>}). \end{aligned}$$

It follows from Proposition 4.2 that the map ξ_M is a left π - C -colinear retraction of ρ_M .

It remains to prove that $\xi = \{\xi_M | M \in {}^{\pi-C}\mathcal{U}(H)_A\}$ is a natural transformation. Let $f : M \rightarrow N$ be a morphism in ${}^{\pi-C}\mathcal{U}(H)_A$. We have to prove that the diagram

$$\begin{array}{ccc} G(M) & \xrightarrow{\xi_M} & M \\ \downarrow G(f) & & \downarrow f \\ G(N) & \xrightarrow{\xi_N} & N \end{array}$$

is commutative. Using that f is right A -linear, we have

$$\begin{aligned} &(f \circ \xi_M)\left(\bigoplus_{\alpha \in \pi} c_\alpha \cdot 1_{<-1,\alpha>} \otimes m_\alpha \cdot 1_{<0,0>}\right) \\ &= f\left(\sum_{\alpha \in \pi} m_{\alpha<0,0>} \cdot 1_{<0,0><0,0>} \right. \\ &\quad \left. \theta_\alpha(c_\alpha \cdot 1_{<-1,\alpha>})(m_{\alpha<-1,\alpha^{-1}>} \cdot 1_{<0,0><-1,\alpha^{-1}>})\right) \\ &= f\left(\sum_{\alpha \in \pi} m_{\alpha<0,0>} \cdot 1_{<0,0><0,0>} \right. \\ &\quad \left. \theta_\alpha(c_\alpha \cdot 1_{<-1,\alpha>})(m_{\alpha<-1,\alpha^{-1}>} \cdot 1_{<0,0><-1,\alpha^{-1}>})\right) \end{aligned}$$

and using that f is left π - C -colinear

$$\begin{aligned} &(\xi_N \circ (G(f)))(\bigoplus_{\alpha \in \pi} c_\alpha \cdot 1_{<-1,\alpha>} \otimes m_\alpha \cdot 1_{<0,0>}) \\ &= \xi_N\left(\bigoplus_{\alpha \in \pi} c_\alpha \cdot 1_{<-1,\alpha>} \otimes f(m_\alpha \cdot 1_{<0,0>})\right) \end{aligned}$$

$$\begin{aligned}
&= \sum_{\alpha \in \pi} f(m_\alpha)_{<0,0>} \cdot 1_{<0,0><0,0>} \theta_\alpha(c_\alpha \cdot 1_{<-1,\alpha>}) \\
&\quad (f(m_\alpha)_{<-1,\alpha^{-1}>} \cdot 1_{<0,0><-1,\alpha^{-1}>}) \\
&= \sum_{\alpha \in \pi} f(m_{\alpha<0,0>}) \cdot 1_{<0,0><0,0>} \\
&\quad \theta_\alpha(c_\alpha \cdot 1_{<-1,\alpha>}) (m_{\alpha<-1,\alpha^{-1}>} \cdot 1_{<0,0><-1,\alpha^{-1}>}).
\end{aligned}$$

I.e., ξ is a natural transformation that splits ρ .

(2) \Rightarrow (3). Assume that for any $M \in {}^{\pi-C}\mathcal{U}(H)_A$. $\rho_M : M \rightarrow G(M)$ splits in the category ${}^{\pi-C}\mathcal{U}$ of left π - C -comodules and the character of the splitting is functorial. In particular,

$$\begin{aligned}
\rho_{G(A)} &: \bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A} \rightarrow \bigoplus_{\beta \in \pi} \bigoplus_{\alpha \in \pi} \overline{C_\beta \otimes C_\alpha \otimes A}, \\
&\rho_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \\
&= \bigoplus_{\beta \in \pi} c_{(1,\beta)} \cdot 1_{<-1,\beta>} \otimes c_{(2,\beta^{-1}\alpha)} \cdot 1_{<0,0><-1,\beta^{-1}\alpha>} \otimes a 1_{<0,0><0,0>} \quad (4.5)
\end{aligned}$$

splits in ${}^{\pi-C}\mathcal{U}$ and let

$$\xi_{G(A)} : \bigoplus_{\beta \in \pi} \bigoplus_{\alpha \in \pi} \overline{C_\beta \otimes C_\alpha \otimes A} \rightarrow \bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A}$$

be the left π - C -colinear retraction of $\rho_{G(A)}$. Using the naturality of ξ , we will prove that ξ is a right π - C -colinear, where $\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A}$ and $\bigoplus_{\beta \in \pi} \bigoplus_{\alpha \in \pi} \overline{C_\beta \otimes C_\alpha \otimes A}$ are right π - C -comodules in section 3.

First, let V be a vector space and $M \in {}^{\pi-C}\mathcal{U}(H)_A$. Then $M \otimes V \in {}^{\pi-C}\mathcal{U}(H)_A$ via the structures arising from the ones of M , i.e.,

$$(m \otimes v) \cdot a = ma \otimes v, \quad \rho_\alpha^{M \otimes V} = \rho_\alpha^M \otimes id_V,$$

for all $\alpha \in \pi, m \in M, a \in A$, and $v \in V$. Let $v \in V$ and $g_v : M \rightarrow M \otimes V$, $g_v(m) = m \otimes v$. Then g_v is a morphism in ${}^{\pi-C}\mathcal{U}(H)_A$. From the naturality of ξ , we obtain that $g_v \circ \xi_M = \xi_{M \otimes V} \circ G(g_v)$. Hence

$$\begin{aligned}
\xi_{M \otimes V}(c \cdot 1_{<-1,\alpha>} \otimes m \cdot 1_{<0,0>} \otimes v) &= g_v(\xi_M(c \cdot 1_{<-1,\alpha>} \otimes m \cdot 1_{<0,0>})) \\
&= \xi_M(c \cdot 1_{<-1,\alpha>} \otimes m \cdot 1_{<0,0>}) \otimes v,
\end{aligned}$$

for all $\alpha \in \pi, c \in C_\alpha$ and $m \in M, v \in V$. Thus we prove that $\xi_{M \otimes V} = \xi_M \otimes id_V$.

In particular, let us take $M = G(A)$ and $V = \bigoplus_{\beta \in \pi} C_\beta$ viewed only as a vector space. Then $G(A) \otimes \bigoplus_{\beta \in \pi} C_\beta = \bigoplus_{\alpha \in \pi} \bigoplus_{\beta \in \pi} \overline{C_\alpha \otimes A} \otimes C_\beta \in {}^{\pi-C}\mathcal{U}(H)_A$ via the

structures:

$$(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>} \otimes d) \cdot b = c \cdot b_{<-1,\alpha>} \otimes ab_{<0,0>} \otimes d, \quad (4.6)$$

$$\begin{aligned} {}^l \rho_\gamma^{G(A) \otimes \bigoplus_{\beta \in \pi} C_\beta} (c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>} \otimes d) \\ = c_{(1,\gamma)} \otimes c_{(2,\gamma^{-1}\alpha)} \cdot 1_{<-1,\gamma^{-1}\alpha>} \otimes a 1_{<0,0>} \otimes d, \end{aligned} \quad (4.7)$$

for all $\alpha, \beta, \gamma \in \pi, c \in C_\alpha, a \in A$ and $d \in C_\beta$. With these structures, the map

$$\begin{aligned} f = {}^r \rho^{G(A)} : G(A) &\rightarrow G(A) \otimes \bigoplus_{\beta \in \pi} C_\beta \\ &{}^r \rho^{G(A)} (c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \\ &= \bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})} \cdot 1_{<-1,\alpha\beta^{-1}>} \otimes a_{<0,0>} 1_{<0,0>} \otimes c_{(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}), \end{aligned}$$

for all $a \in A, \alpha \in \pi$ and $c \in C_\alpha$, is a morphism in ${}^{\pi-C}\mathcal{U}(H)_A$. In fact, for all $a, b \in A, \alpha \in \pi$ and $c \in C_\alpha$, we have

$$\begin{aligned} {}^r \rho^{G(A)} ((c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \cdot b) &= c \cdot b_{<-1,\alpha>} \otimes ab_{<0,0>} \\ &= \bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})} \cdot b_{<-1,\alpha>(1,\alpha\beta^{-1})} 1_{<-1,\alpha\beta^{-1}>} \otimes a_{<0,0>} b_{<0,0><0,0>} 1_{<0,0>} \\ &\quad \otimes c_{(2,\beta)} \cdot b_{<-1,\alpha>(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>} b_{<0,0><-1,\beta^{-1}>}) \\ &\stackrel{(2.16)}{=} \bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})} \cdot b_{<-1,\alpha\beta^{-1}>(1,\alpha)(1,\alpha\beta^{-1})} 1_{<-1,\alpha\beta^{-1}>} \otimes a_{<0,0>} b_{<0,0>} 1_{<0,0>} \\ &\quad \otimes c_{(2,\beta)} \cdot b_{<-1,\alpha\beta^{-1}>(1,\alpha)(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>} b_{<-1,\alpha\beta^{-1}>(2,\beta^{-1})}) \\ &\stackrel{(2.1)}{=} \bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})} \cdot b_{<-1,\alpha\beta^{-1}>(1,\alpha\beta^{-1})} 1_{<-1,\alpha\beta^{-1}>} \otimes a_{<0,0>} b_{<0,0>} 1_{<0,0>} \\ &\quad \otimes c_{(2,\beta)} \cdot b_{<-1,\alpha\beta^{-1}>(2,e)(1,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>} b_{<-1,\alpha\beta^{-1}>(2,e)(2,\beta^{-1})}) \\ &\stackrel{(2.15)}{=} \bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})} \cdot b_{<-1,\alpha\beta^{-1}>(1,\alpha\beta^{-1})} 1_{<-1,\alpha\beta^{-1}>} \otimes a_{<0,0>} b_{<0,0>} 1_{<0,0>} \\ &\quad \otimes c_{(2,\beta)} \cdot \varepsilon_\beta^t(b_{<-1,\alpha\beta^{-1}>(2,e)}) S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}) \\ &\stackrel{(W4)}{=} \bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})} \cdot b_{<-1,\alpha\beta^{-1}>} \otimes a_{<0,0>} b_{<0,0>} \otimes c_{(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}) \\ &\stackrel{(4.6)}{=} (\bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})} \cdot 1_{<-1,\alpha\beta^{-1}>} \otimes a_{<0,0>} 1_{<0,0>} \otimes c_{(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>})) \cdot b. \end{aligned}$$

Thus f is A -linear. We are left to prove that f is π - C -colinear. It is sufficient to check that for all $\gamma \in \pi$, ${}^l \rho_\gamma^{G(A) \otimes \bigoplus_{\beta \in \pi} C_\beta} \circ f = (id_{C_\gamma} \otimes f) \circ {}^l \rho_\gamma^{G(A)}$ holds. As a matter

of fact, for all $\gamma \in \pi$, $a \in A$, $\alpha \in \pi$ and $c \in C_\alpha$, since

$$\begin{aligned} & {}^l\rho^{G(A) \otimes \bigoplus_{\beta \in \pi} C_\beta} \circ f(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \\ = & c_{(1,\gamma)} \otimes f(c_{(2,\gamma^{-1}\alpha)} \cdot 1_{<-1,\gamma^{-1}\alpha>} \otimes a 1_{<0,0>}) \\ = & \bigoplus_{\beta \in \pi} c_{(1,\gamma)} \otimes c_{(2,\gamma^{-1}\alpha)(1,\gamma^{-1}\alpha\beta^{-1})} \cdot 1_{<-1,\gamma^{-1}\alpha\beta^{-1}>} \\ & \otimes a_{<0,0>} 1_{<0,0>} \otimes c_{(2,\gamma^{-1}\alpha)(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}) \end{aligned}$$

and

$$\begin{aligned} & {}^l\rho_\gamma^{G(A) \otimes \bigoplus_{\beta \in \pi} C_\beta} \circ f(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) \\ = & {}^l\rho_\gamma^{G(A) \otimes \bigoplus_{\beta \in \pi} C_\beta} \left(\bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})} \cdot 1_{<-1,\alpha\beta^{-1}>} \right. \\ & \quad \left. \otimes a_{<0,0>} 1_{<0,0>} \otimes c_{(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}) \right) \\ = & \bigoplus_{\beta \in \pi} c_{(1,\alpha\beta^{-1})(1,\gamma)} \otimes c_{(1,\alpha\beta^{-1})(2,\gamma^{-1}\alpha\beta^{-1})} \cdot 1_{<-1,\gamma^{-1}\alpha\beta^{-1}>} \\ & \quad \otimes a_{<0,0>} 1_{<0,0>} \otimes c_{(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}) \\ = & \bigoplus_{\beta \in \pi} c_{(1,\gamma)} \otimes c_{(2,\gamma^{-1}\alpha)(1,\gamma^{-1}\alpha\beta^{-1})} \cdot 1_{<-1,\gamma^{-1}\alpha\beta^{-1}>} \\ & \quad \otimes a_{<0,0>} 1_{<0,0>} \otimes c_{(2,\gamma^{-1}\alpha)(2,\beta)} S_{\beta^{-1}}(a_{<-1,\beta^{-1}>}). \end{aligned}$$

Thus f is π - C -colinear. From the naturality of ξ , we gain the following commutative diagram

$$\begin{array}{ccc} G^2(A) & \xrightarrow{\xi_{G(A)}} & G(A) \\ \downarrow G(f) = id \otimes f & & \downarrow f \\ G(G(A) \otimes \bigoplus_{\alpha \in \pi} C_\alpha) & \xrightarrow{\xi_{G(A)} \otimes id_{\bigoplus_{\alpha \in \pi} C_\alpha}} & G(A) \otimes \bigoplus_{\alpha \in \pi} C_\alpha \end{array} \quad (*)$$

I.e., ξ is also a right π - C -colinear.

(3) \Rightarrow (1). From (2), we have that

$$\begin{aligned} \rho_{G(A)} : G(A) & \rightarrow \bigoplus_{\beta \in \pi} \bigoplus_{\alpha \in \pi} \overline{C_\beta \otimes C_\alpha \otimes A}, \\ \rho_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes a 1_{<0,0>}) & \\ = & \bigoplus_{\beta \in \pi} c_{(1,\beta)} \cdot 1_{<-1,\beta>} \otimes c_{(2,\beta^{-1}\alpha)} \cdot 1_{<0,0>} <-1,\beta^{-1}\alpha> \otimes a 1_{<0,0>} <0,0> \quad (4.8) \end{aligned}$$

splits in ${}^{\pi-C}\mathcal{U}^{\pi-C}$. Let

$$\xi_{G(A)} : \bigoplus_{\alpha \in \pi} \bigoplus_{\beta \in \pi} C_\alpha \otimes C_\beta \otimes A \rightarrow \bigoplus_{\alpha \in \pi} C_\alpha \otimes A,$$

be a split of $\rho_{G(A)}$ in ${}^{\pi-C}\mathcal{U}^{\pi-C}$. In particular,

$$\begin{aligned} & \xi_{G(A)}\left(\bigoplus_{\beta \in \pi} c_{(1,\beta)} \cdot 1_{<-1,\beta>} \otimes c_{(2,\beta^{-1}\alpha)} \cdot 1_{<0,0><-1,\beta^{-1}\alpha>} \otimes a1_{<0,0><0,0>}\right) \\ &= c \cdot 1_{<-1,\alpha>} \otimes a1_{<0,0>}. \end{aligned} \quad (4.9)$$

We define

$$\begin{aligned} \theta_\alpha : C_\alpha &\rightarrow \text{Hom}(C_{\alpha^{-1}}, A), \\ \theta_\alpha(c)(d) &= (\varepsilon_C \otimes id) \circ P_{\overline{C_e \otimes A}} \circ \xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes d \cdot 1_{<0,0><-1,\alpha^{-1}>} \otimes 1_{<0,0><0,0>}), \end{aligned} \quad (4.10)$$

for all $\alpha \in \pi, c \in C_\alpha, d \in C_{\alpha^{-1}}$, where $P_{\overline{C_e \otimes A}}$ is the natural projective map from $\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A}$ onto $\overline{C_e \otimes A}$, we will prove that θ is a total integral, i.e., E.q (4.1) and (4.2) hold. For all $c \in C_e$, we have

$$\begin{aligned} & \sum_{\alpha \in \pi} \theta_\alpha(c_{(1,\alpha)})(c_{(2,\alpha^{-1})}) \\ = & (\varepsilon_C \otimes id) \circ P_{\overline{C_e \otimes A}} \circ \xi_{G(A)}\left(\bigoplus_{\alpha \in \pi} c_{(1,\alpha)} \cdot 1_{<-1,\alpha>} \right. \\ & \quad \left. \otimes c_{(2,\alpha^{-1})} \cdot 1_{<0,0><-1,\alpha^{-1}>} \otimes 1_{<0,0><0,0>}\right) \\ \stackrel{(4.9)}{=} & (\varepsilon_C \otimes id) \circ P_{\overline{C_e \otimes A}}(c \cdot 1_{<-1,\alpha>} \otimes 1_{<0,0>}) = \varepsilon_C(c \cdot 1_{<-1,\alpha>})1_{<0,0>}. \end{aligned}$$

So E.q (4.2) holds.

Now, we are left to check the other E.q(4.1). For all $\alpha, \beta \in \pi$ and $c \in C_{\alpha\beta}, d \in C_{\beta^{-1}}$, the left hand side of E.q (4.1) is

$$\begin{aligned} & c_{(1,\alpha)} \otimes \theta_\beta(c_{(2,\beta)})(d) \\ = & c_{(1,\alpha)} \otimes (\varepsilon \otimes id) P_{\overline{C_e \otimes A}} \xi_{G(A)}(c_{(2,\beta)} \cdot 1_{<-1,\beta>} \\ & \otimes d \cdot 1_{<0,0><-1,\beta^{-1}>} \otimes 1_{<0,0><0,0>}) \\ = & (id_{C_\alpha} \otimes (\varepsilon \otimes id) \circ P_{\overline{C_e \otimes A}}) \rho_\alpha^{G(A)}(\xi_{G(A)}(c \cdot 1_{<-1,\alpha\beta>} \\ & \otimes d \cdot 1_{<0,0><-1,\beta^{-1}>} \otimes 1_{<0,0><0,0>})) \\ = & (P_{\overline{C_\alpha \otimes A}}) \circ \xi_{G(A)}(c \cdot 1_{<-1,\alpha\beta>} \otimes d \cdot 1_{<0,0><-1,\beta^{-1}>} \otimes 1_{<0,0><0,0>}), \end{aligned}$$

where $P_{\overline{C_\alpha \otimes A}}$ is the natural projective map from $\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A}$ onto $\overline{C_\alpha \otimes A}$. In order to compute the right hand side of E.q (4.1) , we adopt the temporary notation

$$\xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes d_{(1,(\alpha\beta)^{-1})} \cdot 1_{<0,0><-1,(\alpha\beta)^{-1}>} \otimes 1_{<0,0><0,0>}) = \bigoplus_{\gamma \in \pi} p_\gamma \otimes q_\gamma.$$

Now,

$$\begin{aligned} & d_{(2,\alpha)} \theta_{\alpha\beta}(c)(d_{(1,(\alpha\beta)^{-1})})_{<-1,\alpha>} \otimes \theta_{\alpha\beta}(c)(d_{(1,(\alpha\beta)^{-1})})_{<0,0>} \\ &= d_{(2,\alpha)} q_{e<-1,\alpha>} \otimes \varepsilon_C(p_e) q_{e<0,0>}. \end{aligned}$$

Hence (4.1) is equivalent to

$$\begin{aligned}
 & (P_{\overline{C_\alpha \otimes A}}) \circ \xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes d \cdot 1_{<0,0><-1,\beta^{-1}>} \otimes 1_{<0,0><0,0>}) \\
 = & (\cdot \otimes id_A)(d_{(2,\alpha)} \otimes id_{H_\alpha} \otimes id_A)(\varepsilon_C \otimes \rho_\alpha^A)(P_{\overline{C_e \otimes A}}) \cdot \xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \\
 & \otimes d_{(1,(\alpha\beta)^{-1})} \cdot 1_{<0,0><-1,(\alpha\beta)^{-1}>} \otimes 1_{<0,0><0,0>}),
 \end{aligned} \tag{4. 11}$$

for all $\alpha, \beta \in \pi$ and $c \in C_{\alpha\beta}, d \in C_{\beta^{-1}}$. Denoting

$$\xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes d \cdot 1_{<0,0><-1,\beta^{-1}>} \otimes 1_{<0,0><0,0>}) = \bigoplus_{\nu \in \pi} D_\nu \otimes A_\nu$$

and evaluating the diagram at(*), we obtain

$$\begin{aligned}
 & \bigoplus_{\gamma \in \pi} \xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \otimes d_{(1,(\gamma)^{-1})} \cdot 1_{<0,0><-1,(\gamma)^{-1}>} \otimes 1_{<0,0><0,0>}) \otimes d_{(2,\gamma)} \\
 = & \bigoplus_{\gamma \in \pi} \bigoplus_{\nu \in \pi} D_{\nu(1,\nu\gamma^{-1})} \cdot 1_{<-1,\nu\gamma^{-1}>} \otimes A_{\nu<0,0>} 1_{<0,0>} \otimes D_{\nu(2,\gamma)} S_{\gamma^{-1}}(A_{\nu<-1,\gamma^{-1}>}),
 \end{aligned}$$

for all $\alpha, \beta \in \pi, c \in C_\alpha$ and $d \in C_{\beta^{-1}}$. Hence, for all $\alpha \in \pi$, we have

$$\begin{aligned}
 & (d_{(2,\alpha)} \otimes id_A)(\varepsilon_C \otimes id_A)(P_{C_e \otimes A}) \circ \xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \\
 & \otimes d_{(1,(\alpha)^{-1})} \cdot 1_{<0,0><-1,(\alpha\beta)^{-1}>} \otimes 1_{<0,0><0,0>}) \\
 = & D_{\alpha(2,\alpha)} S_{\alpha^{-1}}(A_{\alpha<-1,\alpha^{-1}>}) \varepsilon_C(D_{\alpha(1,e)} \cdot 1_{<-1,e>}) \otimes A_{\alpha<0,0>} 1_{<0,0>} \\
 \stackrel{(2.25)}{=} & D_\alpha \cdot \varepsilon_\alpha^t(1_{<-1,e>}) S_{\alpha^{-1}}(A_{\alpha<-1,\alpha^{-1}>}) \otimes A_{\alpha<0,0>} 1_{<0,0>} \\
 \stackrel{(W12)}{=} & D_\alpha \cdot S_{\alpha^{-1}}(1_{<-1,\alpha^{-1}>}) S_{\alpha^{-1}}(A_{\alpha<-1,\alpha^{-1}>}) \otimes A_{\alpha<0,0>} 1_{<0,0>} \\
 \stackrel{(W7,3.1)}{=} & D_\alpha \cdot S_{\alpha^{-1}}(A_{\alpha<-1,\alpha^{-1}>}) \otimes A_{\alpha<0,0>}.
 \end{aligned}$$

Now we apply ρ_α^A to the second factor of both sides, we have

$$\begin{aligned}
 & (id_{C_\alpha} \otimes \rho_\alpha^A) \circ (d_{(2,\alpha)} \otimes id_A)(\varepsilon_C \otimes id_A)(P_{C_e \otimes A}) \circ \xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \\
 & \otimes d_{(1,(\alpha\beta)^{-1})} \cdot 1_{<0,0><-1,(\alpha\beta)^{-1}>} \otimes 1_{<0,0><0,0>}) \\
 = & D_\alpha \cdot S_{\alpha^{-1}}(A_{\alpha<-1,\alpha^{-1}>}) \otimes A_{\alpha<0,0><-1,\alpha>} \otimes A_{\alpha<0,0><0,0>}.
 \end{aligned}$$

Let the second factor act on the first one, we gain

$$\begin{aligned}
 & (\cdot \otimes id_A) \circ (d_{(2,\alpha)} \otimes id_{H_\alpha} \otimes id_A) \circ (id_{C_\alpha} \otimes \rho_\alpha^A) \circ (d_{(2,\alpha)} \otimes id_A) \\
 & (\varepsilon_C \otimes id_A)(P_{C_e \otimes A}) \circ \xi_{G(A)}(c \cdot 1_{<-1,\alpha>} \\
 & \otimes d_{(1,(\alpha\beta)^{-1})} \cdot 1_{<0,0><-1,(\alpha\beta)^{-1}>} \otimes 1_{<0,0><0,0>}) \\
 = & D_\alpha \cdot S_{\alpha^{-1}}(A_{\alpha<-1,e>(1,\alpha^{-1})}) A_{\alpha<-1,e>(2,\alpha)} \otimes A_{\alpha<0,0>} \\
 \stackrel{(2.14)}{=} & D_\alpha \cdot \varepsilon_\alpha^s(A_{\alpha<-1,e>}) \otimes A_{\alpha<0,0>} \stackrel{(2.21)}{=} D_\alpha \otimes A_\alpha.
 \end{aligned}$$

Since

$$\begin{aligned} (\cdot \otimes id_A) \circ (d_{(2,\alpha)} \otimes id_{H_\alpha} \otimes id_A) \circ (id_{C_\alpha} \otimes \rho_\alpha^A) \circ (d_{(2,\alpha)} \otimes id_A) \\ = (\cdot \otimes id_A)(d_{(2,\alpha)} \otimes id_{H_\alpha} \otimes id_A)\rho_\alpha^A, \end{aligned}$$

we can gain E.q (4.11). To sum up, θ is a total integral. \square

From the proof of Theorem 4.4, if there exists

$$\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$$

a total integral, then the natural transformation

$$\xi : F_A \circ G \circ F^C \rightarrow F_A \circ 1_{\pi-C\mathcal{U}(H)_A}$$

splits ρ , so we have

$$\xi_M \circ F_A(\rho_M) = I_{F_A(M)}$$

for any $M \in \pi-C\mathcal{U}(H)_A$, by Rafael's theorem, F^C is F_A -separable. Conversely, if F^C is F_A -separable, then there exists a natural transformation

$$\xi : F_A \circ G \circ F^C \rightarrow F_A \circ 1_{\pi-C\mathcal{U}(H)_A}$$

such that

$$\xi_M \circ F_A(\rho_M) = I_{F_A(M)}$$

for any $M \in \pi-C\mathcal{U}(H)_A$, we have that ξ splits ρ , by Theorem 4.4, there exists $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ a total integral. Immediately, we have the following conclusion.

Corollary 4.5. *Let (H, A, C) be a weak Doi-Hopf π -datum. The following statements are equivalent:*

- (1) F^C is F_A -separable,
- (2) There exists $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ a total integral,

Let $\pi = \{e\}$ be a trivial group, the weak Hopf π -coalgebras is just the weak Hopf algebras. Combining Theorem 4.4 and Corollary 4.5, we have the following result.

Corollary 4.6. *Let (H, A, C) be a weak Doi-Hopf datum. The following statements are equivalent:*

- (1) F^C is F_A -separable,
- (2) There exists $\theta : C \rightarrow \text{Hom}(C, A)$ a total integral,
- (3) the natural transformation

$$\rho : F_A \circ 1_{C\mathcal{U}(H)_A} \rightarrow F_A \circ G \circ F^C$$

splits,

(4) the map

$$\rho_{\overline{C \otimes A}}^l : \overline{C \otimes A} \rightarrow \overline{C \otimes C \otimes A},$$

$$\rho_{\overline{C \otimes A}}(c \cdot 1_{<-1>} \otimes a 1_{<0>}) = c_{(1)} \otimes c_{(2)} \cdot 1_{<-1>} \otimes a 1_{<0>}$$

splits in ${}^C\mathcal{U}^C$, the category of C -bicomodules. Consequently, if one of the equivalent conditions holds, any weak Doi-Hopf module is injective as a left C -comodule.

We shall prove now the main applications of the existence of a total integral.

Theorem 4.7. *Let (H, A, C) be a weak Doi-Hopf π -datum and suppose that there exists $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}$ a total integral. Then for any $M \in {}^{\pi-C}\mathcal{U}(H)_A$, the map*

$$f : \bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A} \otimes M \rightarrow M,$$

$$\begin{aligned} & f\left(\bigoplus_{\alpha \in \pi} c_\alpha \cdot 1_{<-1,\alpha>} \otimes a_\alpha \cdot 1_{<0,0>} \otimes m\right) \\ &= \sum_{\alpha \in \pi} m_{<0,0>} \theta_\alpha(c_\alpha S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>}))(m_{<-1,\alpha^{-1}>}) a_{<0,0>}, \end{aligned}$$

for all $m \in M$ is a k -split epimorphism in ${}^{\pi-C}\mathcal{U}(H)_A$. In particular, $\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A}$ is a generator in the category ${}^{\pi-C}\mathcal{U}(H)_A$.

Proof. $\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A} \otimes M$ is viewed as an object in ${}^{\pi-C}\mathcal{U}(H)_A$ with the structures as follows,

$$(c \cdot 1_{<-1,\alpha>} \otimes a \cdot 1_{<0,0>} \otimes m) \cdot b = c \cdot b_{<-1,\alpha>} \otimes ab_{<0,0>} \otimes m,$$

$$\rho_\beta^{\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A} \otimes M} (c \cdot 1_{<-1,\alpha>} \otimes a \cdot 1_{<0,0>} \otimes m) = c_{(1,\beta)} \otimes c_{(2,\beta^{-1}\alpha)} \cdot 1_{<-1,\beta^{-1}\alpha>} \otimes a \cdot 1_{<0,0>} \otimes m,$$

for all $\alpha, \beta \in \pi$, $c \in C_\alpha$, $a, b \in A$ and $m \in M$. First, we shall prove that f is a split surjection. Let

$$g : M \rightarrow \bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A} \otimes M,$$

$$g(m) = \bigoplus_{\alpha \in \pi} m_{<-1,\alpha>} \cdot 1_{<-1,\alpha>} \otimes 1_{<0,0>} \otimes m_{<0,0>} ,$$

for all $m \in M$. Then g is left π - C -colinear (but is not right A -linear) and for $m \in M$, we have

$$\begin{aligned} (f \circ g)(m) &= f\left(\bigoplus_{\alpha \in \pi} m_{<-1,\alpha>} \cdot 1_{<-1,\alpha>} \otimes 1_{<0,0>} \otimes m_{<0,0>}\right) \\ &= \sum_{\alpha \in \pi} m_{<0,0><0,0>} \theta_\alpha(m_{<-1,\alpha>}) (m_{<0,0><-1,\alpha^{-1}>}) \\ &\stackrel{(2.6)}{=} \sum_{\alpha \in \pi} m_{<0,0>} \theta_\alpha(m_{<-1,e>(1,\alpha)}) (m_{<-1,e>(2,\alpha^{-1})}) \\ &\stackrel{(4.2)}{=} m_{<0,0>} \cdot 1_{<0,0>} \varepsilon_C(m_{<-1,e>} \cdot 1_{<-1,e>}) = m. \end{aligned}$$

Thus g is a left π - C -colinear section of f . For $a, b \in A, c \in C_\alpha, m \in M$, we have

$$\begin{aligned} &f((c \cdot 1_{<-1,\alpha>} \otimes a \cdot 1_{<0,0>} \otimes m) \cdot b) \\ &= f(c \cdot b_{<-1,\alpha>} \otimes ab_{<0,0>} \otimes m) \\ &= m_{<0,0>} \theta_\alpha(c \cdot b_{<-1,\alpha>} S_{\alpha^{-1}}(b_{<0,0><-1,\alpha^{-1}>}) S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>})) \\ &\quad (m_{<-1,\alpha^{-1}>}) a_{<0,0>} b_{<0,0><0,0>} \\ &\stackrel{(2.16, 2.15)}{=} m_{<0,0>} \theta_\alpha(c \cdot \varepsilon_\alpha^t(b_{<-1,e>}) S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>})) (m_{<-1,\alpha^{-1}>}) a_{<0,0>} b_{<0,0>} \\ &\stackrel{(2.22)}{=} m_{<0,0>} \theta_\alpha(c S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>})) (m_{<-1,\alpha^{-1}>}) a_{<0,0>} b \\ &= f(c \cdot 1_{<-1,\alpha>} \otimes a \cdot 1_{<0,0>} \otimes m) \cdot b, \end{aligned}$$

i.e., f is right A -linear. It remains to prove that f is also left π - C -colinear. First, for all $\beta \in \pi, c_\alpha \in C_\alpha, a \in A$ and $m \in M$, we compute

$$\begin{aligned} &(id_C \otimes f) \circ \rho_\beta^{\bigoplus_{\alpha \in \pi} \overline{C_\alpha \otimes A} \otimes M} (c \cdot 1_{<-1,\alpha>} \otimes a \cdot 1_{<0,0>} \otimes m) \\ &= (c_{(1,\beta)} \otimes m_{<0,0>} \theta_{\beta^{-1}\alpha}(c_{(2,\beta^{-1}\alpha)} S_{\alpha^{-1}\beta}(a_{<-1,\alpha^{-1}\beta>})) (m_{<-1,\alpha^{-1}\beta>}) a_{<0,0>}) \end{aligned}$$

and

$$\begin{aligned} &\rho_\beta^M(f(c \cdot 1_{<-1,\alpha>} \otimes a \cdot 1_{<0,0>} \otimes m)) \\ &= \rho_\beta^M(m_{<0,0>} \theta_\alpha(c_\alpha S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>})) (m_{<-1,\alpha^{-1}>}) a_{<0,0>}) \\ &= (m_{<0,0><-1,\beta>} (\theta_\alpha(c_\alpha S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>})) (m_{<-1,\alpha^{-1}>}))_{<-1,\beta>} \\ &\quad a_{<0,0><-1,\beta>} \otimes m_{<0,0><0,0>} (\theta_\alpha(c_\alpha S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}>})) \\ &\quad (m_{<-1,\alpha^{-1}>}))_{<0,0>} a_{<0,0><0,0>}) \\ &\stackrel{(2.6)}{=} (m_{<-1,\alpha^{-1}\beta>(2,\beta)} \theta_\alpha(c_\alpha S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}\beta>(1,\alpha^{-1})}))) \\ &\quad (m_{<-1,\alpha^{-1}\beta>(1,\alpha^{-1})})_{<-1,\beta>} a_{<-1,\alpha^{-1}\beta>(2,\beta)} \otimes m_{<0,0>} \end{aligned}$$

$$\begin{aligned}
& (\theta_\alpha(c_\alpha S_{\alpha^{-1}}(a_{<-1,\alpha^{-1}\beta>(1,\alpha^{-1})}))(m_{<-1,\alpha^{-1}>}))_{<0,0>} a_{<0,0>} \\
\stackrel{(\nu=\alpha^{-1}\beta)}{=} & (m_{<-1,\nu>(2,\beta)} \theta_{\beta\nu^{-1}}(c_{\beta\nu^{-1}} S_{\nu\beta^{-1}}(a_{<-1,\nu>(1,v\beta^{-1})})) \\
& (m_{<-1,\nu>(1,v\beta^{-1})})_{<-1,\beta>} a_{<-1,\nu>(2,\beta)} \otimes m_{<0,0>} \\
& \theta_{\beta\nu^{-1}}(c_{\beta\nu^{-1}} S_{\nu\beta^{-1}}(a_{<-1,\nu>(1,v\beta^{-1})}))(m_{<-1,\nu>(1,v\beta^{-1})})_{<0,0>} a_{<0,0>} \\
\stackrel{(4.1)}{=} & ((c_{\beta\nu^{-1}} S_{\nu\beta^{-1}}(a_{<-1,\nu>(1,v\beta^{-1})}))(1,\beta) a_{<-1,\nu>(2,\beta)} \otimes \\
& m_{<0,0>} \theta_{\nu^{-1}}((c_{\beta\nu^{-1}} S_{\nu\beta^{-1}}(a_{<-1,\nu>(1,v\beta^{-1})}))(2,\nu^{-1}))) (m_{<-1,\nu>}) a_{<0,0>} \\
\stackrel{(W9)}{=} & (c_{\beta\nu^{-1}(1,\beta)} S_{\beta^{-1}}(a_{<-1,\nu>(1,\nu\beta^{-1})(2,\beta^{-1})}) a_{<-1,\nu>(2,\beta)} \otimes \\
& m_{<0,0>} \theta_{\nu^{-1}}(c_{\beta\nu^{-1}(2,\nu^{-1})} S_\nu(a_{<-1,\nu>(1,\nu\beta^{-1})(1,v)})) (m_{<-1,\nu>}) a_{<0,0>} \\
\stackrel{(2.1)}{=} & (c_{\beta\nu^{-1}(1,\beta)} S_{\beta^{-1}}(a_{<-1,\nu>(2,e)(1,\beta^{-1})}) a_{<-1,\nu>(2,e)(2,\beta)} \otimes \\
& m_{<0,0>} \theta_{\nu^{-1}}(c_{\beta\nu^{-1}(2,\nu^{-1})} S_\nu(a_{<-1,\nu>(1,v)})) (m_{<-1,\nu>}) a_{<0,0>} \\
\stackrel{(2.14)}{=} & (c_{\beta\nu^{-1}(1,\beta)} \varepsilon_\beta^s(a_{<-1,\nu>(2,e)}) \otimes m_{<0,0>} \\
& \theta_{\nu^{-1}}(c_{\beta\nu^{-1}(2,\nu^{-1})} S_\nu(a_{<-1,\nu>(1,v)})) (m_{<-1,\nu>}) a_{<0,0>} \\
\stackrel{(W11)}{=} & (c_{\beta\nu^{-1}(1,\beta)} S_{\beta^{-1}}(1_{(2,\beta^{-1})}) \otimes m_{<0,0>} \\
& \theta_{\nu^{-1}}(c_{\beta\nu^{-1}(2,\nu^{-1})} S_\nu(a_{<-1,\nu>1_{(1,\nu)}})) (m_{<-1,\nu>}) a_{<0,0>} \\
\stackrel{(W9)}{=} & ((c_{\beta\nu^{-1}(1,\beta)}) \otimes m_{<0,0>} \theta_{\nu^{-1}}(c_{\beta\nu^{-1}(2,\nu^{-1})} S_\nu(a_{<-1,\nu>})) (m_{<-1,\nu>}) a_{<0,0>}),
\end{aligned}$$

i.e., f is left π - C -colinear. Hence, we proved that f is an epimorphism in ${}^{\pi-C}\mathcal{U}(H)_A$ and has a π - C -colinear section. \square

Remark 4.8. The Theorem 4.7 can be followed in a new way as follows: From the forgetful functors $F_A : {}^{\pi-C}\mathcal{U}(H)_A \rightarrow {}^{\pi-C}\mathcal{U}(H)$ and $F^C : {}^{\pi-C}\mathcal{U}(H)_A \rightarrow \mathcal{U}(H)_A$, we have the functor $\mathfrak{H} = F^C \circ F_A : {}^{\pi-C}\mathcal{U}(H)_A \rightarrow Vect_k$. the natural transformation \mathcal{P} is constructed as follows: for $M, N \in {}^{\pi-C}\mathcal{U}(H)$, we define

$$\mathcal{P}_{M,N} : \text{Hom}_{{}^{\pi-C}\mathcal{U}(H)}(M, N) \rightarrow \text{Hom}_{Vect_k}(M, N), \mathcal{P}_{M,N}(f) = F^C(f).$$

Notice $\mathcal{P} \circ F_A = \mathfrak{H}$, i.e., F_A is \mathfrak{H} -separable functor. From the proof of Theorem 4.4, we know that $F_A(f)$ has a right inverse in ${}^{\pi-C}\mathcal{U}(H)$, by the Maschke's theorem for \mathfrak{H} -separable functor(see[7,Prop.2.4]), $\mathfrak{H}(f)$ has a right inverse in $Vect_k$.

5. The Maschke-type Theorem For weak Doi-Hopf π -modules

In the section, we give the Maschke-type Theorem for weak Doi-Hopf π -modules. First, we need the following Lemma.

Lemma 5.1. *Let (H, A, C) be a weak Doi-Hopf π -datum, $N, M \in {}^{\pi-C} \mathcal{U}(H)_A$ and $u : N \rightarrow M$ a A -linear map. Suppose that there exists $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ an integral such that A -centralising condition, i.e., for all $\alpha \in \pi$, $a \in A, c \in C_\alpha$ and $d \in C_{\alpha^{-1}}$,*

$$a_{<0,0>} <0,0> \theta_\alpha(c \cdot a_{<-1,\alpha>}) (d \cdot a_{<0,0>} <-1,\alpha^{-1}>) = \theta_\alpha(c)(d)a. \quad (5.1)$$

Then the map

$$\tilde{u} : N \rightarrow M, \quad \tilde{u}(n) = \sum_{\alpha \in \pi} u(n_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>}) (u(n_{<0,0>})_{<-1,\alpha^{-1}>}),$$

for all $n \in N$, is left π - C -colinear and A -linear.

Proof. From the Proposition 4.2, \tilde{u} is a left π - C -colinear. We are left to check that \tilde{u} is right A -linear. In fact, for all $n \in N$ and $a \in A$, Since

$$\begin{aligned} \tilde{u}(n \cdot a) &= \sum_{\alpha \in \pi} u(n_{<0,0>} \cdot a_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>} \cdot a_{<-1,\alpha>}) \\ &\quad (u(n_{<0,0>} \cdot a_{<0,0>})_{<-1,\alpha^{-1}>}) \\ &= \sum_{\alpha \in \pi} u(n_{<0,0>})_{<0,0>} \cdot a_{<0,0>} <0,0> \theta_\alpha(n_{<-1,\alpha>} \cdot a_{<-1,\alpha>}) \\ &\quad (u(n_{<0,0>})_{<-1,\alpha^{-1}>} \cdot a_{<0,0>} <-1,\alpha^{-1}>) \\ &= \sum_{\alpha \in \pi} u(n_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>}) (u(n_{<0,0>})_{<-1,\alpha^{-1}>}) a. \end{aligned}$$

So the proof of the lemma is completed. \square

Theorem 5.2. *Let (H, A, C) be a weak Doi-Hopf π -datum, and $N, M \in {}^{\pi-C} \mathcal{U}(H)_A$, and $v : M \rightarrow N$ a morphism in ${}^{\pi-C} \mathcal{U}(H)_A$. Suppose that there exists $\theta = \{\theta_\alpha : C_\alpha \rightarrow \text{Hom}(C_{\alpha^{-1}}, A)\}_{\alpha \in \pi}$ an integral such that A -centralising condition. If the map v is a split injection in \mathcal{U}_A , then u have a retraction in ${}^{\pi-C} \mathcal{U}(H)_A$.*

Proof. Since v is a split injection in \mathcal{U}_A , there exist a morphism $u : N \rightarrow M$ in \mathcal{U}_A such that $uv = id_M$. Define \tilde{u} as follows

$$\tilde{u}(n) = \sum_{\alpha \in \pi} u(n_{<0,0>})_{<0,0>} \theta_\alpha(n_{<-1,\alpha>}) (u(n_{<0,0>})_{<-1,\alpha^{-1}>}).$$

From Lemma 5.1, \tilde{u} is a morphism in ${}^{\pi-C}\mathcal{U}(H)_A$. Next, we shall check $\tilde{u}v = id_M$.

For $m \in M$, one has

$$\begin{aligned}
& \tilde{u}v(m) \\
&= \sum_{\alpha \in \pi} u(v(m)_{<0,0>})_{<0,0>} \theta_\alpha (v(m)_{<-1,\alpha>}) (u(v(m)_{<0,0>})_{<-1,\alpha^{-1}>}) \\
&= \sum_{\alpha \in \pi} u(v(m_{<0,0>}))_{<0,0>} \theta_\alpha (m_{<-1,\alpha>}) (u(v(m_{<0,0>}))_{<-1,\alpha^{-1}>}) \\
&= \sum_{\alpha \in \pi} m_{<0,0>} <0,0> \theta_\alpha (m_{<-1,\alpha>}) (m_{<0,0>} <-1,\alpha^{-1}>) \\
&= \sum_{\alpha \in \pi} m_{<0,0>} \theta_\alpha (m_{<-1,e>(1,\alpha)}) (m_{<-1,e>(2,\alpha^{-1})}) \\
&= m.
\end{aligned}$$

So the theorem is proved. \square

Acknowledgment. The authors would like to thank the referee for the valuable suggestions and comments.

References

- [1] G. Böhm, F. Nill and K. Szlachányi, *Weak Hopf algebras, I. Integral theory and C^* -structure*, J. Algebra, 221 (1999), 385-438.
- [2] G. Böhm, *Doi-Hopf modules over weak Hopf algebras*, Comm. Algebra, 28 (2000), 4687-4698.
- [3] S. Caenepeel, G. Militaru and Zhu Shenglin, *Crossed modules and Doi-Hopf modules*, Israel J. Math., 100 (1997), 221-248.
- [4] S. Caenepeel and M. De Lombaerde, *A categorical approach to Turaev's Hopf group-coalgebras*, Comm. Algebra, 34(7) (2006), 2631-2657.
- [5] S. Caenepeel, G. Militaru and Zhu Shenglin, *A Maschke-type theorem for Doi-Hopf modules. Applications*, J. Algebra, 187 (1997), 388-412.
- [6] S. Caenepeel, Bogdan Ion, G. Militaru and Zhu Shenglin, *Separable functors for the Category of Doi-Hopf modules, Applications*, Adv. Math., 145 (1999), 239-290.
- [7] S. Caenepeel, G. Military, *Maschke functors, semisimple functors and separable functors of the second kind. Applications*, J. Pure Appl. Algebra, 178 (2003), 131-157.
- [8] C. Năstăsescu, M. Van den Bergh and F. Van Oystaeyen, *Separable functors applied to graded rings*, J. Algebra, 123 (1989), 397-413
- [9] Y. Doi, *Unifying Hopf modules*, J. Algebra, 153(2) (1992), 373-385.

- [10] C. Menini and G. Militaru, *Integral, Quantum galois extensions, and the Affineness criterion for quantum Yetter-Drinfeld modules*, J. Algebra, 247 (2002), 467-508.
- [11] D. E. Radford and J. Towber, *Yetter-Drinfeld categories associated to an arbitrary bialgebra*, J. Pure Appl. Algebra, 87(3) (1993), 259-279.
- [12] Shen Bingliang and Wang Shuanhong, *Blattner-Cohen-Montgomery's Duality Theorem for (Weak) Group Smash Products*, Comm. Algebra, 36(6) (2008), 2387-2409.
- [13] M. E. Sweedler, Hopf Algebras, New York: Benjamin, 1969.
- [14] V. G. Turaev, *Homotopy field theory in dimension 3 and crossed group-categories*, Preprint GT/0005291,(2000).
- [15] A. Van Daele, *Multiplier Hopf algebras*, Trans. Amer. Math. Soc., 342(2) (1994), 917-932.
- [16] A. Van Daele, *An algebraic framework for group duality*, Adv. Math., 140 (1998), 323-366.
- [17] A. Van Daele and Wang Shuanhong, *New braided crossed categories and Drinfeld quantum double for weak Hopf π -coalgebras*, Comm. Algebra, 36 (2008), 2341-2386.
- [18] A. Van Daele and Wang Shuanhong, *Multiplier unifying Hopf algebras*. Submitted for publication.
- [19] A. Virelizier, *Hopf group-coalgebras*, J. Pure Appl. Algebra, 171 (2002), 75-122.
- [20] A. Virelizier, *Graded quantum groups and quasitriangular Hopf group-coalgebras*, Comm. Algebra, 33 (9)(2005), 3029-3050.
- [21] Wang Shuanhong, *Group twisted smash products and Doi-Hopf modules for T -coalgebras*, Comm. Algebra, 32(9) (2004), 3437-3458 .
- [22] Wang Shuanhong, *Group entwining structures and group coalgebra Galois extensions*, Comm. Algebra, 32(9) (2004), 3417-3436.
- [23] Wang Shuanhong, *A Maschke-Type Theorem for Hopf π -Comodules*, Tsukuba J. Math., 28(2) (2004), 377-388.
- [24] Wang Shuanhong, *Morita Contexts, π -Galois Extensions for Hopf π -coalgebras*, Comm. Algebra, 34(2) (2006), 521-546.
- [25] D. N. Yetter, *Quantum groups and representations of monoidal categories*, Math. Proc. Cambridge Philos. Soc., 108(2) (1990), 261-290.
- [26] M. Zunino, *Double construction for crossed Hopf coalgebra*, J. Algebra, 278 (2004), 43-75.

- [27] M. Zunino, *Yetter-Drinfeld modules for Turaev crossed structures*, J. Pure Appl. Algebra, 193(1-3) (2004), 313-343.
- [28] M. D. Rafael, *Separable functors revisited*, Comm. Algebra ,18 (1990), 1445-1459.

Chen Quan-guo and Wang Shuan-hong

Department of Mathematics
Southeast University
210096, Nanjing, China
e-mails: cqg211@163.com (C. Quan-guo)
shuanhwang2002@yahoo.com (W. Shuan-hong)