

ANAESTHETIC EFFECTS OF SODIUM BICARBONATE AT DIFFERENT CONCENTRATIONS ON AFRICAN CATFISH (*Clarias gariepinus*) JUVENILES

Cecilia Muthoni GITHUKIA¹, Elijah Migiro KEMBENYA¹, Mary A. OPIYO²

¹Kenya Marine & Fisheries Research Institute, Kegati Aquaculture Research Station, Kisii, Kenya.

²Kenya Marine & Fisheries Research Institute, National Aquaculture Research Development & Training Center, Sagana, Kenya.

Received: 17.06.2015

Accepted: 15.12.2015

Published online: 21.04.2016

Corresponding author:

Cecilia Muthoni GITHUKIA, Kenya Marine & Fisheries Research Institute, Kegati Aquaculture Research Station, P.O. Box 3259-40200, Kisii, Kenya.

E-mail: cmuthoni230@gmail.com

Abstract:

In the present study, anaesthetic effects of different sizes (5, 10, 15, 20, 25, 30, 35 and 40g) of African catfish (*Clarias gariepinus*) juveniles was investigated at different concentrations (0/control, 10, 15, 20, 25, 30, 35, 40, 45, 50 g/L) of sodium bicarbonate. Juveniles were bath immersed in each of the different concentrations of sodium bicarbonate solution for 5 minutes and recovery times in clean aerated water was established. Juveniles took between 0.67 to 5.44 minutes to reach full anaesthesia and 0.66 to 4.25 minutes to recover fully from anaesthesia. An inverse exponential relationship was observed between sodium bicarbonate concentrations and induction time, whereas exponential relationship was observed between sodium bicar-

bonate concentrations and recovery time. Size of *C. gariepinus* juveniles significantly affected the induction time ($P < 0.05$) with smaller juveniles getting to full anaesthesia faster than bigger juveniles at the same concentration of sodium bicarbonate. In addition, recovery time was size dependent where smaller *C. gariepinus* juveniles took significantly longer time ($P < 0.05$) to recover from anaesthesia than bigger juveniles. All juveniles reached anaesthesia and recovered in less than 5 minutes except for 30 g juveniles at 15g/L of sodium bicarbonate which was anaesthetized at 5.44 minutes.

Keywords: Immersion, Induction, Recovery, Anaesthesia

Introduction

Anaesthetics are used to minimize stress associated with handling live fish when marking, tagging, counting, injecting, transporting, weighing, artificial reproduction and surgical operations among others (Hajek et al., 2006, Altun et al., 2009, Yildiz et al., 2013, Opiyo et al., 2013). The need to handle fish without impairing their health or commercial value has led to the development of many techniques to anaesthetize fish (Yildiz et al., 2013). Different anaesthetics act with various intensity driving fish in to general anaesthesia, resulting in loss of consciousness, inhibition of reflex activity, and reduction in skeletal muscle tone (Hajek et al., 2006). Quick induction and recovery from anaesthesia is desirable in most cases (Marking & Meyer, 1985; Stoskopf, 1993). However, long recovery time is desirable while collecting fish from the wild or where fish must be handled for a longer time in the laboratory. An ideal anaesthetic should possess several attributes such as; non-toxic, inexpensive, simple to administer and result in rapid induction and calm recovery (Pawar et al., 2011). It is of paramount importance to identify the effective dose of anaesthetic for specific fish species since response to the same anaesthetic vary depending on the concentration used and the species of fish (Tytler and Hawkins, 1981; Pawar, et al., 2011). Common fish anaesthetics include clove oil, sodium bicarbonate, carbon dioxide gas, metomidate, benzocaine, tricaine, methanesulphonate (MS-222), 2-phenoxyethanol and quinaldine (Massee et al., 1995; Palić et al., 2006). Regardless of the agent, the process of anaesthesia in fish develops in a similar way and runs in a progressive pattern (McFarland, 1959).

Sodium bicarbonate (Na_2CO_3) also known as baking soda, is a white substance that gives carbon dioxide when dissolved in water (McFarland and Klontz, 1969). Its main advantages lie in its low cost, wide availability and safety to both fish and humans (Altun et al., 2009). Sodium bicarbonate has been effectively used as an anaesthetic in common carp (*Cyprinus carpio*) in both cold and warm water conditions (Booke et al., 1978, Altun et al., 2009), Rainbow trout (*Oncorhynchus mykiss*) (Keen et al., 1998) and in Nile tilapia (*Oreochromis niloticus*) (Opiyo et al., 2013).

The African catfish (*Clarias gariepinus*), is found in several countries throughout Africa, where it has a native distributional range as well as in Europe (the Netherlands, Germany and Belgium),

Asia (Indonesia, Thailand) and South America (Brazil) (de Graaf and Janssen, 1996; Brummett, 2008). *C. gariepinus* is the second most important freshwater cultured fish (after tilapia) in Africa (Barasa et al., 2014) with the exception of Nigeria where its production far exceeds that of tilapia production and accounts for 70–80 % of the total freshwater fish production (Ponzoni and Nguyen, 2008). Successful culture of *C. gariepinus* requires artificial propagation whereby there is need of injection of the fish with gonadotropin to spawn and the juveniles are often moved from one place to another during sorting and grading at the larval growth phases (de Graaf and Janssen, 1996). These procedures may induce stress in the fish hence require use of anaesthetics to induce calming effects and reduce the mobility of the aggressive fish. *C. gariepinus* is therefore an ideal species to examine the suitability of sodium bicarbonate as an anaesthetic. The aim of the current study is to determine the efficacy and optimum concentration of sodium bicarbonate in different sizes of *C. gariepinus* which hitherto has not been reported.

Materials and Methods

C. gariepinus juveniles of average weight 10, 15, 20, 25, 30, 35 and 40 g were collected from the nursery ponds at Kenya Marine and Fisheries Research Institute, Kegati Aquaculture Research Station, Kenya. The juveniles were starved for 24 hours to allow for gut evacuation prior to experiment (Weyl et al., 1996). Average water temperature was at $23.5 \pm 1.2^\circ\text{C}$, pH (7.8 ± 0.18), dissolved oxygen (4.5 ± 1.7) mg/L, and Total Ammonia Nitrogen (0.1 ± 0.1) mg/L in the experimental aquaria. To determine the optimal dosage and effect of size on anaesthetic efficacy of sodium bicarbonate, a group of 5 juveniles from each weight class were exposed to different concentrations of sodium bicarbonate by bath immersion. The experiment was carried out in 15 L glass aquaria in triplicates. The concentrations were increased by an interval of 5 based on the ability of the fish to go to stage III anaesthesia as described by Iwama et al., (1989) and Palić et al., (2006). Concentrations of sodium bicarbonate used were (0/control, 10, 15, 20, 25, 30, 35, 40, 45, 50 g/L). The juveniles were observed for three different stages of induction and recovery time (Table 1) as described by Iwama et al., (1989) and Palić et al., (2006).

Table 1. Stages of induction and recovery

Stage of induction	Description
I	Loss of equilibrium
II	Loss of gross body movements but continued opercula movement
III	Same as stage II but opercula movement ceases
Stages of recovery	Description
I	No body movement but opercula movement start
II	Regular opercula movements and body movements start
III	Equilibrium regained with pre-anaesthetic appearance

*Adapted from Iwama et al. (1989) and Palić et al. (2006).

After full anaesthesia, juveniles were removed from the anaesthetic solution and transferred to another aquarium with clean aerated well water for recovery. The time of induction to anaesthesia and recovery were recorded and fish were maintained there for 48 hours in order to observe their behavior and possible mortality.

Differences among means were analyzed by two-way ANOVA followed by Duncan's multiple range tests. All analyses were performed with SPSS version 17.0 statistical package. Differences were considered significant at $P < 0.05$.

Results and Discussion

Induction and recovery time of *C. gariepinus* juveniles exposed to different concentrations of sodium bicarbonate are shown in table 2 and 3. In general juveniles took between 0.67 to 5.44 minutes to reach full anaesthesia in different concentrations of sodium bicarbonate (Table 2). Juveniles exposed to 0 and 10g/L concentration of sodium bicarbonate did not undergo anaesthesia. In addition, 35 and 40 g juveniles did not undergo anaesthesia when exposed to 15g/L concentration of sodium bicarbonate. The recovery time for the juveniles which did not undergo anaesthesia was not recorded. Induction time decreased with increase in the concentration of sodium bicarbonate while recovery time increased with increase in the concentration of sodium bicarbonate (Figure 1 and 2). Concentration of sodium bicarbonate solution had a significant effect ($P < 0.05$) on the induction time to anaesthesia with juveniles attaining full anaesthesia faster at a higher concentration than at a lower concentration. Ju-

veniles weighing 10g took 0.67 minutes to attain anaesthesia at 50 g/L concentration of sodium bicarbonate and 3.04 minutes at 15 g/L of sodium bicarbonate. Size of *C. gariepinus* juveniles significantly affected the induction time to anaesthesia ($P < 0.05$) with smaller juveniles getting to full anaesthesia faster than bigger juveniles at the same concentration of sodium bicarbonate. At a concentration of 20g/L, 10g juveniles took 2.78 minutes to attain anaesthesia while 40 g juveniles took 4.98 minutes to attain anaesthesia.

C. gariepinus juveniles generally took between 0.66 to 4.25 minutes to fully recover from the anaesthetic solution. Sodium bicarbonate concentration significantly ($P < 0.05$) affected the recovery time from anaesthesia with juveniles exposed to a higher concentration taking longer to recover than those exposed to a lower concentration (Table 3). 10 g juveniles took 4.25 minutes to recover from anaesthesia at 50g/L of sodium bicarbonate concentration and 1.66 minutes at 15g/L of sodium bicarbonate concentration. Size of juveniles also significantly ($P < 0.05$) affected the recovery time from anaesthesia with smaller juveniles taking longer time to recover than bigger juveniles at the same concentration of sodium bicarbonate. 10 g juveniles took 1.88 minutes to recover from anaesthesia while 40 g juveniles took 0.66 minutes to recover from anaesthesia when exposed to 20g/L concentration of sodium bicarbonate. No mortality was recorded in any concentration of sodium bicarbonate during the application and post recovery period.

Anaesthetics play a vital role in modern day aquaculture. It helps to reduce the stress which makes fish more susceptible to diseases due to resistance during handling (Ross and Ross, 1999). The use of sodium bicarbonate from baking powder as an anaesthetic in fish is a relatively cheap technique because of its affordability and ease of application (Opiyo et al., 2013). It is safe for human and there is no banning or restriction in its use as it leaves no tissue residues (Altun et al., 2009). Consequently, it can be used on animals entering the human food chain without any withdrawal period (Ross and Ross, 1999). The effectiveness of sodium bicarbonate depends on the weight of the fish and the concentration of administration as reported by Booke et al., (1978) and Altun et al., (2009). Similar phenomenon was realized in the present study.

Table 2. Induction time (minutes) for different sizes of *C. gariepinus* juveniles exposed to different concentrations of sodium bicarbonate

Concentration (g/L)	Fish weight (g)						
	10	15	20	25	30	35	40
0	–	–	–	–	–	–	–
10	–	–	–	–	–	–	–
15	3.04 ± 0.01 ^{aA}	3.42 ± 0.00 ^{bA}	3.81 ± 0.05 ^{cA}	4.38 ± 0.01 ^{dA}	5.44 ± 0.01 ^{eA}	–	–
20	2.78 ± 0.00 ^{aB}	3.06 ± 0.02 ^{bB}	3.14 ± 0.00 ^{cB}	3.34 ± 0.01 ^{dB}	4.12 ± 0.00 ^{eB}	4.89 ± 0.01 ^{fA}	4.98 ± 0.01 ^{gA}
25	2.31 ± 0.01 ^{aC}	1.97 ± 0.01 ^{bC}	2.19 ± 0.01 ^{cC}	2.45 ± 0.00 ^{dC}	3.67 ± 0.00 ^{eC}	4.00 ± 0.00 ^{fB}	4.66 ± 0.00 ^{gB}
30	2.04 ± 0.00 ^{aD}	1.22 ± 0.01 ^{bD}	1.42 ± 0.01 ^{cD}	1.88 ± 0.01 ^{dD}	3.04 ± 0.03 ^{eD}	3.12 ± 0.00 ^{fC}	3.21 ± 0.00 ^{gC}
35	1.16 ± 0.03 ^{aE}	1.08 ± 0.08 ^{bE}	1.27 ± 0.06 ^{cE}	1.35 ± 0.01 ^{dE}	1.99 ± 0.00 ^{eE}	2.35 ± 0.01 ^{fD}	2.77 ± 0.00 ^{gD}
40	1.01 ± 0.03 ^{aF}	0.91 ± 0.01 ^{bF}	1.13 ± 0.01 ^{cF}	1.23 ± 0.01 ^{dF}	1.55 ± 0.00 ^{eF}	1.75 ± 0.01 ^{fE}	2.11 ± 0.01 ^{gE}
45	0.85 ± 0.01 ^{aG}	0.85 ± 0.01 ^{aG}	0.99 ± 0.01 ^{bG}	1.14 ± 0.01 ^{cG}	1.25 ± 0.01 ^{dG}	1.33 ± 0.00 ^{eF}	1.60 ± 0.00 ^{fF}
50	0.67 ± 0.01 ^{aH}	0.76 ± 0.04 ^{bH}	0.90 ± 0.00 ^{cH}	1.01 ± 0.01 ^{dH}	1.06 ± 0.00 ^{eH}	1.12 ± 0.01 ^{fG}	1.23 ± 0.00 ^{gG}

* Data are presented as Mean values ± SE of five determinations

** Means identified by different small letters in the rows or capital letters in the columns were significantly different (P < 0.05).

*** The sign –, means no change in the fish

Table 3. Recovery time (minutes) for different sizes of *C. gariepinus* juveniles exposed to different concentrations of sodium bicarbonate

Concentration (g/L)	Fish weight (g)						
	10	15	20	25	30	35	40
0	–	–	–	–	–	–	–
10	–	–	–	–	–	–	–
15	1.66 ± 0.15 ^{aA}	1.65 ± 0.01 ^{bA}	1.65 ± 0.12 ^{cA}	1.36 ± 0.02 ^{dA}	0.98 ± 0.01 ^{eA}	–	–
20	1.88 ± 0.03 ^{aB}	1.62 ± 0.15 ^{bB}	1.70 ± 0.01 ^{cB}	1.55 ± 0.01 ^{dB}	1.37 ± 0.03 ^{eB}	1.02 ± 0.03 ^{fA}	0.66 ± 0.01 ^{gA}
25	2.33 ± 0.01 ^{aC}	2.27 ± 0.02 ^{bC}	2.04 ± 0.01 ^{cC}	1.87 ± 0.01 ^{dC}	1.78 ± 0.03 ^{eC}	1.58 ± 0.02 ^{fB}	1.04 ± 0.00 ^{gB}
30	2.72 ± 0.01 ^{aD}	2.62 ± 0.05 ^{bD}	2.17 ± 0.02 ^{cD}	2.46 ± 0.02 ^{dD}	2.50 ± 0.01 ^{eD}	1.90 ± 0.01 ^{fC}	1.61 ± 0.02 ^{gC}
35	3.28 ± 0.03 ^{aE}	3.13 ± 0.03 ^{bE}	2.74 ± 0.02 ^{cE}	3.00 ± 0.01 ^{dE}	2.60 ± 0.10 ^{eE}	2.18 ± 0.01 ^{fD}	2.04 ± 0.02 ^{gD}
40	3.62 ± 0.01 ^{aF}	3.47 ± 0.06 ^{bF}	3.24 ± 0.00 ^{cF}	3.13 ± 0.01 ^{dF}	2.94 ± 0.05 ^{eF}	2.40 ± 0.01 ^{fE}	2.28 ± 0.01 ^{gE}
45	3.95 ± 0.01 ^{aG}	3.54 ± 0.16 ^{bG}	3.40 ± 0.02 ^{cG}	3.27 ± 0.01 ^{dG}	3.10 ± 0.02 ^{eG}	2.79 ± 0.01 ^{fF}	2.56 ± 0.01 ^{gF}
50	4.25 ± 0.01 ^{aH}	4.21 ± 0.05 ^{bH}	3.93 ± 0.02 ^{cH}	3.45 ± 0.01 ^{dH}	3.25 ± 0.01 ^{eH}	3.05 ± 0.01 ^{fG}	2.71 ± 0.01 ^{gG}

* Data are presented as Mean values ± SE of five determinations

** Means identified by different small letters in the rows or capital letters in the columns were significantly different (P < 0.05).

*** The sign –, means no change in the fish




Figure 1. Induction of different sizes of *C. gariepinus* juveniles exposed to different concentrations of sodium bicarbonate.




Figure 2. Recovery of different sizes of *C. gariepinus* juveniles exposed to different concentrations of sodium bicarbonate.

It took between 0.67 to 5.44 minutes for *C. gariepinus* juveniles to reach anaesthesia. Fish of the same weight reached anaesthesia faster at higher concentrations of sodium bicarbonate than at lower concentrations. This could be attributed to the fact that more anaesthesia diffused in to the body of the fish at higher concentration than those subjected to a lower concentration. This is consistent with Opiyo et al., (2013) and Altun et al., (2009) who found that higher concentration of sodium bicarbonate resulted to faster anaesthesia in Nile tilapia and Common carp respectively. In addition, smaller juveniles were also anaesthetized faster than bigger ones at the same concentration of sodium bicarbonate. This could be explained by the fact that smaller fish have a bigger surface area to volume ratio than bigger fish and therefore absorbed the anaesthetic faster. The induction time from the present study is in agreement with several authors (Marking and Meyer, 1985; Iversen, 2003; Coyle et al., 2004; Mylonas et al., 2005; Opiyo et al., 2013) who recommended a period of between 3-5 minutes for an effective induction time to anaesthesia.

The concentration of the anaesthetic solution significantly affected recovery time from anaesthesia. Juveniles exposed to higher concentration of sodium bicarbonate took longer to recover from anaesthesia than those at a lower concentration. These results are comparable with previously published work by Solomon et al. (2014) who found that when anaesthetic was in higher concentration, more of it was absorbed and accumulated in the Central Nervous System (CNS) of the fish thus suppressing the activity of the CNS to a greater degree than at lower concentrations and consequently prolonging the recovery time. King et al., (2005) also reported similar results when clove oil was used as anaesthesia in black sea bass (*Centropristis striata*). Moreover, bigger juveniles took shorter period of time to recover from anaesthesia than smaller juveniles at the same concentration. This could be linked to the fact that bigger fish have a smaller surface area to volume ratio than smaller fish and is comparable to (Yildiz et al., 2013). Similar results were recorded by Altun et al., (2009) on Common carp juveniles, Zaikov et al., (2008) on Pike and Opiyo et al., (2013) on Nile tilapia juveniles. Solomon et al., (2014) and Hseu et al., (1998) also noted that higher drug concentration increase recovery time. However, Booke et al., (1978) recorded much lower concentrations of sodium

bicarbonate at 1,000 mg/L and 600 mg/L at pH 6.5 and 7.7 respectively.

Conclusion

In conclusion, 50g/L concentration of sodium bicarbonate is recommended to anaesthetize *C. gariepinus* juveniles since it led to anaesthesia in less than 2 minutes for all the sizes of fish.

Acknowledgements

The authors would wish to thank Kenya Marine and Fisheries Research Institute (KMFRI) through Kenya Agricultural Productivity and Agribusiness Project (KAPAP) grant number KAPAP-CGS/FP/2011/06 for the financial support to undertake this research work. Our sincere gratitude also goes to Laboratory technicians; Mr. Ezekiel Nyarumba and Ms. Susan Mogaka for their technical assistance in carrying out the experiment.

References

- Altun, T., Bilgin, R. & Danabas, D. (2009). Effects of Sodium Bicarbonate on Anaesthesia of Common Carp, *Cyprinus carpio* Juveniles. *Turkish Journal of Fisheries and Aquatic Science*, 9, 29-31.
- Barasa, J. E., Abila, R., Grobler, J.P., Dangasuk, O.G., Njahira, M.N. & Kaunda-Arara, B. (2014). Genetic diversity and gene flow in *Clarias gariepinus* from Lakes Victoria and Kanyaboli, Kenya. *African Journal of Aquatic Science*, 39, 287-293.
- Booke, H.E., Hollender, B. & Lutterbie, G. (1978). Sodium bicarbonate, an inexpensive fish anesthetic for field use, *Progressive Fish Culturist*, 40, 11-13.
- Brummett, R.E. (2008). *Clarias* catfish: Biology, ecology, distribution and biodiversity, pp. 64-72. In: R.W. Ponzoni and N.H. Nguyen (Eds.), 2008. Proceedings of a Workshop on the Development of a Genetic Improvement Program for African catfish, *Clarias gariepinus*. World Fish Center Conference Proceedings Number 1889. The World Fish Center, Penang, Malaysia. 130 pp.
- Coyle, S.D., Durborow, R.M. & Tidwell, J.H. (2004). Anaesthetics in Aquaculture, *SRAC Publication* No. 3900. Texas.
- de Graaf, G. & Janssen, H. (1996). Artificial reproduction and pond rearing of African catfish, *Clarias gariepinus* in Sub-Saharan Af-

- rica - A handbook. FAO Fisheries Technical Paper. No. 362. FAO, Rome, Italy. 73pp.
- Hajek, G.J., Klyszejko B. & Dziaman R. (2006). The anaesthetic effect of clove oil on Common carp, *Cyprinus carpio*. *Acta Ichthyologica*, 36, 93-97.
- Hseu, J.R., Yeh, S.L., Chu, Y.T. & Ting, Y.Y. (1998). Comparison of efficacy of five anaesthetic in goldlined sea bream, *Sparus sarba*. *Acta Zool Taiwan*, 9, 11-18.
- Iversen, M., Finstad, B., McKinley, R.S. & Eliassen R.A. (2003). The efficacy of metomidate, clove oil, Aquist-Mand Benzoak® as anaesthetics in Atlantic salmon, *Salmo salar* smolts, and their potential stress-reducing capacity. *Aquaculture*, 221, 549-566.
- Iwama, G.K., McGeer, J.C. & Pawluk, M.P. (1989). The effects of five fish anaesthetic on acid-base balance, hematocrit, cortisol and adrenaline in Rainbow trout, *Oncorhynchus mykiss*. *Canadian Journal of Zoology*, 67, 2065-2073.
- King, W.V., Hooper, B., Hills Grove, S., Benton, C. & Berlinsky, D. (2005). The use of clove oil, metomidate, tricaine methanesulphonate and PE for inducing anaesthesia and their effect on the cortisol stress response in black sea bass, *Centropristis striata*. *Aquaculture Research*, 36, 1442-1449.
- Marking, L.L. & Meyer, F.P. (1985). Are better anesthetics needed in fisheries? *Fisheries*, 10, 2-5.
- Massee, K.C., Rust, M.B., Hardy, R.W. & Stickney R.R. (1995). The effectiveness of tricaine, quinaldine sulfate and metomidate as anaesthetics for larval fish. *Aquaculture*, 134, 351-359.
- McFarland, W.N. & Klontz G.W. (1969). Anesthesia in fishes. *Federal Proceedings*, 7, 1535-1540.
- McFarland, W.W. (1959). A study of the effects of anaesthetics on the behaviour and physiology of fishes. *Publications of the Institute of Marine Science*, 6, 23-55.
- Mylonas, C.C., Cardinaletti, G., Sigelaki, I. & Polzonetti-Magni, A. (2005). Comparative efficacy of clove oil and 2-phenoxyethanol as anaesthetics in the aquaculture of European sea bass, *Dicentrarchus labrax* and gilthead sea bream, *Sparus aurata* at different temperatures. *Aquaculture*, 246, 467-481.
- Opiyo, M.A., Ogello, E.O. & Charo Karisa H. (2013). Effectiveness of Sodium Bicarbonate as an Anaesthetic for different sizes of Nile tilapia, *Oreochromis niloticus* Juveniles. *International Journal of Aquatic Science*, 4, 14-22.
- Palic, D., Herolt, D.M., Andreason, C.B., Menzel, B.W. & Roth J.W. (2006). Anaesthetic efficacy of tricaine methanesulfonate, metomidate and eugenol: Effects on plasma cortisol concentration and neutrophil function in fathead minnows, *Pimephales promelas*. *Aquaculture*, 254, 675-685.
- Pawar, H.B., Sanaye, S.V., Sreepada, R.A., Harish, V. & Suryavanshi, U., Tanu Ansari, Z.A. (2011). Comparative efficacy of four anaesthetic agents in the yellow seahorse, *Hippocampus kuda*. *Aquaculture*, 311, 155-161.
- Ponzoni, R.W. & Nguyen, N.H. (2008). Proceedings of a Workshop on the Development of a Genetic Improvement Program for African catfish, *Clarias gariepinus*. WorldFish Center Conference Proceedings Number 1889. The WorldFish Center, Penang, Malaysia. 130 pp.
- Ross, L.G. & Ross, B. (1999). Anaesthetic and Sedative Techniques for Aquatic Animals. Blackwell Science, Oxford, 159 pp.
- Solomon, S.G., Cheikyula, J.O. & Anju, D.T. (2014). Behavioural Responses of *Heterobranchus longifilis* Juveniles Exposed to Freeze-dried Bark Extract of *Tephrosia vogelii* as an Anaesthetic. *Journal of Stress Physiology & Biochemistry*, 10, 82-92.
- Stoskopf, M.K. (1993). Anesthesia. In: L. Brown, (Eds.), *Aquaculture for Veterinarians: Farmed Fish Husbandry and Medicine*, Pergamon Press, Oxford, England, pp. 161-168.
- Tytler, P. & Hawkins, A.D. (1981). Vivisection, anaesthetics and minor surgery. In: A.D. Hawkins (Eds.), *Aquarium systems*, Academic Press, New York, USA, pp. 247-278.
- Weyl, O., Kaiser, H. & Hecht T. (1996). The efficacy and mode of action of 2-phenoxyethanol as an anaesthetic for goldfish, *Carassius auratus* at different tempera-

Journal abbreviation: **J Aquacult Eng Fish Res**

- tures and concentrations. *Aquaculture Research*, 27: 757-764.
- Yildiz, M., Kayim, M., Akin S. 2013. The Anesthetic Effects of Clove Oil and 2-Phenoxyethanol on Rainbow Trout, *Oncorhynchus mykiss* at Different Concentrations and Temperatures. *Iranian Journal of Fisheries Sciences*, 12, 947-961.
- Zaikov, A., Iliev, I. & Hubenova, T. (2008). Induction and recovery from anaesthesia in pike, *Esox lucius* exposed to clove oil. *Bulgarian Journal of Agricultural Science*, 14, 165-170.