

Identification of the Solely Time-Dependent Zero-Order Coefficient in a Linear Bi-Flux Diffusion Equation from an Integral Measurement

İbrahim Tekin^{1*} and Mehmet Akif Çetin²¹Department of Fundamental Sciences, Rafet Kayis Faculty of Engineering, Alanya Alaaddin Keykubat University, 07425 Antalya, Türkiye²ALTSO Vocational School, Alanya Alaaddin Keykubat University, 07425 Antalya, Türkiye

*Corresponding author

Article Info

Keywords: Bi-flux equation, Fourier method, Inverse problem**2010 AMS:** 35R30, 35A02**Received:** 7 February 2023**Revised:** 12 July 2023**Accepted:** 28 September 2023**First Online:** 29 September 2023**Published:** 30 September 2023

Abstract

Bi-flux diffusion equation, can be easily affected by the existence of external factors, is known as an anomalous diffusion. In this paper, the inverse problem (IP) of determining the solely time-dependent zero-order coefficient in a linear Bi-flux diffusion equation with initial and homogeneous boundary conditions from an integral additional specification of the energy is considered. The unique solvability of the inverse problem is demonstrated by using the contraction principle for sufficiently small times.

1. Introduction

The classical diffusion model is contingent on Fick's law and, this generally describes some practical problems such as heat and mass diffusion, bacterial infection diffusion, and predator-prey models, [1, 2, 3]. However, a number of sensitive particle systems, for example population systems arise in biology, are known as an anomalous diffusion. Because the diffusion motion of particles can be easily affected by the existence of the disturbing exogenous agents. To model this anomalous diffusion, the Bi-flux approach to diffusion problems was introduced in [4, 5, 6, 7] to deal with bizarre evolutionary processes by using a discrete formulation. The theory leads the flux to bifurcate into two separate currents corresponding to two independent potentials. The 1st and 2nd flows derive from the classical Fick potential and a new potential, respectively. From this theory the Bi-flux diffusion equation is derived as:

$$Z_{\tau}(\chi, \tau) + \beta(1 - \beta)RZ_{\chi\chi\chi\chi}(\chi, \tau) - \beta DZ_{\chi\chi}(\chi, \tau) = H(\chi, \tau, Z), \quad (1.1)$$

where $Z(\chi, \tau)$ is the particle concentration, D is the diffusion coefficient, R is the reaction coefficient, $H(\chi, \tau, Z)$ linear or nonlinear reaction function, and β ($0 \leq \beta \leq 1$) is the fraction of particles displaced regarding to the Fick's law.

If $\beta = 0$, the homogeneous equation (1.1) ($H(\chi, \tau, Z) = 0$) reduces to a stationary equation and $Z(\chi, \tau) = \text{constant}$. If $\beta = 1$, the equation (1.1) reduces to the classical diffusion model which arises in many physical processes such as thermodynamics [8], predator-prey problems [2], bio-heat conduction [9], heat exchangers [10], and mass transport in groundwater [11]. If the diffusion coefficient $D = 0$, the equation (1.1) is Mullin's equation which arises in thermal grooving by surface diffusion [12]. If the reaction function $H(\chi, \tau, Z)$ is linear and in the form $H(\chi, \tau, Z) = k(\tau)Z(\chi, \tau) + h(\chi, \tau)$, then the eq. (1.1) can be rewritten as

Email addresses and ORCID numbers: ibrahim.tekin@alanya.edu.tr, 0000-0001-6725-5663 (I. Tekin), akif.cetin@alanya.edu.tr, 0000-0002-4991-5098 (M.A. Çetin)

Cite as: İ. Tekin, M.A. Çetin, Identification of the solely time-dependent zero-order coefficient in a linear bi-flux diffusion equation from an integral measurement, Fundam. J. Math. Appl., 6(3) (2023), 170-176



$$Z_\tau(\chi, \tau) + \beta(1 - \beta)RZ_{\chi\chi\chi\chi}(\chi, \tau) - \beta DZ_{\chi\chi}(\chi, \tau) = k(\tau)Z(\chi, \tau) + h(\chi, \tau). \tag{1.2}$$

In this work we consider an IP of recovering the time-dependent zero-order coefficient $k(\tau)$ in a linear Bi-flux diffusion equation (1.2) together with the particle concentration $Z(\chi, \tau)$ in the rectangle domain $\Pi_T = \{(\chi, \tau) : (\chi, \tau) \in [0, 1] \times [0, T]\}$ for some fixed $T > 0$, subject to the initial condition (IC)

$$Z(\chi, 0) = z_0(\chi), \chi \in [0, 1], \tag{1.3}$$

and the boundary conditions (BCs)

$$Z(0, \tau) = Z(1, \tau) = Z_{\chi\chi}(0, \tau) = Z_{\chi\chi}(1, \tau) = 0, \tau \in [0, T], \tag{1.4}$$

and the additional condition (AC)

$$\int_0^1 Z(\chi, \tau)d\chi = m(\tau), \tau \in [0, T], \tag{1.5}$$

where, D and R are the positive diffusion and reaction coefficients respectively, $0 < \beta < 1$, $h(\chi, \tau)$ is the source term, $z_0(\chi)$ is the initial particle concentration, and $m(\tau)$ is the extra integral measurement, which often refers specification of the energy/mass, to obtain the solution of the IP.

The IPs for the classical diffusion equation ($\beta = 1$) are satisfactorily studied theoretically and numerically. For instance, IPs of determining the time-dependent heat source are studied in [13, 14, 15, 16], and IPs of reconstructing the time-dependent reaction coefficient are investigated in [17, 18, 19, 20, 21] with different boundary conditions (local, non-local or non-classical conditions).

On contrary to the IPs for the classical diffusion equation, our aim is to study the Bi-flux diffusion equation to identify the time-dependent zero-order coefficient $k(\tau)$ along with the particle concentration $Z(\chi, \tau)$ theoretically, for the first time, in the rectangular domain, using the IC (1.3), homogeneous BCs (1.4) and the AC (1.5).

The paper is organized as follows. In Section 2, the equivalent IP formulation is derived. In Section 3, the equivalent IP reduced to a system of Volterra integral equations with respect to the unknown functions $Z(\chi, \tau)$, and $k(\tau)$. Then the theorem of the existence and uniqueness of the solution of the IP is phrased and proved by means of the contraction principle.

2. Mathematical formulation of the equivalent IP

In this section, first we will give the classical solution of the IP (1.2)-(1.5), and then reduce the considered IP to the equivalent problem.

Definition 2.1. *The classical solution of the IP (1.2)-(1.5) is the pair $\{Z(\chi, \tau), k(\tau)\}$ subject to the following properties:*

- (i) $Z(\chi, \tau) \in C^{4,1}(D_T)$,
- (ii) $k(\tau) \in C[0, T]$,
- (iii) *the pair $\{Z(\chi, \tau), k(\tau)\}$ satisfies the eq. (1.2) and the conditions (1.3)-(1.5) in ordinary sense.*

$Z(\chi, \tau) \in C^{4,1}(\Pi_T)$ means that $Z(\chi, \tau)$ and its partial derivatives with respect to χ upto fourth and τ upto first order are continuous on Π_T .

Let us reduce the IP (1.2)-(1.5) to the equivalent inverse problem to prove the existence and uniqueness theorem for the IP.

Lemma 2.2. *Let $z_0(\chi) \in C^2[0, 1]$, $m(\tau) \in C^1[0, T]$, $m(\tau) \neq 0$ for all $\tau \in [0, T]$ and*

$$z_0(0) = z_0(1) = z_0''(0) = z_0''(1) = 0,$$

$$\int_0^1 z_0(\chi)d\chi = m(0),$$

are satisfied. Then the problem of finding the classical solution of the IP (1.2)-(1.5) is equivalent to the problem of finding the classical solution of the IP from the eq. (1.2) with the IC (1.3), the BCs (1.4) and the AC

$$m'(\tau) + \beta(1 - \beta)R \int_0^1 Z_{\chi\chi\chi\chi}(\chi, \tau)d\chi - \beta D \int_0^1 Z_{\chi\chi}(\chi, \tau)d\chi = k(\tau)m(\tau) + \int_0^1 h(\chi, \tau)d\chi. \tag{2.1}$$

Proof. It is obvious that if the pair $\{Z(\chi, \tau), k(\tau)\}$ is the classical solution of the IP (1.2)-(1.5), then the AC (2.1) is also satisfied.

Let us show that the condition (1.5) is fulfilled if the pair $\{Z(\chi, \tau), k(\tau)\}$ is a classical solution of the IP (1.2)-(1.4) with the AC (2.1).

Integrating the eq. (1.2) with respect to χ from 0 to 1 yields

$$\int_0^1 Z_\tau(\chi, \tau) d\chi + \beta(1-\beta)R \int_0^1 Z_{\chi\chi\chi\chi}(\chi, \tau) d\chi - \beta D \int_0^1 Z_{\chi\chi}(\chi, \tau) d\chi = k(\tau) \int_0^1 Z(\chi, \tau) d\chi + \int_0^1 h(\chi, \tau) d\chi. \quad (2.2)$$

Considering the difference of the equations (2.2) and (2.1) gives the following initial value problem

$$\begin{cases} Y'(\tau) - k(\tau)Y(\tau) = 0, & \tau \in [0, T], \\ Y(0) = 0, \end{cases} \quad (2.3)$$

where $Y(\tau) = \int_0^1 Z(\chi, \tau) d\chi - m(\tau)$. The Cauchy problem (2.3) has only a trivial solution $Y(\tau) = 0$. i.e.

$$\int_0^1 Z(\chi, \tau) d\chi = m(\tau), \quad \tau \in [0, T].$$

Thus the condition (1.5) is fulfilled. \square

3. Unique Solvability of the IP

The elements of the system $\{y_n(\chi)\}_{n=1}^\infty = \{\sqrt{2} \sin(\mu_n \chi)\}_{n=1}^\infty$ are bi-orthonormal on $[0, 1]$ with $\mu_n = n\pi$ for each $n = 1, 2, \dots$, i.e.:

$$(y_n(\cdot), y_m(\cdot)) = \int_0^1 y_n(\chi) y_m(\chi) d\chi = \begin{cases} 1, & m = n \\ 0, & m \neq n \end{cases}.$$

Also the system $\{y_n(\chi)\}_{n=1}^\infty$ is complete and forms a Riesz basis in $L_2[0, 1]$.

The following lemma is useful to prove the unique solvability of the IP:

Lemma 3.1. *Suppose that the assumptions*

$$\begin{aligned} \mathbf{A}_1 \quad & z_0(\chi) \in C^3[0, 1], \quad z_0^{(4)}(\chi) \in L_2[0, 1], \quad z_0(0) = z_0(1) = z_0''(0) = z_0''(1) = 0, \\ \mathbf{A}_2 \quad & h(\chi, \tau) \in C^{3,0}(\Pi_T), \quad h_{\chi\chi\chi\chi}(\cdot, \tau) \in L_2[0, 1], \quad h(0, \tau) = h(1, \tau) = h_{\chi\chi}(0, \tau) = h_{\chi\chi}(1, \tau) = 0, \end{aligned}$$

are satisfied. Then

$$\sum_{n=1}^\infty \mu_n^3 |z_{0n}| \leq c_1 \|z_0^{(4)}\|_{L_2[0,1]}, \quad \sum_{n=1}^\infty \mu_n |z_{0n}| \leq c_2 \|z_0^{(4)}\|_{L_2[0,1]},$$

and

$$\sum_{n=1}^\infty \mu_n^3 |h_n(\tau)| \leq c_1 \|h_{\chi\chi\chi\chi}(\cdot, \tau)\|_{L_2[0,1]}, \quad \sum_{n=1}^\infty \mu_n |h_n(\tau)| \leq c_2 \|h_{\chi\chi\chi\chi}(\cdot, \tau)\|_{L_2[0,1]},$$

hold. Here $z_{0n} = (z_0(\cdot), y_n(\cdot))$, $h_n(\tau) = (h(\cdot, \tau), y_n(\cdot))$, and

$$c_1 = \left(\sum_{n=1}^\infty \frac{1}{\mu_n^2} \right)^{1/2} = \frac{1}{\sqrt{6}}, \quad c_2 = \left(\sum_{n=1}^\infty \frac{1}{\mu_n^6} \right)^{1/2} = \frac{1}{\sqrt{945}}.$$

Proof. Let us prove first one of these estimates. The others can be proved analogously. By applying integration by parts four times we get

$$\mu_n^3 z_{0n} = \mu_n^3 (z_0(\cdot), y_n(\cdot)) = \mu_n^3 \int_0^1 z_0(\chi) y_n(\chi) d\chi = \mu_n^3 \frac{1}{\mu_n^4} \int_0^1 z_0^{(4)}(\chi) y_n(\chi) d\chi = \frac{1}{\mu_n} \int_0^1 z_0^{(4)}(\chi) y_n(\chi) d\chi = \frac{1}{\mu_n} (z_0^{(4)}(\cdot), y_n(\cdot)).$$

By using Cauchy-Schwartz and Bessel inequalities

$$\sum_{n=1}^\infty \mu_n^3 |z_{0n}| = \sum_{n=1}^\infty \frac{1}{\mu_n} \left| (z_0^{(4)}(\cdot), y_n(\cdot)) \right| \leq \left(\sum_{n=1}^\infty \frac{1}{\mu_n^2} \right)^{1/2} \left(\sum_{n=1}^\infty \left| (z_0^{(4)}(\cdot), y_n(\cdot)) \right|^2 \right)^{1/2} \leq c_1 \|z_0^{(4)}\|_{L_2[0,1]}$$

with $c_1 = \left(\sum_{n=1}^\infty \frac{1}{\mu_n^2} \right)^{1/2} = \frac{1}{\sqrt{6}}$. \square

Since the system $\{y_n(\chi)\}_{n=1}^\infty$ forms a Riesz basis, we can seek the solution of the IP (1.2)-(1.4) and (2.1) as

$$Z(\chi, \tau) = \sum_{n=1}^\infty Z_n(\tau)y_n(\chi), \tag{3.1}$$

where

$$Z_n(\tau) = \int_0^1 Z(\chi, \tau)y_n(\chi)d\chi.$$

Applying Fourier method yields $Z_n(\tau)$, $n = 1, 2, \dots$ satisfy the following initial-value problems (IVPs)

$$\begin{cases} Z'_n(\tau) + K_n Z_n(\tau) = H_n(\tau; Z, k), \\ Z_n(0) = z_{0n}, n = 1, 2, \dots, \end{cases} \tag{3.2}$$

where $K_n = \mu_n^4 \beta(1 - \beta)R + \mu_n^2 \beta D$, $H_n(\tau; Z, k) = k(\tau)Z_n(\tau) + h_n(\tau)$.

The solution of the IVPs (3.2) are

$$Z_n(\tau) = z_{0n} \exp(-K_n \tau) + \int_0^\tau H_n(s; Z, k) \exp(-K_n(\tau - s)) ds. \tag{3.3}$$

Substituting (3.3) into (3.1) we get the classical solution of the IP (1.2)-(1.4) (or the first component of the pair $\{Z(\chi, \tau), k(\tau)\}$) as

$$Z(\chi, \tau) = \sum_{n=1}^\infty \left[z_{0n} \exp(-K_n \tau) + \int_0^\tau H_n(s; Z, k) \exp(-K_n(\tau - s)) ds \right] y_n(\chi). \tag{3.4}$$

To derive the equations for the unknown coefficient $k(\tau)$ consider

$$\int_0^1 Z_{\chi\chi\chi\chi}(\chi, \tau) dx = \int_0^1 \sum_{n=1}^\infty \mu_n^4 Z_n(\tau) y_n(\chi) d\chi,$$

and

$$\int_0^1 Z_{\chi\chi}(\chi, \tau) dx = - \int_0^1 \sum_{n=1}^\infty \mu_n^2 Z_n(\tau) y_n(\chi) d\chi,$$

into the AC (2.1). Then we get

$$\begin{aligned} k(\tau) &= \frac{1}{m(\tau)} [m'(\tau) - h_{int}(\tau) \\ &+ \sqrt{2} \sum_{n=1}^\infty \frac{(1 - (-1)^n) K_n}{\mu_n} \left\{ z_{0n} \exp(-K_n \tau) + \int_0^\tau H_n(s; Z, a) \exp(-K_n(\tau - s)) ds \right\}], \end{aligned} \tag{3.5}$$

where $h_{int}(\tau) = \int_0^1 h(\chi, \tau) d\chi$.

The equations (3.4) and (3.5) are Volterra type integral equations with regard to $Z(\chi, \tau)$, and $k(\tau)$ and form the system of integral equations. The inverse problem (1.2)-(1.4) and (2.1) reduced to the system of equations (3.4) and (3.5). Thus, we can conclude that solving the system of integral equations (3.4) and (3.5) and the inverse problem (1.2)-(1.4) and (2.1) are equivalent.

Before setting and proving the existence and uniqueness theorem of the solution of the system (3.4) and (3.5), let us give the following Banach spaces which will be used in the proof of the main theorem:

I

$$\begin{aligned} B_T &= \left\{ Z(\chi, \tau) = \sum_{n=1}^\infty Z_n(\tau) y_n(\chi) : Z_n(\tau) \in C[0, T], \right. \\ &\left. J_T(Z) = \left(\sum_{n=1}^\infty (\mu_n^4 \|Z_n\|_{C[0, T]})^2 \right)^{1/2} < +\infty \right\}, \end{aligned}$$

and the norm of $Z(\chi, \tau)$ defined as $\|Z\|_{B_T} \equiv \left(\sum_{n=1}^\infty (\mu_n^4 \|Z_n\|_{C[0, T]})^2 \right)^{1/2}$.

II $E_T = B_T \times C[0, T]$ is the space of the all pairs $\kappa(\chi, \tau) = \{Z(\chi, \tau), k(\tau)\}$ and the norm is

$$\|\kappa\|_{E_T} = \|Z\|_{B_T} + \|k\|_{C[0, T]}.$$

Theorem 3.2. *Let the assumptions of the Lemma 2.2 and 3.1 are satisfied. Then, the system (3.4) and (3.5) (or the IP (1.2)-(1.4) and (2.1)) has a unique solution for small T .*

Proof. Let $\kappa(\chi, \tau) = \{Z(\chi, \tau), k(\tau)\}$ is an arbitrary element belongs to $E_T = B_T \times C[0, T]$. Then the system of equations (3.4) and (3.5) we can be rewritten into the operator equation form as

$$\kappa = \Phi(\kappa)$$

where $\Phi(\kappa) \equiv \{\phi_1, \phi_2\}$ and

$$\phi_1(\kappa) = \sum_{n=1}^{\infty} \left[z_{0n} \exp(-K_n \tau) + \int_0^{\tau} H_n(s; Z, k) \exp(-K_n(\tau - s)) ds \right] y_n(\chi),$$

and

$$\begin{aligned} \phi_2(\kappa) = & \frac{1}{m(\tau)} [m'(\tau) - h_{im}(\tau) \\ & + \sqrt{2} \sum_{n=1}^{\infty} \frac{(1 - (-1)^n) K_n}{\mu_n} \left\{ z_{0n} \exp(-K_n \tau) + \int_0^{\tau} H_n(s; Z, k) \exp(-K_n(\tau - s)) ds \right\}]. \end{aligned}$$

It is obvious that

$$K_n = \mu_n^4 \beta(1 - \beta)R + \mu_n^2 \beta D > 0, \quad n = 1, 2, \dots$$

Thus

$$\exp(-K_n \tau) \leq 1, \text{ and } \exp(-K_n(\tau - s)) \leq 1 \text{ for } 0 \leq s \leq \tau, 0 \leq \tau \leq T.$$

Let us prove that Φ is a contraction operator in two steps by considering these estimates.

I) Firstly prove that Φ is a continuous onto map on $E_T \rightarrow E_T$. It means that we require to prove $\phi_1(\kappa) \in B_T$ and $\phi_2(\kappa) \in C[0, T]$ for an arbitrary element $\kappa(x, \tau) = \{Z(\chi, \tau), k(\tau)\} \in E_T$ with $Z(\chi, \tau) \in B_T, k(\tau) \in C[0, T]$.

Let us verify that

$$J_T(\phi_1) = \left(\sum_{n=1}^{\infty} (\mu_n^4 \|\phi_1\|_{C[0, T]})^2 \right)^{1/2} < +\infty,$$

where $(\phi_1)_n(\tau) = RHS(Z_n(\tau))$. In other words let us show that $\phi_1(\kappa) \in B_T$. Under the assumptions of the Lemma 3.1 and Theorem 3.2, we obtain

$$J_T(\phi_1) \leq 2 \left\| z_0^{(4)} \right\|_{L_2[0, 1]} + 2T \|h_{\chi\chi\chi\chi}\|_{L_2[0, 1]} + 2T \|k\|_{C[0, T]} \|Z\|_{B_T}. \tag{3.6}$$

Since $Z(\chi, \tau) \in B_T$, and $k(\tau) \in C[0, T]$, the norms $\|k\|_{C[0, T]}, \|Z\|_{B_T}$ are finite. Therefore, $J_T(\phi_1)$ is also finite. Thus $\phi_1(\kappa)$ belongs to the space B_T .

Now let's demonstrate that $\phi_2(\kappa) \in C[0, T]$. From the eq. (3.5), we obtain

$$\begin{aligned} \|\phi_2(\kappa)\|_{C[0, T]} \leq & \frac{1}{\alpha} \left[\|m'\|_{C[0, T]} + \|h_{im}\|_{C[0, T]} \right. \\ & \left. + M \left(\left\| z_0^{(4)} \right\|_{L_2[0, 1]} + T \|h_{\chi\chi\chi\chi}\|_{L_2[0, 1]} + T \|k\|_{C[0, T]} \|Z\|_{B_T} \right) \right] \end{aligned} \tag{3.7}$$

where $\min_{0 \leq \tau \leq T} |m(\tau)| \geq \alpha > 0, M = 2\sqrt{2}(c_1\beta(1 - \beta)R + c_2\beta D)$.

As in the previous part we can conclude that $\|\phi_2(\kappa)\|_{C[0, T]}$ is bounded. So $\phi_2(\kappa) \in C[0, T]$.

From the eqs. (3.6) and (3.7), we can obtain

$$\|\Phi(\kappa)\|_{E_T} \leq L_1(T) + L_2(T) \|k\|_{C[0, T]} \|Z\|_{B_T},$$

where

$$L_1(T) = \left(2 + \frac{M}{\alpha}\right) \|z_0^{(4)}\|_{L_2[0,1]} + T \left(2 + \frac{M}{\alpha}\right) \|h_{\chi\chi\chi\chi}\|_{L_2[0,1]} + \frac{1}{\alpha} \left[\|m'\|_{C[0,T]} + \|h_{int}\|_{C[0,T]}\right],$$

$$L_2(T) = T \left(2 + \frac{M}{\alpha}\right).$$

Since $L_2(T)$ tends to zero as $T \rightarrow 0$, and $L_1(T)$ is a continuous function of T , there exists a sufficiently small $T > 0$ such that

$$(L_1(T) + 1)^2 L_2(T) < 1.$$

Let us define a ball $A := \left\{ \kappa \in E_T : \|\kappa\|_{E_T} \leq L_1(T) + 1 \right\}$ for the fixed T . Then, for every $\kappa \in A$, we get

$$\begin{aligned} \|\Phi(\kappa)\|_{E_T} &\leq L_1(T) + L_2(T) \|k\|_{C[0,T]} \|Z\|_{B_T} \\ &\leq L_1(T) + L_2(T) (L_1(T) + 1)^2 < L_1(T) + 1. \end{aligned}$$

Thus, Φ is an onto continuous map on E_T .

II) This step our aim is to prove that the operator Φ is a contraction mapping operator. Assume that let z_1 and z_2 be any two elements of E_T . According to the definition of the space E_T yields $\|\Phi(\kappa_1) - \Phi(\kappa_2)\|_{E_T} = \|\phi_1(\kappa_1) - \phi_1(\kappa_2)\|_{B_T} + \|\phi_2(\kappa_1) - \phi_2(\kappa_2)\|_{C[0,T]}$. Here $\kappa_i = \{Z^i(\chi, \tau), k^i(\tau)\}$, $i = 1, 2$. consider the following differences

$$\phi_1(\kappa_1) - \phi_1(\kappa_2) = \sum_{n=1}^{\infty} \left[\int_0^{\tau} [H_n(s; Z^1, k^1) - H_n(s; Z^2, k^2)] \exp(-K_n(\tau - s)) ds \right] y_n(\chi),$$

and

$$\phi_2(\kappa_1) - \phi_2(\kappa_2) = \frac{\sqrt{2}}{m(\tau)} \sum_{n=1}^{\infty} \frac{(1 - (-1)^n) K_n}{\mu_n} \int_0^{\tau} [H_n(s; Z^1, k^1) - H_n(s; Z^2, k^2)] \exp(-K_n(\tau - s)) ds.$$

We can obtain from the last equations

$$\|\phi_1(\kappa_1) - \phi_1(\kappa_2)\|_{B_T} \leq 2T \left(\|Z^2\|_{B_T} \|k^1 - k^2\|_{C[0,T]} + \|k^1\|_{C[0,T]} \|Z^1 - Z^2\|_{B_T} \right),$$

and

$$\|\phi_2(\kappa_1) - \phi_2(\kappa_2)\|_{C[0,T]} \leq \frac{TM}{\alpha} \left(\|Z^2\|_{B_T} \|k^1 - k^2\|_{C[0,T]} + \|k^1\|_{C[0,T]} \|Z^1 - Z^2\|_{B_T} \right).$$

From the these inequalities it follows that

$$\|\Phi(\kappa_1) - \Phi(\kappa_2)\|_{E_T} \leq L_2(T) C(k^1, Z^2) \|\kappa_1 - \kappa_2\|_{E_T},$$

where $L_2(T) = T \left(2 + \frac{M}{\alpha}\right)$, and $C(k^1, Z^2) = \max \left\{ \|k^1\|_{C[0,T]}, \|Z^2\|_{B_T} \right\} = L_1(T) + 1$.

Since $L_1(T) + 1 \leq (L_1(T) + 1)^2$,

$$0 < L_2(T) C(k^1, Z^2) = L_2(T) (L_1(T) + 1) \leq L_2(T) (L_1(T) + 1)^2 < 1,$$

i.e. $0 < L_2(T) C(k^1, Z^2) < 1$. Therefore, the operator Φ is contraction mapping operator.

Thus we can conclude that the operator Φ is contraction mapping and it is a continuous onto map on E_T . Then the solution of $\kappa = \Phi(\kappa)$ exists and unique regarding to Banach fixed point theorem. \square

Example 3.3. Consider the IP (1.2)-(1.5) with unknown smooth $k(\tau)$ and inputs:

$$z_0(\chi) = \sin(\pi\chi), \quad m(\tau) = \frac{2}{\pi} e^{\tau}, \quad h(\chi, \tau) = \left[(1 + \pi^4 \beta (1 - \beta) R + \pi^2 \beta D) e^{\tau} - 1 \right] \sin(\pi\chi).$$

These inputs satisfy the conditions of the Lemma 2.2 and Lemma 3.1, i.e.

$$z_0(\chi) \in C^3[0, 1], \quad z_0^{(4)}(\chi) \in L_2[0, 1], \quad z_0(0) = z_0(1) = z_0''(0) = z_0''(1) = 0,$$

$$h(\chi, \tau) \in C^{3,0}(\Pi_T), \quad h_{\chi\chi\chi\chi}(\cdot, \tau) \in L_2[0, 1], \quad h(0, \tau) = h(1, \tau) = h_{\chi\chi}(0, \tau) = h_{\chi\chi}(1, \tau) = 0,$$

and

$$m(\tau) \in C^1[0, T], \quad m(\tau) \neq 0 \text{ for all } \tau \in [0, T] \text{ and } \int_0^1 z_0(\chi) d\chi = m(0) = \frac{2}{\pi}.$$

According to the Theorem 3.2, the solution of the inverse problem (1.2)-(1.5) exists and unique and the solution is

$$\{Z(\chi, \tau), k(\tau)\} = \{e^{\tau} \sin(\pi\chi), e^{-\tau}\}.$$

4. Conclusion

The manuscript studies the IP of obtaining the solely time-dependent zero-order coefficient in a linear Bi-flux diffusion equation from an extra integral observation. The theorem of existence and uniqueness of the solution of the IP is proved for an adequately small time interval by applying the contraction mapping principle. This work is new and has never been studied theoretical or numerical before. The numerical method of the IP will be considered with a suitable scheme as a future work.

Article Information

Acknowledgements: The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

Author's Contributions: The authors contributed equally to the writing of this paper.

Conflict of Interest Disclosure: No potential conflict of interest was declared by the author.

Copyright Statement: Authors own the copyright of their work published in the journal and their work is published under the CC BY-NC 4.0 license.

Supporting/Supporting Organizations: No grants were received from any public, private or non-profit organizations for this research.

Ethical Approval and Participant Consent: It is declared that during the preparation process of this study, scientific and ethical principles were followed and all the studies benefited from are stated in the bibliography.

Plagiarism Statement: This article was scanned by the plagiarism program. No plagiarism detected.

Availability of Data and Materials: Not applicable.

References

- [1] S. Guenneau, D. Petiteau, M. Zerrad, C. Amra, T. Puvirajesinghe, *Transformed Fourier and Fick equations for the control of heat and mass diffusion*, AIP Adv., **5**(5) (2015), 053404.
- [2] D. Shi, S. Zhang, *Unconditional superconvergence of the fully-discrete schemes for nonlinear prey-predator model*, Appl. Numer. Math., **172** (2022), 118-132.
- [3] D. Shi, C. Li, *Superconvergence analysis of two-grid methods for bacteria equations*, Numer. Algorithms, **86**(1) (2021), 123-152.
- [4] L. Bevilacqua, M. Jiang, A. Silva Neto, A.C.R. Galeão, *An evolutionary model of bi-flux diffusion processes*, J. Braz. Soc. Mech. Sci. Eng., **38**(5) (2016), 1421-1432.
- [5] M. Jiang, L. Bevilacqua, A.J.S. Neto, A.C.R. Galeão, J. Zhu, *Bi-flux theory applied to the dispersion of particles in anisotropic substratum*, Appl. Math. Model., **64** (2018), 121-134.
- [6] M. Jiang, L. Bevilacqua, J. Zhu, X. Yu, *Nonlinear Galerkin finite element methods for fourth-order Bi-flux diffusion model with nonlinear reaction term*, Comput. Appl. Math., **39**(3) (2020), 1-16.
- [7] D. Shi, S. Zhang, *Unconditional superconvergence analysis for the nonlinear Bi-flux diffusion equation*, Appl. Math. Comput., **442** (2023), 127771.
- [8] S. Guenneau, C. Amra, D. Veynante, *Transformation thermodynamics: cloaking and concentrating heat flux*, Opt. Express, **20**(7) (2012), 8207-8218.
- [9] E.P. Scott, P.S. Robinson, T.E. Diller, *Development of methodologies for the estimation of blood perfusion using a minimally invasive thermal probe*, Meas. Sci. Technol., **9**(6) (1998), 888.
- [10] L. Marin, L. Elliott, P.J. Heggs, D.B. Ingham, D. Lesnic, X. Wen, *Analysis of polygonal fins using the boundary element method*, Appl. Therm. Eng., **24**(8-9) (2004), 1321-1339.
- [11] E.A. Sudicky, *The Laplace transform Galerkin technique: A time-continuous finite element theory and application to mass transport in groundwater*, Water Resour. Res., **25**(8) (1989), 1833-1846.
- [12] M.I. Ismailov, *Direct and inverse problems for thermal grooving by surface diffusion with time dependent mullins coefficient*, Math. Model. Anal., **26**(1) (2021), 135-147.
- [13] J.R. Cannon, *Determination of an unknown heat source from overspecified boundary data*, SIAM J. Numer. Anal., **5**(2) (1968), 275-286.
- [14] A. Hasanov, *Simultaneous determination of source terms in a linear parabolic problem from the final overdetermination: weak solution approach*, J. Math. Anal. Appl., **330** (2007), 766-779.
- [15] A. Hazanee, D. Lesnic, M.I. Ismailov, N.B. Kerimov, *An inverse time-dependent source problem for the heat equation with a non-classical boundary condition*, Appl. Math. Model., **39**(20) (2015), 6258-6272.
- [16] M.I. Ismailov, F. Kanca, D. Lesnic, *Determination of a time-dependent heat source under nonlocal boundary and integral overdetermination conditions*, Appl. Math. Comput., **218**(8) (2011), 4138-4146.
- [17] J.R. Cannon, Y. Lin, S. Wang, *Determination of a control parameter in a parabolic partial differential equation*, J. Aust. Math. Soc. Series B, **33** (1991), 149-163.
- [18] M.I. Ismailov, I. Tekin, S. Erkovan, *An inverse problem for finding the lowest term of a heat equation with Wentzell-Neumann boundary condition*, Inverse Probl. Sci. Eng., **27**(11) (2019), 1608-1634.
- [19] M.I. Ismailov, F. Kanca, *An inverse coefficient problem for a parabolic equation in the case of nonlocal boundary and overdetermination conditions*, Math. Methods Appl. Sci., **34**(6) (2011), 692-702.
- [20] N.B. Kerimov, M.I. Ismailov, *An inverse coefficient problem for the heat equation in the case of nonlocal boundary conditions*, J. Math. Anal. Appl., **396**(2) (2012), 546-554.
- [21] L. Zhuo, D. Lesnic, M.I. Ismailov, I. Tekin, S. Meng, *Determination of the time-dependent reaction coefficient and the heat flux in a nonlinear inverse heat conduction problem*, Int. J. Comput. Math., **96**(10) (2019), 2079-2099.