

## Comparison of Neck Awareness, Physical and Psychosocial Parameters in Inactive University Students with and without Neck Pain

Boyun Ağrısı Olan ve Olmayan İnaktif Üniversite Öğrencilerinde Boyun Farkındalığı, Fiziksel ve Psikososyal Parametrelerinin Karşılaştırılması

Raziye ERKAN<sup>1\*</sup>, Emine ASLAN TELCİ<sup>2</sup>, Sebahat Yaprak ÇETİN<sup>3</sup>, Fatmanur ALTIN<sup>4</sup>

<sup>1</sup>Burdur Mehmet Akif Ersoy University, Rectorate Department of Physical Education and Sports, Burdur, Türkiye

<sup>2</sup>Pamukkale University, Faculty of Physical Therapy and Rehabilitation, Department of Physical Therapy and Rehabilitation, Denizli, Türkiye

<sup>3</sup>Pamukkale University, Faculty of Health Sciences, Department of Physical Therapy and Rehabilitation, Antalya, Türkiye

<sup>4</sup>Adnan Menderes University, Vocational School of Health Services, Department of Physical Therapy and Rehabilitation, Aydın, Türkiye

**Abstract:** The aim of the study was to compare neck awareness, muscular endurance, mental state and self-efficacy parameters in young adults with and without neck pain and to examine the relationship of awareness with pain intensity, muscular endurance and anxiety and depression in the group with neck pain. A total of 104 students with neck pain (n=41) and without neck pain (n=63) with a mean age of  $22.85 \pm 3.53$  years were included in the study. Visual Analog Scale (VAS), Cervical Deep Flexor Muscle Endurance Test, Hospital Anxiety and Depression Scale (HADS) and Fremantle Neck Awareness Questionnaire (FreBAQ) were used to collect the data. When the groups were compared in terms of physical and psychosocial data, it was determined that the group with neck pain was negatively affected compared to the group without neck pain in terms of all parameters ( $p < 0.05$ ). In the group with neck pain, there was a positive correlation between the FreBAQ and the HAD scale values ( $p < 0.05$ ); there was no correlation between deep flexor neck muscular endurance and neck pain ( $p > 0.05$ ). According to the results of the study, neck pain negatively affects physical and psychological factors in university students and neck awareness is especially associated with anxiety and depression.

**Keywords:** Young Adult, Neck Pain, Neck Awareness, Mental State, Muscular Endurance.

**Öz:** Çalışmanın amacı, boyun ağrısı olan ve olmayan genç yetişkin gruplarda boyun farkındalığı, kassal endurans, ruhsal durum ve öz-yeterlilik parametrelerinin karşılaştırılması ve boyun ağrılı grupta farkındalığın ağrı şiddeti, kassal endurans ve anksiyete depresyon ile ilişkisinin incelenmesidir. Çalışmaya yaş ortalaması  $22,85 \pm 3,53$  yıl olan, boyun ağrılı (n=41) ve boyun ağrısı olmayan (n=63) toplam 104 öğrenci dahil edilmiştir. Çalışmanın verilerinin toplanmasında Görsel Analog Skala (GAS), Servikal Derin Fleksör Kas Endurans Testi, Hastane Anksiyete ve Depresyon Ölçeği (HADÖ) ve Fremantle Boyun Farkındalık Anketi (FreBFA) kullanılmıştır. Gruplar fiziksel ve psikososyal veriler açısından karşılaştırıldıklarında tüm parametreler açısından boyun ağrısı olan grubun boyun ağrısı olmayan gruba göre olumsuz yönde etkilendiği belirlenmiştir ( $p < 0,05$ ). Boyun ağrısı olan grupta, Fremantle Boyun Farkındalık Testi ile HAD ölçeği değerleri arasında pozitif bir korelasyon olduğu ( $p < 0,05$ ); derin fleksör boyun kassal enduransı ile boyun ağrısı arasında ise bir korelasyon olmadığı bulunmuştur ( $p > 0,05$ ). Çalışmanın sonuçlarına göre boyun ağrısının üniversite öğrencilerinde fiziksel ve psikolojik faktörleri olumsuz yönde etkilediği ve boyun farkındalığının özellikle anksiyete ve depresyon ile ilişkili olduğu görülmüştür.

**Anahtar Kelimeler:** Genç Yetişkin, Boyun Ağrısı, Boyun Farkındalığı, Ruhsal Durum, Kassal Endurans.

\*Corresponding author : Raziye ERKAN

e-mail : rerkan@mehmetakif.edu.tr

Geliş tarihi / Received : 09.03.2023

Kabul tarihi / Accepted: 24.04.2023

## Introduction

Chronic neck pain is a common problem affecting 75% of the general population (Fejer et al., 2006). One third of the young adult population complains of neck pain at least once a week (Hogg-Johnson et al., 2008). In a study conducted in university students aged 18- 25 years in our country, it was reported that the lifetime incidence of neck pain in students was 43.70% (Kurtaran et al., 2019). According to the results of studies on chronic pain prevention, neck pain ranks first among the reasons for seeking health care (Fricton et al., 2015). Within a bio-psycho-social framework, a number of factors cause neck pain, and these factors may also contribute to the transition from an acute pain state to a chronic pain state (Nagrle et al., 2010, Lau et al., 2011).

Mechanical causes (Hidelgo et al., 2017), myofascial disorders in the neck/shoulder region (Gallego et al., 2020) are among the factors that cause neck pain. Neck muscle endurance has been found to be associated with chronic neck pain (Schomacher et al., 2013, Kahlaee et al., 2017). Factors such as increased long-term use of phones and computers in inappropriate positions in young populations cause postural disorders and musculoskeletal pain (Sihawong et al., 2011, Kim et al., 2015). It has been reported that the effect of impaired body perception on the severity of the clinical condition makes a more significant contribution than other factors affecting the clinical condition. In case of impaired body perception, an unconscious awareness occurs and conscious perception is impaired (Wand et al., 2016). Studies have shown that improving habitual postural patterns can improve musculoskeletal pain or prevent further deterioration. However, changing habitual patterns requires awareness (Cramer et al., 2018). Researchers state that chronic pain is associated with impaired body awareness. This suggests that mindfulness should be examined in chronic pain (Wand et al., 2016).

How the body perceives and experiences pain has recently attracted attention. The response to pain is related to perceived health, tolerance, and beliefs

about the risk associated with a particular movement or activity (Wiech et al., 2004). Self-efficacy belief is an important factor affecting pain response (Bandura, 1994). In addition, self-efficacy is also effective in the formation and change of patients' health-related behaviors (Yıldırım et al., 2010). There are examples of studies examining patient behaviors related to self-efficacy in the field of health, but there are not many studies on neck pain.

Neck pain may be caused by psychosocial factors as well as physical factors (Linton 2000). Emotional states may affect body positions and functions of extremities. The presence of depression and anxiety is thought to be a factor in the onset and chronicity of neck pain and in increasing the severity of pain (Shahidi et al., 2015). Therefore, it is important to measure the presence of anxiety and depression in patients.

Health-related behaviors acquired in the early period of life affect the risks for lifestyle-related disorders in the future and the incidence of preventable diseases that may cause problems in later ages (Von Bothmer et al., 2005). In our country, university students represent a large portion of the young adult population (Savcı et al., 2010). It is important to identify potential pain factors among young adults and thus prevent the development of neck pain later in life (Kanchanomai et al., 2011).

The primary aim of this study was to compare neck awareness, muscular endurance, anxiety-depression and self-efficacy in young adults with and without neck pain. The secondary aim was to investigate the relationship of mindfulness with pain intensity, muscular endurance, self-efficacy, anxiety and depression in a group with neck pain.

## Materials and Methods

The population of the study was Pamukkale University Physiotherapy and Rehabilitation Faculty students. The sample of the study consisted of volunteer students who met the study criteria in this population.

The study was conducted on inactive volunteer students between the ages of 18-25. In order to carry out the study, all classes were informed by the researchers about the purpose of the study and the inclusion criteria and it was emphasized that the study was voluntary. Individuals who had neck pain for at least 3 months and scored above 0 on the Visual Analog Scale (n=41) and individuals who had not had neck pain for the last 6 months (n=63) were included participate in the study.

Exclusion criteria included a history of previous surgery on the spine, malignant conditions, neurologic and orthopedic diagnoses, musculoskeletal pain apart from cervical region and any condition that posed an obstacle to perform the assessments. At the same time, individuals with improve musculoskeletal pain other than the cervical region were exclusion criteria for both groups. There were no individuals who met the exclusion criteria among the students who volunteered to participate in the study.

Ethical approval was obtained from Pamukkale University Non-interventional Clinical Research Ethics Committee for the conduct of the study with the decision number 60116787- 020/92293 dated 25.12.2019. The study was conducted in accordance with the principles defined in the Declaration of Helsinki. Informed written consent was obtained from all participants.

**Sociodemographic Data Form.** Participants' age, gender, height, body weight, and body mass index were recorded on the socio-demographic form.

**Evaluation of pain intensity.** Pain intensity was assessed with the Visual Analog Scale (VAS). Participants were asked to mark the intensity of pain they felt on a 10 cm long horizontal line. The point marked on the line was measured with a ruler and the GAS value was recorded in cm (0 cm: No pain; 10 cm: Intolerable pain) (Wewers et al., 1990).

**General Health Status Assessment:** To assess general health status, participants were asked the

question "How would you rate your general health status?" and were asked to mark one of the answers "excellent, very good, fair, poor, poor" (Cavlak et al., 2009)

**Evaluation of cervical region deep flexor muscle endurance:** The 'Cervical Region Deep Flexor Muscle Endurance Test' was used for the evaluation. The test was performed in supine hook position. Individuals were asked to maximally retract the chin (*chin tuck* position) and lift the head and neck approximately 2.5 cm from the lying position while maintaining the retraction isometrically. The students were asked to maintain this position for as long as possible and the time to maintain the position was recorded in seconds. The test was terminated if the test position disappeared, if there was a sudden and severe increase in pain or if the person did not want to continue the test (Harris et al., 2005)

**Assessment of Neck Awareness:** Neck awareness was assessed with the Fremantle Neck Awareness Questionnaire. The Turkish validity and reliability of the questionnaire developed by Wand et al. was performed by Onan et al. (ICC0.711). The questionnaire is a Likert-type scale that evaluates individual-specific altered perception. The 9 statements that make up the scale are graded from 0 = Never/never feel this way, 1 = Rarely feel this way, 2 = Sometimes, or some times feel this way, 3 = Often feel this way, 4 = Always or most of the time feel this way. The total score ranges from 0-36, with an increase in score indicating a poor prognosis (Wand et al., 2016, Onan et al., 2019).

**Assessment of Self-Efficacy:** General Self-Efficacy Scale was used in the study. The Turkish validity and reliability of the General Self-Efficacy Scale developed by Sherer et al. (1982) was conducted by Yıldırım and İlhan (2010) (Yıldırım 2010). Items 2, 4, 5, 6, 7, 10, 11, 12, 14, 16 and 17 in the 17-item Likert format scale are reverse scored and the score of each question varies between 1-5. The total score of the scale can vary between 17-85; the higher the score, the higher the self-efficacy belief (Yıldırım 2010).

**Assessment of anxiety and depression:** The Hospital Anxiety and Depression Scale (HAD) developed by Zigmond et al. in 1983 was used in the study (Zigmond et al., 1983). The HAD consists of 14 items and assesses 2 different psychological states: anxiety and depression. The questionnaire is scored between 0 and 21 and the higher the score, the more mood disturbance. The Turkish validity and reliability of the scale was conducted by Aydemir et al. in 1997 (Aydemir et al., 1997).

### Data Analysis

Data analysis was performed with 22.0 SPSS package program. In the analysis of the demographic characteristics of the participants, % distributions, mean and standard deviations were given. According to whether the other data were normally distributed or not, the data of the groups

were compared with the T test in independent groups from parametric tests or Mann-Whitney U test from non-parametric tests.  $p < 0.05$  was accepted as significant. Pearson correlation analysis was used if the data were normally distributed, and Spearman correlation analysis was used when the data were not normally distributed.

### Results

The study included 44 female and 30 male students with a mean age of  $22.85 \pm 3.53$  years. The mean pain duration of the students in the neck pain group was 20.53 months, ranging from 3 months to 72 months, and the mean pain intensity was 4.86 according to the VAS pain scale. There was no difference between the socio-demographic data of individuals with and without neck pain ( $p > 0.05$ ) (Table 1).

**Table1. Comparison of socio-demographic data of individuals with and without neck pain**

	Neck pain (n=41)		No neck pain (n=63)		p
	$\bar{x}$	S.S.	$\bar{x}$	S.S.	
Age (years)	23,17	3,87	22,63	3,29	0,450
Body weight (kg)	64,87	11,91	64,42	14,52	0,867
Height (cm)	168	8	167	9	0,720
BMI (kg/m) <sup>2</sup>	22,84	3,17	22,75	3,35	0,891
Duration of pain (months)	20,53	11,64	-	-	
	<b>n</b>	<b>%</b>	<b>n</b>	<b>%</b>	
Gender	30	73,2	44	69,8	0,717
Woman	11	26,8	19	30,2	

BMI: Body Mass Index \* $p < 0.05$  \*Independent samples T test

**Table2. Comparison of general health status of individuals with and without neck pain**

General Health Status Questionnaire	Neck pain (n=41)		No neck pain (n=63)		p
	n	%	n	%	
Weak	1	2,4	0	0	<b>0,009*</b>
Middle	19	46,3	15	23,8	
Very good	18	43,9	40	63,5	
Perfect	3	7,3	8	12,7	

\* $p < 0.05$  \* Independent samples T test.

The general health status of the group without neck pain was found to be better than the group with neck pain ( $p < 0.05$ ), (Table 2).

When the groups were compared in terms of physical and psychosocial data, a significant difference was found in favor of the group without neck pain according to the results of deep flexor neck endurance test, HADE anxiety, depression sub-parameters and total value and FreBAQ questionnaire ( $p < 0.05$ ). There was no statistically

significant difference between the two groups in terms of self-efficacy scores ( $p > 0.05$ ) (Table 3).

In the neck pain group, a positive correlation was found between FreBAQ value and HADE-anxiety, HADE-depression and HADE-total values ( $p < 0.05$ ). There was no statistically significant relationship between FreBAQ value and deep neck flexor muscle endurance and neck pain ( $p > 0.05$ ) (Table 4).

**Table 3. Comparison of Physical and Psychosocial Data of Students with and without Neck Pain**

General Health Status Questionnaire	Neck pain (n=41)		No neck pain (n=63)		p
	n	%	n	%	
Weak	1	2,4	0	0	<b>0,009*</b>
Middle	19	46,3	15	23,8	
Very good	18	43,9	40	63,5	
Perfect	3	7,3	8	12,7	

$p < 0.05$  \*Independent groups T test; \*\*Mann Whitney-U, HADE: Hospital Anxiety and Depression Scale.

**Table 4. The Relationship Between Fremantle Neck Awareness Questionnaire Value and Pain Severity, Anxiety and Depression, Self-Efficacy and Muscular Endurance in Individuals with Neck Pain.**

	VAS Activity	HADE anxiety	HADE depression	HADE total	Neck endurance	Self- efficacy
Neck Awareness Scale	0,310	0,001**	0,015*	0,003**	0,912	-0,081

$p < 0.05$  HADE: Hospital Anxiety and Depression Scale, VAS: Visual Analog Scale.

## Discussion

The first results of this study showed that neck pain negatively affected physical, psychological parameters and neck awareness in young adults compared to young adults without neck pain. Another result we obtained from our study was that there was a relationship between neck awareness and psychological parameters in young adults with neck pain.

In recent years, there has been an increase in researches on how neck pain affects the cervical

motor system, posture and movement (Blomgren et al., 2018). Most studies have examined the relationship between neck pain and flexor muscle endurance and generally found that flexor muscle endurance is low in people with neck pain (Piper et al., 2009, Edmondston et al., 2011, Parazza et al., 2014). Falla et al. demonstrated the loss of endurance of deep cervical flexors using surface electromyography in patients with chronic neck pain and found that this condition caused pain (Falla et al., 2004). In the recent study, in parallel with the literature, it was determined that cervical

region flexor muscle endurance was lower in young individuals with neck pain compared to those without neck pain. We thought that the inclusion of training for cervical muscle endurance, which decreased with neck pain especially in young individuals, in the rehabilitation program may be protective against physical effects such as postural problems that may be seen in later ages.

In the literature, it is known that one of the underlying causes of chronic neck pain is psychosocial reasons and it is stated that these reasons should be investigated (Blozik et al., 2009 Carroll et al., 2008). Studies have shown that anxiety and depression symptoms are more common and severe in individuals with neck pain (Demyttenaere et al., 2007, Liu et al., 2014, Dimitriadis et al., 2015). Carroll et al. found a strong correlation between depressive symptoms and the onset of a pain episode and reported that the most depressed group had a fourfold increased risk of neck pain compared to the least depressed group (Carroll et al., 2008). Studies on psychological factors in neck pain have emphasized that psychological factors may cause neck pain or neck pain may cause psychological problems (Linton et al., 2000). In a study examining the relationship between neck pain, anxiety and depression in 448 patients, it was shown that depression and anxiety were closely related to recurrent neck pain (Blozik et al., 2009). In our study, anxiety and depression symptoms were found to be higher in the group with neck pain ( $p < 0.05$ ). The recent results are consistent with the literature. This may be due to the fact that patients with prolonged and severe neck pain are negatively affected in terms of mood.

It has been reported that self-efficacy is associated with patients' ability to continue daily activities in the face of obstacles such as pain; therefore, it is an important predictive factor for pain (Nicholas et al., 2007). Self-efficacy has been found to significantly mediate the relationship between pain and disability (Lee et al., 2015). Greater self-efficacy may enable individuals to recognize and reduce some of the underlying causes of

musculoskeletal pain, such as poorly coordinated posture and movement habits, excessive muscle tension, and associated psychological distress (Woodman et al., 2018). Studies have shown that neck pain and self-efficacy are interrelated (Chiarotto et al., 2018, Monticone et al., 2021). The recent , no significant difference was found between the groups with and without neck pain and the general self-efficacy scale. The reasons for this may be explained by the fact that the individuals who make up the population of my study are young university students; in other words, apart from the pain, they are young individuals, and the opportunities provided by university education, such as academic and social opportunities, are equivalent, and different situations enable them to have the same level of self-efficacy.

Studies have reported that chronic pain is associated with the deterioration of the perceived body image of painful body parts and that this deterioration leads to an increase in chronic pain severity and prolongation of pain duration, and that pain severity decreases as awareness increases (Moreira et al., 2017, Cramer et al., 2018). In the first of two different studies examining the contribution of neck awareness to the relationship between neck awareness and chronic neck pain in the literature, it was concluded that body awareness was affected in individuals with chronic neck pain (Özel 2022). The recent, it was determined that the awareness levels of students with neck pain were lower. Another study found that pain intensity was weakly associated with neck awareness in individuals with chronic neck pain (Şimşek et al., 2022). The recent, it was found that there was no difference between awareness level and pain intensity. It was thought that the reason for the difference between the two studies in terms of pain intensity may be due to the sociodemographic characteristics of the subjects participating in the study.

When the literature is examined, it is seen that there are few studies examining the relationship between muscle endurance and neck awareness and different opinions have been put forward on

this issue. In Dere's study, it was found that there was a relationship between neck awareness and cervical flexor, trunk, upper extremity and scapular region muscular endurance in individuals with neck pain (Dere 2021). Şimşek et al. found a weak positive correlation between neck pain and decreased neck awareness (Şimşek et al., 2022). In a different study conducted with patients with chronic neck pain, no relationship was found between neck awareness and cervical muscle endurance in individuals with chronic neck pain (Özel 2022). Similarly, in our study, there was no relationship between muscular endurance and awareness. It is thought that this may be related to the evaluation of not only cervical endurance but also body parts adjacent to this region with neck awareness.

Anxiety can exacerbate chronic pain, while reducing anxiety and anxiety levels can mediate recovery (Linton 2000). In the study by Özel et al. a weak positive relationship was found between neck awareness and anxiety, while no relationship was found between neck awareness and depression (Özel 2022). The researchers associated this situation with the borderline anxiety and depression of the individuals with neck pain who participated in the study. Onan et al. stated that chronic neck pain contributes to the deterioration of neck awareness and perception in individuals (Onan et al., 2019). According to the results of our study, neck awareness was found to be positively associated with anxiety and depression symptoms. The relationship we found between awareness and mental status in young individuals reveals the importance of psychological support with a multidisciplinary approach as well as interventions for physical parameters in individuals with neck pain.

The most important limitation of our study was that possible postural errors such as head posture, which may be related to neck awareness in individuals with neck pain, and severity of disability were not examined. In the literature, interest in examining neck awareness, especially with patient-reported questionnaires, has recently increased. In order to determine the effect of neck

pain on awareness, future studies should examine various factors that may affect neck awareness in populations with different characteristics such as age and gender. The strength of our study was that it was a study examining the physical and psychosocial effects of neck pain on neck awareness in young adult groups. It was thought that defining neck awareness and psychosocial status in young adult groups will help to determine the necessary treatment programs to take precautions for neck pain later in life.

## Conclusion

According to the results of this study, neck endurance and neck awareness levels were lower and anxiety and depression levels were higher in students with neck pain compared to students without neck pain. In addition, neck awareness was associated with anxiety and depression in young people with neck pain. The results of this study revealed that muscular endurance, emotional state and neck awareness should be included in the evaluation parameters of young individuals with neck pain and the importance of developing appropriate treatment strategies with a multidisciplinary approach.

## References

- Aydemir, Ö., Güvenir, T., Küey, L., Kültür, S., 1997. Reliability and Validity of the Turkish version of Hospital Anxiety and Depression Scale. *Türk Psikiyatri Dergisi* 8(4), 280 -287.
- Bandura A., 1994. Self-efficacy. *Encyclopedia of Human Behavior*, New York. Academic Press, pp. 71-81.
- Blomgren, J., Strandell, E., Jull, G., Vikman, I., Röijezon, U., 2018. Effects of deep cervical flexor training on impaired physiological functions associated with chronic neck pain: a systematic review. *BMC musculoskeletal disorders*, 19(1):415.
- Blozik, E., Laptinskaya, D., Herrmann-Lingen, C., Schaefer, H., Kochen, M. M., Himmel, W., Scherer, M., 2009. Depression and anxiety as major determinants of neck pain: a cross-sectional study in general practice. *BMC musculoskeletal disorders* 10, 13. <https://doi.org/10.1186/1471-2474-10-13>

- Carroll, L. J., Hogg-Johnson, S., van der Velde, G., Haldeman, S., Holm, L. W., Carragee, E. J., Hurwitz, E. L., Côté, P., Nordin, M., Peloso, P. M., Guzman, J., Cassidy, J. D., 2008.** Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders Course and prognostic factors for neck pain in the general population: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine*, 33(4Suppl),S75-S82. <https://doi.org/10.1097/BRS.0b013e31816445be>
- Cavlak, U., Yagci N., Aslan U.B., Ekici G., 2009.** A new tool measuring health-related quality of life (HRQOL): The effects of musculoskeletal pain in a group of older Turkish people. *Archives of Gerontology and Geriatrics* 49, 298-303.
- Chiarotto, A., Falla, D., Polli, A., Monticone, M., 2018.** Validity and Responsiveness of the Pain Self-Efficacy Questionnaire in Patients With Neck Pain Disorders. *The Journal of orthopaedic and sports physical therapy*, 48(3), 204-216. <https://doi.org/10.2519/jospt.2018.7605>
- Cramer, H., Mehling, W.E., Saha, F. J., Dobos, G., Lauche, R., 2018.** Postural awareness and its relation to pain: validation of an innovative instrument measuring awareness of body posture in patients with chronic pain. *BMC musculoskeletal disorders* 19(1),109
- Demyttenaere, K., Bruffaerts, R., Lee, S., Posada-Villa, J., Kovess, V., Angermeyer, M.C., 2007.** Mental disorders among persons with chronic back or neck pain: results from the World Mental Health Surveys. *Pain* 129(3), 332-42.
- Dere, T., 2021.** Servikal disk hernili bireylerde kassal enduransın ağrı, boyun farkındalığı ve kinezyofobi ile ilişkisinin incelenmesi. Yüksek lisans tezi, Hacettepe Üniversitesi Sağlık Bilimleri Enstitüsü, Ankara.
- Dimitriadis, Z., Kapreli, E., Strimpakos, N., Oldham, J., 2015.** Do psychological states associate with pain and disability in chronic neck pain patients?. *Journal Back Musculoskelet Rehabilitaion* 28(4),797-802.
- Edmondston, S.J., Björnsdóttir, G., Pálsson, T., Solgard, H., Ussing, K., Allison, G., 2011.** Endurance and fatigue characteristics of the neck flexor and extensor muscles during isometric tests in patients with postural neck pain. *Manuel Therapy*, 16(4), 332-338. [10.1016/j.math.2010.12.005](https://doi.org/10.1016/j.math.2010.12.005).
- Falla, D.L., Jull, G.A., Hodges, P.W., 2004.** Patients with neck pain demonstrate reduced electromyographic activity of the deep cervical flexor muscles during performance of the craniocervical flexion test. *Spine* 29(19):2108-14.
- Fejer, R., Kyvik, K. O., Hartvigsen, J., 2006.** The prevalence of neck pain in the world population: a systematic critical review of the literature. *European spine journal: official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society* 15(6), 834-848. <https://doi.org/10.1007/s00586-004-0864-4>
- Fricton, J., Anderson, K., Clavel, A., Fricton, R., Hathaway, K., Kang, W., Jaeger, B., Maixner, W., Pesut, D., Russell, J., Weisberg, M. B., Whitebird, R., 2015.** Preventing Chronic Pain: A Human Systems Approach-Results From a Massive Open Online Course. *Global advances in health and medicine*, 4(5), 23-32. <https://doi.org/10.7453/gahmj.2015.048>
- Gallego-Sendarrubias, G.M., Rodríguez-Sanz, D., Calvo-Lobo, C., Martín, J.L., 2020.** Efficacy of dry needling as an adjunct to manual therapy for patients with chronic mechanical neck pain: a randomised clinical trial. *Acupuncture in Medicine* 38(4),244-254. doi:10.1136/acupmed-2018-011682
- Harris,K.D., Heer, M.D., Roy,T.C., Santos, D.M., Whitman,J.M., Wainne, S.R., 2005.** Reliability of a Measurement of Neck Flexor Muscle Enduranc. *Physical Therapy* 85 (12), 1349-1355.
- Hidalgo, B., Hall, T., Bossert, J., Dugeny, A., Cagnic, B., Pitance, L., 2017.** The efficacy of manual therapy and exercise for treating non-specific neck pain: A systematic review. *Journal of back and musculoskeletal rehabilitation* 6,30(6),1149-1169. doi: 10.3233/BMR-169615. PMID: 28826164; PMCID: PMC5814665.
- Hogg-Johnson, S., Van Der Velde, G., Carroll, L.J., Holm, L.W., Cassidy, J.D., Guzman, J., 2008.** The burden and determinants of neck pain in the general population *Spine* 17(1), 39-51.
- Kahlacae, A. H., Rezasoltani, A., Ghamkhar, L., 2017.** Is the clinical cervical extensor endurance test capable of differentiating the local and global muscles?. *The spine journal : official journal of the North American Spine Society* 17(7), 913-921. <https://doi.org/10.1016/j.spinee.2017.01.014>
- Kanchanomai, S., Janwantanakul P., Pensri, P., Jiamjarasrangsi, W., 2011.** Risk factors for the onset and persistence of neck pain in undergraduate students: 1-year prospective cohort study. *BMC Public Health* 11, 566-574.
- Kim, H.J., Kim, J.S. 2015.** The relationship between smartphone use and subjective musculoskeletal symptoms and university students. *Journe Physical Therapy Science*27, 575-579.



- Kurtaran, M., Baktır, S., Şeker Abanoz, E., Yeldan, İ., 2019.** Evaluation of Pain, Alexithymia, Depressive Symptom Prevalence and Quality of Life in University Students Sakarya Tıp Dergisi 9(3),433-41.
- Lau, H. M., Wing Chiu, T. T., Lam, T. H., 2011.** The effectiveness of thoracic manipulation on patients with chronic mechanical neck pain - a randomized controlled trial. Manual therapy 16(2), 141-147. <https://doi.org/10.1016/j.math.2010.08.003>
- Lee, H., Hübscher, M., Moseley, G. L., Kamper, S. J., Traeger, A. C., Mansell, G., McAuley, J. H., 2015.** How does pain lead to disability? A systematic review and meta-analysis of mediation studies in people with back and neck pain. Pain 156(6), 988-997. <https://doi.org/10.1097/j.pain.0000000000000146>
- Linton, S.J., 2000.** A review of psychological risk factors in back and neck pain. Spine 25(9), 1148-56.
- Liu, F., Fang, T., Zhou, F., Zhao, M., Chen, M., You, J., 2018.** Association of Depression/Anxiety Symptoms with Neck Pain: A Systematic Review and Meta-Analysis of Literature in China. Pain research & management. 3259431.
- Monticone, M., Giordano, A., Franchignoni, F., 2021.** Scale Shortening and Decrease in Measurement Precision: Analysis of the Pain Self-Efficacy Questionnaire and Its Short Forms in an Italian-Speaking Population With Neck Pain Disorders. Physical Therapy 101(6), pزاب039. <https://doi.org/10.1093/ptj/pزاب039>
- Moreira, C., Bass, A.R., Brandão, M.P., Silva, A.G., 2017.** Do patients with chronic neck pain have distorted body image and tactile dysfunction? European Journal of Physiotherapy 19(4):215-21.
- Nagrle, A. V., Glynn, P., Joshi, A., Ramteke, G., 2010.** The efficacy of an integrated neuromuscular inhibition technique on upper trapezius trigger points in subjects with non-specific neck pain: a randomized controlled trial. The Journal of manual & manipulative therapy 18(1), 37-43. <https://doi.org/10.1179/106698110X12595770849605>.
- Nicholas, M.K., 2007.** The pain self-efficacy questionnaire: taking pain into account. European journal of pain 11(2), 153-163. <https://doi.org/10.1016/j.ejpain.2005.12.008>
- Onan, D., Gokmen, D., Ulger, O. 2019.** The Fremantle Neck Awareness Questionnaire in Chronic Neck Pain Patients: Turkish Version, Validity and Reliability Study. Spine 45(3), E163-E169. doi: 10.1097/BRS.00000000000003207
- Özel, M., 2022.** Kronik boyun ağrısında boyun farkındalığı ile eklem pozisyon hissi, baş postürü ve kas dayanıklılığı arasındaki ilişki. Yüksek lisans tezi, Hacettepe Üniversitesi Sağlık Bilimleri Enstitüsü, Ankara.
- Parazza, S., Vanti, C., O'Reilly, C., Villafañe, J. H., Tricás Moreno, J. M., Estébanez De Miguel, E., 2014.** The relationship between cervical flexor endurance, cervical extensor endurance, VAS, and disability in subjects with neck pain. Chiropractic & manual therapies 22(1), 10. <https://doi.org/10.1186/2045-709X-22-10>.
- Piper, A., 2009.** Comparison of endurance capacity of deep cervical flexors between healthy and subjects with cervical pain. Manuelle Therapie 13, 202-211.
- Savcı, S., Öztürk, M., Arıkan, H., İnal İnce, D., Tokgözoğlu, L., 2006.** Physical activity levels of university students. Archives of the Turkish Society of Cardiology 34(3), 166-172.
- Schomacher, J., Falla, D., 2013.** Function and structure of the deep cervical extensor muscles in patients with neck pain. Manual Therapy 18(5),360-366. <https://doi.org/10.1016/j.math.2013.05.009>
- Shahidi, B., Curran-Everett, D., Maluf, K.S., 2015.** Psychosocial, Physical, and Neurophysiological Risk Factors for Chronic Neck Pain: A Prospective Inception Cohort Study. The Journal of Pain 16(12),1288-1299.
- Sherer, M., Adams, C.H., 1983.** Construct validation of the Self-Efficacy Scale. Psychol Rep, 53:899-902.
- Sihawong, R., Janwantanakul, P., Sitthipornvorakul, E., Pensri, P., 2011.** Exercise therapy for office workers with nonspecific neck pain: a systematic review. Journal Manipulative Physiol Therapy 34, 62-71
- Şimşek, Ş. , Yağcı, N. , Oymak Soysal, A. N. , Kaş Özdemir, A., Bergin, M., 2022.** The relation between pain, functional status, and neck awareness in individuals with chronic neck pain. Türk Fizyoterapi ve Rehabilitasyon Dergisi 33 (2) , 106-113. DOI: 10.21653/tjpr.979737
- Von Bothmer, M.I., Fridlund, B., 2005.** Gender differences in health habits and in motivation for a healthy lifestyle among Swedish university students. Nursing Health Science 7, 107-18.
- Wand, B.M., Catley, J., Rabey, M., O'Sullivan, B.P., O'Connell, E. N., Smith, J., 2016.** Disrupted Self-Perception in People With Chronic Low Back Pain. Further Evaluation of the Fremantle Back

Awareness Questionnaire. The Journal of Pain 17 (9), 1011- 1012.

**Wewers, M.E., Lowe, N.K., 1990.** A critical review of visual analogue scales in the measurement of clinical phenomena. Research in Nursing & Health 13, 227-236.

**Wiech, K., 2016.** Deconstructing the sensation of pain: the influence of cognitive processes on pain perception. Science 354 (6312), 584–587.

**Woodman, J., Ballard, K., Hewitt, C., MacPherson, H., 2018.** Self-efficacy and self-care-

related outcomes following Alexander Technique lessons for people with chronic neck pain in the ATLAS randomised, controlled trial. European Journal of Integrative Medicine 17, 64–71. <https://doi.org/10.1016/j.eujim.2017.11.006>

**Yıldırım, F., İlhan, İ.Ö., 2010.** The Validity and Reliability of the General Self-Efficacy Scale-Turkish Form Türk Psikiyatri Dergisi 21(4), 301-308.

**Zigmond, A. S., Snaith, R. P., 1983.** The hospital anxiety and depression scale. Acta psychiatrica Scandinavica 67(6), 361–370. <https://doi.org/10.1111/j.1600-0447.1983.tb0971>