



Araştırma Makalesi

www.ziraat.selcuk.edu.tr/ojs
Selçuk Üniversitesi
Selçuk Tarım ve Gıda Bilimleri Dergisi
25 (4): (2011) 11-14
ISSN: 1309-0550



Kahramanmaraş'ta Sebze Bitkilerinde Cicadellidae (Hemiptera) Türleri

Tülin ÖZSİSLİ^{1,2}

¹Kahramanmaraş Sütçü İmam Üniversitesi, Ziraat Fakültesi, Bitki Koruma Bölümü, Kahramanmaraş/Türkiye

(Geliş Tarihi: 11.08.2011, Kabul Tarihi: 10.10.2011)

Özet

Kahramanmaraş'ta yetişirilen sebze bitkilerinden Cicadellidae (Hemiptera) türlerini tanımlamak için surveyler yapılmıştır. Sebzeler üzerindeki cicadellid türleri 1997–1998 yıllarında toplanmıştır. Sebzelerden Aconurella prolixa (Lethierry), Arboridia sp., Circulifer haematoceps (M.-R.), Edwardsiana sp., Empoasca decipiens Paoli, Empoasca sp., Euacelidius sp., Macrosteles quadripunctulatus (Kirschbaum), Neoaliturus fenestratus (Herrich-Schäffer), Psammotettix provincialis (Ribaut), Psammotettix sitriatus L., Recilia schimidtgeni (Wagner), Zyginidia pullula (Boheman), Zyginidia sp. türleri elde edilmişdir. Sebze alanlarındaki en yaygın cicadellid türü E. decipiens olmuştur.

Key words: Cicadellid, survey, sebzeler, Kahramanmaraş, Türkiye.

Cicadellidae (Hemiptera) Species of Vegetable Plants from Kahramanmaraş, Turkey

Abstract

Surveys were conducted to identify leafhopper family, Cicadellidae (Hemiptera) species from vegetable plants in Kahramanmaraş, Turkey. Cicadellid species on vegetables were collected during 1997–1998. Aconurella prolixa (Lethierry), Arboridia sp., Circulifer haematoceps (M.-R.), Edwardsiana sp., Empoasca decipiens Paoli, Empoasca sp., Euacelidius sp., Macrosteles quadripunctulatus (Kirschbaum), Neoaliturus fenestratus (Herrich-Schäffer), Psammotettix provincialis (Ribaut), Psammotettix sitriatus L., Recilia schimidtgeni (Wagner), Zyginidia pullula (Boheman), Zyginidia sp. were obtained from vegetables. E. decipiens was the most common cicadellid species in the vegetable fields.

Key words: Cicadellid, survey, vegetables, Kahramanmaraş, Turkey

Introduction

Family Cicadellidae (Hemiptera) is one of the pest groups of vegetables. There are few data on the cicadellid species on vegetables in the other provinces from different regions of Turkey. Karsavuran et al. (1992) studied that investigations on the population trends of *Asymmetrasca decedens* (Paoli) and *Empoasca decipiens* Paoli on processing tomatoes in Yenişehir (Bursa). Additionally, the pests of processing tomatoes were also studied in Turkey during 1988-89 (Öncüer et al., 1992). Başpinar (1994) reported that the population changes of *A. decedens* and *E. decipiens* on eggplant and tomato in Adana province in 1991. Akkaya and Uygun (1996) determined *E. decipiens* as one of the harmful species on summer vegetables areas. Akkaya and Uygun (1999) also reported that the study was carried out to determine the harmful and beneficial insect fauna of cucurbit vegetables, in Diyarbakır and Şanlıurfa provinces in 1993-94. Yiğit and Uygun (1982) determined cicadellid species only in apple orchards in Kahramanmaraş. The object of the present faunistic paper is to determine the cicadellids on different vegetable plants in Kahramanmaraş as a basis for future studies.

Material and Methods

Vegetable growing areas of Kahramanmaraş (Centre, Pazarcık, and Türkoğlu counties) were surveyed randomly from July to August in 1997–1998.

Surveys were performed in eggplant (*Solanum melongena* L.), melon (*Cucumis melo* L.), watermelon (*Citrullus lanatus* (Thunb.)), tomato (*Lycopersicon esculentum* L.), cucumber (*Cucumis sativus* L.; *Cucumis melo* var. *flexuosus*), okra (*Hibiscus esculentus* L.), pepper (*Capsicum annuum* L.), zucchini (*Cucurbita pepo* L.), and legume (*Phaseolus vulgaris* L.) fields. However, most of samples were taken from *C. annuum*, *C. sativus*, and *S. melongena* fields. Samplings were conducted in 14 and 24 fields in 1997 and 1998, respectively. In total, 25 leaves and 25 sweeping net (size: 30x75cm) samples were taken randomly from each vegetable fields. The samples were placed in separate plastic bags and were brought to the laboratory and cicadellid adults were prepared for the identification. Method of mounting minute insects was used for mounting of cicadellid adults (Borror et al., 1992).

²Sorumlu Yazar: tulin@ksu.edu.tr

Results and Discussion

Family Cicadellidae (Hemiptera)

Aconurella prolixa (Lethierry)

Material examined: Center 568 m: Hacımustafa, 18.VI.1998, (1 individual), on *Solanum melongena* L. leaves.

Arboridia sp.

Material examined: Center: Hacımustafa, 18.VI.1998, (1), on *Phaseolus vulgaris* L. leaves.

Circulifer haematoceps (M.-R.)

Material examined: Center: Hacımustafa, 18.VI.1998, (1), on *Solanum melongena* L. leaves; Center: Hacımustafa, 18.VI.1998, (1), on *Cucurbita pepo* L. collected with a sweeping net; Türkoğlu 490 m: Akçalı, 15.VII.1998, (1), *Cucumis sativus* L. collected with a sweeping net; Pazarcık 850 m: Alibeyuşağı, 21.VII.1998, (1), *Capsicum annuum* L. collected with a sweeping net.

Edwardsiana sp.

Material examined: Türkoğlu: Ceceli, 30.VII.1997, (1), *Lycopersicon esculentum* L. collected with a sweeping net; Pazarcık: Emiroğlu, 18.VI.1998, (5), on *Lycopersicon esculentum* L. collected with a sweeping net.

Empoasca decipiens Paoli

Material examined: Pazarcık: Karaçay, 27.VII.1997, (1), on *Solanum melongena* L. leaves; Center: Hacımustafa, 29.VII.1997, (1), *Solanum melongena* L. collected with a sweeping net; Pazarcık: Bölkçam, 15.VIII.1997, (1), on *Phaseolus vulgaris* L. leaves; Center: Hacımustafa, 18.VI.1998, (1), on *Solanum melongena* L. leaves; Center: Hacımustafa, 09.VII.1998, (11), *Cucurbita pepo* L. collected with a sweeping net; Türkoğlu: Ceceli, 15.VII.1998, (9), *Lycopersicon esculentum* L. collected with a sweeping net; Türkoğlu: Ceceli, 15.VII.1998, (2), *Cucumis sativus* L. collected with a sweeping net; Türkoğlu: Akçalı, 15.VII.1998, (2), on *Cucumis sativus* L. by sweeping net; Pazarcık: Alibeyuşağı, 21.VII.1998, (9), *Capsicum annuum* L. collected with a sweeping net; Pazarcık: Bölkçam, 24.VII.1998, (3), *Capsicum annuum* L. collected with a sweeping net; Pazarcık: Bölkçam, 24.VII.1998, (3), *Lycopersicon esculentum* L. collected with a sweeping net; Center: Şahinkayası, 29.VII.1998, (11), *Lycopersicon esculentum* L. collected with a sweeping net; Center: Şahinkayası, 29.VII.1998, (18), *Capsicum annuum* L. collected with a sweeping net; Türkoğlu: Beyoğlu: 03.VIII.1998, (1), *Capsicum annuum* L. collected with a sweeping net; Pazarcık: Çınarlı, 03.VIII.1998, (1), *Capsicum annuum* L. collected with a sweeping net; Center: Çığlı, 03.VIII.1998, (6), *Capsicum annuum* L. collected with a sweeping net; Center: Tevekkeli,

11.VIII.1998, (8), *Hibiscus esculentus* L. collected with a sweeping net; Center: Küpeli, 11.VIII.1998, (8), *Solanum melongena* L. collected with a sweeping net.

Empoasca sp.,

Material examined: Türkoğlu: 30.VII.1997, (1), *Solanum melongena* L. collected with a sweeping net; Center: Hacımustafa, 18.VI.1998, (3), on *Phaseolus vulgaris* L. leaves; Center: Hacımustafa, 18.VI.1998, (1), on *Solanum melongena* L. leaves.

Euacelidius sp.,

Material examined: Center: Hacımustafa, 18.VI.1998, (1), on *Solanum melongena* L. leaves.

Macrosteles quadripunctulatus (Kirschbaum)

Material examined: Türkoğlu: Akçalı, 15.VII.1998, (1), *Cucumis sativus* L. collected with a sweeping net; Pazarcık: Karaçay, 27.VII.1998, (1), on *Solanum melongena* L. leaves.

Neoaliturus fenestratus (Herrich-Schaffer)

Material examined: Pazarcık: Karaçay, 30.VII.1998, (1), *Lycopersicon esculentum* L. collected with a sweeping net.

Psammotettix provincialis (Ribaut)

Material examined: Center: Hacımustafa, 18.VI.1998, (2), on *Cucurbita pepo* L. collected with a sweeping net; Türkoğlu: Akçalı, 15.VII.1998, (4), *Cucumis sativus* L. collected with a sweeping net.

Psammotettix sitriatus L.

Material examined: Center: Hacımustafa, 18.VI.1998, (1), on *Phaseolus vulgaris* L. leaves; Center: Hacımustafa, 18.VI.1998, (3), on *Solanum melongena* L. leaves; Center: Hacımustafa, 18.VI.1998, (6), on *Phaseolus vulgaris* L. leaves; Türkoğlu: 15.VII.1998, (6), *Cucumis sativus* L. collected with a sweeping net. Center: Ceceli, 15.VII.1998, (8), *Cucumis sativus* L. collected with a sweeping net. Pazarcık: Alibeyuşağı, 21.VII.1998, (1), *Capsicum annuum* L. collected with a sweeping net; Pazarcık: Karaçay, 27.VII.1998, (1), on *Solanum melongena* L. leaves; Center: Şahinkayası, 29.VII.1998, (10), *Capsicum annuum* L. collected with a sweeping net; Center: Çığlı, 30.VIII.1998, (2), *Capsicum annuum* L. collected with a sweeping net.

Recilia schimidtgeni (Wagner)

Material examined: Center: Ceceli, 15.VII.1998, (1), *Cucumis sativus* L. collected with a sweeping net.

Zyginidia pullula (Boheman)

Material examined: Pazarcık: Alibeyuşağı, 21.VII.1998, (9), *Capsicum annuum* L. collected with a sweeping net.

Zyginidia sp.

Material examined: Türkoğlu: 30.VII.1997, (1), *Cucumis sativus* L. collected with a sweeping net; Türkoğlu: 30.VII.1997, (1), *Lycopersicon esculentum* L. collected with a sweeping net; Center: Hacimustafa, 18.VI.1998, (3), on *Phaseolus vulgaris* L. leaves; Center: Hacimustafa, 18.VI.1998, (1), on *Solanum melongena* L. leaves; Center: Şahinkayası, 29.VII.1998, (1), *Lycopersicon esculentum* L. collected with a sweeping net.

In conclusion, 9 different cicadellid species and 5 genus were obtained from vegetable fields of Centre, Pazarcık and Türkoğlu counties of Kahramanmaraş Province. Cicadellid species were not found on *C. melo*, *C. lanatus* and *C. melo* var. *flexuosus* in this study. *E. decipiens* was obtained from *S. melongena*, *P. vulgaris*, *C. pepo*, *L. esculentum*, *C. sativus*, *H. esculentus* and *C. annuum* in 1998. *E. decipiens* was the most common cicadellid species of sweeping net samples of vegetables in Kahramanmaraş. *P. sitriatus* followed *E. decipiens* (Table 1).

Table 1. The total number of cicadellid species obtained from vegetable plants in Kahramanmaraş (1998).

Cicadellid species	Vegetables							
	Sm	Pv	Cp	Le	Cs	He	Ca	Total
<i>Aconurella prolixa</i>	A*	1						1
	F**	1						1
<i>Arboridia</i> sp	A		1					1
	F		1					1
<i>Circulifer haematoceps</i>	A	1		1		1		4
	F	1		1		1		4
<i>Edwardsiana</i> sp.	A			5				5
	F			1				1
<i>Empoasca decipiens</i>	A	9		11	23	4	8	26
	F	2		1	3	2	1	15
<i>Empoasca</i> sp.	A	1	3					4
	F	1	1					2
<i>Euacelidius</i> sp.	A	1						1
	F	1						1
<i>Macrosteles quadripunctulatus</i>	A	1				1		2
	F	1				1		2
<i>Neoaliturus fenestratus</i>	A			1				1
	F			1				1
<i>Psammotettix provincialis</i>	A		2		4			6
	F		1		1			2
<i>Psammotettix sitriatus</i>	A	3	7	6	8		13	37
	F	2	1	1	1		3	8
<i>Recilia schimidtgeni</i>	A				1			1
	F				1			1
<i>Zyginidia pullula</i>	A						1	1
	F						1	1
<i>Zyginidia</i> sp.	A	1	3		1			5
	F	1	1		1			3
Total no sampels		4	3	-	1	7	-	24

Sm, *Solanum melongena*; Pv, *Phaseolus vulgaris*; Cp, *Cucurbita pepo*; Le, *Lycopersicon esculentum*; Cs, *Cucumis sativus*; He, *Hibiscus esculentus*; Ca, *Capsicum annuum*.

*A: Abundance, number of individuals; **F: Frequency, number of sampels.

E. decipiens was reported to be harmful on many culture plants especially in the South and The Western Anatolia regions of Turkey by Lodos, (1986). *E. decipiens* also was determined as one of the most common species on processing tomatoes, squash and cucumber by the other researchers (Delrio et al., 1989; Karsavuran et al., 1992; Öncür et al., 1992; Akkaya

and Uygun, 1999). Previously, Yiğit and Uygun (1982) reported *P. sitriatus*, *Edwardsiana rosae* (L.), *E. decipiens*, *Neoaliturus haematoceps* (M.R.), *Psammotettix* sp. Prob. *provincialis* (Ribaut) from apple orchards in Adana and Kahramanmaraş. The cicadellid fauna can also be considered in terms of integrated pest management for the other field areas

and orchards of Kahramanmaraş in addition these studies.

Acknowledgements

I would like to express my sincere to Prof. Dr. Hüseyin Başpinar (Adnan Menderes University, Agricultural Faculty, Plant Protection Department, Aydın) for his help in identification of cicadellids. I would like to thank Kahramanmaraş Sütçü İmam University for their financial support of this study.

References

- Akkaya A. and Uygun N., 1996. Studies on the insect fauna associated to summer vegetables areas in Diyarbakır and Şanlıurfa provinces. *Proceedings of the Third Turkish National Congress of Entomology*, 423-430, 24-28 Eylül 1996, Ankara.
- Akkaya A. and Uygun N., 1999. Faunistic studies on harmful and beneficial insects on cucurbit vegetables in the Southeastern Anatolian region of Turkey. *Acta Horticulturae* (492): 335-340.
- Başpinar H., 1994. Some observations on dominant structure and population changes of *Asymmetrasca decedens* and *Empoasca decipiens* (Hom., Cicadellidae) on different crops in Adana. *Turk. J. Entomol.*, 18 (2): 71-76.
- Borror, D.J., Triplehorn, C.A. and Johnson, N.F., 1992. An introduction to the study of insects, 6th edition. Harcourt Brace College Publishers, 875 p. Orlando, FL.
- Delrio, G., Luciano, P., Floris, I., Cabitza, F., Cubeddu, M. and Cabras, P., 1989. Control trials against the pests of processing tomatoes in Sardinia. *Difesa delle Piante*, 12 (1-2): 97-106.
- Karsavuran, Y., Öncüler, C., Civelek, H.S. and Gümüş, M., 1992. *Proceedings of the Second Turkish National Congress of Entomology*, İzmir: Ege University, 541-548.
- Lodos N., 1986. *Türkiye Entomolojisi II.*, Genel, uygulamalı ve faunistik. E. Ü. Ziraat Fakültesi Yayınları, No: 429, Bornova, İzmir, 580 pp.
- Öncüler, C., Karsavuran, Y., Yoldaş, Z. and Durmuşoğlu, E., 1992. *Proceedings of the Second Turkish National Congress of Entomology* Izmir: Ege University, 705-713.
- Yiğit, A. and Uygun, N., 1982. Studies on the determination of beneficial and injurious fauna of apple orchards in Adana, İçel and Kahramanmaraş Provinces. *Plant Protec. Bull.*, 22 (4): 163-178.