



In Vitro Vitamin C Equivalent Antioxidant Capacity, Cytotoxicity and Anti-Cancer Activity of Methanolic *Urtica dioica* L. Leaf Extract as a Food Supplement

Serol Korkmaz^{1,3,*}, Hanife Banu Aydın², Burcu Irem Omurtag Korkmaz²

¹ Pendik Veterinary Control Institute, Istanbul, Türkiye

² Marmara University, Faculty of Health Sciences, Department of Nutrition and Dietetics, Istanbul, Türkiye

³ Marmara University, Faculty of Health Sciences, Department of Basic Health Sciences, Istanbul, Türkiye

HIGHLIGHTS

- This in vitro study was conducted on some cell cultures originating from vital mammalian tissues.
- The safest concentrations of *Urtica dioica* L. leaf extract were determined as 15.71 mg/ml and 5.14 mg/ml of CC₅₀.
- The extract showed anti-cancer activity on human hepatocellular carcinoma cells with a 2.46 mg/ml of IC₅₀.

Abstract

Antioxidant capacity, cytotoxicity on two vital cell lines and the anti-cancer activity of methanolic *Urtica dioica* L. leaf extract (UDE) collected from Duzkoy, Giresun, Turkey were studied by determining safe concentration. The antioxidant capacity of the extract was expressed as vitamin C equivalency by spectrophotometric MTT assay. The cytotoxic concentration 50% was measured by the linearity between UDE concentrations (CC₅₀) and the cell viability of non-cancer kidney cell lines (BHK-21, MDBK). The anti-cancer activity was conducted on human hepatocellular carcinoma cells (HepG2) by determining inhibition concentration (IC₅₀) on cell proliferation. The vitamin C equivalence of UDE increased linearly by increasing the concentration. The cytotoxic and non-toxic concentrations of UDE were determined on BHK-21 and MDBK with 15.71 mg/ml and 5.14 mg/ml of CC₅₀ respectively. The extract inhibited the proliferation of human hepatocellular carcinoma cells with a 2.46 mg/ml of IC₅₀. In conclusion, the present study tried to explain in detail the dose-dependent activity of *Urtica dioica* L. leaf extract. The dose-response results showed that *Urtica dioica* L. leaf extract could have low cytotoxicity, but potential anti-cancer activity at safe concentrations.

Keywords: anti-cancer; cytotoxicity; nettle; *Urtica dioica* L.

1. Introduction

Urtica dioica L. belongs to the Urticaceae family. It has been identified and widely distributed worldwide and is considered native to Europe, North Africa, Asia, and North America (Upton 2013; Dhouibi et al. 2020; European Commission 2023). It and its extracts are used in both pharmaceutical and food industries as a

Citation: Korkmaz S, Aydın HB, Omurtag Korkmaz BI (2024). In Vitro Vitamin C Equivalent Antioxidant Capacity, Cytotoxicity and Anti-Cancer Activity of Methanolic *Urtica dioica* L. Leaf Extract as a Food Supplement. *Selcuk Journal of Agriculture and Food Sciences*, 38 (1), 140-146. <https://doi.org/10.15316/SJAfS.2024.013>

Corresponding Author E-mail: serolkorkmaz@yahoo.com

Received date: 29/09/2023

Accepted date: 25/04/2024

Author(s) publishing with the journal retain(s) the copyright to their work licensed under the CC BY-NC 4.0.

<https://creativecommons.org/licenses/by-nc/4.0/>

supplement and a food additive for extending shelf-life, ensuring microbial safety of foods and higher consumer acceptability (Alp and Aksu 2010). Phytochemical studies have mainly focused on its bioactive compounds and activities related to antioxidant contents (Körpe et al. 2013; Dhouibi et al. 2020; Veiga et al. 2020). Therefore, a wide range of plants and their extracts were applied as food additives and preservatives during food production and service to improve safety and quality by presenting antioxidant activities at their safe concentrations (Alp and Aksu, 2010; Körpe et al. 2013; Dhouibi et al. 2020).

In addition to antioxidant capacity, several *in vitro* and *in vivo* studies of plants aimed to determine the non-toxic doses for preventing adverse effects before using them including *Urtica dioica* L. as a potential therapeutic agent in modern and traditional medicine (Özkol et al. 2012; Sayhan et al. 2012; Dhouibi et al. 2020; Veiga et al. 2020). Recently, the aerial and subsoil parts of *Urtica dioica* L. were experimentally used to treat many cancer types by inhibiting cell proliferation (Konrad et al. 2000; Gözümlü et al. 2003). Also, the cancer treatment has a risk of toxicity in different tissues including the kidney and liver by inducing oxidative stress and free radical generation (Özkol et al. 2012; Dhouibi et al. 2020). Therefore, the present study aimed to assess dose-dependently the antioxidant capacity, cytotoxicity on non-cancer kidney cell culture lines and anti-cancer activity on liver cancer cells of methanolic *U. dioica* L. leaf extract (UDE).

2. Materials and Methods

2.1. Extraction of *U. dioica* L.

The sampling and extraction methods used in this study were adapted from Aydın et al. (2023). Briefly, *U. dioica* L. leaf samples were collected from Duzkoy province (40°56'59.1"N. 38°36'06.5"E), the city of Giresun on the Black Sea coast of Turkey in May 2021. They were identified and authenticated as "*Urtica dioica* L." at the Department of Pharmaceutical Botany, Faculty of Pharmacy in Marmara University, and a herbarium record was created with the code "MARE 23334". The samples were dried in cool room conditions avoiding direct sunlight.

Solid/liquid extraction and evaporation methods were used for the extraction of dried samples. Dried samples were ground by a water-cooled miller. Fifty grams of ground leaves were macerated in 500 ml of methanol (reagent grade, $\geq 99.7\%$) on a magnetic stirrer for 24 hours. The alcoholic suspension was filtered twice through filter paper (FilterLab-50 g/m²). The solvent in the remaining filtrate was evaporated in a rotary evaporator (Heidolph, Germany) at 40-45°C, 150 mbar, and 135 rpm. The extract was completely concentrated in a vacuum oven (Nüve EV 018, Turkey) at 45°C and -1 bar pressure. A UDE stock solution of 40 mg/ml was prepared in ultra-distilled water and stored at +4 °C for further analysis.

2.2. Handling of Cell Lines

Baby hamster kidney fibroblast (BHK-21, CCL-10), bovine kidney epithelial (MDBK, CCL-22) and human hepatocellular carcinoma (HepG2, HB-8065) cell lines were from ATCC, USA. All cell lines were cultured with Eagle's Minimum Essential Medium (EMEM) in the incubator with the standard condition (SC) of 37 °C and 5% CO₂. The medium contained foetal bovine serum (10%), L-alanyl-L-glutamine (200 mM) and 1% penicillin (10,000 unit/ml)-streptomycin (10 mg/ml)-amphotericin B (0.025 mg/ml). For the anti-cancer and cytotoxicity assay, the stock viable cell suspension (3×10^5 cell/ml) (100 μ l) was transferred in each well (3×10^4 cell/well) and kept in SC for 24 h to achieve the confluence of at least 90% in 96-well microplates.

2.3. Vitamin C equivalent antioxidant capacity

Vitamin C (l-ascorbic acid, HPLC grade) was two-fold diluted as 600, 300, 150, 75 and 37.5 μ mol with ultra-distilled water. UDE was two-fold diluted as 0.62, 1.25, 2.5, 5, 10 and 20 mg/ml with ultra-distilled water. MTT (2,2-diphenyl-1H-tetrazolium bromide, 1 mg/ml) (380 μ l) and each Vitamin C or extract dilution (20 μ l) were mixed in an Eppendorf tube and incubated at 37 °C for 4 h. After incubation, DMSO (400 μ l) was added to all tubes and mixed well to solve the blue formazan salt formed during the incubation. A 100 μ l of each mixture was added quadruplicated to the U-bottom 96-well microplate. The microplate was read at 570 nm.

2.4. Anti-cancer and cytotoxicity assays by MTT

The extract was two-fold serially diluted with the maintaining medium at the concentrations of 0.62, 1.25, 2.5, 5, 10 and 20 mg/ml. Then, each dilution (100 µl) was added to the microplates with monolayer cell culture at six-replicated wells. The cell control wells contained only fresh medium and cells. The blank wells contained the medium without cells. The microplates were incubated in SC for 24 h. 10 µl MTT in PBS (5 mg/ml) was added to each well. After a 4 h incubation in SC. The supernatant was discarded and DMSO (100 µl) was added to wells. The microplate was gently shaken to solubilize the formazan crystals and read at a wavelength of 570 nm (Absorbance 96. Byonoy. Germany).

2.5. Data analysis

The percentage of cell inhibition was calculated using the equation (1) as follows,

$$\text{Cell viability (\%)} = (\text{OD sample} - \text{OD blank}) / (\text{OD control} - \text{OD blank}) \times 100'' (1)$$

The 50% cytotoxic concentration (CC₅₀) was calculated from concentration-based-curves after non-linear regression analysis ($y = me^{bx}$). The vitamin C standard curve was generated with optic density (OD, nm) values of five vitamin C dilutions by linear regression analysis (vitamin C equivalence = $m \times \text{OD}_{570} + b$, R²). The equivalence to vitamin C of extract dilutions was calculated concerning the standard curve (Figure 1). Statistical analyses were conducted by using SPSS version 15 software (IBM. USA). The graphs were generated by using Office Excel 2016 (Microsoft. USA).

3. Results

3.1. Vitamin C equivalent antioxidant capacity of UDE

The natural antioxidants in herbal additives such as carotenoids, tocopherol (vitamin E), some phenolic compounds, and ascorbic acid inhibit oxidative damage by free-radical scavenging (Dini 2019; El-Saber Batiha et al. 2021). Previous research suggested that fresh *U. dioica* L. leaves contained various amounts of total ascorbic acid content from 16 to 112.8 mg/100 g fresh weight at different harvest times (Skalozubova and Reshetova 2013; Shonte et al. 2020). The highest vitamin C was measured in August while the lowest was in September (Paulauskienė et al. 2021). As the vegetation period and drying methods influenced the vitamin C content and antioxidant capacity of *U. dioica* L. leaves, it was not varied by meteorological conditions. The highest vitamin C content was determined with low-temperature drying compared to oven drying (Shonte et al. 2020; Garcia et al. 2021; Paulauskienė et al. 2021). In this study, *U. dioica* L. leaves were harvested in May, dried under cool-dry room conditions and extracted by methanol. The vitamin C equivalence of UDE was calculated concerning the standard curve by linear regression analysis (vitamin C equivalence. µmol = $1191.6 \times \text{OD of UDE} - 1.4723$, R² = 0.998). The vitamin C equivalence of UDE increased by increasing the concentrations of UDE with a linearity of R² = 0.690 (vitamin C equivalence, µmol = $29.578 \times \text{UDE mg/ml} + 501.51$) (Figure 1). In harmony with previous studies, the average vitamin C content was calculated as approximately 42.3 µg/mg in methanolic extract of *U. dioica* L. leaves collected from Giresun region in Turkiye.




Figure 1. The vitamin C equivalency antioxidant capacity of UDE based on concentration.

3.2. Cytotoxicity of UDE

The cytotoxicity and safety levels of plant extracts vary related to the extraction method and are generally tested with cell lines of mammalian kidney and liver. A previous study suggested that methanolic extract (CC₅₀: 0.702-0.803 mg/ml) was safer than aqueous extract (CC₅₀: 0.37-0.49 mg/ml) of *U. dioica* L. on BHK-21 (Flores-Ocelotl et al. 2018). Also, the ethanolic extracts of *U. dioica* L. showed higher cytotoxicity than the aqueous extracts (Mannila et al. 2022). In this study, the cytotoxic effects of methanolic *U. dioica* L. leaf extract experimented on the viabilities of non-cancer (BHK-21 and MDBK) cell lines with concentration-response curves (Table 1). The relation between the cell viability and the concentration was the cell viability % = $99.474e^{-0.044 \times \text{UDE (mg/ml)}}$ (R² = 0.936) for BHK-21 and the cell proliferation % = $163.76e^{-0.23 \times \text{UDE (mg/ml)}}$ (R² = 0.913) for MDBK (Figure 2). The CC₅₀ was calculated as 15.71 mg/ml and 5.14 mg/ml for BHK-21 and MDBK respectively (Table 1). The Higher CC₅₀ than the previous study indicated that the methanolic extract of *U. dioica* L. leaves might be a safer and potential food supplement and additive.

Table 1. The dose-response effects of UDE on % cell viability.

UDE Conc. (mg/ml)	Non-Cancer Cells		Cancer Cell
	BHK-21	MDBK	HepG2
	Mean±SD (%)	Mean±SD (%)	Mean±SD (%)
20	44.66±5.00c	1.13±4.44e	2.37±3.38d
10	56.36±6.50bc	43.21±4.30d	22.65±2.30c
5	73.59±9.15b	61.73±8.35c	34.49±5.25bc
2.5	93.45±4.25ab	79.42±4.01b	44.05±2.30b
1.25	99.82±8.35ab	101.54±2.02a	60.05±6.61a
0.62	99.95±5.51a	102.82±1.51a	60.13±5.45a
P value	<0.01	<0.01	<0.01
CC ₅₀ (mg/ml)	15.71	5.14	2.46

3.3. Anti-cancer activity of UDE

Various studies have recently demonstrated the cytotoxic and anti-cancer properties of *U. dioica* L. in particular against colon, gastric, lung, prostate and breast cancers (Esposito et al. 2019). Aqueous extract of *U. dioica* L. leaf cultured in Iran inhibited the proliferation of human breast cancer (MCF-7) at 2 mg/ml after 72h treatment (Fattahi et al. 2013, 2018). Its aqueous extract (from 5 to 30 µg/mL) harvested in Turkey has shown a dose-dependent inhibition effect on three breast cancer cell lines (MCF-7, MDA-MB-468, and MDA-MB-231) with IC₅₀s of 14-18 µg/ml (Karakol et al. 2022). For Its dichloromethane extract against both mouse and human breast cancer cells, IC₅₀ was determined between 31.37 and 38.14 mg/ml. Similarly, its anti-cancer activity was measured by inhibiting the metastasis of breast cancer with 20 mg/kg daily injection treatment in vivo mouse models (Mansoori et al. 2017; Mohammadi et al. 2017). For its anti-cancer activities on human prostate cancer, the dichloromethane and aqueous extracts inhibited the proliferation of PC3 and LNCaP cells with 15.54 µg/ml and 42-50 µg/ml of IC₅₀, respectively (Durak et al. 2004; Levy et al. 2013; Mohammadi et al. 2016). Meanwhile, no cytotoxicity was determined in non-cancer human prostate stromal cells (Konrad et al. 2000). *U. dioica* L. hydroalcoholic (50:50 v/v) extract inhibited 25.4% of human hepatocellular carcinoma (HepG2) cell viability at the concentration of 0.35 mg/ml (Carvalho et al. 2017). In this study. The anti-cancer effect of UDE on the proliferation of HepG2 cells was determined by spectrophotometric MTT assay with the concentration-response curve (cell proliferation % = $74.941e^{-0.163 \times \text{UDE (mg/ml)}}$, R² = 0.965) (Figure 2). The IC₅₀ was calculated as 2.46 mg/ml for human hepatocellular carcinoma (Table 1). These IC₅₀ values present the potential anti-cancer activity through hepatoprotective and reducing tumorigenesis activities of methanolic extract of *Urtica dioica* L. leaf used at safe concentrations.




Figure 2. The cell viability and inhibition with CC₅₀ (mg/ml) of UDE

4. Conclusions

In conclusion, the present study gives information about the bioactivity of *U. dioica* L. leaf harvested during the growing season in Giresun City, Turkey. The extract had dose-dependently antioxidant capacity expressed as vitamin C equivalency. Non-toxic and safe concentrations of the extract were determined in detail on vital cell lines originating from kidney tissue, which are generally used in food and drug research for consumer safety. The dose-response results indicated that the extract was less toxic to kidney cells while inhibiting the proliferation of liver cancer cells at a lower concentration.

Author Contributions: The authors have an equal contribution. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding

Acknowledgements: The authors thank Prof. Dr. Zafer Odabas, Prof. Dr. Hasan Kilic, and Prof. Dr. Aysen Gargili Keles (Head of the Basic Health Sciences Department and Research Laboratory, Marmara University). The extraction method used in this study is from a part of the MSc thesis of Hanife Banu Aydin supervised by B. Irem Omurtag Korkmaz.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Alp E, Aksu MI (2010). Effects of water extract of *Urtica dioica* L. and modified atmosphere packaging on the shelf life of ground beef. *Meat Science*, 86(2): 468–473. Doi:10.1016/j.meatsci.2010.05.036
- Aydin HB, Korkmaz S, Omurtag Korkmaz BI (2023). Antibacterial and antiviral activities of stinging nettle (*Urtica dioica* L.) leaf extract on norovirus and campylobacter jejuni as foodborne pathogens. *Journal of Microbiology, Biotechnology and Food Sciences*, 13(1):9768. Doi:10.55251/jmbfs.9768
- Carvalho AR, Costa G, Figueirinha A, Liberal J, Prior JAV, Lopes MC, Cruz MT, Batista MT (2017). *Urtica* spp.: Phenolic composition, safety, antioxidant and anti-inflammatory activities. *Food Research International*, 99: 485–494. Doi:10.1016/j.foodres.2017.06.008
- Dhouibi R, Affes H, Salem M Ben, Hammami S, Sahnoun Z, Zeghal KM, Ksouda K (2020). Screening of pharmacological uses of *Urtica dioica* and others benefits. *Progress in Biophysics and Molecular Biology*, 150: 67–77. Doi:10.1016/j.pbiomolbio.2019.05.008
- Dini I (2019). An overview of functional beverages. Woodhead Publishing, MA, USA. p.1–40. Doi:10.1016/B978-0-12-816397-9.00001-7
- Durak I, Biri H, Devrim E, Sözen S, Avci A (2004). Aqueous extract of *urtica dioica* makes significant inhibition on adenosine deaminase activity in prostate tissue from patients with prostate cancer. *Cancer Biology & Therapy*, 3(9): 855–857. Doi:10.4161/cbt.3.9.1038
- El-Saber Batiha G, Hussein DE, Algammal AM, George TT, Jeandet P, Al-Snafi AE, Tiwari A, Pagnossa JP, Lima CM, Thorat ND, Zahoor M, El-Esawi M, Dey A, Alghamdi S, Hetta HF, Cruz-Martins N (2021). Application of natural antimicrobials in food preservation: Recent views. *Food Control*, 126. Doi:10.1016/j.foodcont.2021.108066
- Esposito S, Bianco A, Russo R, Maro A Di, Isernia C, Pedone PV (2019). Therapeutic Perspectives of Molecules from *Urtica dioica* Extracts for Cancer Treatment. *Molecules*, 24(15). Doi:10.3390/molecules24152753
- European Commission (2023) Novel Food Catalogue. https://webgate.ec.europa.eu/fip/novel_food_catalogue/ (accessed date: 28.09.2023).
- Fattahi S, Ardekani AM, Zabihi E, Abedian Z, Mostafazadeh A, Pourbagher R, Akhavan-Niaki H (2013). Antioxidant and apoptotic effects of an aqueous extract of *Urtica dioica* on the MCF-7 human breast cancer cell line. *Asian Pacific Journal of Cancer Prevention*, 14(9): 5317–5323
- Fattahi S, Ghadami E, Asouri M, Ardekanid AM, Akhavan-Niaki H (2018). *Urtica dioica* inhibits cell growth and induces apoptosis by targeting Ornithine decarboxylase and Adenosine deaminase as key regulatory enzymes in adenosine and polyamines homeostasis in human breast cancer cell lines. *Cellular and Molecular Biology*, 64(3): 97–102
- Flores-Ocelotl MR, Rosas-Murrieta NH, Moreno DA, Vallejo-Ruiz V, Reyes-Leyva J, Domínguez F, Santos-López G (2018). *Taraxacum officinale* and *Urtica dioica* extracts inhibit dengue virus serotype 2 replication in vitro. *BMC complementary and alternative medicine*, 18(1): 95. Doi:10.1186/s12906-018-2163-3
- García LM, Ceccanti C, Negro C, Bellis L De, Incrocci L, Pardossi A, Guidi L (2021). Effect of drying methods on phenolic compounds and antioxidant activity of *Urtica dioica* L. leaves. *Horticulturae*, 7(1): 1–12. Doi:10.3390/horticulturae7010010
- Gözüm S, Tezel A, Koc M (2003). Complementary alternative treatments used by patients with cancer in eastern Turkey. *Cancer nursing* 26(3): 230–236. Doi:10.1097/00002820-200306000-00010
- Karakol P, Saraydin SU, Bozkurt M, Hepokur C, Inan ZDS, Turan M (2022). Anticancer Effects of *Urtica Dioica* in Breast Cancer. *Asian Pacific Journal of Cancer Prevention*, 23(2): 673–681. Doi:10.31557/APJCP.2022.23.2.673

- Konrad L, Müller HH, Lenz C, Laubinger H, Aumüller G, Lichius JJ (2000). Antiproliferative effect on human prostate cancer cells by a stinging nettle root (*Urtica dioica*) extract. *Planta Medica*, 66(1): 44–47. Doi:10.1055/s-2000-11117
- Körpe DA, Işeri ÖD, Sahin FI, Cabi E, Haberal M (2013). High-antibacterial activity of *Urtica* spp. seed extracts on food and plant pathogenic bacteria. *International Journal of Food Sciences and Nutrition*, 64(3): 355–362. Doi:10.3109/09637486.2012.734290
- Levy A, Sivanesan D, Murugan R, Quinonez Y, Jaffe M, Rathinavelu A, Fattahi S, Ardekani AM, Zabihi E, Abedian Z, Mostafazadeh A, Pourbagher R, Akhavan-niaki H (2013). *Urtica dioica* Induces Cytotoxicity in Human Prostate Carcinoma LNCaP Cells : Involvement of Oxidative Stress, Mitochondrial Depolarization and Apoptosis. *Asian Pacific Journal of Cancer Prevention*, 13(5): 711–717
- Mannila E, Marti-Quijal FJ, Selma-Royo M, Calatayud M, Falcó I, la Fuente B de, Barba FJ, Collado MC, Linderborg KM (2022). In Vitro Bioactivities of Food Grade Extracts from Yarrow (*Achillea millefolium* L.) and Stinging Nettle (*Urtica dioica* L.) Leaves. *Plant Foods for Human Nutrition*, 78: 132–138. Doi:10.1007/s11130-022-01020-y
- Mansoori B, Mohammadi A, Hashemzadeh S, Shirjang S, Baradaran A, Asadi M, Doustvandi MA, Baradaran B (2017). *Urtica dioica* extract suppresses miR-21 and metastasis-related genes in breast cancer. *Biomedicine & Pharmacotherapy*, 93: 95–102. Doi:10.1016/j.biopha.2017.06.021
- Mohammadi A, Mansoori B, Baradaran PC, Khaze V, Aghapour M, Farhadi M, Baradaran B (2017). *Urtica dioica* Extract Inhibits Proliferation and Induces Apoptosis and Related Gene Expression of Breast Cancer Cells In Vitro and In Vivo. *Clinical Breast Cancer*, 17(6): 463–470. Doi:10.1016/j.clbc.2017.04.008
- Mohammadi A, Mansoori B, Goldar S, Shanehbandi D, Khaze V, Mohammadnejad L, Baghbani E, Baradaran B (2016). Effects of *Urtica dioica* dichloromethane extract on cell apoptosis and related gene expression in human breast cancer cell line (MDA-MB-468). *Cellular and molecular biology (Noisy-le-Grand, France)* 62(2): 62–67
- Özkol H, Musa D, Tuluce Y, Koyuncu I (2012). Ameliorative influence of *Urtica dioica* L against cisplatin-induced toxicity in mice bearing Ehrlich ascites carcinoma. *Drug and Chemical Toxicology*, 35(3): 251–257. Doi:10.3109/01480545.2011.598531
- Paulauskienė A, Tarasevičienė Ž, Laukagalis V (2021). Influence of harvesting time on the chemical composition of wild stinging nettle (*Urtica dioica* l.). *Plant,s* 10(4). Doi:10.3390/plants10040686
- Sayhan MB, Kanter M, Oguz S, Erboga M (2012). Protective effect of *Urtica dioica* L. on renal ischemia/reperfusion injury in rat. *Journal of Molecular Histology*, 43(6): 691–698. Doi:10.1007/s10735-012-9436-9
- Shonte TT, Duodu KG, Kock HL de (2020). Effect of drying methods on chemical composition and antioxidant activity of underutilized stinging nettle leaves. *Heliyon*, 6(5). Doi:10.1016/j.heliyon.2020.e03938
- Skalozubova TA, Reshetova VO (2013). Leaves of common nettle (*Urtica dioica* L.) as a source of ascorbic acid (vitamin C). *World Applied Sciences Journal*, 28(2): 250–253. Doi:10.5829/idosi.wasj.2013.28.02.13792
- Upton R (2013). Stinging nettles leaf (*Urtica dioica* L.): Extraordinary vegetable medicine. *Journal of Herbal Medicine*, 3(1): 9–38. Doi:10.1016/j.hermed.2012.11.001
- Veiga M, Costa EM, Silva S, Pintado M (2020). Impact of plant extracts upon human health: A review. *Critical Reviews in Food Science and Nutrition*, 60(5): 873–886. Doi:10.1080/10408398.2018.1540969