

THE CHECK-LIST OF THE SUBFAMILY CRYPTOCEPHALINAE  
(COLEOPTERA, CHRYSOMELIDAE) IN TURKEY

İrfan ASLAN<sup>(1)</sup>

Hikmet ÖZBEK<sup>(1)</sup>

**SUMMARY:** In this study 75 species and 9 subspecies of 3 genera in 3 tribes have been recorded up to date in Turkey. Eighteen species are endemic. The host plant of some species were indicated.

TÜRKİYE' NİN CRYPTOCEPHALINAE (COLEOPTERA,  
CHRYSOMELIDAE) TÜRLERİ LİSTESİ

**ÖZET:** Bu çalışmada, 3 tribus'e ait 3 cins'e bağlı 75 tür ve 9 alttür'ün ülkemizde bulunduğu belirlenmiştir. Bunlardan 36 tür ve 2 alttür incelenmiş, geri kalan 39 tür ve 7 alttür ise literatür değerlendirilerek verilmiştir. Tespit edilen türlerden 18 tür endemiktir. Ayrıca, bir çok türlerin de konukçuları tespit edilmiştir.

INTRODUCTION

Chrysomelidae known as leaf beetles, is a large family of phytophagous Coleoptera, comprising approximately 50000 species in over 2000 genera and 19 subfamilies in world fauna (Lopatin, 1977).

The subfamily Cryptocephalinae has a world-wide distribution. The members are usually small, 10 mm or less in length, more or less cylindirical, compact, globrous beetles, with a flat hypognathous head, deeply sunk into the prothorax, and a large exposed pygidium (Booth et al., 1990).

Turkey is a large country, has various climates and topographic structure. Therefore it is expected that the Cryptocephalinae fauna of Turkey should be very rich. In spite of world fauna of this subfamily has been well studied (Clavareau, 1913; Mohr, 1966; Lopatin, 1977; Gruev and Tomov, 1984) particularly native researchers have not studied the Turkish fauna. Except Güll-Zümreoğlu (1972) and Tuatay et al. (1972) indicated presence of two species everyone of them, Selmi (1982) noted eight species with their host plants. Foreign scientists as Pic (1908), Clavareau (1913), Medvedev (1970, 1975), Warchałowski (1976) and Gruev and Tomov (1979) made faunal studies and described some new taxa. But all records were from the western part of Turkey, only one species from Trabzon, two species from Mardin and three species from Malatya were recorded (Clavareau, 1913; Medvedev, 1970).

(1) Atatürk University, Agricultural Faculty, Department of Plant Protection, 25240 Erzurum, Turkey.

This study based upon the material collected mostly from the eastern part of the country. The species have been already recorded were reviewed.

## RESULTS

Present study and previous records reveal that 75 species and 9 subspecies of 3 genera in 3 tribes occur in Turkey.

### Tribe: CRYPTOCEPHALINI

Genus: *Cryptocephalus* Geoffroy, 1762

*Cryptocephalus amasiensis* Weise, 1894

Distribution in Turkey: Amasya (Weise, 1894), Turkey, no locality (Clavareau, 1913), Turkey, no locality (Warchałowski, 1976). It is an endemic species in Turkey.

*Cryptocephalus angorensis* Pic, 1908

Distribution in Turkey: Ankara (Pic, 1908), Anatolia (Clavareau, 1913), Turkey, no locality (Warchałowski, 1976). It is an endemic species in Turkey.

*Cryptocephalus apicalis* Gebler, 1830

Distribution in Turkey: Turkey, no locality (Clavareau, 1913), West Black Sea Region (Warchałowski, 1991).

Distribution in the world: Hungary, Russia, Siberia, Anatolia (Clavareau, 1913), European part of the USSR, northern Caucasus, Kazakhstan (Lopatin, 1977), Bulgaria, Crimea (Warchałowski, 1976), Europe, Asia Minor, Kazakhstan (Gruev and Tomov, 1984).

*Cryptocephalus aureolus* Suffrian, 1847

Material examined: Dikyar, Uzundere (Erzurum), 1500 m, 2.VII.1994, 1♀, 1♂, *Centaurea solstitialis*; Palandöken, Erzurum, 2300 m, 28.VII.1992, 4♂♂, 10.VIII.1993, 2♂♂, 13.VII.1993, 2♀♀, 1♂, 19.VIII.1993, 1♂, *C. solstitialis* and some other Compositae; Altıparmak, Yusufeli (Artvin), 1700 m, 6.VII.1994, 2♀♀, 5♂♂, *C. solstitialis* and some other Compositae; Akarsu, Ardanuç (Artvin), 1600 m, 6.VII.1994, 1♀, 7♂♂, Compositae; Sarıgöl, Yusufeli, 1200 m, 5.VII.1994, 1♂, Compositae; Ferhatlı, Ardanuç, 1500 m, 7.VII.1994, 1♀, *C. solstitialis*; Pehlivanhı, Tortum (Erzurum), 1300 m, 12.VII. 1993, 1♂, *C. solstitialis*; Sandıkköy, Kemaliye (Erzincan), 21.VII.1984, 1♂; Belkız (Antalya), 200 m, 3.VIII.1994, 1♂; Abdurrahmangazi, Erzurum, 2200 m, 17.VIII.1993, 1♂, *C. solstitialis*; Oğuzkent, Olur (Erzurum), 2200 m, 23.VIII.1993, 2♂♂, *C. solstitialis*. and some other Compositae; Kireçli, Tortum (Erzurum), 1700 m, 8.VIII.1995, 2♂♂.

Distribution in Turkey: Turkey, no locality (Warchałowski, 1976).

Distribution in the world: Europe (Clavareau, 1913), Mid Europe (Mohr, 1966), European part of the USSR, Caucasus, Kazakhstan, Europe, Asia Minor (Lopatin, 1977), Bulgaria, Poland, Venezuela, Albania, Yugoslavia, Greece (Warchałowski, 1976).

***Cryptocephalus bicolor*** Eschscholtz, 1818

Distribution in Turkey: Oltu, Tortum, Uzundere (Erzurum) (Aslan and Özbek, 1997).

Distribution in the world: Caucasus, Hungary, Trans-Slovakia, South Russia (Clavareau, 1913), Bulgaria, Crimea, Caucasus, Azerbaijan, Armenia (Warchałowski, 1976) Europe, Central Asia and Kazakhstan (Gruev and Tomov, 1984).

***Cryptocephalus biguttatus*** (Scopoli, 1763)

***Buprestis biguttata*** Scopoli, 1763 (Gruev and Tomov, 1984).

Distribution in Turkey: Ankara (Tomov and Gruev, 1975).

Distribution in the world: Europe, Siberia (Clavareau, 1913), European part of the USSR, southern Urals, southern Siberia, Kazakhstan, Europe (Lopatin, 1977), Europe (Gruev and Tomov, 1984).

***Cryptocephalus biledjekensis*** Pic, 1909

Distribution in Turkey: Bilecik (Pic, 1909), Anatolia, no locality (Clavareau, 1913), Asia Minor, no locality (Gruev and Tomov, 1984). It is an endemic species in Turkey.

***Cryptocephalus bilineatus*** (Linnaeus, 1767)

***Chrysomela bilineata*** Linnaeus, 1767 (Gruev and Tomov, 1984).

Distribution in Turkey: Turkey, no locality (Warchałowski, 1976).

Distribution in the world: Europe, Anatolia, Siberia, Japan (Clavareau, 1913), Mid Europe (Mohr, 1966), Poland, Crimea, Caucasus, Azerbaijan, Armenia (Warchałowski, 1976), Bulgaria, Europe (Gruev and Tomov, 1984), North Caucasus (Yaroshenko, 1985)

***Cryptocephalus bipunctatus*** (Linnaeus, 1758)

***Chrysomela bipunctata*** Linnaeus, 1758 (Gruev and Tomov, 1984).

Material examined: Penek, Şenkaya (Erzurum), 1200m, 19.V.1996, 1♀, 1♂, *Populus* spp.; Madenköprübaşı, İspir (Erzurum), 1500 m, 18.VI.1994, 1♂, *Salix* spp.; Abdurrahmangazi, Erzurum, 2200 m, 14.VII.1992, 1♂; Dikyar, Uzundere (Erzurum), 1550 m, 2.VII.1994, 1♀, 2♂♂; 30.VI.1996, 2♀♀, 3♂♂, *Salix* and *Populus* spp.; Rabat, Pasinler (Erzurum), 2300m, 11.VIII.1996, 4♀♀, 6♂♂, *Populus* spp.; Pişkidağ, Erzincan, 1200 m, 8.VII.1993, 1♀, *Salix* spp.; Ankara, 900 m, 2.VII. 1992, 1♀, 1♂; Altıparmak,

Yusufeli (Artvin), 1700 m, 5.VII.1994, 1♀, *Salix* spp.; Akarsu, Ardanuç (Artvin), 1500 m, 7.VII.1994, 2♂♂, *Salix* and *Populus* spp..

Distribution in Turkey: Kaynarca (Iznik), between Elmalık and Bakacak, between Amasya and Ezinepazarı (Medvedev, 1970), Amasya, Ankara, İçel, Sinop (Tomov and Gruev, 1975).

Distribution in the world: Europe, Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Crimea, Venezuela, Albania, Yugoslavia (Warchałowski, 1976), European part of the USSR, Caucasus, Siberia, Kazakhstan, northern Africa, central and southern Europe, northern China, Korea (Lopatin, 1977), Europe, Greece, Central Asia (Gruev and Tomov, 1984).

***Cryptocephalus bodemeyeri* Weise, 1900**

Material examined: Turnalı, Şenkaya (Erzurum), 1750 m, 15.V.1994, 1♂; Sarıkamış (Kars), 2000 m, 22.VI.1995, 1♀; Sutkans, Oltu (Erzurum), 1800 m, 18.VI.1996, 1♀; Tepeköy, Aşkale (Erzurum), 8.VI.1996, 1♀.

Distribution in Turkey: Bilecik, Anatolia (Clavareau, 1913), Kavak (Samsun) (Medvedev, 1970), Turkey (Warchałowski, 1976). It is an endemic species in Turkey.

***Cryptocephalus chrysopus* Gmelin, 1788**

Distribution in Turkey: Turkey (Warchałowski, 1976), Kavak (Samsun) (Medvedev, 1970).

Distribution in the world: Mid and North Europe (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Crimea, Caucasus, Azerbaijan, Armenia, Albania, Yugoslavia (Warchałowski, 1976), Europe, Asia, Bulgaria (Gruev and Tomov, 1984), North Caucasus (Yaroshenko, 1985).

***Cryptocephalus concolor* Suffiran, 1847**

Distribution in Turkey: Amasya, Ankara, Artvin, Erzurum, Gümüşhane, Samsun, İçel, Sinop, Sivas, Trabzon (Tomov and Gruev, 1975), Turkey, no locality (Warchałowski, 1976).

Distribution in the world: Caucasus, Anatolia, South Russia (Clavareau, 1913), Bulgaria, Greece, Caucasus, Azerbaijan, Armenia, Albania, Yugoslavia (Warchałowski, 1976), Southern caucasus and Turkmenia (Lopatin, 1977), Bulgaria, Mid Asia (Gruev and Tomov, 1984).

***Cryptocephalus connexus* Olivier, 1807**

Material examined: Very common in Pasinler (Erzurum), *Onobrychis sativa*; Dumlu, Erzurum, 1900 m, 2.VII.1994, 2♀♀; Kinalıçam, Yusufeli (Artvin), 850 m, 4.VII.1994, 5♀♀,

6♂♂, *Ulmus* spp.; Anzavderesi, Oltu (Erzurum), 900m, 5.VII.1995, 10♀♀, 13♂♂, *O. sativa*; Bilecik, 800m, 15.VII.1995, 2♀♀, 3♂♂; Palandöken, Erzurum, 2300 m, 19.VIII.1993, 6♀♀, 2♂♂; Dikyar, Uzundere (Erzurum), 1450m, 8.IX.1996, 5♀♀, 10♂♂, *Ulmus* spp.; Kandilli, Erzurum, 1750m, 22.VIII. 1996, 3♀♀, 2♂♂, *O. sativa*.

Distribution in Turkey: İzmir (Gül-Zümreoglu, 1972), All Turkey (Warchalowski, 1991).

Distribution in the world: Austria, Hungary, Rumania and Russia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Crimea, Venezuela, Albania, Yugoslavia (Warchalowski, 1976), European part of the USSR, Caucasus, Trans-Caucasus, Central Asia, western Turkmenia, Kazakhstan, Mediterranean zone and eastern Europe (Lopatin, 1977), North Caucasus, East Europe, North Kazakhstan (Yaroshenko, 1985).

***Cryptocephalus corithius* Pic, 1914**

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976). It is an endemic species in Turkey.

***Cryptocephalus creticus* Suffiran, 1847**

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Crete, Dalmatica (Clavareau, 1913), Albania, Yugoslavia Greece (Warchalowski, 1976).

***Cryptocephalus cibratus* Suffiran, 1847**

Material examined: Selale, Uzundere (Erzurum), 1350 m, 9.VI.1996, 3♀♀, 2♂♂, *Salix* and *Populus* spp.; Dikyar, Uzundere, 1450m, 17.VI. 1995, 4♀♀, 3♂♂, 30.VI.1996, 5♀♀, 4♂♂, 31.VII.1993, 1♀, 2.VII.1994, 1♂, *Salix* and *Populus* spp.; Karagöl, Borça (Artvin), 1050 m, 18.VII.1994, 1♂, *Salix alba*.

Distribution in Turkey: Gümüşhane (Tomov and Gruev, 1975), Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Turkey, Caucasus (Clavareau, 1913), Caucasus, Azerbaijan, Armenia (Warchalowski, 1976), All Caucasus, Turkmenia, Asia Minor and Iran (Lopatin, 1977).

***Cryptocephalus curda* Jacobson, 1897**

Distribution in Turkey: Dinar (Afyon), Palandöken, University Field (Erzurum) (Aslan and Özbek, 1997).

Distribution in the world: Russia (Clavareau, 1913), Caucasus, Azerbaijan, Armenia (Warchalowski, 1976).

*Cryptocephalus duplicatus* Suffiran, 1847

Material examined: Very common in Erzurum and Artvin provinces (from 600 m up to 2000m), *Centaurea solstitialis* and some other Compositae; Derinöz, Digor (Kars), 2000 m, 21.VI.1995, 8♀♀, 10♂♂, *C. solstitialis*; Sarıkamış (Kars), 1900 m, 13.VII.1996, 7♀♀, 6♂♂, Compositae.

Distribution in Turkey: Belgrat Forest, İstanbul, *Salix cinerea* and *Populus tremula* (Selmi, 1982).

Distribution in the world: Central Caucasus (Clavareau, 1913), Caucasus, Iran (Berti and Rapilly, 1973), Caucasus, Azerbaijan, Armenia (Warchałowski, 1976).

*Cryptocephalus elegantulus* Gravenhorst, 1807

Distribution in Turkey: Turkey, no locality (Warchałowski, 1976).

Distribution in the world: Mid Europe, Anatolia, West Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Venezuela, Crimea, Caucasus, Azerbaijan, Armenia (Warchałowski, 1976), European part of the USSR, Caucasus, Transcaucasus, Siberia, Kazakhstan, Kirgizia, Mongolia, Korea (Lopatin, 1977), Europe, Central Asia, Mongolia (Gruev and Tomov, 1984).

*Cryptocephalus exiguum* Schneider, 1792

Distribution in Turkey: Turkey, no locality (Warchałowski, 1976), Kaynarca (İznik) (Medvedev, 1970), Asia Minor, no locality (Gruev and Tomov, 1984).

Distribution in the world: Mid Europe, West Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Venezuela (Warchałowski, 1976), Europe, Asia Minor, Bulgaria (Gruev and Tomov, 1984).

*Cryptocephalus flavipes* Fabricius, 1781

Material examined: Madenkörprübaşı, İspir (Erzurum), 1400 m, 1♀, 2♂♂; Dikyar, Uzundere (Erzurum), 1450 m, 9.VI.1996, 1♀, 2♂♂; Turnalı, Şenkaya (Erzurum), 1750 m, 25.VII.1996, 1♂; Eğerti, İlca (Erzurum), 1800 m, 28.VI.1997, 2♀♀, 1♂; Sutkans, Oltu (Erzurum), 1900 m, 10.VI.1997, 2♀♀, 3♂♂; Ormanlı, Şenkaya (Erzurum), 20.V.1997, 1♀; Yusufeli (Artvin), 11.VI.1997, 1♀; Altınparmak, Yusufeli (Artvin), 12.VI.1997, 1♀, 1♂.

Distribution in Turkey: Taşdelen (Malatya), Kaynarca (İznik), between Bolu and Yeniçağ, 1000 m, beween Elmalık and Bakacak, 850m, Mataracı, Bafra (Samsun), Vakfıkebir (Trabzon) (Medvedev, 1970), Ordu (Tuatay et al., 1972), İstanbul, Samsun (Tomov and Gruev, 1975), Turkey, no locality (Warchałowski, 1976).

Distribution in the world: Europe, Anatolia, Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Crimea, Venezuela, Caucasus, Azerbaijan, Armenia, Albania, Yugoslavia (Warchałowski, 1976), European part of the USSR, Caucasus, southern Urals, Siberia, Kazakhstan, southwestern Turkmenia, northern Iran (Lopatin, 1977), Europe, Greece, Central Asia, Caucasus (Gruev and Tomov, 1984).

*Cryptocephalus fulvus* Goeze, 1777

Material examined: Erzincan, 14.VI.1994, 2♂♂, *Artemisia* spp.; Köse (Gümüşhane), 900 m, 10.VII.1995, 1♀, 2♂♂, *Artemisia* spp.; Pasinler (Erzurum) 1600m, 15.VII.1995, 6♀♀, 7♂♂, 4.VIII.1993, 3♂♂, *Artemisia* spp.; Rizekent, İlica (Erzurum), 1900 m, 18.VIII.1995, 3♀♀, 2♂♂, *Artemisia* spp.; Turnalı, Şenkaya (Erzurum), 1750m, 8.IX. 1994, 2♀♀, 1♂; Tece (İçel), 2.IX.1992, 4♀♀, 3♂♂.

Distribution in Turkey: Efes (İzmir), Antalya, Ovacık, Silifke (Mersin) (Medvedev, 1975).

Distribution in the world: Europe, West Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Venezuela, Albania, Yugoslavia (Warchałowski, 1976), Europe, Asia Minor, Bulgaria (Gruev and Tomov, 1984), North Caucasus (Yaroshenko, 1985).

*Cryptocephalus glriosus* Mulsant, 1853

Distribution in Turkey: Karaman (Clavareau, 1913), Turkey, no locality (Warchałowski, 1976). It is an endemic species in Turkey.

*Cryptocephalus hypochoeridis* (Linnaeus, 1758)

*Chrysomela hypochoeridis* Linnaeus, 1758 (Gruev and Tomov, 1984).

Distribution in Turkey: Oltu, Tortum, Uzundere (Erzurum), Borçka, Yusufeli (Artvin), Sarıkamış (Kars) *Centaurea solstitialis* and some other Compositae, Rize (Aslan and Özbek, 1997).

Distribution in the world: Europe, Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Venezuela, Albania, Yugoslavia (Warchałowski, 1976), East Europe, North Caucasus, Crimea, North Kazakhstan (Yaroshenko, 1985), Europe and Central Asia (Gruev and Tomov, 1984).

*Cryptocephalus hypochoeridis praticola* Weise, 1901

Distribution in Turkey: Distribution in Turkey: Oltu, Uzundere (Erzurum), Yusufeli (Artvin) *Centaurea solstitialis* and some other Compositae (Aslan and Özbek, 1997).

Distribution in the world: North Kazakhstan and North Caucasus (Yaroshenko, 1985).

*Cryptocephalus hypochoeridis ruguliventris* Franz, 1949

Distribution in Turkey: Artvin (Tomov and Gruev, 1975). It is an endemic species in Turkey.

*Cryptocephalus ilicis* Olivier, 1808

Distribution in Turkey: Turkey, no locality (Warchałowski, 1976), Gönen (Balıkesir), Alemdağ (İstanbul), Bahçeköy (İstanbul), *Quercus coccifera* and *Salix babylonica* (Selmi, 1982).

Distribution in The world: South Europe (Clavareau, 1913), Greece, Italia, Seilia, France, Spain (Burlini, 1955), Albania, Yugoslavia, Greece (Warchałowski, 1976),

*Cryptocephalus ilicis* var. *oblitteratipennis* Pic, 1900

Distribution in Turkey: İstanbul (Clavareau, 1913). It is an endemic species in Turkey.

*Cryptocephalus ilicis* var. *principipennis* Pic, 1908

Distribution in Turkey: İstanbul (Clavareau, 1913). It is an endemic species in Turkey.

*Cryptocephalus imperialis* Laicharting, 1781

Material examined: Aşağı Meydanlar, Tortum (Erzurum), 1700m, 14.VII.1991, 1♀, 1♂, *Quercus* sp.; Şellale, Uzungere (Erzurum), 9.VI.1996, 1♂.

Distribution in Turkey: Turkey, no locality (Warchałowski, 1976), Amasya, 420-900m (Medvedev, 1975).

Distribution in the world: South Europe (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Venezuela, Albania, Yugoslavia, Greece (Warchałowski, 1976), South and North Europe, Asia (Gruev and Tomov, 1984).

*Cryptocephalus inaequalis* Fairmaire, 1902

Distribution in Turkey: Ankara (Clavareau, 1913). It is an endemic species in Turkey.

*Cryptocephalus labiatus* (Linnaeus, 1761)

*Chrysomela labiatus* Linnaeus, 1761 (Gruev and Tomov, 1984).

Material examined: Eğerti, İlica (Erzurum), 1800 m, 28.VI.1997, 1♀, 1♂.

Distribution in Turkey: Taşdelen (Malatya) (Medvedev, 1970), Ordu (Tuatay et al., 1972).

Distribution in the world: Europe, Mongolia, East Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Europe, Asia, Bulgaria (Gruev and Tomov, 1984), North Caucasus (Yaroshenko, 1985).

*Cryptocephalus janthinus* Germar, 1824

Distribution in Turkey: İçel (Tomov and Gruev, 1975), Kars, Ardahan (Warchalowski, 1991).

Distribution in the world: Mid and South Europe, Siberia (Clavareau, 1913), European part of the USSR, Caucasus, Siberia, western Kazakhstan, Central and southern Europe (Lopatin, 1977), Europe, Central Asia (Gruev and Tomov, 1984).

*Cryptocephalus laevicollis* Gebler, 1830

Distribution in Turkey: Amasya, Çorum (Tomov and Gruev, 1975), Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Turkey, South Russia, Siberia (Clavareau, 1913), Slovakia (Mohr, 1966), Bulgaria, Poland, Crimea, Caucasus, Azerbaijan, Armenia (Warchalowski, 1976), European part of the USSR, Kazakhstan, Central and eastern Europe (Lopatin, 1977), Europe, Asia, Bulgaria (Gruev and Tomov, 1984).

*Cryptocephalus macellus* Suffrian, 1860

Material examined: Sükans, Oltu (Erzurum), 1800 m, 10.VI.1997, 1♀, 1♂.

Distribution in Turkey: Bilecik (Tomov and Gruev, 1975), Turkey, no locality (Warchalowski, 1976), Taşdelen Malatya, Belğirat Forest (İstanbul), Samsun, between Turhal and Ezinepazarı, between Amasya and Ezinepazarı, Madenli, Urabat (Medvedev, 1970), Biga (Çanakkale), Adapazarı, Belğirat Forest (İstanbul), Kalkım (Balıkesir), *Populus alba*, *Quercus petraea*, *Q. robur* (Selmi, 1982).

Distribution in the world: Mid and South Europe (Clavareau, 1913), Mid Europe, Slovakia (Mohr, 1966), Spain, Austria, Cyprus, Crete, Italia, Germany, northwestren Iran (Roubal, 1937-1941), Bulgaria, Venezuela, Albania, Yugoslavia, Greece (Warchalowski, 1976), Europe, Asia, Bulgaria (Gruev and Tomov, 1984).

*Cryptocephalus moehringi* Weise, 1884

Distribution in Turkey: Amasya (Clavareau, 1913), Turkey, no locality (Warchalowski, 1976), between Amasya and Ezinepazarı, Ezinepazarı (Medvedev, 1970), Bolu (Tomov and Gruev, 1975).

Distribution in the world: Greece, Turkey (Warchalowski, 1976).

*Cryptocephalus moraei* (Linnaeus, 1758)

*Chrysomela moraei* Linnaeus, 1758 (Gruev and Tomov, 1984).

Material examined: Silvan (Diyarbakır), 800 m, 17.IV.1995, 8♀♀, 7♂♂, *Hypericum* sp.; Sutkans, Oltu, 1600 m, 18.VI.1996, 3♂♂, *Hypericum* sp.; Dikyar, Uzundere (Erzurum), 1450m, 25.VI.1996, 3♀♀, 4♂♂, *Hypericum* sp.; Palandöken, Erzurum, 2300 m, 13.VII.1994, 1♂; Çamlıbel, Oltu (Erzurum), 1750m, 14.VII.1996, 3♀♀, 2♂♂, *Hypericum* sp.; Kelkit (Gümüşhane), 850 m, 9.VII.1995, 3♂♂, *Hypericum* sp.; Ordu, 100 m, 16.VII.1991, 1♂; Sandık Köyü, Kemaliye (Erzincan), 1100 m, 21.VII.1984, 1♀; Üzümlü (Erzincan), 1600 m, 8.VII.1993, 3♂♂, *Hypericum* sp.; Yollarüstü (Erzincan), 1300 m, 9.VII.1994, 2♀♀, 2♂♂, *Hypericum* sp.; Üzümlü, 1450 m, 13.VII.1994, 2♀♀, *Hypericum* sp..

Distribution in Turkey: İzmir (Gül-Zümreoğlu, 1972), Samsun (Tomov and Gruev, 1975), Turkey, no locality (Warchałowski, 1976), Uludağ (Bursa) (Gruev and Tomov, 1979).

Distribution in the world: Europe, West Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Crimea, Venezuela, Albania, Yugoslavia, Greece (Warchałowski, 1976), Europe, Central Asia, Caucasus (Gruev and Tomov, 1984), North Caucasus (Yaroshenko, 1985)..

*Cryptocephalus ocellatus* Drapiez, 1819

Material examined: Çerli Köyü, Perşembe (Ordu), 100 m, 29.VI.1991, 1♂; Demirkent, Yusufeli (Artvin), 700 m 6.VII.1994, 2♂♂, *Salix* sp.; Karagöl, Borçka (Artvin), 1100 m, 7.VII.1994, 6♀♀, 8♂♂, *Alnus* sp.; Yollarüstü (Erzincan), 1300 m, 9.VII.1993, 1♀; Rizekent, İlica (Erzurum), 1800 m, 18.VIII.1995, 2♀♀, 1♂, *Salix* sp.; Köşkköyü, Erzurum, 2000 m, 20.VI.1996, 1♂.

Distribution in Turkey: Rize (Tomov and Gruev, 1975), Turkey, no locality (Warchałowski, 1976), Kepsut (Balıkesir), İnegöl (Bursa), Bahçekök (İstanbul), Bayramiç (Çanakkale), *Salix alba*, *S. fragilis* and *Alnus glutinosa* (Selmi, 1982).

Distribution in the world: South and Mid Europe (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Crimea, Caucasus, Azerbaijan, Armenia, Venezuela, Albania, Greece, Yugoslavia (Warchałowski, 1976), European part of the USSR, Kazakhstan, Europe except north (Lopatin, 1977), Europe, Asia Minor, Caucasus, Kazakhstan (Gruev and Tomov, 1984).

*Cryptocephalus octacosmus* Bedel, 1891

Material examined: Çerkezköy, Tekirdağ, 10.VI.1995, 2♂♂; Bilecik, 800 m, 15.VI.1995, 10♀♀, 13♂♂; Aydıntepe (Bayburt), 1500 m, 30.VI.1991, 3♂♂; Pasinler (Erzurum), 1600 m, 17.VII.1980, 1♂; Demirözü (Bayburt), 1700 m, 23.VII.1991, 1♀;

Kopdağı, Bayburt, 2300 m, 10.VIII.1991, 1♀, 1♂; Bayburt, 1600 m, 11.VIII.1991, 1♂;  
Anzavderesi, Oltu (Erzurum), 28.VIII.1996, 1♀, 1♂.

Distribution in Turkey: Ankara, Tokat (Tomov and Gruev, 1975), Jalomba (İzmit) (Medvedev, 1975), Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Mid and South Europe, West Siberia (Clavareau, 1913), Southern Mid Europe (Mohr, 1966), Bulgaria, Poland, Crimea, Venezuela, Albania, Yugoslavia, Greece (Warchalowski, 1976), European part of the USSR, Caucasus, western Siberia, Kazakhstan, Uzbekistan, Kirgizia, Northwestren China (Lopatin, 1977), Europe, Central Asia, Caucasus (Gruev and Tomov, 1984).

*Cryptocephalus paradisiacus* Weise, 1900

Distribution in Turkey: Mardin (Clavareau, 1913), Turkey, no locality (Warchalowski, 1976). It is an endemic species in Turkey.

*Cryptocephalus parvulus* Müller, 1776

Distribution in Turkey: Samsun (Tomov and Gruev, 1975), Uludağ (Bursa) ,800-1750m (Medvedev, 1975).

Distribution in the world: Europe, Siberia, Japan (Clavareau, 1913), Europe (Mohr, 1966), Poland, Venezuela (Warchalowski, 1976), European part of the USSR, Urals, Siberia, Kazakhstan, Europe (Lopatin, 1977), Europe, Central Asia, Kazakhstan (Gruev and Tomov, 1984).

*Cryptocephalus peyroni* Marseul, 1875

Distribution in Turkey: Borçka (Artvin), 1050 m (Aslan and Özbek, 1997).

Distribution in the world: Lebanon (Beirut) ( Clavareau, 1913).

*Cryptocephalus prope macellus* Suffrian, 1860

Distribution in Turkey: Kanlıdivane (Samsun) (Medvedev, 1970). It is an endemic species in Turkey.

*Cryptocephalus prusias* Suffrian, 1853

Material examined: Tepeköy, Aşkale (Erzurum), 8.VI.1997, 1♀, 2♂♂.

Distribution in Turkey: Bilecik (Bodameyer, 1900), Amasya, Bursa (Clavareau, 1913), between Amasya and Ezinepazarı (Medvedev, 1970), Amasya, Gümüşhane (Tomov and Gruev, 1975), Sındırıcı (Balıkesir), Bandırma (Balıkesir), Demirköy (Kırklareli), Kestel (Bursa), *Salix triandra*, *S.alba* and *Quercus petraea* (Selmi, 1982).

Distribution in the world: Turkey (Clavareau, 1913), Bulgaria, Greece (Warchalowski, 1976),

*Cryptocephalus pusillus* Fabricius, 1777

*Chrysomela minuta* Herbst, 1783; *Cryptocephalus verticalis* Boheman, 1851 (Warchalowski, 1991).

Distribution in Turkey: Aegean Region (Warchalowski, 1991).

Distribution in the world: Germany, Spain, All Europe, Bulgaria, Poland, Venezuela (Warchalowski, 1991).

*Cryptocephalus pygmaeus* Fabricius, 1792

Material examined: Dinar (Afyon), 14.VIII. 1990, 2♀, 3♂.

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976; 1991).

Distribution in the world: Germany, Spain (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Venezuela (Warchalowski, 1976), Europe, Asia, Caucasus, Bulgaria (Gruev and Tomov, 1984), North Caucasus, Kazakhstan (Yaroshenko, 1985)

*Cryptocephalus pygmaeus* var. *lugubris* Demaison, 1904

Distribution in Turkey: Anatolia (Clavareau, 1913). It is an endemic subspecies in Turkey.

*Cryptocephalus pygmaeus* var. *orientalis* Weise, 1882

Distribution in Turkey: Anatolia (Clavareau, 1913). It is an endemic subspecies in Turkey.

*Cryptocephalus quatuordecimmaculatus coloratus* Fabricius, 1798

Distribution in Turkey: Turkey, no locality (Clavareau, 1913).

Distribution in the world: Austria, Turkey, Russia, Siberia (Clavareau, 1913).

*Cryptocephalus reitteri* Weise, 1882

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Dalmatia, Croatia (Clavareau, 1913), Crimea, Caucasus, Azerbaijan, Armenia, Venezuela, Albania, Yugoslavia, Greece (Warchalowski, 1976),

*Cryptocephalus rufipes* Goeze, 1777

Distribution in Turkey: İstanbul (Medvedev, 1970).

Distribution in the world: Europe (Clavareau, 1913), Mid Europe (Mohr, 1966), Crimea (Warchalowski, 1976).

*Cryptocephalus rugicollis* Olivier, 1807

Distribution in Turkey: İzmir, Amasya (Gül-Zümreoglu, 1972), Antalya, Muğla (Tomov and Gruev, 1975), Efes (İzmir), Aydın (Medvedev, 1975).

Distribution in The world: South Europe (Clavarau, 1913).

*Cryptocephalus schaefferi* Schrank, 1789

Material examined: Tepeköy, Aşkale (Erzurum), 8.VI.1997, 3♀♀, 2♂♂.

Distribution in Turkey: Turkey, no locality (Clavareau, 1913), Edirne, Kırklareli (Warchalowski, 1991).

Distribution in the world: France, Germany, Italia, Russia, West Siberia (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Crimea, Azerbaijan, Caucasus, Armenia (Warchalowski, 1976), Europe, Greece, Caucasus, Kazakhstan (Gruev and Tomov, 1984), European part of the USSR, Caucasus, Transcaucasus, western Siberia, Central Kazakhstan, central and southern Europa (Lopatin, 1977)

*Cryptocephalus sericeus* (Linnaeus, 1758)

*Chrysomela sericea* Linnaeus, 1758; (Gruev and Tomov, 1984).

Material examined: Karagöbek, Erzurum, 1900 m, 5.VII.1969, 1♂; Çiftlik Köyü (Tortum (Erzurum), 1500 m, 14.VII.1994, 1♀; Aksuyaylaşı, Tortum (Erzurum), 2250m, 14.VII.1996, 3♀♀, 6♂♂, *Centaurea solstitialis* and some other Compositae; Beytepe, Ankara, 900 m, 20.VII.1994, 1♂; Dikyar, Uzundere (Erzurum), 1500 m, 31.VII.1993, 1♂; 9.VII.1996, 7♀♀, 10♂♂, 6.VIII.1996, 5♀♀, 9♂♂, *C. solstitialis* and some other Compositae; Oğuzkent, Olur (Erzurum), 2100 m, 23.VII.1993, 1♀, *C. solstitialis*; Palandöken, Erzurum, 2300 m, 10.VII.1993, 1♂; 23.VII.1992, 2♀♀, 5♂♂, *C. solstitialis* and some other Compositae; Süngübeyir, Olur, 1850 m, 20.VII.1994, 1♀, 3♂♂; 28.VII.1993, 2♂♂, *C. solstitialis*, and some other Compositae; Turnalı, Şenkaya (Erzurum), 1700 m, 6.VIII.1988, 1♀; Sarıkamış (Kars), 2100 m, 13.VII.1996, 7♀♀, 3♂♂, *C. solstitialis* and some other Compositae.

Distribution in Turkey: Kars (Tomov and Gruev, 1975), Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Europe, Anatolia, Siberia (Clavareau, 1913), Austria (Mohr, 1966), Bulgaria, Poland, Crimea, Venezuela, Albania, Greece, Yugoslavia (Warchalowski, 1976), European part of the USSR, Caucasus, Siberia, Kazakhstan, Central and Southern Europe (Lopatin, 1977).

*Cryptocephalus signatifrons* Suffrian, 1847

Material examined: Sutkans, Oltu (Erzurum), 10.VI.1997, 3♀♀, 2♂♂.

Distribution in Turkey: Taşdelen (Malatya) (Medvedev, 1970).

Distribution in the world: Bulgaria, Venezuela, Albania, Yugoslavia (Warchalowski, 1976).

*Cryptocephalus sultani* Pic, 1920

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976). It is an endemic specimens in Turkey.

*Cryptocephalus tappesi* Marseul, 1868

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Lebanon, Syria (Clavareau, 1913).

*Cryptocephalus tappesi* var. *disconiger* Pic, 1904

Distribution in Turkey: Tarsus (Clavareau, 1913). It is an endemic subspecies in Turkey.

*Cryptocephalus trapezensis* Tappes, 1871

Material examined: Rabat, Pasinler (Erzurum), 2400 m, 11.VI.1996, 2♀♀, 2♂♂; 13.VI.1996, 1♀, 2♂♂.

Distribution in Turkey: Trabzon (Clavareau, 1913), Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Turkey (Clavareau, 1913), Caucasus, Azerbaijan, Armenia (Warchalowski, 1976),

*Cryptocephalus trimaculatus* Rossi, 1790

Material examined: Şelale, Uzungöl (Erzurum), 1300 m, 9.VI.1996, 1♂; Paşalı, Şenkaya (Erzurum), 1300 m, 12.VI.1996, 1♀; Demirkent, Yusufeli (Artvin), 900 m, 6.VII.1994, 1♂; Ferhatlı, Ardanuç (Artvin), 1100 m, 7.VII.1994, 1♀; Anlı, Tortum (Erzurum), 1600 m, 24.VIII.1993, 1♀; Çamlıbel, Oltu (Erzurum), 1700 m, 14.VIII.1996, 2♂♂; Paşalı, Şenkaya (Erzurum), 1600 m, 12.VI.1996, 1♂♂; Şellale, Uzungöl (Erzurum), 1800 m, 9.VI.1996, 1♂.

Distribution in Turkey: Amasya, Ankara, Gümüşhane, Bayburt, İçel (Tomov and Gruev, 1975), Turkey, no locality (Warchalowski, 1976).

Distribution in the world: South Europe, Germany, Siberia (Clavareau, 1913), Mid Europe, Slovakia (Mohr, 1966), Bulgaria, Venezuela, Albania, Greece, Yugoslavia (Warchałowski, 1976), Europe, Asia, Caucasus (Gruev and Tomov, 1984).

*Cryptocephalus turcicus* Suffrian, 1847

Distribution in Turkey: Eskişehir (Bodameyer, 1900), Turkey (Warchałowski, 1976), Kaynarca (İznik), Soğuksu Millet Parkı, 1000m, Mataracı (Medvedev, 1970), Bilecik (Tomov and Gruev, 1975), Sındırğı (Balıkesir), Çanakkale, Armutlu, Gemlik (Bursa), Bahçeköy, Belgrat Ormanları (İstanbul), *Quercus petraea*, *Q. robur* and *Q. ilex* (Selmi, 1982).

Distribution in the world: Dalmatica, Turkey, Greece, Corsica (Clavareau, 1913), Mid Europe (Mohr, 1966), Sout France, Italia, Austria, Scilia, Greece (Burlini, 1955), Bulgaria, Venezuela, Albania, Greece, Yugoslavia (Warchałowski, 1976), Europe, Crentral Asia, Bulgaria, Greece (Gruev and Tomov, 1984).

*Cryptocephalus variceps* Weise, 1884

Material examined: Gölbaşı Yaylası, Uzundere (Erzurum), 2000 m, 16.VI.1995, 3♀, 2♂♂.

Distribution in Turkey: Turkey, no locality (Clavareau, 1913) Turkey, no locality (Warchałowski, 1976). It is an endemic species in Turkey.

*Cryptocephalus violaceus* Laicharting, 1781

Material examined: Sütkans, Oltu (Erzurum), 2000 m, 25.VI.1996, 2♀, 3♂♂, *Centaurea solstitialis* and some other Compositae; Dikyar, Uzundere (Erzurum), 1450 m, 10.VII.1994, 1♀, *Centaurea solstitialis*; Sarıgöl, Yusufeli (Artvin), 1100 m, 5.VII.1994, 1♂, Compositae; Kireçli, Tortum (Erzurum), 1700 m, 8.VIII.1995, 2♂♂, Compositae; Belkiz (Antalya), 100 m, 3.VIII.1994, 1♂.

Distribution in Turkey: Ordu (Tuatay et al., 1972), Turkey, no locality (Warchałowski, 1976).

Distribution in the world: Europe (Clavareau, 1913), South and Mid Europe (Mohr, 1966), Bulgaria, Greece, Poland, Venezuela, Albania, Greece, Yugoslavia, Caucasus, Azerbaijan, Armania (Warchałowski, 1976), Mid and North Europe, Greece (Gruev and Tomov, 1984).

*Cryptocephalus vittula* Suffrian, 1848

Material examined: İlica, Erzurum, 14.VI.1988, 1♂; Dinar (Afyon), 14.VIII.1990, 1♀, 2♂♂.

Distribution in Turkey: Turkey, no locality (Warchałowski, 1976).

Distribution in the world: Bulgaria, Venezuela, Albania, Greece, Yugoslavia, Caucasus, Azerbaijan, Armenia (Warchalowski, 1976),

***Cryptocephalus wehnckeii* Weise, 1882**

Distribution in Turkey: Tarsus (İçel) (Clavareau, 1913), Turkey, no locality (Warchalowski, 1976). It is an endemic species in Turkey

**Tribe: PACHYBRACHINI**

**Genus: *Pachybrachis* Chevrolat, 1837**

***Pachybrachis adaliensis* Weise, 1886**

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Anatolia (Clavareau, 1913), Greece (Warchalowski, 1976),

***Pachybrachis bodemeyeri* Weise, 1906**

Material examined: Köşk Köyü, Erzurum, 2000 m, 20.VI.1971, 9♂♂, 5♀♀, 20.VI.1996, 16♀♀, 20♂♂, *Salix* spp..

Distribution in Turkey: Anatolia (Clavareau, 1913), Turkey, no locality (Warchalowski, 1976). It is an endemic species in Turkey.

***Pachybrachis excisus* Weise, 1897**

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976), Elmalı (Gruev and Tomov, 1979).

Distribution in the world: Anatolia (Clavareau, 1913), Asia, Europe (Gruev and Tomov, 1984).

***Pachybrachis fimbriolatus* Suffrian, 1848**

Material examined: Very common in Bayburt, Pasinler (Erzurum), İğdır and Kağızman (Kars), Papilionacea; Gölbaşı Yayla, Uzundere (Erzurum), 2000 m, 16.VI.1995, 2♀♀, 5♂♂, Papilionacea.

Distribution in Turkey: Sabanözü (Çankırı), Kızılıcahamam (Ankara) (Gruev and Tomov, 1979).

Distribution in the world: South Europe, Siberia, Germany, France (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Crimea, Albania, Greece, Yugoslavia (Warchalowski, 1976), North Caucasus and Kazakhstan (Yaroshenko, 1985)

***Pachybrachis flexuosus* Weise, 1882**

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Sicilia, Greece, Austria (Clavareau, 1913), Bulgaria, Venezuela, Albania, Greece, Yugoslavia (Warchalowski, 1976),

***Pachybrachis glycyrrhizae*** Olivier, 1808

Distribution in Turkey: İğdir, Kağızman (Kars) (Aslan and Özbek, 1997).

Distribution in the world: Iran (Clavareau, 1913), Azerbaijan, Armenia, Caucasus (Warchalowski, 1976), Trans-Caucasus, southern Turkmenia, Southern Uzbekistan, Southern Tadzhicistan, Iran, Iraq, Syria, Jordan and Afghanistan (Lopatin, 1977)

***Pachybrachis hieroglyphicus*** (Laicharting, 1781)

***Cryptocephalus hieroglyphicus*** Laicharting, 1781 (Gruev and Tomov, 1984)

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Mid Europe, Swidin, Siberia (Clavareau, 1913), Bulgaria, Poland, Venezuela, Albania, Yugoslavia, Greece (Warchalowski, 1976), Europe, Asia Minor, Kazakhstan (Gruev and Tomov, 1984).

***Pachybrachis instabilis*** Weise, 1887

Distribution in Turkey: Akyazı (Adapazarı), Sındırı (Balıkesir), Armutlu, Gemlik (Bursa), Demirköy (Kırklareli), *Quercus petraea* and *Q. robur* (Selmi, 1982).

Distribution in the world: Turkestan (Weise, 1887), Kyndyr-Tau (Clavareau, 1913).

***Pachybrachis limbatus*** (Ménétriés, 1836)

***Cryptocephalus limbatus*** Ménétriés, 1836 (Gruev and Tomov, 1984).

Distribution in Turkey: İçel (Tomov and Gruev, 1975), Turkey, no locality (Warchalowski, 1976), Kaynarca (İznik), Bolu ve Yeniçağa arası, Soğuksu Millet Parkı, 1000m, Kavak (Samsun), (Medvedev, 1970), Sındırı (Balıkesir), Armutlu, Gemlik (Bursa), Alemdağ (İstanbul), Kalkım (Balıkesir), Çanakkale, *Quercus cerris*, *Q. frainetto* and *Q. coccifera* (Selmi, 1982).

Distribution in the world: Turkey, Greece (Clavareau, 1913), Bulgaria, Venezuela, Albania, Yugoslavia (Warchalowski, 1976), Bulgaria, Greece, Central Asia (Gruev and Tomov, 1984).

***Pachybrachis mardinensis*** Weise, 1900

Distribution in Turkey: Mardin (Clavareau, 1913), Turkey, no locality (Warchalowski, 1976). It is an endemic species in Turkey.

***Pachybrachis mendax*** Suffrian, 1860

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Crimea, Azerbaijan, Armenia, Caucasus (Warchalowski, 1976).

***Pachybrachis misellus*** Weise, 1900

Distribution in Turkey: Mardin (Clavareau, 1913), Turkey, no locality (Warchalowski, 1976). It is an endemic species in Turkey.

***Pachybrachis oertzeni*** Weise, 1896

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Greece (Clavareau, 1913).

***Pachybrachis scriptidorsum*** Marseul, 1875

Material examined: Silvan (Diyarbakır), 1000 m, 17.IV.1995, 2♂♂, *Salix* sp.; Pasinler (Erzurum), 11.VI.1996, 1♀.

Distribution in Turkey: Gümüşhane (Tomov and Gruev, 1975).

Distribution in the world: South Russia, Caucasus, West Siberia (Clavareau, 1913), Bulgaria (Warchalowski, 1976), Europe, Central Asia, Kazakhstan (Gruev and Tomov, 1984), East Europe, North Kazakhstan, Crimea (Yarosenko, 1985).

***Pachybrachis sinuatus*** Mulsant, 1859

***Pachybrachis haliciensis*** Mill, 1868 (Claveareu, 1913).

Material examined: Dikyar, Uzundere (Erzurum), 1450m, 2.VII.1994, 2♂♂; 17.VI.1995, 16♀♀, 22♂♂; 30.VI.1996, 6♀♀, 10♂♂, *Salix* spp..

Distribution in Turkey: Soğuksu, Kızılcahamam (Ankara) (Medvedev, 1975), Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Germany, Hungary, South France (Clavareau, 1913), Mid Europe (Mohr, 1966), Bulgaria, Poland, Albania, Yugoslavia (Warchalowski, 1976), Europe, Asia, Bulgaria (Gruev and Tomov, 1984).

***Pachybrachis tesellatus*** (Olivier, 1791)

***Cryptocephalus tesellatus*** Olivier, 1791 (Gruev and Tomov, 1984).

Material examined: Very common in Erzurum, Pasinler (Erzurum) and Bayburt, *Quercus* sp.; Erzincan, 1300 m, 8.VII.1993, 3♀♀, 2♂♂, *Quercus* sp..

Distribution in Turkey: Turkey, no locality (Warchalowski, 1976).

Distribution in the world: Germany, France, Austria, South Europe (Clavereau, 1913) South and Mid Europe (Mohr, 1966), Bulgaria, Poland, Venezuela, Albania, Yugoslavia, Greece, Azerbaijan, Armenia, Caucasus (Warchałowski, 1976), Europe, Bulgaria (Gruev and Tomov, 1984), North Caucasus (Yaroshenko, 1985).

***Pachybrachis tesellatus orientalis* Weise, 1900**

Material examined: Dikyar, Uzundere (Erzurum), 1500 m, 7.VI.1995, 6♀♀, 7♂♂, *Quercus* sp.; Şelale, Tortum (Erzurum), 1400 m, 9.VI.1996, 3♀♀, 2♂♂, *Quercus* sp.; Çamlıbel, Oltu (Erzurum), 1750 m, 14.VII.1996, 7♀♀, 4♂♂, *Quercus* sp.; Madenköprübaşı, İspir (Erzurum), 1300 m, 7.VII.1996, 3♀♀, 4♂♂, *Quercus* sp.; Erzincan, 1300 m, 7.VII.1993, 3♀♀, 4♂♂, *Quercus* sp..

Distribution in Turkey: Kaynarca (İznik), Soğuksu Millî Parkı, 1000m (Medvedev, 1970), Amasya, Ankara, Gümüşhane, Bayburt (Tomov and Gruev, 1975).

Distribution in the world: Asia Minor (Gruev and Tomov, 1984). (It is an endemic subspecies in Turkey).

**Tribe: STYLOSOMINI**

**Genus: *Stylosomus* Suffiran, 1848**

***Stylosomus subelongatus* Pic, 1899**

Distribution in Turkey: Torul (Gümüşhane) (Aslan and Özbek, 1997). In an endemic species in Turkey.

***Stylosomus tamaricis* (Herrich-Schaffer, 1838)**

Material examined: Dutlu, Oltu (Erzurum), 1200 m, 27.VI.1996, 6♀♀, 7♂♂, *Tamarix* sp..

Distribution in Turkey: Aegean, West Black Sea, Mediterranean Region (Warchałowski, 1991).

Distribution in the world: All Europe (Warchałowski, 1991).

**Acknowledgements**

We would like to express our sincere thanks to Prof. Dr. V. Tomov (Plovdiv, Bulgaria) and Prof. Dr. A.Warchałowski (Wrocław, Poland) for determination of the reference material.

## LITERATURE

- Aslan, İ. and H. Özbek, 1997. Some new records of the subfamilies Cassidinae and Cryptocephalinae (Coleoptera, Chrysomelidae) for the Turkish fauna. *Zoology in the Middle East*, 14, 87-91.
- Berti, N. and M. Rapilly, 1973. Contribution a la faune de l'Iran voyages de M.M.R. Naviaux et M. Rapilly (Coleoptera, Chrysomelidae). *Ann. Soc. Entom. France*, 9(4): 861-894.
- Bodemeyer, H.E., 1900. Quer durch Klein-Asien in der Bulghar Dagh. Eine Naturwissenschaftliche Studienreise,. 169 p.
- Burlini, M., 1955. Revisione dei *Cryptocephalus* italiani e della maggior parte delle specie di Europe (Coleoptera, Chrysomelidae). *Mem. Soc. Ent. Ital.*, 34: 1-287.
- Clavareau, H., 1913. Coleopterorum Catalogus Chrysomelidae: 8 .Cryptocephalinae in: W. Junk, S.Schenkling. Berlin W. 15., Pars 53, 85-209.
- Gruev, B.A. and V.Tomov, 1979. Zur Kennetnis einiger in der Türkei, Jugoslawien und Griechenland Workommender Arten der Familie Chrysomelidae (Coleoptera) aus der Zoologischen Staatssammlung München. *Spixiana* 2(3): 259-267.
- Gruev, B. A. and V. Tomov, 1984. Fauna Bulgarica. 13, Coleoptera, Chrysomelidae, Part I, Orsodacninae, Zeugophorinae, Donaciinae, Cricerinae, Clytrinae, Cryptocephalinae, Lamprosamatinae, Eumolpinae. In Aedibus Academia Scientiarum Bulgaricae, 220 p.
- Gül-Zümreoğlu, S., 1972. İzmir Bölge Zirai Mücadele Araştırma Enstitüsü Böcek ve Genel Zararlılar Kataloğu, 1928-1969, 1.Kısim. T.C. Tarım Bakanlığı Zirai Mücadele ve Zirai Karantina Genel Müdürlüğü Yayınları, 48-52.
- Jolivet, P., E.Petitpierre and T.H.Hasiao, 1988. Biology of Chrysomelidae. Kluwer Academic Publishers, 615 p.
- Lopatin, I., 1977. Leaf-beetles (Chrysomelidae) of Middle Asia and Kazakhstan. Nauka, Leningrad, 268 p.
- Medvedev, L.N., 1970. A list of Chrysomelidae collected by Dr. W. Wittmer in Turkey (Coleoptera). *Rev. Suis. Zool.*, 77, 2 (22): 309-319.
- Medvedev, L.N., 1975. Chrysomelidae collected by Dr.W.Wittemer in Turkey and Iran. *Ent. Gaselsch. Basel*, 25(1):12-19.
- Mohr, K., 1966. Chrysomelidae In: H. Freude, K.W. Harde and G.A. Lohse (Goecke and Evers Ed.), Die Kater Miteuropas. Krefeld, 9: 95-299.
- Pic,M., 1908. Descrizioni di forme nuove e note varie. *Echange*, 24: 13-14.
- Pic, M., 1909. Descriptions, diagnoses et notes diverses. *Echange*, 25: 145-186.
- Roubal, J., 1937-41. Catalogue des Coleoptères de la Slovaquie et des Carpathes orientales. Tom. III, 363 p.
- Selmi, E., 1982. Marmara Bölgesi Ormanlarında Zarar yapan Chrysomelidae (Coleoptera) Türleri ve Bazı Önemli Türlerin Biyolojileri. İstanbul Üniversitesi Orman Fakültesi Yayınları, 122 s. (Unpublished theses).

- Tuatay, N., Kalkandelen A. ve Aysev N., 1972. Nebat Koruma Müzesi Kataloğu, 1961-1971. T.C. Tarım Bakanlığı Zirai Mücadele ve Zirai Karantina Genel Müdürlüğü Yayınları, 55-57.
- Tomov, V. and B.A.Gruiev, 1975. Chrysomelidae (Coleoptera) Collected by K.M. Guichard in Turkey, Greece and Yugoslavia. *Trav. Sci. Univ. Plovdiv, Bulgaria, Biol.*, **13 (4)**: 133-151.
- Warchałowski, A., 1976. Biogeographiscche Studien über die Blattkäfer der Pontischen Provinz (Coleoptera: Chrysomelidae). *Polskie Pismo Entomologiczne*, **46**: 29-94.
- Warchałowski, A., 1991. Chrysomelidae stonkowate (Insecta: Coleoptera). Fauna Polski, Tom 13. Polska Akademia Nauk Instytut Zoologii 247 p.
- Weise, J., 1887. 9. Beitrage zur Coleopteren Fauna von Turkestan von Major Z.D.Dr.L. von Heyden, unter Mitwirkung der Herrn Abaille de Perrin, Dr. Eppelsheim, Faust und Weise. *Zeitshrift*, **31**: 303-336.
- Weise, J., 1894. Chrysomeliden aus Amasia. *Zeitshrift*, **31**: 91-92
- Yaroshenko, V.A., 1985. Ecological-Faunistical Characteristic of the Flea-beetles (Chrysomelidae, Cryptocephalinae) of the northern Caucasus. *Ecologia, Zoological Büll., UDK*, **595. 768.12 (470.6)**, 52-55.