



The Effects of Anatolian Black Pine Cone Extracts on Carbonic Anhydrase Enzymes

Berfin VAPOR , Ekrem TUNCA * 

Kütahya Dumlupınar University, Faculty of Arts and Sciences, Department of Biochemistry, 43100 Kütahya, Türkiye

Highlights

- Anatolian black pine cone extracts were prepared using solvents of different polarities.
- The effects of the extracts on hCA I and hCA II activities were examined.
- The extracts had remarkable inhibition effects on hCA I and hCA II.

Article Info

Received: 09 Jan 2024
Accepted: 13 Aug 2024

Keywords

Anatolian black pine
Cone
Carbonic anhydrase
Esterase activity
Inhibition

Abstract

Nowadays, there is an increasing trend in the use of natural products to treat diseases or alleviate their effects. This situation has emerged as a result of the side effect problem seen in the use of synthetic drugs. In this study, Anatolian black pine cone extracts were prepared using solvents of different polarities (water, methanol, ethanol, ethyl acetate, *n*-hexane). Then, the effects of these extracts on the esterase activities of hCA I and hCA II isoenzymes, which are associated with many physiological disorders, were examined *in vitro*. IC₅₀ values were in the range of 5.10 to 49.50 µg/mL for hCA I and 2.57 to 15.07 µg/mL for hCA II. It can be said that these results are promising in terms of the potential of Anatolian black pine cone as a natural pharmacological agent.

1. INTRODUCTION

Meldrum and Roughton, who were investigating the transport of gases in the blood of vertebrates, discovered carbonic anhydrase (CA), which catalyzes the reversible hydrolysis of carbon dioxide (CO₂) and water (H₂O) to bicarbonate (HCO₃⁻) and proton (H⁺), in 1933 [1]. The reaction of two simple molecules (CO₂ and H₂O) produces a weak base (HCO₃⁻) and a strong acid (H⁺), which induces a pH change [2]. This system is used as an effective means of pH regulation by many organisms from Prokaryotes to Eukaryotes [3]. To date, eight genetically different CA families (α -, β -, γ -, δ -, ζ -, η -, θ - and ι -) have been defined [4-8]. Mammalian CAs, which belong to the class of α -CAs, have 16 different isoenzymes in mammals. As in all organisms encoding CA enzymes, human CA enzymes (hCAs) play a role not only in pH regulation but also respiration, CO₂ and HCO₃⁻ transport, electrolyte secretion, pH and CO₂ homeostasis, gluconeogenesis, adipogenesis, ureagenesis, bone resorption and calcification [3,9-11]. It is inevitable that CA isoenzymes, which are involved in many processes, are associated with multitude of diseases. Glaucoma, retinal and cerebral edema, epilepsy, obesity, hypoxic tumors, inflammation, neuropathic pain, Alzheimer, oxidative stress and vertigo are diseases associated with hCA isoenzymes [12-14]. Interfering with the activities of hCAs using modulators (activators or inhibitors) is pharmacologically important for the management of CA-related diseases. Despite tremendous strides over the past decade to understand the molecular interactions associated with CAs, the development of selective CA inhibitors (CAIs) is still an ongoing process. Over the past years, various research groups have worked to design and synthesize new CAIs effective against a variety of pathological conditions in which the activity of these enzymes is dysregulated. Sulfonamides, phenols, sulfamates, sulfamides, coumarins, polyamines, dithiocarbamates,

*Corresponding author, e-mail: ekrem.tunca@dpu.edu.tr

xanthates, benzoxaboroles, phosphoramidates, selenols, and ninhydrins are classes of CAIs discovered from 1940 to 2020 [9]. Compounds of completely natural origin as well as synthetic or natural-synthetic hybrid CAIs have been studied as CAIs. In the literature, there are studies in which CAIs of natural origin are used directly as extracts and the compounds obtained from the extract by chromatographic techniques are examined as CAIs. Some plants in which hCA inhibition effects were investigated as total extracts can be listed as *Foeniculum vulgare* Mill., *Alcea rosea* L., *Laurus azorica* (Seub.) Franco, *Elettaria cardamomum* L., *Lavandula stoechas* L. [15], *Artemisia dracuncululus* L. [16], *Leucas cephalotes* (Roth) Spreng. [17], and *Abrus precatorius* L [18].

Anatolian black pine (*Pinus nigra* Arnold.) is distributed in the Balkans, Southern Carpathians, Crimea, Turkey, Cyprus and Syria. It has a smooth cylindrical body that can reach up to 40 m in length and exceed 1 m in diameter. Needle leaves are 4–8 cm long, dark green and hard. The ovoid-conical cones are symmetrical and have almost non-existent short stems. Cone length varies between 3.5 cm – 10 cm [19]. The production of pine cone extracts, whose medicinal and protective properties have been discovered, has gained importance in recent years. It has been reported that some pine cone extracts have toxic activities against fungi. It has also been observed that essential oils obtained from pine cones protect the plant against fungal and insect attacks. The cones, leaves and resins of the *Pinaceae* family are believed to have healing effects among the public and are used in the treatment of various disorders [20]. In East Asia, cones, cortex, needles, and pollen, which are various parts of pine trees, are widely consumed as dietary supplements to support immune system. In these countries, it is believed that consumption of pine tree fractions positively affects neuronal and gastrointestinal problems and prevents diabetes, hypertension and atherosclerosis. In studies conducted on the use of antioxidant components found in the structure of pine bark and cones in foods, it has been proven that the present components prevent protein oxidation at rates ranging from 42% to 64%. In traditional medicine, pine cones have been used to relieve cough, moisten the lungs and reduce fever [21,22].

In last two decades, there has been an increase in the use of natural compounds/extracts instead of synthetic compounds for the treatment or prevention of diseases, as they have fewer possible side effects. In this context, various groups have studied the CA inhibition effects of some of the extracts mentioned above. In this study, extracts of pine cones were prepared using solvents of different polarities (water, methanol, ethanol, ethyl acetate and hexane). The inhibitory effects of the prepared extracts on hCA I and hCA II isoenzymes were determined as *in vitro*.

2. MATERIAL METHOD

The pine cones used in the study were collected from Anatolian black pine trees in Kütahya Dumlupınar University Evliya Çelebi Campus (Türkiye). Extraction solvents (methanol, ethanol, ethyl acetate and *n*-hexane) were purchased from Fischer Scientific International Inc. (New Hampshire, USA). TRIS and H₂SO₄ used to prepare buffer solution were purchased from Merck (Darmstadt, Germany). 4-nitrophenyl acetate, which was used as a substrate in kinetic measurements, was purchased from Sigma-Aldrich (Missouri, USA). DMSO, which was used for preparation of stock inhibitor solutions, was purchased from Fischer Scientific International Inc. (New Hampshire, USA). Enzyme activity measurements were performed with SHIMADZU UV-1700 PharmaSpec UV-Vis spectrophotometer (Kyoto, Japan). Buffer solutions were prepared with the pH meter (Mettler Toledo SevenDirect SD-20 Kit, Ohio, USA). Grinding operations were carried out using an IKA A11 basic grinder (Staufen, Germany).

2.1. Preparation of Pinecone Extracts

The unopened cones collected from *Pinus nigra* Arnold. were kept in a sunless environment at room temperature for a few days. 50 mL of solvent (water, methanol, ethanol, ethyl acetate and *n*-hexane) was added to 5 g of the sample taken from the milled pinecone and mixed for 24 h at 25 °C. The suspension was filtered and the filtrate was centrifuged at 10⁵ g for 20 min to obtain clear solution. Then the supernatant was concentrated at low temperature and low pressure in a rotary evaporator. The solid residue was weighed and stored in -20 °C until used [15,16,23,24].

catalysis mechanism of these isoenzymes, approximately half of the active site consists of hydrophobic amino acid residues, while the other half consists of hydrophilic amino acid residues. Therefore, since substances of different polarities are obtained in extraction processes with solvents of different polarities, the possibility of obtaining extracts that can inhibit hCA isoenzymes has increased.

Table 1. Amount of the extracts obtained from larch pinecones

Solvent	Extract amount (g)
Water	0.82
Methanol	0.61
Ethanol	0.42
Ethyl acetate	0.27
<i>n</i> -Hexane	0.13

The best extraction efficiency was obtained in the presence of water, which is the most polar solvent. As expected, this is due to the polyphenols present in large amounts in pinecones. As the polarity decreased, the amount of extract obtained also decreased (Table 1). This shows that the oily substances in the pinecone are in very small amounts. The reason why DMSO was chosen as the extract solvent for inhibition studies is that it can dissolve all extracts and does not inhibit or activate CA enzymes.

Table 2. Esterase IC₅₀ values of the *Pinus nigra* pinecone extracts for hCA isoenzymes

Extracts	IC ₅₀ (µg.mL ⁻¹)	
	hCA I	hCA II
Acetazolamide ^a	1.652±0.033	0.016±0.0001
Water	11.18±0.56	7.37±0.37
Methanol	5.10±0.26	2.57±0.13
Ethanol	12.60±0.63	10.19±0.51
Ethyl acetate	30.13±1.51	11.36±0.57
<i>n</i> -Hexane	49.50±2.48	15.07±0.75

^aControl compound, and the IC₅₀ values of this compound were taken from [16]

Among the extracts used in the study, methanolic extract had the strongest inhibitory effect (hCA I IC₅₀ value is 5.10 µg/mL, and hCA II IC₅₀ value is 2.57 µg/mL) (Figure 2). However, it is important for a pharmacological agent to be selective among CA isoenzymes as well as being a strong CA inhibitor. When the IC₅₀ values of the extracts are examined, it is seen that all of them are relatively more selective against the hCA II isoenzyme (Table 2).




Figure 2. Enzyme activity vs. inhibitor concentration graphs of methanolic extract (A: hCA I, B: hCA II)

When the amino acid sequences in active sites of hCA I and hCA II are compared, it is seen that there are differences mostly between the amino acid residues in the hydrophobic part (Table 3) [34]. It is possible, then, that these differences enable the extracts to interact more tightly with hCA II. The fact that the extract prepared using *n*-hexane inhibited the hCA II isoenzyme approximately 3.5 times more strongly than hCA I indicates that the selectivity may be due to the hydrophobic part of the isoenzymes. However, to make a definitive comment, the compounds in the extract should be determined, and molecular docking studies and even X-ray diffraction analyses should be performed (Figure 3).

Table 3. Hydrophobic amino acid residues in the active sites of hCA I and hCA II*

Residue	hCA I	hCA II
91	Phe	Ile
121	Ala	Val
131	Leu	Phe
135	Ala	Val
141	Leu	Leu
143	Val	Val
204	Tyr	Leu
207	Val	Val
209	Trp	Trp

* Data was taken from [34]




Figure 3. IC₅₀ values and isoenzyme selectivities according to extract polarity ($E_T(30)$ values were taken from [35])

Some studies in the literature on the subject and the results obtained are explained below. Kaya et al. (2019) prepared aqueous, ethanolic and methanolic extracts of *Foeniculum vulgare*, *Elettaria cardamomum*, *Alcea rosea*, *Lavandula stoechas* and *Laurus azorica* plants and examined their inhibition effects on hCA I and hCA II in terms of esterase activity inhibition. They determined that methanolic extracts of all plants were the most effective CA inhibitors [15]. In a study in which dichloromethane, *n*-hexane, methanol and ethanol extracts of *Artemisia dracunculus* L. leaves were prepared and their inhibition effects on hCA I and hCA II esterase activity were examined, it was determined that dichloromethane extract was the strongest inhibitor (hCA I IC₅₀ value was 0.020 µg/mL) [16]. In another paper, methanol and oil extracts of *Cucumis melo* L. seeds were reported to show interesting biological activities for human erythrocyte CAs. While these extracts activated the hCA I, they inhibited the hCA II isoform. hCA II IC₅₀ values were determined as 0.497 ng/mL for the oil extract and 10.98 µg/mL for the methanol extract [36]. Aydin et al. (2021) prepared extracts of the *Satureja cuneifolia* plant with petroleum ether, chloroform and methanol, examined their effects on hCA I and II esterase activities, and found that the methanol extract showed the strongest inhibitory effect. Researchers had determined the hCA I esterase IC₅₀ value of the direct methanol extract as 31 µg/mL, and the hCA II IC₅₀ value of the methanol extract as 12 µg/mL [37].

The IC₅₀ values determined in our study were in the range of 5.10 – 49.50 µg/mL for hCA I and 2.57 – 15.07 µg/mL for hCA II (Table 2) and were comparable to other studies in the literature. Additionally,

similar to some studies in the literature, the strongest inhibitory effect was observed in the methanolic extract in our study.

4. CONCLUSION

In recent years, the use of natural compounds both as food supplements and for therapeutic purposes has become widespread. Intensive efforts are being made to discover therapeutic agents of natural origin. Within the scope of this study, the effects of Anatolian black pine cone extracts, whose many benefits are known to the public and reported in the literature, on hCA I and hCA II isoenzymes, which are known to be associated with various disorders, were examined *in vitro*. While methanolic extract showed the strongest inhibitory effect among the extracts, *n*-hexane extract showed the most selective inhibitory effect on hCA II isoenzyme. These promising results should be supported by further studies including *in vivo* tests and toxicity tests.

ACKNOWLEDGEMENTS

This study was supported by TUBITAK (grant number 1919B012105233).

CONFLICTS OF INTEREST

No conflict of interest was declared by the authors.

REFERENCES

- [1] Meldrum, R.U., Roughton, F.J., "Carbonic anhydrase. Its preparation and properties", The Journal of Physiology, 80: 113-142, (1933).
- [2] Krebs, H.A., Roughton, F.J., "Carbonic anhydrase as a tool in studying the mechanism of reactions involving H_2CO_3 , CO_2 or HCO_3^- ", Biochemical Journal, 43: 550-555, (1948).
- [3] Supuran, C.T., "Carbonic anhydrases: novel therapeutic applications for inhibitors and activators", Nature Reviews Drug Discovery, 7: 168-181, (2008).
- [4] Capasso, C., Supuran, C.T., "An overview of the alpha-, beta-, and gamma-carbonic anhydrases from Bacteria: can bacterial carbonic anhydrases shed new light on evolution of bacteria?", Journal of Enzyme Inhibition and Medicinal Chemistry, 30: 325-332, (2015).
- [5] Bertucci, A., Moya, A., Tambutté, S., Allemand, D., Supuran, C.T., Zoccola, D., "Carbonic anhydrases in anthozoan corals – A review", Bioorganic & Medicinal Chemistry, 21: 1437-1450, (2013).
- [6] Del Prete, S., Vullo, D., Fisher, G.M., Andrews, K.T., Poulsen, S.A., Capasso, C., Supuran, C.T., "Discovery of a new family of carbonic anhydrases in the malaria pathogen *Plasmodium falciparum* – The η -carbonic anhydrases", Bioorganic & Medicinal Chemistry Letters, 24: 4389-4396, (2014).
- [7] Kikutani, S., Nakajima, K., Nagasato, J., Tsuji, Y., Miyatake, A., Matsuda, Y., "Thylakoid luminal theta-carbonic anhydrase critical for growth and photosynthesis in the marine diatom *Phaeodactylum tricorinitum*", PNAS, 113: 9828-9833, (2016).
- [8] Nocentini, A., Supuran, C.T., Capasso, C., "An overview on the recently discovered iota-carbonic anhydrases", Journal of Enzyme Inhibition and Medicinal Chemistry, 36: 1988-1995, (2021).
- [9] Supuran, C.T., "A simple yet multifaceted 90 years old, evergreen enzyme: Carbonic anhydrase, its inhibition and activation", Bioorganic & Medicinal Chemistry Letters, 93: 129411, (2023).

- [10] Aspatwar, A., Supuran, C.T., Waheed, A., Sly, W.S., Parkkila, S., "Mitochondrial carbonic anhydrase VA and VB: properties and roles in health and disease", *The Journal of Physiology*, 601: 257-274, (2023).
- [11] Lynch, C.J., Fox, H., Hazen, S.A., Stanley, B.A., Dogson, S., Lanoue, K.F., "Role of hepatic carbonic anhydrase in de novo lipogenesis", *Biochemical Journal*, 310: 197-202, (1995).
- [12] Alterio, V., Di Fiore, A., D'Ambrosio, K., Supuran, C.T., De Simone, G., "Multiple binding modes of inhibitors to carbonic anhydrases: How to design specific drugs targeting 15 different isoforms?", *Chemical Reviews*, 112: 4421-4468, (2012).
- [13] Jang, B.G., Yun, S.M., Ahn, K., Song, J.H., Jo, S.A., Kim, Y.Y., Kim, D.K., Park, M.H., Han, C., Koh, Y.H., "Plasma carbonic anhydrase II protein is elevated in Alzheimer's disease", *Journal of Alzheimer's Disease*, 21: 939-945, (2010).
- [14] Sluch, I.M., Elliott, M.S., Dvorak, J., Ding, K., Farris, B.K., "Acetazolamide: A new treatment for visual vertigo", *Neuro-Ophthalmology*, 41: 315-320, (2017).
- [15] Kaya, E.D., Erğun, B., Demir, Y., Alım, Z., Beydemir, Ş., "The *in vitro* impacts of some plant extracts on carbonic anhydrase I, II and paraoxonase-1", *Hacettepe Journal of Biology and Chemistry*, 47: 51-59, (2019).
- [16] Aydın, T., Yurtvermez, B., Şentürk, M., Kazaz, C., Çakır, A., "Inhibitory effects of metabolites isolated from *Artemisia drancunculus* L. against the human carbonic anhydrase I (hCA I) and II (hCA II)", *Records of Natural Products*, 13: 216-225, (2019).
- [17] Sethi, K.K., Verma, S.M., Kumar, P.M., Mishra, R., Supuran, C.T., "Carbonic anhydrase I and II inhibition with natural products: *Leucas cephalotes*", *Pharmacognosy Communications*, 1: 41-46, (2011).
- [18] Sethi, K.K., Suresh, P., Mondal, S., "Carbonic anhydrase I and II inhibition with natural products: *Abrus precatorius*", *Current Enzyme Inhibition*, 7: 119-124, (2011).
- [19] Hazer, O., Akbbik, M., Güngör, K., "Simultaneous determination of the main organic acids in Anatolian black pine by HPLC with DAD detector", *Eurasian Journal of Analytical Chemistry*, 11: 181-195, (2016).
- [20] Said, O., Aksut, Y., Yardıncı, R.E., Karatas, S., Yigit, H., Tel, A.Z., Arda, N., "Antioxidant and antibacterial activities of methanol extracts from various plant parts of pomegranate and Anatolian black pine", *Current Perspectives on Medicinal and Aromatic Plants*, 5: 50-62, (2022).
- [21] Iglesias, J., Pazos, M., Lois, S., Medina, I., "Contribution of galloylation and polymerization to the antioxidant activity of polyphenols in fish lipid systems", *Journal of Agricultural and Food Chemistry*, 58: 7423-7431, (2010).
- [22] Frontela-Saseta, C., Lopez-Nicolas, R., Gonzalez-Bermudez, C.A., Peso-Echarri, P., Ros-Berruezo, G., Martinez-Gracia, C., Virgili, F., "Evaluation of antioxidant activity and antiproliferative effect of fruit juices enriched with Pycnogenol® in colon carcinoma cells. The effect of in vitro gastrointestinal digestion", *Phytotherapy Research*, 25: 1870-1875, (2011).
- [23] Dilek, E., Erol, H.S., Cakir, A., Koc, M., Halici, M.B., "Natural product inhibitors of carbonic anhydrase I and II isoenzymes: osajin and pomiferin", *Archives of Physiology and Biochemistry*, 123: 219-224, (2017).

- [24] Demir, T., Gunesli, Z., Sonmez, F., Bilen, C., Yavuz, E., Gencer, N., "Inhibition of carbonic anhydrase I and II with total anthocyanins extracted from sweet cherry cultivars", *Environmental Engineering and Management Journal*, 14: 935-941, (2015).
- [25] İlkimen, H., Yenikaya, C., Sarı, M., Bülbül, M., Tunca, E., Süzen, Y., "Synthesis and characterization of a proton transfer salt between dipicolinic acid and 2-amino-6-methylbenzothiazole and its complexes, and their inhibition studies on carbonic anhydrase isoenzymes", *Polyhedron*, 61: 56-64, (2013).
- [26] İlkimen, H., Yenikaya, C., Sarı, M., Bülbül, M., Tunca, E., Dal, H., Baş, M., "Synthesis and characterization of complexes of a novel proton transfer salt and their inhibition studies on carbonic anhydrase isoenzymes", *Journal of Enzyme Inhibition and Medicinal Chemistry*, 30: 195-203, (2015).
- [27] Al-Dhrub, A.H.A., Sahin, S., Ozmen, I., Tunca, E., Bulbul, M., "Immobilization and characterization of human carbonic anhydrase I on amine functionalized magnetic nanoparticles", *Process Biochemistry*, 57: 95-104, (2017).
- [28] Tunca, E., Bülbül, M., İlkimen, H., Saygılı Canlıdınç, R., Yenikaya, C., "Investigation of the effects of the proton transfer salts of 2-aminopyridine derivatives with 5-sulfosalicylic acid and their Cu(II) complexes on cancer-related carbonic anhydrases: CA IX and CA XII", *Chemical Papers*, 74: 2365-2374, (2020).
- [29] Okumuş, A., Elmas, G., Binici, A., Tunca, E., Hökelek, T., Kılıç, Z., "Phosphorus-nitrogen compounds: part 70. Syntheses of tetraaminomono/bis(4-fluorobenzyl)spiro(N/N)cyclotriphosphazanes: structural characterization, Hirshfeld surface analysis and comparative evaluation of esterase activities of hCA I and hCA II isoenzymes", *New Journal of Chemistry*, 47: 8578-8588, (2023).
- [30] Verpoorte, J.A., Mehta, S., Edsall, J.T., "Esterase activities of human carbonic anhydrases B and C", *Journal of Biological Chemistry*, 242: 4221-4229, (1967).
- [31] İlkimen, H., Yenikaya, C., Sarı, M., Bülbül, M., Tunca, E., Dal, H., "Synthesis and characterization of a proton transfer salt between 2,6-pyridinedicarboxylic acid and 2-aminobenzothiazole, and its complexes and their inhibition studies on carbonic anhydrase isoenzymes", *Journal of Enzyme Inhibition and Medicinal Chemistry*, 29: 353-361, (2014).
- [32] Karaçam, H., Tunca E., Kaygısız, Y., İmdat, G., Bülbül, M., "Investigation of the effects of some edible mushroom extracts on human carbonic anhydrase isoenzymes", *Hacettepe Journal of Biology and Chemistry*, 43: 187-193, (2015).
- [33] Alkan Alkaya, Z., İlkimen, H., Yenikaya, C., Tunca, E., Bülbül, M., Tunç, T., Sarı, M., "Synthesis and characterization of Cu(II) complexes of 2-amino-6-sulfamoylbenzothiazole and their inhibition studies on carbonic anhydrase isoenzymes", *Polyhedron*, 151: 199-205, (2018).
- [34] Bhatt, A., Mahon, B.P., Cruzeiro, V.W.D., Cornelio, B., Laronze-Cochard, M., Ceruso, M., Sapi, J., Rance, G.A., Khlobystov, A.N., Fontana, A., Roitberg, A., Supuran, C.T., McKenna, R., "Structure-activity relationships of benzenesulfonamide-based inhibitors towards carbonic anhydrase isoform specificity", *ChemBioChem*, 18: 213-222, (2017).
- [35] Marcus, Y., "The effectiveness of solvents as hydrogen bond donors", *Journal of Solution Chemistry*, 20: 929-944, (1991).

- [36] Akkemik, E., Aybek, A., Felek, I., “Effects of cefan melon (*Cucumis melo*) seed extracts on human erythrocyte carbonic anhydrase I-II enzymes”, Applied Ecology and Environmental Research, 17: 14699-14713, (2019).
- [37] Aydın, F.G., Türkoğlu, E.A., Kuzu, M., Taşkın, T., “*In vitro* carbonic anhydrase inhibitory effects of the extracts of *Satureja cuneifolia*”, Turkish Journal of Agricultural and Natural Sciences, 8: 1146-1150, (2021).

EARLY VIEW