

Growth and Immune Responses, Oxidative Stress Biomarkers, and Antioxidative Enzymes of Broilers Fed with Supplementation of *Chenopodium ambrosioides* L. and *Crassocephalum crepidioides* leaf meals

Andrew Bamidele FALOWO^{1,*} 

¹Adekunle Ajasin University, Faculty of Agriculture, Department of Animal Science, Akungba-Akoko, Nigeria.

Article History

Received: 08 June 2023

Accepted: 20 December 2023

Corresponding Author

E-mail: andrew.falowo@aaua.edu.ng

Anddele2013@gmail.com

Tel: +2348065320285

Keywords

Broiler

Immune system

Leaf meal

Supplement

Stress biomarkers

Abstract

This study assessed the effect of dietary supplementation of *Crassocephalum crepidioides* (CCLM) and *Chenopodium ambrosioides* L. (CALM) leaf meals on relative growth rate, oxidative stress biomarkers, immune response, serum and meat antioxidative enzymes of broilers. A total of 120 one-day-old Cobb 500 chicks were randomly assigned to four dietary treatments (three replicates of 10 birds/treatment) as follows, Diet 1 (basal/control diet), Diet 2 (basal +200mg/kgVitC), Diet 3 (basal +2500mg/kgCCLM) and Diet 4 (basal +2500mg/kgCALM) for 42 days. The result revealed that supplementation of CCLM and CALM significantly reduced ($P < 0.05$) the concentration of heat shock protein and hydroxy-2-deoxyguanosine in broilers compared to the control group. Broilers fed diet containing CCLM and CALM exhibited higher ($P < 0.05$) serum glutathione peroxidase content compared to the control group. The level of immunoglobulins were significantly higher in birds fed diet containing CCLM, CALM and Vit C compared to control group. However, supplementation of CCLM and CALM did not influence ($P > 0.05$) the relative growth rate and antioxidant content of breast muscle compared to control group. It's concluded that CCLM and CALM could be used as natural additive to alleviate oxidative damage, improve immune system and serum antioxidant content of broilers.

Introduction

Commercial poultry is often subject to a range of stressors, from environmental to nutritional factors, that could impair performance during growing period, despite advances in management, genetic selection, and biosecurity [1]. In terms of nutrition, poultry production requires a balanced diet to support the immune system and inhibit cell damage. However, stresses that occur due to nutritional imbalances or deficiency may induce the production of free radicals, which in turn could weaken the immune system, create health problems, and result in significant production-related financial losses [1-3]. In order to avoid this, the inclusion of natural antioxidants in diets has been encouraged [4-5]. This is because natural antioxidants contain inherent bioactive compounds that can

activate the production of immunoglobulin [6], increase endogenous antioxidants [7], and inhibit the oxidative process that is caused by an imbalance between the production of reactive oxygen species (ROS) and endogenous antioxidant defense mechanisms in the body [4,8]. In addition, studies have shown that natural antioxidants can be used to improve feed intake, enhance body weight gain, and improve meat quality and the overall performance of poultry during production [7, 9, 10].

Interestingly, most natural antioxidants are obtained from plant materials, such as herbs (flowers, leave, fruits, seed, stem, root, etc.), vegetables, and spices, as well as their extracts such as essential oils [4, 11]. Its utilization has been encouraged in animal diets because they are easily affordable, accessible and safe when used at low concentrations compared to

antibiotic growth promoters [12]. Natural antioxidants possessed enormous phytochemicals such as phenols, flavonoid, alkaloid, tannin, saponin, and other bioactive compounds. *Chenopodium ambrosioides* L. and *Crassocephalum crepidioides* plants are among the natural antioxidants with a lot of potential that can be utilized as feed additives in poultry production.

Chenopodium ambrosioides L. is an aromatic herbaceous medicinal/vegetable plant that belongs to the family of Chenopodiaceae and the genus *Chenopodium*. The plant is extensively found in West Africa, particularly in Nigeria, and is generally known as Mexican tea [13]. In traditional medicine, the leaf of the plant is used to treat uterine fibroids as well as bacterial, fungal, parasitic, and viral diseases [14-15]. According to research by Maldonado-Garcia *et al.* [16] and Reyes-Becerril *et al.* [17], fish fed a diet supplemented with *C. ambrosioides* L. at 0.5, 1.0, and 2.0% (w/w) had stronger immune cell and antioxidant enzyme activities than fish without the supplements. *Crassocephalum crepidioides* is a succulent annual leafy vegetable and herb that belongs to the family Asteraceae (Compositae). It is widely distributed in tropical and subtropical regions of the world, including Africa, Asia and Australia [18]. Every part of a plant is known to have abundant secondary metabolites with the ability to elicit antibacterial, hypoglycemic, and antioxidant actions [18-19]. According to a preliminary investigation by Falowo *et al.* [19], both *Chenopodium ambrosioides* L. and *Crassocephalum crepidioides* plants are readily available and their leaves possessed moderate nutritional content, high bioactive compounds and antioxidant properties, which qualifies them as potential natural additives. In fact, the dietary supplementation of leaf meal of *Crassocephalum crepidioides* plant has been reported to improve the performance, immune systems and meat quality of broiler chicken [20]. However, to our knowledge the dietary utilization of *Chenopodium ambrosioides* L. leaf meal as natural additives on boiler performance has not been studied. Therefore, this study was designed to examine the effect of *Chenopodium ambrosioides* L. and *Crassocephalum crepidioides* leaf meals on oxidative stress biomarkers, immune response, serum and meat antioxidative enzymes of broiler chickens.

Materials and Methods

Plant collection and extract preparation

Freshly harvested leaves of *Chenopodium ambrosioides* L. and *Crassocephalum crepidioides* were purchased from the commercial markets in southwestern Nigeria. The leaves were cleaned and air-dried in an open shade. The dried leaves were ground using an electric blending machine and the powdered samples were packed into a black polyethylene bag for further analysis.

Experimental site

The study was conducted at the Poultry Unit of the Teaching and Research Farm, Federal College of Agriculture, Akure, Nigeria. The experimental site is located at 7°25' N and 5°19' E with average annual temperature and annual rainfall of 25.3°C and 1455 mm, respectively. The entire study was carried out for six weeks following the research ethics and guidelines of the Animal Health and Production Technology Department of the institution.

Experimental diets and animals

Two basal diets [starter (0-21 days) and finisher phase (22-42days)] were formulated to meet the broiler's nutritional requirement as recommended by NRC [21] (Table 1). At each phase, the experimental diets were divided into four treatments and designated as Diet 1 (basal/control diet), Diet 2 (basal + 200mg/kg Vitamin C as positive control), Diet 3 (basal + 2500mg/kg CALM) and Diet 4 (basal + 2500mg/kg CCLM). One hundred and twenty 1-day-old Cobb 500 unsexed broiler chicks were randomly distributed to four dietary treatments. Each treatment was replicated three times. Thirty birds were assigned to each treatment (10 birds/replicate) in a completely

Table 1. The Experimental Basal Diets' Composition.

Ingredients	Starter feed	Finisher diet
Maize	52.33	59.32
Maize bran	7.02	0.00
Rice bran	0.00	6.03
Fish meal	3.00	3
Soybean meal	30	24
Premix	0.30	0.30
Bone meal	3.00	3.00
Soy oil	3.00	3.00
Methionine	0.30	0.30
Limestone	0.50	0.50
Salt	0.30	0.30
Lysine	0.25	0.25
<i>Analyzed composition (g/kg)</i>		
Crude fibre	3.55	3.63
Crude fat	4.47	3.94
Crude protein	22.19	20.09
<i>Calculated composition (g/kg)</i>		
Calcium	1.02	0.97
Available phosphorus	0.44	0.41
Methionine	0.68	0.65
Lysine	1.36	1.24
Metabolizable energy	3018.93	3108.16
(Kcal/kg)		

randomized design (CRD). The birds were housed in their respective pen (200 x 100 cm) with the floor covered with wood shavings. The temperature of the house was maintained within $31\text{oC} \pm 2$ for the first 7 days and reduced by 2°C after each consecutive 7 days until the house temperature was 26 ± 2 oC. The feed and water were provided ad libitum throughout the six weeks feeding trial. The lights were left on for 24 hours on the first day, and then reduced by 1 hour until 7th days, and later reduced to 20 hours per day til the end of the feeding trial.

Growth, Oxidative Stress Biomarkers, Immune and antioxidant enzymes analysis

The body weights of broilers were measured every seven days. The relative growth rate (RGR) was calculated using the following formula [22]:

$$\text{RGR} = [(w_2 - w_1) / ((w_1 + w_2) / 2)] * 100$$

Where w_1 = Bodyweight when the trial began, and w_2 = Bodyweight at the conclusion of the study

At 42 weeks of age, three birds per treatment were randomly selected and humanely slaughtered. Before slaughter, feeds were withdrawn from the birds overnight. During slaughtering, blood samples were obtained from jugular veins into a plain blood sample bottle for antioxidant enzymes (catalase and glutathione peroxidase). The blood sample in each of the plain bottles was centrifuged for 10 min at 3000 rpm to obtain clean supernatant serum. The harvested supernatant serum was kept at -20°C before analysis. The serum enzymatic activities of the catalase (CAT) and glutathione peroxidase (GPx) activities were determined according to the method of Aebi [21,3] and Rotruck *et al.* [22,4], respectively. Also, the fresh centrifuged serum was used to determine the concentration of immunoglobulin A (IgA), immunoglobulin G (IgG), immunoglobulin M (IgM) and immunoglobulin E (IgE) of the broiler chickens. The concentrations of IgE, IgM, and IgA were determined by mixing serum with buffer solution and antibody reagents to form an antigen/antibody complexes which, following agglutination, were measured turbidimetrically, using commercial ELISA quantitation kits (Fortress Diagnostics Limited, United Kingdom): IgG (Catalog No BXC0721A), IgM (Cat No BXC0731A), IgE (Cat No BXC0751A) and IgA (Cat No BXC0701A).

The levels of oxidative stress biomarkers were analyzed in the collected serum. The concentration of

8-hydroxydeoxyguanosine (8-OHdG) was determined in the blood serum using a commercial ELISA Kit (Elabscience Biotechnology Inc, USA) with catalog No E-EL-0028 96T, based on the manufacturer's instructions. The level of heat shock protein (HSP)/heat shock factor (HSP) was also determined using ELISA kit, according to the manufacturer's instructions (Bioassay Technology Laboratory, Shanghai, China). For antioxidant meat enzyme analysis, about 100g of fresh meat samples were excised from the breast muscle of broiler chicken after slaughter and evisceration for determination of catalase activity and superoxide dismutase, by the methods described by Hadwan and Khabt [23, 5] and Marhlund and Marklund [24, 6]

Statistical analysis

Data obtained on growth and immune responses, oxidative stress biomarkers and antioxidative enzymes of broilers were analyzed using PROC GLM procedures of Statistical Analysis System (SAS, version 9.1.3 of 2007). Differences in mean values were computed using Duncan's Multiple Range Test for multiple comparisons. For all statistical tests, significance was determined at $P < 0.05$.

Results

The result of the effect of CCLM and CALM supplements on serum oxidative biomarkers of broilers is presented in Table 2. The result revealed significant ($P < 0.05$) differences in heat shock protein (HSP) and 8-Hydroxy-2-deoxyguanosine (8OHdG) content of broilers among the treatments. Broilers fed the diet supplemented with CCLM (0.50pg/mL), CALM (0.46pg/mL) and Vit C (0.48 pg/mL) recorded lower heat shock protein content compared to the control group (0.65 pg/mL). Similarly, the 8-Hydroxy-2-deoxyguanosine content was significantly lowered in broiler chicken fed diet supplemented with CCLM (95.06 ng/mL), CALM (83.01 ng/mL) and Vit C (81.82 ng/mL) compared to the control group (122.06 ng/mL).

Table 3 shows the result of the serum antioxidant enzymes of broiler chickens fed diet supplemented with CCLM and CALM. The analysis revealed that broiler chickens fed diet containing CCLM (45.79 mg/ml), CLAM (32.20 mg/ml) and Vit C (46.01 mg/ml) had a higher concentration of glutathione peroxidase

Table 2: Serum oxidative stress biomarkers of broiler chicken fed diets supplemented with *Chenopodium ambrosioides L.* and *Crassocephalum crepidioides leaf meal*.

Parameters	Diet 1 Control	Diet 2 200mg/kg Vit C	Diet 3 2500mg/kg CCLM	Diet 3 2500mg/kg CALM	SEM	P value
HSP (pg/mL)	0.65 ^a	0.48 ^b	0.50 ^b	0.46 ^b	0.02	0.00
8OHdG (ng/mL)	122.06 ^a	81.82 ^b	95.06 ^{ab}	83.01 ^b	9.01	0.03

Means within a row with different letters and significantly different ($P < 0.05$). SEM Standard error. CALM = *Chenopodium ambrosioides leaf meal*. HSP = Heat shock protein. 8OHdG= 8-Hydroxy-2-deoxyguanosine

Table 3: Serum antioxidant enzymes of broiler chicken fed diets supplemented with *Chenopodium ambrosioides* L. and *Crassocephalum crepidioides* leaf meal

Parameters	Diet 1 Control	Diet 2 200mg/kg Vit C	Diet 3 2500mg/kg CCLM	Diet 3 2500mg/kg CALM	SEM	P value
GPX (mg/ml)	29.72 ^b	46.01 ^a	45.79 ^a	32.20 ^b	2.13	0.01
Catalase (mg/g)	135.01	160.78	160.94	170.53	9.27	0.13

Means within a row with different letters and significantly different ($P < 0.05$). SEM Standard error. CALM = *Chenopodium ambrosioides* leaf meal, GPX = Glutathione peroxidase, LDH = Lactate dehydrogenase

compared to the control group (29.72 mg/ml) ($P < 0.05$). The concentration of lactate dehydrogenase was significantly higher ($P < 0.05$) in broiler chickens fed diet containing CALM compared to other treatments. However, there was no significant influence ($p > 0.05$) of supplemented diets (CCLM and CALM) on the catalase content of the broiler chicken compared to the control group.

Table 4 shows the result of the immune response of broiler chicken fed the diet supplemented with CCLM and CALM. The results revealed significant differences ($P < 0.05$) in the value of immunoglobulin A (IgA) and M (IgM) across the experimental treatments. Broiler chicken fed diet supplemented with CCLM (277.55 mg/dl), CALM (259.65 mg/dl) and Vit C (253.79 mg/dl) had higher IgA content compared to the control group (209.58). Similarly, broiler chicken fed a diet containing CCLM, CALM and Vit C had higher IgM content at 382.92, 456.41 and 365.12 mg/dl respectively, compared to the control group (357.51 mg/dl). However, the result did not show any significant effect of supplemented diets (CCLM and CALM) on immunoglobulin E (IgE) and G (IgG) of the broiler chicken across the experimental treatments.

As presented in Table 5, supplementation of CCLM and CALM did not significantly influence the meat antioxidant content (catalase and superoxide dismutase) of the broiler chicken ($P > 0.05$) across treatments. The result of the effect of CCLM and CALM supplements on relative growth rate of the broilers is shown in Figure 1. The result did not show any significant ($P > 0.05$) effect of CCLM and CALM supplements on relative growth rate of the broilers compared to control group in both the starter and finisher phases of the feeding trial.

Discussion

Oxidative stress biomarkers

Natural antioxidants are commonly used in animal nutrition to promote growth, enhance endogenous antioxidants, inhibit oxidative stress, and boost the immune response of broilers [4, 6]. In this study, supplementation of CCLM and CALM significantly decreased the expression of heat shock protein in the broilers compared to the control. HSPs are specific proteins that are produced by cells in response to a variety of oxidative stressors such as oxidants, toxins, heavy metals, free radicals, and microbes [27]. They are produced as molecular chaperones to ameliorate the adverse effect of oxidative stress during animal production [28]. HSPs have been implicated in a diverse range of cellular activities, including senescence, endoplasmic stress, cell death signaling and inflammation [29]. The result of this study is suggesting that CCLM and CALM possessed the ability to attenuate the production of HSPs and alleviate oxidative damage in the cell. Different studies have shown that the dietary supplementation of natural antioxidants that is rich in phytochemicals can be used to modulate or inhibit the expression of heat shock protein in broilers under heat stress [30-32]. Conversely, the observed reduction in serum concentration of 8-Hydroxy-2-deoxyguanosine (8OHdG) of broiler chicken fed the diet supplemented with CCLM and CALM is further buttressing the ability of the CCLM and CALM to inhibit oxidative damages in poultry during production. Many 8-Hydroxy-2'-deoxyguanosine (8-OhdG) are produced in the cells as a product of DNA damage or oxidation, which is caused

Table 4: Immune response of broiler chicken fed diets supplemented with *Chenopodium ambrosioides* L. and *Crassocephalum crepidioides* leaf meal.

Parameters (mg/dl)	Diet 1 Control	Diet 2 200mg/kg Vit C	Diet 3 2500mg/k g CCLM	Diet 3 2500mg/kg CALM	SEM	P value
IgA	209.58 ^b	253.79 ^a	277.55 ^a	259.65 ^a	11.70	0.001
IgE	1082.15	1075.88	1077.37	1100.25	25.67	0.88
IgG	303.24	318.50	344.67	318.58	20.91	0.73
IgM	357.51 ^b	365.12 ^b	382.92 ^b	456.41 ^a	9.57	0.00

Means within a row with different letters and significantly different ($P < 0.05$). SEM Standard error. CALM = *Chenopodium ambrosioides* leaf meal, IgA =Immunoglobulin A, IgE= =Immunoglobulin E, IgG =Immunoglobulin G, IgM =Immunoglobulin M




Figure 1: The effects of *Chenopodium ambrosioides L.* and *Crassocephalum crepidioides leaf meal* on the relative growth rate of broilers.

by hydroxyl radical, superoxide, hydrogen peroxide, singlet oxygen or direct photodynamic action [33]. This result is similar to the findings of Prima *et al.* [34] who found that broiler chickens fed the diet supplemented with olive leaves and marigold petal extract significantly reduce the plasma concentration of 8-Hydroxy-2-deoxyguanosine compared to the control. Liu *et al.* [35] in their study also found that the dietary inclusion of *Macleaya cordata* extracts significantly reduced hepatic contents of 8-hydroxy-2'-deoxyguanosine in broiler chickens compared to the control group.

Antioxidant enzymes

The result of serum antioxidant enzymes revealed that supplementation of CCLM and CALM significantly increased the concentration of glutathione peroxidase compared to control. Glutathione peroxidase is one of the endogenous enzymatic antioxidants that help to protect the cell against oxidative damage by inactivating or scavenging or removing free radicals and other reactive oxygen species in cells [36]. This result is in agreement with the findings of Adeyeye *et al.* [36] who reported a significant increase in glutathione peroxidase content of broiler chickens fed the diet supplemented with wild sunflower and goat weed leaf

meals composite-mix compared to the control group. The increase in GPX in diets supplemented with CCLM, CALM and Vit C is indicating that they could exert higher antioxidative activity against oxidative damage in broiler chicken compared to the control. However, the observed similarity in the value of catalase of the broiler chicken across experimental treatments is in contrast with the report of Adu *et al.* [7] who found a significant increase in catalase content of broiler chicken fed the diet supplemented with *Syzygium aromaticum* leaf meal compared to control.

Immune response

Besides the potential of plant leaf meal as a natural antioxidant to inhibit oxidative damage and increase endogenous antioxidants, several studies have reported their ability to induce or modulate the immune response and increase immunoglobulin secretion in broiler chicken during production [37-38]. Immunoglobulins (IgA, IgM, IgE and IgG) are produced by plasma cells as part of the body's adaptive humoral immune response against a foreign pathogen [39]. In this study, the observed increase in the immune responses of the broiler chickens has shown that supplementation of CCLM and CALM could be used to stimulate the production of immunoglobulin M and A

Table 5: Meat antioxidant content of broiler chickens fed diets supplemented with *Chenopodium ambrosioides L.* and *Crassocephalum crepidioides leaf meal*.

Parameters	Diet 1 Control	Diet 2 200mg/kg Vit C	Diet 3 2500mg/kg CCLM	Diet 3 2500mg/kg CALM	SEM	<i>P value</i>
Catalase (mM/ml/min)	48.53	48.58	46.91	46.17	0.57	0.15
Superoxide dismutase (%)	91.22	89.75	93.66	94.63	1.88	0.31

Means within a row with different letters and significantly different ($P < 0.05$). SEM Standard error.

during production. Immunoglobulin (Ig) M is known as the first antibody isotype that helps to protect the host against infections and regulate other immune responses and tolerance [39]. While IgA plays an important role in inhibiting macromolecule absorption or binding of allergens to mucosal target cells, neutralizing bacterial toxins and enhancing nonspecific defense mechanisms in animals [41-42]. The result is similar to the findings of Osman *et al.* [43] and Cheng *et al.* [44] who found that broiler chickens fed the diet supplemented with turmeric powder and lotus leaf extract as natural antioxidants, respectively, had higher IgA and IgM concentrations compared to control. On the contrary, this result agreed with the report of Adeyemi *et al.* [20] who found no significant difference in the value of serum IgM of broiler chicken supplement with *Crassocephalum crepidioides* leaf meal at 1000 and 2000 ppm compared to the control. However, the observed similarity in concentration of IgE and IgG is indicating that the dietary inclusion of CCLM and CALM at 2500mg/kg could not improve their content compared to the control. This result is in contrast with the findings of Osman *et al.* [43] and Yao *et al.* [45] who reported a significant improvement in Immunoglobulins (IgA, IgM, IgE and IgG) of chicken supplemented with turmeric powder and sea buckthorn extract as a natural antioxidant, respectively, compared to control.

Meat antioxidant content

The observed similarity in the meat catalase and superoxide dismutase across experimental treatments is indicating that the inability of dietary inclusion of CCLM and CALM at 2500mg/kg to meat antioxidant content of the broiler chicken during production. This result is in line with the report of Adu *et al.* [7] and Gbore *et al.* [46] who found that dietary supplementation of *Syzygium aromaticum* and moringa leaf meal at 2500mg/kg each, respectively did not increase the catalase content of breast meat of broiler chicken compared to control.

Relative growth rate

The observed similarity in the relative growth rate between the broilers fed leaf meals and control diet is indicating that the utilization of CCLM and CALM at 2500mg/kg could not improve the growth performance of broilers during production. This result is in contrast with report of Oloruntola *et al.* [22] and Adu *et al.* [7] who found that dietary inclusion of plant leaf meal as feed additives at 2500mg/kg significantly increased the growth performance (body weight) of broilers compared to control group.

Conclusion

The findings of the present study reveal the potential of *Chenopodium ambrosioides* L. and *Crassocephalum crepidioides* leaf meal to inhibit

oxidative damage, enhance endogenous antioxidant enzymes, improve the immune system and maintain the meat quality of broiler chickens when used as a natural antioxidant in the diet. It was shown that the inclusion of CCLM and CALM at 2500mg/kg reduced the production of heat shock protein and Hydroxy-2-deoxyguanosine content, increased the level of serum glutathione peroxidase and catalase content, enhance the production of immunoglobulin IgA and IgM, and maintain the breast meat antioxidant of broiler chicken during production. Further research is required to assess CCLM and CALM effects on the meat fatty acid profile of broiler chickens.

Conflict of Interest

The author(s) declare that they have no known competing financial or non-financial, professional, or personal conflicts that could have appeared to influence the work reported in this paper.

References

- Bacou E, Walk C, Rider S, Litta G, Perez-Calvo E (2021). Dietary oxidative distress: A review of nutritional challenges as models for poultry, swine and fish. *Antioxidants*, 10(4):525.
- Ponnampalam EN, Kiani A, Santhiravel S, Holman B, Lauridsen C, Dunshea FR (2022). The Importance of Dietary Antioxidants on Oxidative Stress, Meat and Milk Production, and Their Preservative Aspects in Farm Animals: Antioxidant Action, Animal Health, and Product Quality—Invited Review. *Animals*. 12(23):3279.
- Surai PF, Fisinin VI (2016). Vitagenes in poultry production: Part 3. Vitagene concept development. *World's Poultry Science Journal*. 72:751.
- Falowo AB, Fayemi PO, Muchenje V (2014). Natural antioxidants against lipid-protein oxidative deterioration in meat and meat products: A review. *Food Research International* 64: 171–181
- Mousa MA, Osman AS, Hady HA (2017). Performance, immunology and biochemical parameters of *Moringa oleifera* and/or *Cichorium intybus* addition to broiler chicken ration. *Journal of veterinary medicine and animal health*, 9(10):255-263.
- Gharechopogh AM, Fakhraei J, Hosseini SA, Yarahmadi HM, Lotfollahian H (2021). Performance, immune responses, and blood biochemistry of broiler chickens fed with plant growth compound. *Tropical Animal Science Journal*, 44(1):62-70.
- Adu OA, Gbore FA, Oloruntola OD, Falowo AB, Olarotimi OJ (2020). The effects of *Myristica fragrans* seed meal and *Syzygium aromaticum* leaf meal dietary supplementation on growth performance and oxidative status of broiler chicken. *Bulletin of the National Research Centre*, 44(1):1-10.
- Oloruntola OD, Agbede JO, Ayodele SO, Oloruntola DA (2018). Neem, pawpaw, and bamboo leaf meal dietary supplementation in broiler chickens: Effect on performance and health status. *Journal of Food Biochemistry*, 2723

- Ayodele SO, Oloruntola OD, Ayeyeye SA, Jimoh OA, Falowo AB, Omoniyi IS (2021). Supplementation value of Mucuna seed powder on performance, antioxidant enzymes, meat cholesterol and peroxidation, and serum metabolites of broiler chickens. *Malaysian Journal of Animal Science*, 24(1):11-22.
- Falowo AB (2022). Beneficial effect of phytobiotic and synbiotic feed additives on growth performance, carcass characteristics and meat quality of broiler chicken. *Annals of Animal and Biological Research*, 2(1):61-70
- Reis JH, Gebert RR, Barreta M, Baldissera MD, Dos Santos ID, Wagner R, Campigotto G, Jaguezeski AM, Gris A, de Lima JL, Mendes RE (2018). Effects of phytogenic feed additive based on thymol, carvacrol and cinnamic aldehyde on body weight, blood parameters and environmental bacteria in broilers chickens. *Microbial pathogenesis*, 125:168-176
- Falowo AB (2022). Potential of medicinal plants as Hypocholesterolemic agents in chicken meat production. *Science Letter*, 10(1):24-31
- Kasali FM, Tusiimire J, Kadima JN, Agaba AG (2021). Ethnomedical uses, chemical constituents, and evidence-based pharmacological properties of *Chenopodium ambrosioides* L.: extensive overview. *Future Journal of Pharmaceutical Sciences*, 7:1-36.
- Kumar S, Pandey AK (2013). Chemistry and Biological Activities of Flavonoids: An Overview. *Science World Journal*, 2013 | Article ID 162750
- Diroff T (2008). Epazote and Holistic Health in Mexico, Senior Thesis Projects, University of Tennessee. Available at: http://trace.tennessee.edu/utk_interstp5/9
- Maldonado-Garcia M, Angulo C, Vazquez-Martinez J, Sanchez V, Lopez MG, Reyes-Becerril M (2019). Antioxidant and immunostimulant potentials of *Chenopodium ambrosioides* L. in Pacific red snapper (*Lutjanus peru*). *Aquaculture*, 513:734414.
- Reyes-Becerril M, Angulo C, Sanchez V, Vázquez-Martínez J, López MG (2019). Antioxidant, intestinal immune status and anti-inflammatory potential of *Chenopodium ambrosioides* L. in fish: In vitro and in vivo studies. *Fish and Shellfish Immunology*, 1, 86:420-428
- Can N M, Thao DTP (2020). Wound Healing Activity of *Crassocephalum crepidioides* (Benth.) S. Moore. Leaf Hydroethanolic Extract. *Oxidative medicine and cellular longevity*, 2020: 1-10
- Falowo AB, Oloruntola OD, Akinmoladun OF (2023). Assessment of Nutritional Composition and Antioxidant Properties of *Dysphania ambrosioides* (L.) Mosyakin & Clemants and *Crassocephalum crepidioides* Leaf Meal as Potential Feed Additives. *Turkish Journal of Agriculture - Food Science and Technology*, 11(2): 274-279, 2023
- Adeyemi KD, Sola-Ojo FE, Ajayi DO, Banni F, Isamot HO, Lawal MO (2021). Influence of dietary supplementation of *Crassocephalum crepidioides* leaf on growth, immune status, caecal microbiota, and meat quality in broiler chickens. *Tropical Animal Health Production*, 53(1):125.
- NRC (1994). Nutrient Requirements of Poultry (19th rev. ed.), *The National Academies Press*, Washington, DC (1994)
- Oloruntola OD, Ayodele SO, Adeyeye SA, Oloruntola DA, Osowe CO, Fasuhami OS (2022). Broiler Chickens' Growth, Haematological Indices, Guts Microbiota, Carcass and Meat Analysis in Response to Dietary Supplementation with *Anacardium occidentale* Leaf Powder and A Mix of Prebiotic, Probiotic and Acidifier. *Journal of Poultry Research*, 19(2):52-59.
- Aebi H (1974). Catalase estimation. In: Bergmeyer HV (ed) *Methods of enzymatic analysis*. Verlag Chemic, New York Academic Press, New York.
- Rotruck JT, Pope AL, Ganther HE, Swanson AB, Hafeman DG, Hoekstra WG (1973). Selenium: biochemical role as a component of glutathione peroxidase. *Science*, 179: 588-590.
- Hadwan M, Khabt H (2018). Simple spectrophotometric method for analysis of serum catalase activity. *Journal of Clinical and Diagnostic Research*, 12(Suppl 9):13-16
- Marklund S, Marklund G (1974). Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European Journal of Biochemistry*, 47: 469-474
- Ponomarenko M, Stepanenko I, Kolchanov N (2013). Heat Shock Proteins. Editor(s): Stanley Maloy, Kelly Hughes, *Brenner's Encyclopedia of Genetics* (Second Edition), Academic Press, 2013, Pages 402-405, ISBN 9780080961569, <https://doi.org/10.1016/B978-0-12-374984-0.00685-9>. (<https://www.sciencedirect.com/science/article/pii/B9780123749840006859>)
- Shehata AM, Saadeldin IM, Tukur HA, Habashy WS (2020). Modulation of Heat-Shock Proteins Mediates Chicken Cell Survival against Thermal Stress. *Animals* (Basel), 10(12):2407.
- Shankar K, Mehendale, HM (2014). Heat-Shock Proteins, Editor(s): Philip Wexler, *Encyclopedia of Toxicology* (Third Edition), Academic Press, 2014, Pages 830-831, ISBN 9780123864550, <https://doi.org/10.1016/B978-0-12-386454-3.00320-1>. (<https://www.sciencedirect.com/science/article/pii/B9780123864543003201>)
- Wan X, Ahmad H, Zhang L, Wang Z, Wang T (2018). Dietary enzymatically treated *Artemisia annua* L. improves meat quality, antioxidant capacity and energy status of breast muscle in heat-stressed broilers. *Journal of the Science of Food and Agriculture*, 98(10): 3715-3721.
- Balakrishnan KN, Ramiah SK, Zulkifli I (2023). Heat Shock Protein Response to Stress in Poultry: A Review. *Animals*, 13 (2):317.
- Sugito S, Etriwati E, Akmal M, Rahmi E, Delima M, Muchlisin ZA, Hasan DI (2021). Immunohistochemical Expression of AQP2 and HSP70 in Broiler Kidney Tissue Treated with *Salix tetrasperma* Roxb. Extract under Heat Exposure. *The Scientific World Journal*, 18;2021.
- Zhang L, Bai K, Zhang J, Xu W, Huang Q, Wang T (2017). Dietary effects of *Bacillus subtilis* fmbj on the antioxidant capacity of broilers at an early age. *Poultry science*, 96(10):3564-73.
- Priman T, Rezar V, Vrecl M, Salobir J, Levart A (2021). Effect of Olive leaves or Marigold petal extract on oxidative stress, gut fermentative activity, and mucosa morphology in broiler chickens fed a diet rich in n-3 polyunsaturated fats. *Journal of Poultry Science* 58 (2):119-130
- Liu Y, Li Y, Niu J, Liu H, Jiao N, Huang L, Jiang S, Yan L, Yang W (2022). Effects of Dietary *Macleaya cordata* extract containing isoquinoline alkaloids supplementation as an alternative to antibiotics in the diets on growth performance and liver health of broiler chickens. *Frontiers in Veterinary Science*, 9:950174

- Adeyeye SA, Oloruntola OD, Ayodele SO, Falowo AB, Agbede JO (2020). Wild sunflower and goat weed leaf meals composite-mix supplementation in broiler chickens: effects on performance, health status and meat. *Acta Fytotechnica et Zootechnica*, 23 (4): 205–212
- Dhama K, Latheef SK, Mani S, Samad HA, Karthik K, Tiwari R, Khan RU, Alagawany M, Farag MR, Alam GM, Laudadio V (2015). Multiple beneficial applications and modes of action of herbs in poultry health and production-a review. *International Journal of pharmacology*, 11(3):152-176.
- Mahfuz S, Piao XS (2019). Application of Moringa (*Moringa oleifera*) as natural feed supplement in poultry diets. *Animals*, 9(7):431.
- Sathe A, Cusick JK (2023). *Biochemistry, Immunoglobulin M*. In: Stat Pearls Treasure Island (FL): StatPearls Publishing: <https://www.ncbi.nlm.nih.gov/books/NBK555995/>
- Liu J, Wang Y, Xiong E, Hong R, Lu Q, Ohno H, Wang JY (2019). Role of the IgM Fc receptor in immunity and tolerance. *Frontiers in immunology*, 10:529. |
- Zhao K, Rong G, Hao Y, Yu L, Kang H, Wang X, Wang X, Jin Z, Ren Z, Li Z (2016). IgA response and protection following nasal vaccination of chickens with Newcastle disease virus DNA vaccine nanoencapsulated with Ag@ SiO₂ hollow nanoparticles. *Scientific Reports*, 6(1):1-2.
- Merino-Guzmán R, Latorre JD, Delgado R, Hernandez-Velasco X, Wolfenden AD, Teague KD, Graham LE, Mahaffey BD, Baxter MF, Hargis BM, Tellez G (2017). Comparison of total immunoglobulin A levels in different samples in Leghorn and broiler chickens. *Asian Pacific Journal of Tropical Biomedicine*, (2):116-120.
- Osman AH, El-Far AH, Sadek KM, Abo-Ghanema II, Abdel-Latif MA (2017). Immunity, Antioxidant Status, and Performance of Broiler Chickens Fed Turmeric (*Curcuma Longa*) Rhizome Powder. *Alexandria Journal for Veterinary Sciences*, 54(2) 19-28
- Cheng L, Zhang W, Jin Q, Zhu Y, Chen R, Tian Q, Yan N, Guo L (2021). The effects of dietary supplementation with lotus leaf extract on the immune response and intestinal microbiota composition of broiler chickens. *Poultry Science*, 100(3):100925.
- Yao BN, Liao FY, Yang JY, Liu A, Wang J, Zhu BG, Feng G, Yang SL (2023). Effect of sea buckthorn extract on production performance, serum biochemical indexes, egg quality, and cholesterol deposition of laying ducks. *Frontiers in Veterinary Science*, 10:185.
- Gbore FA, Oloruntola OD, Adu OA, Olarotimi OJ, Falowo AB, Afolayan EO (2021). Serum and meat antioxidative status of broiler chickens fed diets supplemented with garlic rhizome meal, moringa leaf meal and their composite. *Tropical Animal Health and Production*, 53:1-10.