

## Assessment of wound cultures in an oncology hospital

### *Onkoloji hastanesindeki hastaların yara kültürlerinin değerlendirilmesi*

Ferzan Arslan, Esra Tavukcu, Buket Demirhan, İpek Mumcuoğlu, Turgay Ulaş, Serap Süzük Yıldız, Ayşe Semra Güreşer, Neşe İnan, Gülşen İskender, Tuba Dal

Posted date:23.01.2024

Acceptance date:05.06.2024

#### Abstract

**Purpose:** The aim of this study is to evaluate the patient's demographic, clinical and laboratory data to determine whether the bacteria isolated from wound cultures are causative agents or colonization, and to determine their antimicrobial susceptibilities. This study aims to assess the demographic, clinical, and laboratory data of patients to distinguish between pathogenic bacteria and colonization in wound cultures, while also determining their antimicrobial susceptibilities.

**Materials and methods:** This retrospective research was conducted in Dr. Abdurrahman Yurtaslan Ankara Oncology Training and Research Hospital between January 1, 2021 and December 31, 2022. Two hundred thirty six isolates from 186 patients wound cultures were included in the study. Demographic data, clinical data and laboratory results of the patients were evaluated. The isolated bacteria and their antimicrobial susceptibilities were determined. The Q score system was used to evaluate the microbiological quality of wound samples.

**Results:** One hundred fifty nine cases (85%) were inpatients. Totally 119 (63.9%) patients were diagnosed with infection. The Q score for 136 samples (85.5%) was assessed as Q3. The most common isolated microorganisms were coagulase negative-staphylococci (CoNS) (19%), Escherichia coli (14.8%), and Staphylococcus aureus (13.1%), respectively in wound bacterial cultures. The methicillin resistance rate was 55.5% in CoNS and 54.1% in Staphylococcus aureus. Gram-negative bacteria were isolated in 81 (59.9%) infected patients.

Among patients with infected wounds, 39 (32.7%) patients had surgical site infections, 25 (21%) prosthesis infections, and diabetic foot infections 3 (2.5%). Infection rates were statistically significantly higher in patients with surgery, prosthesis, and diabetic foot ( $p=0.054$ ).

**Conclusion:** The Q score serves as a strong indicator for identifying the causative agent in wound infection and distinguishing it from colonization, thus aiding in the prevention of unnecessary antibiotic use. Regular review of local antibiotic susceptibility data is crucial in the clinical treatment of specific patient groups with oncological conditions.

**Keywords:** Wound culture, Q score, oncology patient, antimicrobial susceptibility.

Arslan F, Tavukcu E, Demirhan B, Mumcuoglu I, Ulas T, Suzuk Yildiz S, Gureser AS, Inan N, Iskender G, Dal T. Assessment of wound cultures in an oncology hospital. Pam Med J 2024;17:654-663.

#### Öz

**Amaç:** Bu çalışmanın amacı, yara kültürlerinden izole edilen bakterilerin etken/kolonizasyon ayrımının yapılmasında; hastaya ait demografik, klinik ve laboratuvar verilerinin değerlendirilmesi ve etken bakterilerin antimikrobiyal duyarlılıklarının belirlenmesidir.

**Gereç ve yöntem:** Bu retrospektif araştırma, 1 Ocak 2021-31 Aralık 2022 tarihleri arasında Dr. Abdurrahman Yurtaslan Ankara Onkoloji Eğitim ve Araştırma Hastanesi'nde gerçekleştirildi. Çalışmaya, 186 hastaya ait 236 yara kültürü dahil edildi. Hastalara ait demografik veriler, klinik veriler ve laboratuvar sonuçları değerlendirildi. İzole edilen bakteriler ve antimikrobiyal duyarlılıkları belirlendi. Yara örneklerinin mikrobiyolojik kalitesini değerlendirmek için Q skor sistemi kullanıldı.

Ferzan Arslan, Ph.D. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Medical Microbiology, Ankara, Türkiye, e-mail: md.ferzanarslan@gmail.com (<https://orcid.org/0009-0002-3934-3150>) (Corresponding Author)

Esra Tavukcu, Ph.D. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Medical Microbiology, Ankara, Türkiye, e-mail: md.esratavukcu@gmail.com (<https://orcid.org/0009-0005-6638-7492>)

Buket Demirhan, Ph.D. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Infectious Diseases and Clinical Microbiology, Ankara, Türkiye, e-mail: buket.demirhan@hotmail.com (<https://orcid.org/0009-0008-6072-3995>)

İpek Mumcuoğlu, Prof. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Medical Microbiology, Ankara, Türkiye, e-mail: ipekmumcuoglu@gmail.com (<https://orcid.org/0000-0002-6392-8880>)

Turgay Ulaş, Prof. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Hematology, Ankara, Türkiye, e-mail: turgayulas@yahoo.com (<https://orcid.org/0000-0001-9332-663X>)

Serap Süzük Yıldız, Assoc. Prof. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Medical Microbiology, Ankara, Türkiye, e-mail: serapsuzuk@gmail.com (<https://orcid.org/0000-0002-4820-6986>)

Ayşe Semra Güreşer, Assoc. Prof. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Medical Microbiology, Ankara, Türkiye, e-mail: asemragureser@hitit.edu.tr (<https://orcid.org/0000-0002-6455-5932>)

Neşe İnan, M.D. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Medical Microbiology, Ankara, Türkiye, e-mail: neseurdogan@yahoo.com (<https://orcid.org/0000-0002-1559-6244>)

Gülşen İskender, Assoc. Prof. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Infectious Diseases and Clinical Microbiology, Ankara, Türkiye, e-mail: golshan1669@hotmail.com (<https://orcid.org/0000-0001-7619-1366>)

Tuba Dal, Prof. Health Sciences University, Dr. Abdurrahman Yurtaslan Oncology Training and Research Hospital, Medical Microbiology, Ankara, Türkiye, e-mail: tuba\_dal@yahoo.com (<https://orcid.org/0000-0001-7045-1462>)

**Bulgular:** Vakaların 159'u (%85) yatan hastalardı. Toplam 119 (%63,9) hastada etken olarak kabul edildi. Q puanı 136 örnek için (%85,5) Q3 olarak değerlendirildi. Yara bakteri kültürlerinde en sık izole edilen mikroorganizmalar sırasıyla koagülaz negatif stafilokoklar (KNS) (%19), E. coli (%14,8) ve S. aureus (%13,1) oldu. Metisilin direnci oranı KNS'lerde %55,5; S. aureus'ta ise %54,1 olarak belirlendi. Enfeksiyöz hastaların 81'inde (%59,9) Gram negatif bakteri izole edildi. Enfekte yarası olan hastaların 39'unda (%32,7) cerrahi alan enfeksiyonu, 25'inde (%21) protez enfeksiyonu, 3'ünde (%2,5) diyabetik ayak enfeksiyonu vardı. Ameliyatlı, protezli ve diyabetik ayaklı hastalarda enfeksiyon oranları istatistiksel olarak anlamlı derecede yüksekti ( $p=0,054$ ).

**Sonuç:** Q skorlaması yara enfeksiyonunda etkenin saptanması ve kolonizasyonun dışlanmasında güçlü bir belirteçtir ve gereksiz antibiyotik kullanımının önlenmesine yardımcı olur. Onkolojik hastalar gibi özel hasta gruplarının ampirik tedavilerinin verilmesinde lokal antibiyotik duyarlılık verilerinin güncel olarak incelenmesi gereklidir.

**Anahtar kelimeler:** Yara kültürü, Q skoru, onkoloji hastaları, antimikrobiyal duyarlılık.

Arslan F, Tavukcu E, Demirhan B, Mumcuoğlu İ, Ulaş T, Süzük Yıldız S, Güreser AS, İnan N, İskender G, Dal T. Onkoloji hastanesindeki hastaların yara kültürlerinin değerlendirilmesi. Pam Tıp Derg 2024;17:654-663.

## Introduction

Wound infections are one of the most common causes of healthcare-associated infections (HCAIs) and lead to high mortality and morbidity. Timely and accurate evaluation of wound infections is vital. Determining the causative pathogens and their antimicrobial susceptibility increases the effectiveness of treatment and reduces mortality and morbidity. Management of wound infections will also contribute to the Sustainable Development Goals. According to 2017 data from the National Healthcare-Associated Infections Surveillance Network (USHIESA), 8,194 cases (1.3%) out of 617,745 total healthcare infections were attributed to surgical site infections (SSIs) [1].

The aim of this study is to differentiate whether the bacteria isolated from wound cultures are pathogens or colonization, to determine the antimicrobial susceptibility of bacteria interpreted as causative agents, and to evaluate the demographic, clinical and laboratory data of the patients.

## Material and methods

Cultures of 186 patients were included in the study between January 1, 2021 and December 31, 2022. Demographic data, clinical data and laboratory results of the patients (including C-reactive protein-CRP levels, procalcitonin levels and leukocyte counts) were evaluated. Clinical samples sent to the Medical Microbiology Laboratory were stained with Gram stain and microscopic examination was performed. Culture samples were simultaneously inoculated into 5% sheep blood agar and eosin methylene blue agar and evaluated after the appropriate

incubation period. The isolated bacteria were identified using traditional microbiological methods and the VITEK® 2 automated system (BioMérieux, France). Antibiotic susceptibility test results were determined according to EUCAST standards. Antibiotic susceptibility tests were performed both with the disc diffusion method and the VITEK® 2 automatic system (BioMérieux, France). Sensitive (S) and sensitive increasing exposure (I) results were considered sensitive.

In this retrospective study, demographic and clinical findings of the patients were obtained from the hospital data system. CRP and procalcitonin level were determined by AU5800 (Beckman Coulter INC) and Centaur XP (Siemens Healthcare), and leukocyte count levels were evaluated using the Automated Hematology Analyzer (Mindray BC-6200, Shenzhen Mindray Bio-Medical Electronics Co., Ltd., Shenzhen, China).

Colonization was characterized by the isolation of microorganisms from the wound without local and/or systemic signs and symptoms of infection. Local infection was defined by the presence of signs and symptoms of infection, which included erythema, local warmth, swelling, purulent discharge, delayed wound healing beyond expected timelines, the appearance of new or intensified pain, and increased foul odor [2]. Surgical site infections (SSI) are defined as infections that affect the incisional wound created during the surgical procedure or occur near the surgical site or organ. SSI was diagnosed according to the criteria of the Center for Disease Control and Prevention (CDC). Infections that occurred

within 30 days after surgery and 90 days when an implant (e.g., hip prosthesis) was used were designated as SSIs [3].

The Q score system was used to evaluate sample quality and determine the required extent of culture investigation for potential pathogens (PP). The Q score assigns positive values to the number of polymorphonuclear cells (PMNs) and negative values to the number of squamous epithelial cells (SECs) observed directly in the Gram-stained smear. The number resulting from the addition of these values creates the "Q score". Starting with a maximum value of 3, the score then continues to decrease values, maintaining the lower limit of zero; Negative numbers are always rounded to zero in the final calculation of the Q score [4, 5].

### Statistical analysis

Statistical data were analysed using SPSS (IBM Corp. Released 2019. IBM SPSS Statistics for Windows, Version 26.0. Armonk, NY: IBM Corp). The categorical data were expressed as percentage, and numbers; continuous variables were expressed as median, minimum

and maximum. The Chi-square test was used to compare the categorical data. *P* value <0.05 was considered to be significant statistically.

This research was approved by the Dr. Abdurrahman Yurtaslan Ankara Oncology Training and Research Hospital Non-Interventional Clinical Research Ethics Committee.

### Results

Of the 186 patients with positive wound cultures, 71 (38.2%) were male and 115 (61.8%) were female. The average age of the patients was 61.72 years. One hundred twenty-nine (69.4%) of the cases were inpatients and 27 (14.5%) were outpatients. Thirty (16.1%) patients were from intensive care unit. One hundred forty-two (76.3%) of the samples consisted of wound swabs, and 44 (23.7%) consisted of tissue, debridement, and drainage fluids. According to CDC criteria, 119 (64%) of the bacteria isolated from wound cultures were identified as causative agents and 42 (22.6%) were identified as colonization (Table 1).

**Table 1.** Demographic and clinical characteristics of patients

<b>Demographic characteristics</b>	
<b>The average age (years)</b>	61.7
<b>Gender</b>	
Female	115 (61.8%)
Male	71 (38.2%)
<b>Hospital Unit</b>	
Inpatient	129 (69.4%)
Outpatient	27 (14.5%)
Intensive care patients	30 (16.1%)
<b>Distribution of wound samples</b>	
Wound swab	142 (76.3%)
Tissue, debridement	44 (23.7%)
<b>Infection</b>	119 (64%)
<b>Colonization</b>	42 (22.6%)
<b>Undetermined sample</b>	25 (13.4%)

Out of 154 samples with available Gram stain results, in 53 samples (34.4%), the leukocyte count was  $\geq 25$ , and no epithelium was observed under x10 magnification microscopy. The Q score for these 53 samples was assessed as Q3. In 41 samples (26.6%), the leukocyte count

during the x10 scan fell within the range of 1-9, and epithelium was absent. The Q score for these 41 samples was also Q3. For 42 samples (27.2%) where no cells were detected, the Q Score was designated as Q3. The Q score for these 136 samples (88.2%) was assessed as

Q3. In 11 samples (7.2%), only 1-9 squamous epithelial cells were visible in each x10 scan area, and no leukocytes were detected so the Q score was determined as Q0.

Bacterial or yeast cells were observed in only seven samples (4.6%) by direct microscopy. The most frequently isolated microorganisms from wound cultures were coagulase-negative staphylococci (CoNS) (19%), *Escherichia coli* (14.8%) and *Staphylococcus aureus* (13.1%), respectively (Table 2). A single agent was isolated in 155 of the patients included in the study, two agents were isolated in 29 patients, and three or more different microorganisms were isolated in two patients. It was observed that the microorganism isolated from wound cultures was simultaneously isolated from non-

wound samples in 25 of the patients. Especially *E. coli* (8 cases), *Staphylococcus* spp. (5 cases) and *Candida* spp. (5 cases) were the most common isolates in these concurrent samples.

The methicillin resistance rate was 55.5% in CoNS and 54.1% in *S. aureus* (Table 3). Antimicrobial susceptibility of *Enterobacterales* to cephalosporins and carbapenem was observed 49.9% and 78.3%, respectively. In non-fermenter Gram-negative bacteria, ceftazidime and carbapenem susceptibility was 32% and 52%, respectively (Table 4). In particular, ceftazidime-avibactam susceptibility was performed for multidrug resistance 42 Gram-negative isolates and 88% (37/42) was susceptible.

**Table 2.** The distribution of microorganism species isolated in wound cultures

Microorganism	Number (n=236)	Percent (%)
Coagulase negative staphylococci	46	19.4
<i>Escherichia coli</i>	35	14.8
<i>Staphylococcus aureus</i>	31	13.1
<i>Klebsiella</i> spp.	26	11
<i>Enterococcus faecalis/faecium</i>	18	7.6
<i>Acinetobacter baumannii</i>	13	5.5
<i>Pseudomonas aeruginosa</i>	12	5
<i>Enterobacter</i> spp.	11	4.6
<i>Candida</i> spp.	8	3.3
<i>Proteus</i> spp.	6	2.5
Others	30	13.2

**Table 3.** Distribution of antibiotic susceptibility of Gram-positive microorganisms isolated in wound culture

	AM (%)	GN (%)	GNHR (%)	CIP (%)	LEV (%)	E (%)	DA (%)	LNZ (%)	VA (%)	TEC (%)	FA (%)	SXT (%)	FOX (%)
<b>CoNS (n=46)</b>	NA	NA	NA	23.2*	27.2*	26.6	97	97.8	100	100	41.3	71.7	44.5
<b><i>S. aureus</i> (n=31)</b>	NA	NA	NA	62.5*	NA	67.7	77.4	96.7	100	100	89.2	90.3	45.9
<b><i>Enterococcus</i> spp. (n=18)</b>	50	75	40	36.6	50	IR	IR	100	72.2	77.7	IR	IR	NA

AM: Ampicillin, GN: Gentamicin, GNHR: Gentamicin high dose resistance, CIP: Ciprofloxacin, E: Erythromycin, DA: Clindamycin, LNZ: Linezolid  
 VA: Vancomycin, TEC: Teicoplanin, FA: Fusidic acid, SXT: Trimethoprim-Sulphamethoxazole, FOX: Cefoxitin, NA: Not applicable  
 (\*): Susceptible, increased exposure, IR: Intrinsic Resistance

**Table 4.** Distribution of antibiotic susceptibility of Gram-negative microorganisms isolated in wound cultures

	CN (%)	AK (%)	AM (%)	TPZ (%)	CXM (%)	CAZ (%)	CRO (%)	FEP (%)	ETP (%)	MEM (%)	IMP (%)	CIP (%)	LEV (%)	TGC (%)	SCF (%)	SXT (%)
<i>E. coli</i> (n=35)	84	100	22.8	60.6	6.2	54.8	51.5	52.9	88.5	100	100	50	61.5	81.4	89.9	54.2
<i>Klebsiella spp.</i> (n=26)	46.1	54.1	IR	36.3	5.8	30.7	30.7	36	57.6	57.6	66.6	38.4	25	80	50	42.3
<i>Enterobacter spp.</i> (n=11)	70	100	IR	72.7	0	63.6	54.5	70	90.9	100	100	72.7	NA	NA	77.7	72.7
<i>Proteus spp.</i> (n=6)	40	100	16.6	100	0	100	83.3	100	100	100	100	50	100	16.6	100	33.3
<i>Pseudomonas aeruginosa</i> (n=12)	16.6	100	NA	33.3	NA	66.6	NA	100	NA	90.9	66.6	50	66.6	IR	100	NA
<i>Acinetobacter baumannii</i> (n=13)	36.3	58.3	NA	15.3	NA	27.2	NA	NA	NA	23	50	9	0	87.5	33.3	41.6

CN: Gentamicin, AK: Amikacin, AM: Ampicillin, TPZ: Piperacillin-tazobactam, CXM: Cefuroxime, CAZ: Ceftazidime, CRO: Ceftriaxone, FEP: Cefepime, ETP: Ertapenem  
MEM: Meropenem, IMP: Imipenem, CIP: Ciprofloxacin, LEV: Levofloxacin, TGC: Tigecycline, SCF: Cefoperazone/sulbactam, NA: Not applicable IR: Intrinsic Resistance  
SXT: Trimethoprim/sulfamethoxazole

In our study, we evaluated the clinical, laboratory data and treatment schedule and divided the patients into two groups: infection and colonization; (Table 5). Since we could not access the data of 25 patients, we evaluated infection or colonization in a total of 161 patients (Table 5). There was no significant difference in the presence of fever, local infection symptoms, CRP level, procalcitonin level and leukocyte count in patients with infection and colonization ( $p>0.05$ ) (Table 5). However, in

77.6% (125/161) of the cases, CRP levels were above normal limits, with an average of 98.4. Regarding procalcitonin, high levels were observed in 17.4% (28/161) of the patients tested. Empirical treatment and post-culture treatment was significantly compatible ( $p\leq 0.05$ ). Wound isolates identified as pathogens were diagnosed with cancer in 65 cases (54.6%), while among those considered as colonization, 22 cases (52.3%) were diagnosed with cancer. Patients with infectious wounds, 39 (32.8%)

had surgical site infection, 25 (21%) had prosthesis infection, and 3 (2.5%) had diabetic foot infection. Infection rates were higher in patients with surgery, prosthesis and diabetic feet ( $p=0.054$ ). While gram-negative bacteria

were isolated in 81 (60%) of the patients with infectious wounds, gram-positive bacteria were isolated in 30 (71.4%) of the patients with colonization.

**Table 5.** Clinical and laboratory findings of patients with wound infection/colonization

	Infection n=119, (%)	Colonization n=42, (%)	Values/p
<b>Clinical findings</b>			
Fever	26 (21.8)	5 (11.9)	1.932/0.381
<b>Local signs of infection</b>			
Fever, erythema, pain, tenderness	13 (10.9)	4 (9.5)	1.560/0.906
Serous discharge	22 (18.5)	6 (14.2)	
Fistula	3 (2.5)	1 (2.3)	
Purulent discharge	23 (19.3)	7 (16.6)	
Local signs of infection $\geq 1$	27 (22.7)	8 (19)	
<b>Laboratory findings</b>			
Increased CRP (C-reactive protein)	98 (82.3)	27 (64.2)	92.497/0.437
Increased Procalcitonin	19 (15.9)	9 (21.4)	2.356/0.502
Increased Leukocyte count	41 (34.4)	15 (35.7)	15.585/0.792
Receiving empirical treatment	90 (75.6)	28 (66.6)	4.316/0.497
<b>Compliance with empirical therapy and post-culture therapy</b>			
The same with post-culture treatment	38 (32)	21 (50)	1.142/0.378
Narrowed Post-culture therapy	14 (11.8)	0	0.798/0.508
Extended post-culture therapy	43 (36.1)	11 (26.2)	1.326/0.315
Antibiotic started patients for the first time after culture	16 (13.4)	2 (4.8)	1.762/0.184
Patients not given antibiotics	8 (6.7)	8 (19)	3.735/0.053
<b>The presence of cancer</b>	65 (54.6)	22 (52.3)	0.063/0.802
<b>The presence of prosthesis</b>	28 (23.5)	8 (19)	1.744/0.187
<b>Clinic/followed unit</b>			
Outpatient	16 (13.5)	6 (14.3)	0.030/0.985
Inpatient	81 (68)	28 (66.7)	
Intensive care unit	22 (18.5)	8 (19)	
<b>Wound types</b>			
Surgical side	39 (32.8)	8 (19)	7.693/0.103
Prosthesis	25 (21)	6 (14.3)	
Diabetic foot	3 (2.5)	2 (4.8)	
Other	52 (43.7)	26 (61.9)	
<b>History of hospitalization in the last three months</b>	51 (42.8)	14 (33.3)	0.366/0.545
<b>Culture result</b>			
Gram negative isolation	n=135 81 (60)	n=42 7 (16.6)	9.704/0.018
Gram positive isolation	51 (37.8)	30 (71.4)	4.892/0.026
<i>Candida</i> isolation	3 (2.2)	5 (12)	6.071/0.103

The Pearson Chi-Square test was used for the categorical variables

## Discussion

The human skin hosts a wide variety of microorganisms, many of which play a crucial role in defending against harmful pathogens through a phenomenon known as bacterial interference. These microorganisms, constituting the skin's flora, can be categorized as either resident or transient. Resident bacteria refer to the naturally occurring microorganisms that inhabit an individual's skin. These bacteria make their home on visible skin areas and within the skin's accessory structures. Transient bacteria are acquired when individuals come into contact with others or are exposed to surfaces teeming with bacterial presence. Among the diverse array of bacteria present on human skin, notable species include *Staphylococcus*, *Micrococcus*, *Peptococcus*, *Corynebacterium*, *Brevibacterium*, *Propionibacterium*, *Streptococcus*, *Neisseria*, and *Acinetobacter* species. Additionally, *Candida* spp. and the mites also take up residence on the skin. The quantity of bacteria within the stratum corneum is regulated to a certain extent by the continuous shedding of squames from the uppermost skin layer. The research findings highlighted that a significant portion of wound infections are caused by microorganisms commonly found in the body's natural flora and were seen in hospitalized patients. The skin's microbial community can generate biofilm, potentially leading to colonization and subsequent infection [6-11]. Similarly, in our study, the majority of the patients (>85%) were hospitalized patients. In our study, gram-negative bacteria isolation was detected in 60% of patients with infectious wounds, and gram-positive bacteria in 71.4% of patients with colonization. This data suggested that Gram-negative bacterial wound infections were more frequent in our hospital.

The colonized wounds' progress into infections is determined by several crucial factors. These factors encompass the concentration of bacteria per Gram of tissue and the host's immune system. In cases with appropriate wound care and management, the infection can escalate into septicemia, potentially leading to fatal outcomes. Wounds that have not progressed through the normal healing process and are open for  $\geq 1$  month are classified as chronic wounds. The most common risk factors of chronic wound infections were reported as

metabolic disruptions (e.g., diabetes), vascular deficits (e.g., venous or arterial insufficiency), or mechanical impacts. A breach in the skin integrity heals uneventfully with time and is defined as acute wounds. Acute wounds are injuries that occur suddenly and typically heal within a predictable time frame. They are often caused by external trauma, such as cuts, burns, abrasions, or surgical incisions. Advanced age, inadequate nutrition, obesity, diabetes, prolonged use of steroids, and compromised immune function were the factors for wound infections [7, 8]. In a retrospective study from China, 815 patients were analyzed. Microbial culture positivity was most pronounced in the wound tissue of ulcers resulting from infections (87.6%), with pressure-related ulcers following closely at (77.1%), followed by diabetes-related ulcers at (68.3%), and venous diseases at (67.7%). Within this patient group, (63.9%) of the tested samples exhibited microbial growth, comprising (13.4%) polymicrobial infections and (86.6%) monomicrobial infections. [9]. The surgery, presence of prosthesis, and diabetes mellitus were the most common risk factors for wound infections. Similarly, in our study, monomicrobial isolation was frequent in wound infections. We found that the patients with surgery, prosthesis, diabetes mellitus, advanced age, and, immunosuppression were at risk for the development of wound infections. Immunosuppression is very common in our study group due to 54.6% of the patients had cancer.

Within acute care settings, surgical wounds constitute the most prevalent wound type and they can cause potential complications like bleeding and wound reopening. SSI primarily manifest at the location of the surgical procedure, encompassing both the deep regions within the surgical zone contiguous to the operated organ (such as the hip, colon, pelvis, or brain) and the point of incision (the fascia, subcutaneous tissue, or skin). In conditions where a surgical site infection develops following a joint replacement procedure, the source of the infectious agent may be the nearby skin or the operating room. In an international study, the incidence of surgical site infections was estimated to occur in 1.9% to 40% of surgical procedures [6, 10]. In our study, among 119 patients evaluated as infected, 32.8% were surgical site infections. It was showed that surgical site infections were problematic in our hospital. We suggested

avoiding improper decontamination procedures (inappropriate antibiotic selection, compromised sterility practices) to decrease the incidence of post-surgical complications.

When evaluated clinically, infection in both acute and chronic wounds is typically characterized by an exaggerated inflammatory response surrounding the wound, elevated body temperature, pain, cellulitis, wound dehiscence, foul-smelling discharge, presence of pus, swelling, and warmth. Infection involves the infiltration of bacteria into tissue, while colonization is generally limited to the surface of the wound [7]. Our study indicated that, among the clinical findings, fever (21.8%), local signs of infection (10.9%), serous discharge (18.5%), and purulent discharge (19.3%) rates were higher in patients with infectious wounds than patients with colonization, although not statistically significant. We recommended evaluation of local and systemic signs, Gram-staining and culture results together for the diagnosis of the infection and colonization findings.

In a study conducted with 249 patients who had cesarean delivery, serum PCT, CRP levels, and WBC counts were measured at the postoperative 6<sup>th</sup>, 12<sup>th</sup>, and 24<sup>th</sup> hours. SSI assessments were conducted on the patients on the 2<sup>nd</sup>, 4<sup>th</sup>, and 7<sup>th</sup> days postoperatively. The study reported that 6% of the patients developed surgical site infections. The area under the curve (AUC) for PCT in predicting the SSI was 0.912 (95% CI: 0.79-1) with a sensitivity of 93.3% and specificity of 92.3% ( $p < 0.001$ ). The AUC for CRP was 0.854 and with a sensitivity of 80%, and specificity of 82.4%. Serum procalcitonin levels proved to be a more sensitive and specific indicator for the early diagnosis of SSIs following cesarean operation compared to others [12]. In our study, laboratory findings including elevated levels of CRP (82.3%), procalcitonin (15.9%), and higher numbers of leukocytes (34.4%) were seen in patients with infectious wounds. Although these results were not statistically significant, we recommended evaluating clinical and laboratory findings together in the diagnosis and following of wound infections. On the other hand, due to the majority of our patients having comorbidity and immunosuppression, there might be no statistically significant difference between infection and colonization.

Antibiotic resistance is a significant public health problem. In a study evaluating wound cultures, 600 isolates were analyzed, with 46.2% identified as Gram-positive bacteria, 51.3% as Gram-negative bacteria, and 2.5% as *Candida* spp. The most common isolates included *S. aureus* (29.2%), *E. coli* (11.5%), *P. aeruginosa* (11%), *Proteus mirabilis* (8%), and *Klebsiella pneumoniae* (5.8%). In a study evaluating wound cultures, 600 isolates were analyzed, with 46.2% identified as gram-positive bacteria, 51.3% as gram-negative bacteria, and 2.5% as *Candida* spp. The most common isolates were *S. aureus* (29.2%), *E. coli* (11.5%), *P. aeruginosa* (11%), *Proteus mirabilis* (8%), and *Klebsiella pneumoniae* (5.8%). Susceptibility tests revealed that 116 of the cultured bacteria exhibited resistance to multiple drugs, indicating the presence of multidrug-resistant strains. The resistance rates of *S. aureus* were >50% to methicillin, 92% to penicillin, 58.3% to erythromycin, and 50.9% to clindamycin. The resistance rates of *E. coli* were 68.1% to ampicillin, 68.1% to ciprofloxacin, 60.9% to levofloxacin, 3.9% to tigecycline, and 3.6% to amikacin [9]. In a study conducted with 5409 wound swabs in Saudi Arabia, a total of 14 different bacterial species were isolated and 9 of them were determined to be Gram negative bacteria. The most common isolates were *Klebsiella pneumoniae*, followed by *Pseudomonas aeruginosa*, *Escherichia coli*, *Acinetobacter baumannii*, methicillin-resistant *S. aureus* (MRSA), vancomycin-resistant *Enterococci* (VRE), and vancomycin-resistant *S. aureus* (VRSA). Multidrug resistant strains were determined as follows: *A. baumannii*, 97%; *K. pneumoniae*, 81%; *E. coli*, 71%; MRSA, 60%; *P. aeruginosa*, 33%; VRE, 22%; and VRSA, 2% [13]. In our study, the most common infection/colonization wound culture isolates were CoNS (19.4%), *E. coli* (14.8%), and *S. aureus* (13.1%). However, the Gram-negative isolation rate was higher in patients with infectious wounds than in patients with colonization (60% vs 16.6%). The rate of methicillin resistance was >50% in both CoNS and *S. aureus*. Within enteric bacilli, the resistance rate for 3rd generation cephalosporin was 55.1% and carbapenem 21.7%. In nonfermenter Gram-negative bacteria, resistance for ceftazidime was 68% and for carbapenem 48%. Notably, ceftazidime-avibactam was assessed in 42 Gram-negative

isolates with a resistance rate of 22%. In our study, 75.6% of patients with infectious wounds and 66.6% of patients with colonization received empirical treatment. Narrowed post-culture antimicrobial therapy was applied for 11.8%, and extended post-culture antimicrobial therapy for 36.1% of the patients with infectious wounds. Additionally, 13.4% of the patients with infectious wounds received antimicrobials for the first time after culture. Our study data suggested that every hospital should know the pathogenic agents and their antibiotic susceptibility patterns. The culture and antibiogram results had an important role in the management of wound infections and infection control. When choosing an empirical antimicrobial treatment option by clinicians, it should be kept in mind that Gram-negative bacterial wound infections are at the forefront in our hospital.

Our presented study, literature data [4, 5] indicated that the presence of microorganisms in Gram staining was not a good evaluating criteria for diagnosis of wound infection. The Q score for 136 samples (85.5%) was assessed as Q3. It was shown that Q scoring which assigns positive values to the count of PMNL cells and negative values to the count of squamous epithelial cells was a powerful marker for the determination of sample quality. In our study, it was determined that 76.3% of the wound samples were swab samples. It was reported that samples should be taken with at least two swabs for culture and Gram staining. The swab should be placed in 1-2 ml physiological saline or liquid medium, vortexed, and then inoculated into the medium, then the preparation is prepared for Gram staining [14]. We suggested that for an accurate diagnosis, appropriate and timely collection of swab samples and the application of laboratory sending criteria were necessary. If the samples dry out, the probability of bacterial isolation decreases. Given that clinical samples are not typically submitted to the laboratory in sets of two swabs, the reliability of our study Gram staining process becomes a concern.

In conclusion, wound infection rates, especially SSIs were common in our hospital among oncological patients. The most commonly isolated organisms from wound cultures were CoNS, *E. coli* and *S. aureus*. Gram negative isolation rate was higher in

patients with infectious wounds than patients with colonization. Gram positive isolation rate was higher in colonized patients than patients with infectious wounds. The patients with surgery, prosthesis, diabetes mellitus, and old age, immunosuppression were prone wound infection. Among the clinical findings, the presence of fever, local signs of infection, serous discharge, purulent discharge, elevated levels of CRP, procalcitonin, and higher numbers of leukocyte contributed to the diagnosis of wound infection. We recommended to evaluate clinical and laboratory findings together in the diagnose of wound infections. The culture and antibiogram results had an important role for the management of wound infections and infection control. When choosing an empirical antimicrobial treatment option by clinicians, it should be kept in mind that Gram-negative pathogen rates isolated from wound infections are common in our hospital. Q scoring was a powerful marker for diagnosis of wound infection and exclude of colonization. The appropriate and timely collection of swab samples were necessary. Avoiding improper decontamination procedures (inappropriate antibiotic selection, compromised sterility practices) will decrease the incidence of post-surgical complications.

**Conflict of interest:** No conflict of interest was declared by the authors.

## References

1. Cerrahi Alan Enfeksiyonu Sürveyansı. Available at: [https://hsgm.saglik.gov.tr/depo/birimler/bulasici-hastaliklar-ve-erken-uyari-db/Dokumanlar/Rehberler/CERRAHI\\_ALAN\\_ENFEKSIYONU\\_SURVEYANSI.pdf](https://hsgm.saglik.gov.tr/depo/birimler/bulasici-hastaliklar-ve-erken-uyari-db/Dokumanlar/Rehberler/CERRAHI_ALAN_ENFEKSIYONU_SURVEYANSI.pdf). Accessed December 14, 2023
2. Li S, Renick P, Senkowsky J, Nair A, Tang L. Diagnostics for wound infections. *Adv Wound Care (New Rochelle)* 2021;10:317-327. <https://doi.org/10.1089/wound.2019.1103>
3. Reeves N, Torkington J. Prevention of surgical site infections. *Surgery (Oxford)* 2021;40:20-24. <https://doi.org/10.1016/j.mpsur.2021.11.008>
4. Matkoski C, Sharp SE, Kiska DL. Evaluation of the Q score and Q234 systems for cost-effective and clinically relevant interpretation of wound cultures. *J Clin Microbiol* 2006;44:1869-1872. <https://doi.org/10.1128/JCM.44.5.1869-1872.2006>
5. McCarter YS, Sharp SE. Best laboratory practices for respiratory cultures. *Clinical Microbiology Newsletter* 2013;35:35-43. <https://doi.org/10.1016/j.clinmicnews.2013.02.001>

6. Alverdy JC, Hyman N, Gilbert J. Re-examining causes of surgical site infections following elective surgery in the era of asepsis. *Lancet Infect Dis* 2020;20:38-43. [https://doi.org/10.1016/S1473-3099\(19\)30756-X](https://doi.org/10.1016/S1473-3099(19)30756-X)
7. Wysocki AB. Evaluating and managing open skin wounds: colonization versus infection. *AACN Clin Issues* 2002;13:382-397. <https://doi.org/10.1097/00044067-200208000-00005>
8. Sen CK. Human Wound and Its Burden: Updated 2020 Compendium of Estimates. *Advances in Wound Care* 2021;10:281-292. <https://doi.org/10.1089/wound.2021.0026>
9. Guan H, Dong W, Lu Y, et al. Distribution and antibiotic resistance patterns of pathogenic bacteria in patients with chronic cutaneous wounds in China. *Front Med* 2021;8:609584. <https://doi.org/10.3389/fmed.2021.609584>
10. Li T, Zhang H, Chan PK, Fung WC, Fu H, Chiu KY. Risk factors associated with surgical site infections following joint replacement surgery: a narrative review. *Arthroplasty (London, England)* 2022;4:11(e1-8). <https://doi.org/10.1186/s42836-022-00113-y>
11. Percival SL, Emanuel C, Cutting KF, Williams DW. Microbiology of the skin and the role of biofilms in infection. *Int Wound J* 2012;9:14-32. <https://doi.org/10.1111/j.1742-481X.2011.00836.x>
12. Kanza Gül D. Procalcitonin and C-reactive protein measurements in the early diagnosis of surgical site infections after cesarean section. *CBUI SBED* 2021;8:232-240. <https://doi.org/10.34087/cbusbed.794037>
13. Al Said HM, Alghamdi A, Ashgar SS, et al. Isolation and Detection of Drug-Resistant Bacterial Pathogens in Postoperative Wound Infections at a Tertiary Care Hospital in Saudi Arabia. *Saudi J Med Med Sci* 2023;11:229-234. [https://doi.org/10.4103/sjmms.sjmms\\_405\\_22](https://doi.org/10.4103/sjmms.sjmms_405_22)
14. Gizzie N, Adukwu E. Evaluation of Liquid-Based Swab Transport Systems against the New Approved CLSI M40-A2 Standard. *J Clin Microbiol* 2016;54:1152-1156. <https://doi.org/10.1128/JCM.03337-15>

**Ethics committee approval:** Permission was obtained from University of Health Sciences, Dr. Abdurrahman Yurtaslan Ankara Oncology Training and Research Hospital Non-Interventional Clinical Research Ethics Committee for the study (permission date: 11.01.2024, permission number: 2023-12/125).

### Authors' contributions to the article

F.A., I.M., T.D. constructed the main idea and hypothesis of the study. F.A., I.M., T.D., G.I., N.I., S.S.Y. developed the theory and arranged/edited the material and method section. F.A., E.T., B.D., T.U. have done the evaluation of the data in the Results section. Discussion section of the article written by F.A., I.M., T.D., S.G., reviewed, corrected and approved. In addition, all authors discussed the entire study and approved the final version.