



# Double phase variable exponent problems with nonlinear matrices diffusion

Abderrahim Charkaoui<sup>1</sup> , Jinlan Pan<sup>\*2</sup> 

<sup>1</sup> *Interdisciplinary Research Laboratory in Sciences, Education and Training, Higher School of Education and Training Berrechid (ESEFB), Hassan First University Route de Casablanca Km 3.5, BP 539, 26100, Berrechid, Morocco*

<sup>2</sup> *Center for Applied Mathematics of Guangxi, and Guangxi Colleges and Universities Key Laboratory of Complex System Optimization and Big Data Processing, Yulin Normal University, Yulin 537000, Guangxi, P.R. China*

## Abstract

This work tackles a class of *double phase* elliptic problems with variable exponents and matrices diffusion. Under suitable assumptions on the data, we use critical point theory to establish both the existence and uniqueness of weak solutions to the *double phase* problem under consideration.

**Mathematics Subject Classification (2020).** Primary 35K55, 35A02; Secondary 35R11, 68U10, 65J15

**Keywords.** double phase problem, variable exponent, weak solution, matrices diffusion

## 1. Introduction

In this work, we are concerned with a nonlinear elliptic problem whose prototype reads as

$$\begin{cases} -\operatorname{div} \left( \langle A \nabla u, \nabla u \rangle^{\frac{p(x)-2}{2}} A \nabla u + \omega(x) \langle B \nabla u, \nabla u \rangle^{\frac{q(x)-2}{2}} B \nabla u \right) = f(x, u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where  $\Omega$  is an open and bounded subset of  $\mathbb{R}^N$ , with smooth boundary  $\partial\Omega$ . The variable exponents  $p(\cdot), q(\cdot) : \bar{\Omega} \rightarrow [2, +\infty)$  are two continuous functions satisfying  $2 \leq p(x) < N$  and  $p(x) < q(x)$  for all  $x \in \bar{\Omega}$ . The function  $\omega(\cdot)$  is nonnegative function belonging to a certain Lebesgue space. The source term  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function which belongs to a suitable Lebesgue type space to be discuss later. We assume that

\*Corresponding Author.

Email addresses: abderrahim.charkaoui@uhp.ac.ma (A. Charkaoui), jinlanpanylnu@163.com (J. Pan)

Received: 02.02.2024; Accepted: 14.05.2024

$A = (a_{i,j})_{1 \leq i,j \leq N}$ ,  $B = (b_{i,j})_{1 \leq i,j \leq N}$  are two symmetric diffusion matrices, i.e.  $a_{i,j} = a_{j,i}$  and  $b_{i,j} = b_{j,i}$  satisfying

$$a_{i,j}, b_{i,j} \in L^\infty(\Omega) \cap C^1(\Omega), \quad (1.2)$$

$$\langle A\xi, \xi \rangle = \sum_{i,j=1}^N a_{i,j}(x) \xi_i \xi_j \geq \lambda_A |\xi|^2, \quad (1.3)$$

$$\langle B\xi, \xi \rangle = \sum_{i,j=1}^N b_{i,j}(x) \xi_i \xi_j \geq \lambda_B |\xi|^2, \quad (1.4)$$

for all  $\xi \in \mathbb{R}^N$  and for almost  $x \in \Omega$ , with  $\lambda_A$  and  $\lambda_B$  being nonnegative constants, and  $\langle \cdot, \cdot \rangle$  stands for the scalar product on  $\mathbb{R}^N$ .

To effectively highlight the novelty and originality of our work, we aim to review pertinent existing literature on the same subject. We initiate this process by establishing the connection between the operator-driven problem (1.1) and certain well-known operators from earlier literature. It is worth noting that when  $\omega \equiv 0$ , the operator involved problem (1.1) reduces to

$$\mathcal{A}_{p(x)}(u) := -\operatorname{div} \left( \langle A\nabla u, \nabla u \rangle^{\frac{p(x)-2}{2}} A\nabla u \right), \quad (1.5)$$

which is recognized as the  $\mathcal{A}_{p(x)}$ -Laplace operator. It constitutes a generalization of the so-called  $p(x)$ -Laplace operator, i.e.  $\Delta_{p(x)}u = \operatorname{div} \left( |\nabla u|^{p(x)-2} \nabla u \right)$  with the latter obtained as a special case when  $A$  equals the identity operator  $I_d$  in (1.5). The investigation of partial differential equations involving the  $p(x)$ -Laplace operator has garnered escalating attention in recent years [2, 3, 13–15, 17, 24]. The heightened interest in studying these operators has been spurred by their diverse applications in areas such as image restoration [1, 18], mathematical biology [26], elastic mechanics [46, 47], electrorheological and thermorheological fluids [35, 39, 41]. Conversely, partial differential equations featuring the  $\mathcal{A}_{p(x)}$ -Laplace operator have undergone thorough examination in recent years. In [4], Alvino et al. delved into the eigenvalue problem for an elliptic equation involving the  $\mathcal{A}_p$ -Laplace operator with constant exponents ( $p(x) \equiv p$ ). The authors elucidated certain properties of the first eigenvalue and its corresponding eigenfunction by establishing a Payne-Rayner-type inequality in the process. For further exploration of PDEs governed by the  $\mathcal{A}_p$ -Laplace operator with constant exponent, we refer the readers to [23, 40], along with references provided therein. In a recent development, Mihăilescu and Repovš [36] have tackled a nonlinear elliptic equation with  $\mathcal{A}_{p(x)}$ -Laplace operator. Under a suitable assumption on the variable exponent  $p(x)$ , the authors combined Schauder's fixed point theorem with variational arguments to investigate the existence of solutions. In another scenario where  $A \equiv B \equiv I_d$  and  $\omega : \bar{\Omega} \rightarrow [0, \infty)$  is a measurable function, the problem (1.1) governed by the operator taking the form

$$\mathcal{L}(u) := -\operatorname{div} \left( |\nabla u|^{p(x)-2} \nabla u + \omega(x) |\nabla u|^{q(x)-2} \nabla u \right). \quad (1.6)$$

We would like to emphasize that the operator  $\mathcal{L}$  is commonly known as the *double phase* operator, a term that has gained significant attention in recent years. The term "*double phase*" was originally introduced in [19, 20], indicating that the flux function  $\left( |\nabla u|^{p(x)-2} + \omega(x) |\nabla u|^{q(x)-2} \right) \nabla u$  oscillates between two distinct elliptic scenarios. To phrase it differently, its behavior depends on the values assumed by the function  $\omega : \bar{\Omega} \rightarrow [0, \infty)$ . Specifically, over the subset  $\{x \in \Omega : \omega(x) = 0\}$ , the operator is governed by the gradient of  $p(x)$ -growth, whereas in scenarios where  $\{x \in \Omega : \omega(x) \neq 0\}$ , it is influenced by the gradient of  $q(x)$ -growth. This dual behavior is the fundamental reason for designating it as the "*double phase*" operator.

This type of operator has demonstrated its efficacy in modeling a diverse range of complex physical phenomena, including those arising in nonlinear elasticity, highly anisotropic materials, and the Lavrentiev phenomenon. For a deeper understanding of the real-world applications of such operators, we encourage interested readers to explore the following works [5, 6, 46–48]. It is crucial to highlight that the impetus for investigating this operator’s advantages can be traced back to Zhikov’s work [46], where the author delved into the behavior of highly anisotropic materials through the analysis of the following functional:

$$u \mapsto \int_{\Omega} (|\nabla u|^p + \omega(x)|\nabla u|^q) dx. \quad (1.7)$$

The inclusion of the modulating coefficient  $\omega(\cdot)$  serves to regulate the balance between two distinct materials, each characterized by power hardening rates of  $p$  and  $q$  respectively. In the existing literature, comprehensive investigations have explored the well-posedness of the functional (1.7). This scrutiny is conducted in various works, examining the functional in its original form [9, 27, 28, 33, 34, 37, 44], as well as in its role as a differential operator with variable growth behaviors, characterized by functions  $p(\cdot)$  and  $q(\cdot)$  [8, 10, 29, 42, 43, 45].

An important line of research, led by Mingione and his collaborators, has delved into the regularity of minimizers in variational problems and solutions to differential equations involving *double phase* operators, as documented in [7, 19, 20]. In a study by Cencelj-Rădulescu-Repovš [11], a class of *double phase* variational integrals was investigated, governed by non-uniform potentials with variable exponents  $p(\cdot)$  and  $q(\cdot)$ . Their analysis of the corresponding Euler equation revealed the existence of two distinct Rayleigh quotients. One of these quotients is linked to the existence of an infinite range of eigenvalues, while the second one is associated with the nonexistence of eigenvalues. The study [12] presented a novel parabolic *double phase* model which was applied in the field of image processing. The authors not only proved the existence and uniqueness of solutions but also illustrated the practicality of *double phase* operators in reducing image noise. In the subsequent work [16], the same authors extended their research from [12] by considering a reaction-diffusion parabolic system with variable exponents, governed by the *double phase* operator. They examined the existence and uniqueness of weak solutions to the proposed problem and subsequently validated the robustness of their model in the context of removing noise from Magnetic Resonance Images. Their research marks a significant milestone as it represents the first instance of utilizing parabolic *double phase* problems for image denoising and decomposition.

In the case where  $p(\cdot) \equiv p$  and  $q(\cdot) \equiv q$  are constants, Liu and Dai [31] successfully derived a sign-changing ground-state solution for a *double phase* problem, employing the Nehari manifold approach. Their subsequent work [32] extended the investigation, demonstrating the existence of at least three ground-state solutions through the application of the strong maximum principle. In a recent contribution [22], the authors established the first Schauder-type results for minima of nonuniformly elliptic integrals with nearly linear growth, encompassing the *double phase* integral operator as a special case. Furthermore, the paper by Ho and Winkert [28] introduced new embedding results for Musielak-Orlicz Sobolev spaces of *double phase* type. Their methodology involved the use of De Giorgi iteration and localization arguments, yielding appropriate boundedness results for corresponding weak solutions to two types of *double phase* problems, one with Dirichlet conditions and the other with nonlinear boundary conditions.

The main contribution of this work is to study the existence and uniqueness of solutions to the innovative *double phase* problem (1.1). The key novelty of our paper lies in the integration of a *double phase* operator with matrices diffusion. To the best of our knowledge, this is the first instance of a study encompassing both of these concepts. Specifically, problem (1.1) encapsulates two noteworthy phenomena. Firstly, the operator involved in

(1.1) is the *double phase* operator, characterized by its ability to switch between two distinct elliptic situations. Secondly, our work introduces matrices  $A$  and  $B$ , which serve to generalize not only the *double phase* operator (1.6) but also the well-known  $\mathcal{A}_{p(x)}$ -Laplace operator (1.5).

The remainder of this paper is structured as follows: In Section 2, we commence by revisiting key definitions and notable properties of Lebesgue, Sobolev, and Musielak-Orlicz spaces with variable exponents. Section 3 is devoted to presenting our main findings. Using variational calculus theory, we establish two results concerning the existence and uniqueness of weak solutions to problem (1.1). The first pertains to the scenario when the source term is independent of the solution. Conversely, the second is dedicated to the case of strong nonlinear source terms.

## 2. Mathematical preliminaries

In this section, we offer a concise overview of fundamental properties of variable exponent Lebesgue-Sobolev spaces. For a more comprehensive understanding, we direct the reader to see the book [38]. Throughout this paper, we will consistently employ the following notation:

$$\mathcal{M}(\Omega) := \{u/u : \Omega \rightarrow \mathbb{R} \text{ measurable}\}.$$

### 2.1. Lebesgue and Sobolev spaces with variable exponent

The objective of this section is to establish the necessary foundation for our forthcoming analysis, centered around the Lebesgue and Sobolev spaces with variable exponents. Let us denote:

$$\mathcal{C}^+(\overline{\Omega}) = \left\{ p \in \mathcal{C}(\overline{\Omega}) : \inf_{x \in \overline{\Omega}} p(x) \geq 1 \right\}.$$

We define the Lebesgue space with a variable exponent as  $L^{p(x)}(\Omega)$ , where

$$L^{p(x)}(\Omega) = \{u \in \mathcal{M}(\Omega) \mid \mathfrak{R}_{p(x)}(u) < \infty\},$$

here  $\mathfrak{R}_{p(\cdot)}$  stands for the following convex modular

$$\mathfrak{R}_{p(x)}(u) = \int_{\Omega} |u(x)|^{p(x)} dx.$$

We endow the Lebesgue space  $L^{p(x)}(\Omega)$  with the Luxemburg norm

$$\|u\|_{L^{p(x)}(\Omega)} = \inf \left\{ \delta > 0 \mid \mathfrak{R}_{p(x)}\left(\frac{u}{\delta}\right) \leq 1 \right\}.$$

For each  $p \in \mathcal{C}^+(\overline{\Omega})$  given, we introduce the pair  $(p^-, p^+)$  as follows

$$p^- = \inf_{x \in \overline{\Omega}} p(x) \quad \text{and} \quad p^+ = \sup_{x \in \overline{\Omega}} p(x).$$

**Proposition 2.1** ([25], Theorems 1.2 and 1.3). *Let  $u \in L^{p(x)}(\Omega)$  and  $\{u_n\} \subset L^{p(x)}(\Omega)$ . Then, we have*

(i) *the inequalities are satisfied*

$$\min \left\{ \|u\|_{L^{p(x)}(\Omega)}^{p^-}, \|u\|_{L^{p(x)}(\Omega)}^{p^+} \right\} \leq \mathfrak{R}_{p(x)}(u) \leq \max \left\{ \|u\|_{L^{p(x)}(\Omega)}^{p^-}, \|u\|_{L^{p(x)}(\Omega)}^{p^+} \right\}, \quad (2.1)$$

$$\min \left\{ \mathfrak{R}_{p(x)}^{\frac{1}{p^-}}(u), \mathfrak{R}_{p(x)}^{\frac{1}{p^+}}(u) \right\} \leq \|u\|_{L^{p(x)}(\Omega)} \leq \max \left\{ \mathfrak{R}_{p(x)}^{\frac{1}{p^-}}(u), \mathfrak{R}_{p(x)}^{\frac{1}{p^+}}(u) \right\}. \quad (2.2)$$

(ii) *if  $u_n \rightarrow u$  in  $L^{p(x)}(\Omega)$ , then the succeeding statements are equivalent:*

(a)  $\lim_{n \rightarrow +\infty} \|u_n - u\|_{L^{p(x)}(\Omega)} = 0.$

$$(b) \lim_{n \rightarrow +\infty} \mathfrak{R}_{p(x)}(u_n - u) = 0.$$

$$(c) \lim_{n \rightarrow +\infty} \mathfrak{R}_{p(x)}(u_n) = \mathfrak{R}_{p(x)}(u) \text{ and } u_n \rightarrow u \text{ in measure in } \Omega.$$

**Proposition 2.2** ([25], Theorems 1.6 and 1.10). *Let  $p \in \mathcal{C}^+(\overline{\Omega})$ . Then the space  $(L^{p(x)}(\Omega), \|\cdot\|_{L^{p(x)}(\Omega)})$  is separable. Moreover, when  $p^- > 1$ , the generalized Lebesgue space  $L^{p(x)}(\Omega)$  becomes uniform convex and thus is reflexive.*

For any  $p \in \mathcal{C}^+(\overline{\Omega})$ , we define the conjugate exponent of  $p$  as  $p'$  which meets  $\frac{1}{p(x)} + \frac{1}{p'(x)} = 1$  for all  $x \in \overline{\Omega}$ , with the convention that  $\frac{1}{\infty} = 0$ . The subsequent proposition is commonly known as the generalized Hölder inequality.

**Proposition 2.3** ([30], Theorem 2.1). *Let  $p \in \mathcal{C}^+(\overline{\Omega})$  with  $p^- > 1$ . For any  $(u, v) \in L^{p(x)}(\Omega) \times L^{p'(x)}(\Omega)$ , we have*

$$\left| \int_{\Omega} uv \, dx \right| \leq 2 \|u\|_{L^{p(x)}(\Omega)} \|v\|_{L^{p'(x)}(\Omega)}.$$

The following proposition emphasizes notable embedding results for Lebesgue spaces with variable exponents.

**Proposition 2.4** ([25], Theorem 1.11). *Let  $p_1, p_2 \in \mathcal{C}^+(\overline{\Omega})$  with  $1 \leq p_1(x) \leq p_2(x)$  almost everywhere in  $\Omega$ . Then, we have the following continuous embedding  $L^{p_2(x)}(\Omega) \hookrightarrow L^{p_1(x)}(\Omega)$ .*

We define the Sobolev space with a variable exponent as  $W^{1,p(x)}(\Omega)$ , where

$$W^{1,p(x)}(\Omega) = \left\{ u \in L^{p(x)}(\Omega) \mid |\nabla u| \in L^{p(x)}(\Omega) \right\}.$$

We equip  $W^{1,p(x)}(\Omega)$  with the norm  $\|\cdot\|_{W^{1,p(x)}(\Omega)}$  defined as

$$\|u\|_{W^{1,p(x)}(\Omega)} = \|u\|_{L^{p(x)}(\Omega)} + \|\nabla u\|_{L^{p(x)}(\Omega)}.$$

It is not hard to prove that the above norm is equivalent to the one

$$\|u\|_{W^{1,p(x)}(\Omega)} = \inf \left\{ \delta > 0 \mid \int_{\Omega} \left( \left| \frac{\nabla u(x)}{\delta} \right|^{p(x)} + \left| \frac{u(x)}{\delta} \right|^{p(x)} \right) dx \leq 1 \right\}.$$

Hereafter, we say that the function  $p(x)$  satisfies the log-Hölder continuity condition, if there exists a constant  $C > 0$  such that

$$|p(x_1) - p(x_2)| \leq \frac{C}{-\log |x_1 - x_2|}, \text{ for all } x_1, x_2 \in \Omega, \text{ with } |x_1 - x_2| < \frac{1}{2}. \quad (2.3)$$

Assumption (2.3) ensures the density of the space of smooth functions  $\mathcal{C}_c^\infty(\Omega)$  within  $W^{1,p(x)}(\Omega)$ . For added convenience, we introduce the space  $W_0^{1,p(x)}(\Omega) := \overline{\mathcal{C}_c^\infty(\Omega)}^{W^{1,p(x)}(\Omega)}$ , with its dual space denoted as  $(W_0^{1,p(x)}(\Omega))^*$ . Furthermore, for any  $u \in W_0^{1,p(x)}(\Omega)$ , the generalized Poincaré inequality holds:

$$\|u\|_{L^{p(x)}(\Omega)} \leq C \|\nabla u\|_{L^{p(x)}(\Omega)}, \quad (2.4)$$

where  $C$  is a nonnegative constant depending only on  $p(x)$  and  $\Omega$ . Taking advantage of generalized Poincaré inequality (2.4), we can establish the following equivalent norm on the space  $W_0^{1,p(x)}(\Omega)$ :

$$\|u\|_{W_0^{1,p(x)}(\Omega)} = \|\nabla u\|_{L^{p(x)}(\Omega)}.$$

In a manner akin to  $L^{p(x)}(\Omega)$ , when  $p^- > 1$ , both the Banach spaces  $W^{1,p(x)}(\Omega)$  and  $W_0^{1,p(x)}(\Omega)$  are also separable and reflexive. The ensuing propositions succinctly outline noteworthy embedding results that establish connections between the spaces  $W^{1,p(x)}(\Omega)$  and  $L^{p(x)}(\Omega)$ .

**Proposition 2.5** ([25], Theorem 2.3). *Let  $(p(\cdot), q(\cdot)) \in \mathcal{C}^+(\bar{\Omega}) \times \mathcal{C}^+(\bar{\Omega})$  with  $p^- > 1$  and  $1 \leq q(x) < p^*(x)$  for almost  $x \in \bar{\Omega}$ . Then the embedding  $W_0^{1,p(x)}(\Omega) \hookrightarrow L^{q(x)}(\Omega)$  is continuous and compact, where*

$$p^*(x) := \begin{cases} \frac{Np(x)}{N-p(x)}, & \text{if } p(x) < N \\ +\infty, & \text{if } p(x) \geq N. \end{cases}$$

## 2.2. Orlicz-Sobolev spaces with variable exponent

This subsection provides a basic overview of Orlicz-Sobolev spaces, introducing key concepts. For a more in-depth exploration, readers are encouraged to see the work [21]. Let  $(p(\cdot), q(\cdot)) \in \mathcal{C}^+(\bar{\Omega}) \times \mathcal{C}^+(\bar{\Omega})$  be such that  $1 < p(x) < N$  and  $p(x) < q(x)$  for all  $x \in \bar{\Omega}$ , we define  $\mathcal{H} : \Omega \times [0, \infty) \rightarrow [0, \infty)$  as

$$\mathcal{H}(x, r) := r^{p(x)} + \omega(x)r^{q(x)}, \quad \text{for all } (x, r) \in \Omega \times [0, \infty),$$

where  $\omega$  is a nonnegative function belonging to  $L^\infty(\Omega)$ . It is straightforward to confirm that  $\mathcal{H}$  is a generalized  $N$ -function and is locally integrable. We proceed to introduce the Musielak-Orlicz space, denoted as  $L^{\mathcal{H}}(\Omega)$

$$L^{\mathcal{H}}(\Omega) = \{u \in \mathcal{M}(\Omega) \mid \mathfrak{R}_{\mathcal{H}}(u) < \infty\},$$

where the convex modular  $\mathfrak{R}_{\mathcal{H}}$  is introduced as

$$\mathfrak{R}_{\mathcal{H}}(u) = \int_{\Omega} \mathcal{H}(x, |u|) dx.$$

and endowed  $L^{\mathcal{H}}(\Omega)$  with the norm

$$\|u\|_{L^{\mathcal{H}}(\Omega)} = \inf \left\{ \delta > 0 \mid \mathfrak{R}_{\mathcal{H}}\left(\frac{u}{\delta}\right) \leq 1 \right\}.$$

Also, we introduce the Musielak-Orlicz Sobolev space  $W^{1,\mathcal{H}}(\Omega)$  as follows

$$W^{1,\mathcal{H}}(\Omega) = \left\{ u \in L^{\mathcal{H}}(\Omega) \mid |\nabla u| \in L^{\mathcal{H}}(\Omega) \right\}.$$

equipped  $W^{1,\mathcal{H}}(\Omega)$  with the norm

$$\|u\|_{W^{1,\mathcal{H}}(\Omega)} = \|u\|_{L^{\mathcal{H}}(\Omega)} + \|\nabla u\|_{L^{\mathcal{H}}(\Omega)}.$$

Moreover, we denote by  $W_0^{1,\mathcal{H}}(\Omega)$  the Musielak-Orlicz Sobolev spaces with zero traces, which, actually, represents the closure of  $\mathcal{C}_c^\infty(\Omega)$  with respect to  $W^{1,\mathcal{H}}(\Omega)$  norm, that is

$$W_0^{1,\mathcal{H}}(\Omega) = \overline{\mathcal{C}_c^\infty(\Omega)}^{W^{1,\mathcal{H}}(\Omega)}.$$

With these norms, the spaces  $L^{\mathcal{H}}(\Omega)$ ,  $W^{1,\mathcal{H}}(\Omega)$ , and  $W_0^{1,\mathcal{H}}(\Omega)$  attain the properties of being separable, reflexive, and uniformly convex Banach spaces. The subsequent proposition elucidates the relationships between the modular  $\mathfrak{R}_{\mathcal{H}}(\cdot)$  and the norm  $\|\cdot\|_{L^{\mathcal{H}}(\Omega)}$ .

**Proposition 2.6** ([21]). *Let  $u$  be an element of  $L^{\mathcal{H}}(\Omega)$ . Then, we have*

(i) *the inequalities hold*

$$\min \left\{ \|u\|_{L^{\mathcal{H}}(\Omega)}^{p^-}, \|u\|_{L^{\mathcal{H}}(\Omega)}^{q^+} \right\} \leq \mathfrak{R}_{\mathcal{H}}(u) \leq \max \left\{ \|u\|_{L^{\mathcal{H}}(\Omega)}^{p^-}, \|u\|_{L^{\mathcal{H}}(\Omega)}^{q^+} \right\}. \quad (2.5)$$

(ii) if  $p(x) < q(x) < p^*(x)$  for all  $x \in \Omega$  and  $0 \leq \omega(\cdot) \in L^\infty(\Omega)$ , then, we have

$$W^{1,\mathcal{H}}(\Omega) \hookrightarrow L^{\mathcal{H}}(\Omega) \text{ is compact.}$$

(iii) there exists a nonnegative constant  $C$  such that

$$\|u\|_{L^{\mathcal{H}}(\Omega)} \leq C \|\nabla u\|_{L^{\mathcal{H}}(\Omega)}, \quad \text{for all } u \in W_0^{1,\mathcal{H}}(\Omega). \quad (2.6)$$

From (2.6), we can see that the following norm is an equivalent norm of  $W_0^{1,\mathcal{H}}(\Omega)$

$$\|u\|_{W_0^{1,\mathcal{H}}(\Omega)} = \|\nabla u\|_{L^{\mathcal{H}}(\Omega)}, \quad \text{for all } u \in W_0^{1,\mathcal{H}}(\Omega).$$

Notably, it is worth observing that the Lebesgue space with a variable exponent  $L^{p(x)}(\Omega)$  is a particular instance of the Musielak-Orlicz space. In relation to the Musielak-Orlicz Sobolev space  $W^{1,\mathcal{H}}(\Omega)$ , we present the following set of embedding theorems.

**Proposition 2.7** ([21]). *Let  $s(\cdot) \in \mathcal{C}(\overline{\Omega})$ . Then, we have*

- (i) if  $1 \leq s(x) \leq p(x)$ , for all  $x \in \Omega$ , then the embedding  $W^{1,\mathcal{H}}(\Omega) \hookrightarrow W^{1,s(x)}(\Omega)$  is continuous.
- (ii) if  $1 \leq s(x) \leq p^*(x)$ , for all  $x \in \Omega$ , then the embedding  $W^{1,\mathcal{H}}(\Omega) \hookrightarrow L^{s(x)}(\Omega)$  is continuous.
- (iii) if  $1 \leq s(x) < p^*(x)$ , for all  $x \in \Omega$ , then the embedding  $W^{1,\mathcal{H}}(\Omega) \hookrightarrow L^{s(x)}(\Omega)$  is compact.

Throughout this paper, we assume that  $(p, q) \in \mathcal{C}^+(\overline{\Omega}) \times \mathcal{C}^+(\overline{\Omega})$  take values within the intervals  $(p^-, p^+)$  and  $(q^-, q^+)$  respectively, satisfying log-Hölder continuity condition (2.3) such that

$$2 \leq p(x) < N \quad \text{and} \quad p(x) < q(x). \quad (2.7)$$

Assumption (2.7) leads to employing the aforementioned properties of Lebesgue and Sobolev spaces with variable exponent. Furthermore, we assume that  $\omega : \Omega \rightarrow [0, +\infty)$  is a measurable function belonging to  $L^\infty(\Omega)$ .

### 3. Main results

We begin this section by clarifying the notion of weak solutions used to solve problem (1.1).

**Definition 3.1.** A measurable function  $u : \Omega \rightarrow \mathbb{R}$  is said to be a *weak* solution to problem (1.1) if it satisfies

$$\int_{\Omega} \langle A \nabla u, \nabla u \rangle^{\frac{p(x)-2}{2}} \langle A \nabla u, \nabla \varphi \rangle dx + \int_{\Omega} \omega(x) \langle B \nabla u, \nabla u \rangle^{\frac{q(x)-2}{2}} \langle B \nabla u, \nabla \varphi \rangle dx = \int_{\Omega} f(x, u) \varphi dx, \quad (3.1)$$

for all  $\varphi \in W_0^{1,\mathcal{H}}(\Omega)$ .

In the following theorem, we present our first existence result, which pertains to the case where the source term  $f(x, u)$  is independent of the steady-state solution.

**Theorem 3.2.** *Under the assumptions (1.2), (1.3), (1.4) and (2.7), we assume that  $f(x, r) = f(x)$  with  $f \in L^{\theta(x)}(\Omega)$ , where  $\theta \in \mathcal{C}^+(\overline{\Omega})$  is such that  $\frac{1}{\theta(x)} + \frac{1}{p^*(x)} = 1$ . Then, problem (1.1) admits a unique weak solution.*

**Proof.** Let us start by introducing the energy functional  $\mathcal{J} : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$  associated to problem (1.1) as follows

$$\mathcal{J}(u) = \int_{\Omega} \frac{1}{p(x)} \langle A \nabla u, \nabla u \rangle^{\frac{p(x)}{2}} dx + \int_{\Omega} \frac{\omega(x)}{q(x)} \langle B \nabla u, \nabla u \rangle^{\frac{q(x)}{2}} dx - \int_{\Omega} f(x) u dx. \quad (3.2)$$

By employing standard arguments, we can ensure that  $\mathcal{J}$  belongs to  $\mathcal{C}^1(W_0^{1,\mathcal{H}}(\Omega); \mathbb{R})$  and its derivative in the weak sense is given by

$$\langle \mathcal{J}'(u), \varphi \rangle = \int_{\Omega} \langle A \nabla u, \nabla u \rangle^{\frac{p(x)-2}{2}} \langle A \nabla u, \nabla \varphi \rangle dx + \int_{\Omega} \omega(x) \langle B \nabla u, \nabla u \rangle^{\frac{q(x)-2}{2}} \langle B \nabla u, \nabla \varphi \rangle dx - \int_{\Omega} f(x) \varphi dx,$$

for all  $\varphi \in W_0^{1,\mathcal{H}}(\Omega)$ . Therefore, the critical points of  $\mathcal{J}$  precisely correspond to the weak solutions of problem (1.1). By analyzing (3.2) and taking into account (1.3) and (1.4), it follows that

$$\begin{aligned} \mathcal{J}(u) &\geq \int_{\Omega} \frac{\lambda_A}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{\lambda_B}{q(x)} \omega(x) |\nabla u|^{q(x)} dx - \|f\|_{L^{\theta(x)}(\Omega)} \|u\|_{L^{p^*(x)}(\Omega)}, \\ &\geq \frac{\lambda_A}{p^+} \int_{\Omega} |\nabla u|^{p(x)} dx + \frac{\lambda_B}{q^+} \int_{\Omega} \omega(x) |\nabla u|^{q(x)} dx - C \|f\|_{L^{\theta(x)}(\Omega)} \|u\|_{W_0^{1,\mathcal{H}}(\Omega)}, \end{aligned}$$

where  $C$  is the Sobolev embedding constant of  $W_0^{1,\mathcal{H}}(\Omega) \hookrightarrow L^{p^*(x)}(\Omega)$ . We mention that the later inequality is obtained via the use of Hölder inequality with the help of Proposition (2.7). Then, we have

$$\mathcal{J}(u) \geq \min \left\{ \frac{\lambda_A}{p^+}, \frac{\lambda_B}{q^+} \right\} \int_{\Omega} \left( |\nabla u|^{p(x)} + \omega(x) |\nabla u|^{q(x)} \right) dx - C \|f\|_{L^{\theta(x)}(\Omega)} \|u\|_{W_0^{1,\mathcal{H}}(\Omega)}. \quad (3.3)$$

By taking advantage of (2.5), it results from (3.3) that for any  $u \in W_0^{1,\mathcal{H}}(\Omega)$  with  $\|u\|_{W_0^{1,\mathcal{H}}(\Omega)} > 1$ , one has

$$\mathcal{J}(u) \geq \min \left\{ \frac{\lambda_A}{p^+}, \frac{\lambda_B}{q^+} \right\} \|u\|_{W_0^{1,\mathcal{H}}(\Omega)}^{p^-} - C \|f\|_{L^{\theta(x)}(\Omega)} \|u\|_{W_0^{1,\mathcal{H}}(\Omega)}. \quad (3.4)$$

Therefore, inequality (3.4) proves that  $\mathcal{J}$  is coercive on  $W_0^{1,\mathcal{H}}(\Omega)$ . But, from the definition of  $\mathcal{J}$ , it can see that  $\mathcal{J}$  is bounded and continuous. In addition, with the help of the Clarkson's type inequality (see p. 449 of [4]), it indicates that for all  $r, s \geq 2$  and for all  $\xi_1, \xi_2 \in \mathbb{R}^N$ , one has

$$\frac{\langle A \xi_1, \xi_1 \rangle^{\frac{r}{2}} + \langle A \xi_2, \xi_2 \rangle^{\frac{r}{2}}}{2} \geq \left\langle A \left( \frac{\xi_1 + \xi_2}{2} \right), \frac{\xi_1 + \xi_2}{2} \right\rangle^{\frac{r}{2}} + \left\langle A \left( \frac{\xi_1 - \xi_2}{2} \right), \frac{\xi_1 - \xi_2}{2} \right\rangle^{\frac{r}{2}}, \quad (3.5)$$

$$\frac{\langle B \xi_1, \xi_1 \rangle^{\frac{s}{2}} + \langle B \xi_2, \xi_2 \rangle^{\frac{s}{2}}}{2} \geq \left\langle B \left( \frac{\xi_1 + \xi_2}{2} \right), \frac{\xi_1 + \xi_2}{2} \right\rangle^{\frac{s}{2}} + \left\langle B \left( \frac{\xi_1 - \xi_2}{2} \right), \frac{\xi_1 - \xi_2}{2} \right\rangle^{\frac{s}{2}}. \quad (3.6)$$

The estimates (3.5) and (3.6) illustrates that  $\mathcal{J}$  is convex and therefore weakly lower semi-continuous. Consequently, through the application of classical critical point theory, we establish the existence of  $u \in W_0^{1,\mathcal{H}}(\Omega)$  as a minimizer of  $\mathcal{J}$ . This, in turn, implies that  $u$  constitutes a weak solution to problem (1.1). Furthermore, the strict convexity property of  $\mathcal{J}$  allows us to affirm the uniqueness of  $u$ . This completes the proof of Theorem 3.2.  $\square$

Now, let us comeback to study the existence of solutions to (1.1) when the source term is nonlinear. To this end, we shall assume that there are two nonnegative constants  $C_1$  and  $C_2$  such that

$$|f(x, r)| \leq C_1 + C_2 |r|^{\delta-1}, \quad (3.7)$$

for a.e  $x \in \Omega$  and for all  $r \in \mathbb{R}$ , where  $1 \leq \delta < p^-$ . We have the following existence result.

**Theorem 3.3.** *Assuming that (1.2), (1.3), (1.4), (2.7) and (3.7) are satisfied. Then, problem (1.1) has a weak solution  $u$  in the sense of Definition 3.1.*



**Proof.** We consider the energy functional  $\mathcal{K} : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$  associated to problem (1.1) as follows

$$\mathcal{K}(u) = \int_{\Omega} \frac{1}{p(x)} \langle A \nabla u, \nabla u \rangle^{\frac{p(x)}{2}} dx + \int_{\Omega} \frac{\omega(x)}{q(x)} \langle B \nabla u, \nabla u \rangle^{\frac{q(x)}{2}} dx - \int_{\Omega} F(x, u) dx, \quad (3.8)$$

where  $F(x, r) = \int_0^r f(x, s) ds$ . From (3.7), we can see that

$$|F(x, r)| \leq C \left(1 + |r|^\delta\right), \quad (3.9)$$

for a.e  $x \in \Omega$  and for all  $r \in \mathbb{R}$ . Furthermore, by setting  $\Psi(u) = \int_{\Omega} F(x, u) dx$ , it follows

that  $\Psi' : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \left(W_0^{1,\mathcal{H}}(\Omega)\right)^*$  is completely continuous, which means that if  $(u_n) \rightharpoonup u$  in  $W_0^{1,\mathcal{H}}(\Omega)$ , then  $\Psi(u_n) \rightarrow \Psi(u)$ . Consequently, the functional  $\Psi$  is weakly continuous. It is obvious that  $\mathcal{K}$  belongs to  $\mathcal{C}^1(W_0^{1,\mathcal{H}}(\Omega); \mathbb{R})$ . Hence, the weak solutions of (1.1) correspond precisely to the critical points of  $\mathcal{K}$ . Analyzing (3.8) while considering (1.3), (1.4) and (3.9), we can deduce that

$$\begin{aligned} \mathcal{K}(u) &\geq \min \left\{ \frac{\lambda_A}{p^+}, \frac{\lambda_B}{q^+} \right\} \int_{\Omega} \left( |\nabla u|^{p(x)} + \omega(x) |\nabla u|^{q(x)} \right) dx - C \int_{\Omega} |u|^\delta dx - C_3 \\ &\geq \min \left\{ \frac{\lambda_A}{p^+}, \frac{\lambda_B}{q^+} \right\} \int_{\Omega} \left( |\nabla u|^{p(x)} + \omega(x) |\nabla u|^{q(x)} \right) dx - C_4 \|u\|_{W_0^{1,\mathcal{H}}(\Omega)}^\delta - C_3. \end{aligned} \quad (3.10)$$

The latter inequality is derived through the use of Proposition (2.7). Employing (2.5), it follows from (3.10) that for any  $u \in W_0^{1,\mathcal{H}}(\Omega)$  with  $\|u\|_{W_0^{1,\mathcal{H}}(\Omega)} > 1$ , the following relationship holds:

$$\mathcal{K}(u) \geq \min \left\{ \frac{\lambda_A}{p^+}, \frac{\lambda_B}{q^+} \right\} \|u\|_{W_0^{1,\mathcal{H}}(\Omega)}^{p^-} - C_4 \|u\|_{W_0^{1,\mathcal{H}}(\Omega)}^\delta - C_3.$$

The fact that  $1 \leq \delta < p^-$  enables us to establish that

$$\mathcal{K}(u) \rightarrow +\infty \quad \text{as} \quad \|u\|_{W_0^{1,\mathcal{H}}(\Omega)} \rightarrow +\infty.$$

Thus,  $\mathcal{J}$  is coercive on  $W_0^{1,\mathcal{H}}(\Omega)$ . As  $\mathcal{K}$  is weakly lower semicontinuous, we conclude that  $\mathcal{K}$  possesses a minimum point, denoted as  $u$  in  $W_0^{1,\mathcal{H}}(\Omega)$ . In consequence, we infer that  $u$  is a weak solution to (1.1). This completes the proof.  $\square$

## Acknowledgements

The authors thank the anonymous reviewers for constructive and helpful remarks.

**Author contributions.** The authors contributed equally to this paper.

**Conflict of interest statement.** The authors read and approved the final manuscript. The authors have no relevant financial or non-financial interests to disclose.

**Funding.** Jinlan Pan is partially supported by the Natural Science Foundation of Guangxi Grant Nos. 2021GXNSFFA196004 and GKAD23026237, the NNSF of China Grant No. 12371312, the Startup Project of Doctor Scientific Research of Yulin Normal University No. G2023ZK13, and the project cooperation between Guangxi Normal University and Yulin Normal University. It is also supported by the European Union's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie grant agreement No. 823731 CONMECH.

**Data availability.** This paper has no associated data and material.

## References

- [1] H. Alaa, N. E. Alaa, A. Bouchriti and A. Charkaoui, *An improved nonlinear anisotropic model with  $p(x)$ -growth conditions applied to image restoration and enhancement*, Math. Meth. Appl. Sci. **47**(9), 7546–7575, 2024.
- [2] H. Alaa, N. E. Alaa and A. Charkaoui, *Time periodic solutions for strongly nonlinear parabolic systems with  $p(x)$ -growth conditions*, J. Elliptic Parabol. Equ. **7**, 815–839, 2021.
- [3] N. E. Alaa, A. Charkaoui, M. El Ghabi and M. El Hathout, *Integral Solution for a Parabolic Equation Driven by the  $p(x)$ -Laplacian Operator with Nonlinear Boundary Conditions and  $L^1$  Data*, Mediterr. J. Math. **20**(5), 244, 2023.
- [4] A. Alvino, V. Ferone and G. Trombetti, *On the properties of some nonlinear eigenvalues*, SIAM J. Math. Anal. **29**, 437–451, 1998.
- [5] A. Bahrouni, V. D. Rădulescu and D.D. Repovš, *Double phase transonic flow problems with variable growth: nonlinear patterns and stationary waves*, Nonlinearity, **32**, 2481–2495, 2019.
- [6] A. Bahrouni, V. D. Rădulescu and D. D. Repovš, *A weighted anisotropic variant of the Caffarelli-Kohn-Nirenberg inequality and applications*, Nonlinearity, **31**, 1516–1534, 2018.
- [7] P. Baroni, M. Colombo and G. Mingione, *Harnack inequalities for double phase functionals*, Nonlinear Anal. **121**, 206–222, 2015.
- [8] P. Baroni, M. Colombo and G. Mingione, *Regularity for general functionals with double phase*, Calc. Var. PDEs **57**, 48 pp. 2018.
- [9] S. S. Byun, J. Ok, K. Song, *Hölder regularity for weak solutions to nonlocal double phase problems*, J. Math. Pures Appl. **168**, 110–142, 2022.
- [10] J. X. Cen, S. J. Kim, Y. H. Kim and S. Zeng, *Multiplicity results of solutions to the double phase anisotropic variational problems involving variable exponent*, Adv. Differential Equ. **28**, 467–504, 2023.
- [11] M. Cencelj, V. D. Rădulescu and D. D. Repovš, *Double phase problems with variable growth*, Nonlinear Anal. **177**, 270–287, 2018.
- [12] A. Charkaoui, A. Ben-Loghfyry and S. Zeng, *A Novel Parabolic Model Driven by Double Phase Flux Operator with Variable Exponents: Application to Image Decomposition and Denoising*, Available at SSRN 4682810.
- [13] A. Charkaoui, *Periodic solutions for nonlinear evolution equations with  $p(x)$ -growth structure*, Evol. Equ. Control Theory **13**(3), 877–892, 2024.
- [14] A. Charkaoui and N. E. Alaa, *Existence and uniqueness of renormalized periodic solution to a nonlinear parabolic problem with variable exponent and  $L^1$  data*. J. Math. Anal. Appl. **506**(2), 125674, 2022.
- [15] A. Charkaoui and N. E. Alaa, *An  $L^1$ -theory for a nonlinear temporal periodic problem involving  $p(x)$ -growth structure with a strong dependence on gradients*, J. Evol. Equ. **23**, 73, 2023.

- [16] A. Charkaoui, A. Ben-Loghfyry and S. Zeng, *Nonlinear Parabolic Double Phase Variable Exponent Systems with Applications in Image Noise Removal*, Applied Mathematical Modelling, **132**, 495–530, 2024.
- [17] A. Charkaoui, H. Fahim and N. E. Alaa, *Nonlinear parabolic equation having nonstandard growth condition with respect to the gradient and variable exponent*, Opuscula Math. **41**, 25–53, 2021.
- [18] Y. Chen, S. Levine and M. Rao, *Variable exponent linear growth functionals in image restoration*, SIAM J. Appl. Math. **66**, 1383–1406, 2006.
- [19] M. Colombo and G. Mingione, *Regularity for double phase variational problems*, Arch. Ration. Mech. Anal. **215**, 443–496, 2015.
- [20] M. Colombo and G. Mingione, *Bounded minimisers of double phase variational integrals*, Arch. Ration. Mech. Anal. **218**, 219–273, 2015.
- [21] A. Crespo-Blanco, L. Gasiński, P. Harjulehto and P. Winkert, *A new class of double phase variable exponent problems: existence and uniqueness*, J. Differential Equ. **323**, 182–228, 2022.
- [22] C. De Filippis and G. Mingione, *Regularity for double phase problems at nearly linear growth*, Arch. Ration. Mech. Anal. **247**, 50 pp. 2023.
- [23] A. El Khalil, P. Lindqvist and A. Touzani, *On the stability of the first eigenvalue of the problem:  $A_p u + \lambda g(x)|u|^{p-2}u = 0$  with varying  $p$* , Rend. Mat. **24**, 321–336, 2004.
- [24] H. Fahim, A. Charkaoui and N. E. Alaa, *Parabolic systems driven by general differential operators with variable exponents and strong nonlinearities with respect to the gradient*. J Elliptic Parabol. Equ. **7**, 199–219, 2021.
- [25] X. Fan and D. Zhao, *On the spaces  $L^{p(x)}(\Omega)$  and  $W^{m,p(x)}(\Omega)$* , J. Math. Anal. Appl. **263**, 424–446, 2001.
- [26] G. Fragnelli, *Positive periodic solutions for a system of anisotropic parabolic equations*, J. Math. Anal. Appl. **367**, 204–228, 2010.
- [27] L. Gasiński and P. Winkert, *Existence and uniqueness results for double phase problems with convection term*, J. Differ. Equ. **268**, 4183–4193, 2020.
- [28] K. Ho and P. Winkert, *New embedding results for double phase problems with variable exponents and a priori bounds for corresponding generalized double phase problems*, Calc. Var. PDEs **62**, 38 pp. 2023.
- [29] I. H. Kim, Y. H. Kim, M. W. Oh and S. Zeng, *Existence and multiplicity of solutions to concave-convex-type double-phase problems with variable exponent*, Nonlinear Anal. **67**, 103627, 2022.
- [30] O. Kováčik and J. Rákosník, *On spaces  $L^{p(x)}(\Omega)$  and  $W^{1,p(x)}(\Omega)$* , Czechoslovak Math. J. **41**, 592–618, 1991.
- [31] W. Liu and G. Dai, *Existence and multiplicity results for double phase problem*, J. Differ. Equ. **265**, 4311–4334, 2018.
- [32] W. Liu and G. Dai, *Three ground state solutions for double phase problem*, J. Math. Phys. **59**, 121503, 2018.
- [33] P. Marcellini, *Regularity of minimizers of integrals of the calculus of variations with non standard growth conditions*, Arch. Ration. Mech. Anal. **105**, 267–284, 1989.
- [34] P. Marcellini, *Regularity and existence of solutions of elliptic equations with  $p, q$ -growth conditions*, J. Differ. Equ. **90**, 1–30, 1991.
- [35] M. Mihăilescu and V. Rădulescu, *A multiplicity result for a nonlinear degenerate problem arising in the theory of electrorheological fluids*, Proc. R. Soc. A **462**, 2625–2641, 2006.
- [36] M. Mihăilescu and D. Repovš, *On a PDE involving the  $A_{p(\cdot)}$ -Laplace operator*, Nonlinear Anal. **75**, 975–981, 2012.
- [37] N. S. Papageorgiou, V. D. Rădulescu and D. D. Repovš, *Double-phase problems and a discontinuity property of the spectrum*, P. Am. Math. Soc. **147**, 2899–2910, 2019.

- [38] V. D. Rădulescu and D. D. Repovš, *Partial Differential Equations with Variable Exponents: Variational Methods and Qualitative Analysis*, CRC Press Taylor and Francis Group, 2015.
- [39] K. Rajagopal, *Mathematical modelling of electrorheological fluids*, Contin. Mech. Thermodyn. **13**, 59–78, 2001.
- [40] Y. G. Reshetnyak, *Set of singular points of solutions of certain nonlinear elliptic equations*, Sibirsk. Mat. Zh. **9**, 354–368, 1968. (in Russian).
- [41] M. Ruzicka, *Electrorheological Fluids: Modeling and Mathematical Theory*, Springer Science and Business Media, 2000.
- [42] S. Zeng, N. S. Papageorgiou and P. Winkert, *Inverse problems for double-phase obstacle problems with variable exponents*, J. Optim. Theory Appl. **196**, 666–699, 2023.
- [43] S. Zeng, V. D. Rădulescu and P. Winkert, *Double phase obstacle problems with variable exponent*, Adv. Differential Equ. **27**, 611–645, 2022.
- [44] S. Zeng, V. D. Rădulescu and P. Winkert, *Double phase implicit obstacle problems with convection and multivalued mixed boundary value conditions*, SIAM J. Math. Anal. **54**, 1898–1926, 2022.
- [45] Q. Zhang and V. D. Rădulescu, *Double phase anisotropic variational problems and combined effects of reaction and absorption terms*, J. Math. Pures Appl. **118**, 159–203, 2018.
- [46] V. V. Zhikov, *Averaging of functionals of the calculus of variations and elasticity theory*, Izv. Ros. Akadem. Nauk. **50**, 675–710, 1986.
- [47] V. V. Zhikov, *On Lavrentiev's phenomenon*, Russian J. Math. Phys. **3**, 2 pp. 1995.
- [48] V. V. Zhikov, *On some variational problems*, Russian J. Math. Phys. **5**, 105 pp. 1997.