

## Salmonellae in the air environment: A review

Abdel Hameed A. Awad

Air Pollution Research Department, Environment and Climate Change Research Institute, National Research Centre, Dokki, Giza, EGYPT  
e-mail: [abed196498@yahoo.com](mailto:abed196498@yahoo.com), ORCID: [orcid.org/0000-0003-0784-9835](https://orcid.org/0000-0003-0784-9835)

### Cite this article as:

Awad, A.H.A. 2024. Salmonellae in the air environment: A review. *Trakya Univ J Nat Sci*, 25(2): 211-227, DOI: 10.23902/trkjinat.1433216

Received: 07 February 2024, Accepted: 01 July 2024, Online First: 06 August 2024, Published: 15 October 2024

**Abstract:** *Salmonella* bacteria, a zoonotic pathogen, are frequently transmitted through food and water, causing foodborne outbreaks and illnesses. Bioaerosols are a growing concern as pathogenic microorganisms could be transmitted to the indoor and ambient air environments. The airborne transmission of pathogenic microorganisms is considered a risk of contamination or a route of infection. *Salmonella* have been found in rare numbers in the air, but their detection indicate their ability to survive in the air environment. Physical, biological and environmental stressors affect the survival of airborne microorganisms. The infectivity of airborne *Salmonella* is determined by its pathogenicity, infective dose and individual health conditions. The accurate assessment of *Salmonella* in aerosols is a problem due to the synergistic influence of many uncontrollable environmental conditions and a lack of standardized analysis and sampling protocols. Knowledge of the airborne transmission of *Salmonella* and factors influencing their viability is critical to understanding their potential health risk and the related control measures. This review provides evidence for the transmission of *Salmonella* in different air environments, focusing on the presence of *Salmonella* in the air as a risk of biocontamination. The sampling, detection and enumeration methodologies of *Salmonella* in the air are discussed with recommended mitigation and control strategies.

**Edited by:**  
Mustafa Yamaç

**Key words:**  
Air environment  
Salmonella  
Survival  
Waste applications  
Livestock house  
Survivability  
Bio-indicator

**Özet:** Zoonotik bir patojen olan *Salmonella* cinsi bakteriler sıklıkla gıda ve su yoluyla bulaşarak gıda kaynaklı salgınlara ve hastalıklara neden olmaktadır. Patojenik bakterilerin hava ortamına bulaşabilmesine aracılık ettikleri için biyo-aerosoller giderek artan bir sorun olarak ele alınmaktadır. Patojenik mikroorganizmaların hava yoluyla bulaşması, kontaminasyon veya enfeksiyon riski olarak kabul edilir. *Salmonella*'nın havada az sayılarda bulunması, hava ortamında hayatta kalma yeteneklerini göstermektedir. Fiziksel, biyolojik ve çevresel stres etkenleri havadaki mikroorganizmaların hayatta kalmasını etkileyen faktörlerdir. Hava ortamında bulunan *Salmonella* üyelerinin bulaşıcılığı patojeniteleri, enfektif doz ve bireylerin sağlık koşullarınca belirlenir. Aerosollerle taşınan *Salmonella* üyelerinin doğru bir şekilde değerlendirilmesi, kontrol edilemeyen birçok çevresel koşulun sinerjik etkisine ve standartlaştırılmış analiz ve numune alma protokollerinin eksikliğine bağlı bir sorun olarak görülmektedir. *Salmonella* üyelerinin hava yoluyla bulaşması ve canlılıklarını etkileyen faktörlerin bilinmesi, potansiyel sağlık risklerinin ve ilgili kontrol önlemlerinin anlaşılması açısından kritik öneme sahiptir. Bu derleme, biyolojik kontaminasyon riski olarak havadaki *Salmonella* varlığına odaklanarak *Salmonella* üyelerinin farklı hava ortamlarında bulaştığına dair kanıtlar sunmaktadır. Hava ortamında bulunan *Salmonella* üyelerinin örnekleme, tespit ve sayımı metodolojileri, önerilen azaltma ve kontrol stratejileriyle birlikte tartışılmıştır.

### Introduction

Aerosols are ubiquitous in the earth's atmosphere and they are central to many environmental issues and public health (Colbeck & Lazaridis 2010, Zhang 2020). Atmospheric aerosols are suspensions of liquid, solid or mixed particles with highly variable chemical composition and size distribution (Putaud *et al.* 2010). Bioaerosols are particles of biological origin (e.g.

bacteria, viruses, fungi, algae, biological fragments and pollen) suspended in the air and are an important part of aerosols (Wéry 2014, Smets *et al.* 2016). Bioaerosols, which considerably vary in composition and size (0.2-100 µm) (Stetzenbach 2009), are produced in the environment from a variety of natural and anthropogenic sources (Kim *et al.* 2018, Xie *et al.* 2021), affecting living organisms



OPEN ACCESS

through infectivity, allergenicity and toxicity (Cox & Wathes 1995) (Fig. 1). Moreover, bioaerosols could be a source of pollution for plants, animals and surface water (Michalkiewicz, 2019). Biological particles are transported up in the air as free (single cells, spores or aggregates) or attached to non-biological particles (Jones & Harrison 2004), thus leading to considerable differences in their stability, survivability, composition and dispersal mechanisms (Cambrá-López *et al.* 2010). Sewage treatment plants, biosolid landfills, spray irrigation (untreated / or insufficiently treated), wastewater (Brooks *et al.* 2004), composting, livestock facilities and herb processing have been considered as potential sources of bioaerosols and pathogenic microorganisms (Hickey & Reist 1975, Skórska *et al.* 2005, Zhang *et al.* 2019, Dai *et al.* 2020). Transmission of pathogenic microorganisms is of a great concern due to their ability to affect worker's and the nearby residents' health.


The transmission of pathogenic microorganisms in the atmosphere has to be paid attention, as their transmission is attributed to the initial health symptoms resembling enteric diseases among workers and population living near sewage treatment plants, biosolid landfills, composting and livestock facilities (Chinivasagam *et al.* 2009). The detection of pathogenic microorganisms in the air environment indicates their ability to persist harsh atmospheric conditions. Nowadays, the potential of aerosolization of pathogenic microorganisms has become a debated issue. The available information on emission, source apportionment and transmission of pathogenic microorganisms into the air environment is scarce (Xie *et al.* 2021). This review aims to highlight the transmission of *Salmonella* in the air environment, factors affecting

their survivability, sampling and analysis methods and control strategies.

### Salmonella bacteria

Salmonellae belong to *Enterobacteriaceae*, a family of Gram-negative bacteria represented with facultative anaerobic bacilli with 2-5 µm long and 0.5-1.5 µm wide and are motile by peritrichous flagella (Andino & Hanning 2015). *Salmonella* grow at temperatures in the range of 5-45°C, with ideal temperatures between 35-37°C, but some species can grow at temperatures as high as 54°C and as low as 2°C (Gray & Fedorka-Cray 2002) and at optimum pH range of 6.5 and 7.5 (Shaji *et al.* 2023). *Salmonella* can be distinguished from other bacterial species by their biochemical and antigenic features. Salmonellae are a complex group containing ≥2600 serovars based on somatic (O), flagellar (H) and surface capsule (Vi) antigens (Mumy 2014).

Salmonellae are ubiquitous human and animal pathogens and can be divided into 2 groups, typhoidal *Salmonella* (TS) and Non-typhoidal *Salmonella* (NTS) (Wang *et al.* 2023a). *Salmonella enterica* ser. Enteritidis (*S. Enteritidis*) and *Salmonella enterica* ser. Typhimurium (*S. Typhimurium*), belonging to NTS group, are responsible for the majority of human salmonellosis (Ashurst, *et al.* 2022). NTS group is responsible for ~ 93 million cases of gastroenteritis and 155,000 fatalities annually and is frequently zoonotic (Gordon 2011, Cosby *et al.* 2015). The natural habitat of *Salmonella* is the gastrointestinal tract of humans and animals. Historically, transmission of *Salmonella* and enteric zoonotic infections (e.g. Q-fever, brucellosis, and avian and swine influenza) via aerosols has been neglected (Kallapura *et al.* 2014).



**Fig. 1.** Diagram of bioaerosols emission sources and fate (Krishnamoorthy *et al.* 2020).

Transmission of *Salmonella* aerosols is less pathogenic and rarely occurs (Shuval *et al.* 1986). The possibility of transmission and survival of *Salmonella* (Oliveira *et al.* 2006, López *et al.* 2012) in aerosols should be considered.

*Salmonella* aerosol is a concern route for vegetables contamination and foodborne outbreaks. *Salmonella* directly enter water and agricultural environments via waste and sewage irrigation (Heaton & Jones 2008) or indirectly via *Salmonella* aerosols. *Salmonella*, pathogenic *Escherichia coli* and *Listeria monocytogenes* have been linked to bacterial outbreaks of foodborne diseases associated with ready-to-eat fruit and vegetables (Thomas *et al.* 2024).

A low number of *Salmonella* cells may be sufficient to cause disease in a large number of people (Werber *et al.* 2005). For instance, ~ 13 CFU/g is enough to cause salmonellosis outbreaks. Infectious dose of outbreaks of salmonellosis is found between 10 and 1,000 cells (Blaser & Lee 1982, Vought & Tatini 1998). The infectious dose of *Salmonella* via respiratory pathway is lower than the oral route (Darlow *et al.* 1961). Inhalation of *S. Typhimurium* by mice was reported to cause disease in animals in a dose dependent manner, where the lowest dose was reported as ~150 CFU that could produce a disease (Wathes *et al.* 1988).

#### Sources of *Salmonella* aerosols

##### Wastewater and sludge applications

The primary concern of wastewater treatment plants (WWTPs) is to remove contaminants and inactivate pathogenic organisms to protect environment and human health. The enteric bacteria, viruses, protozoa and helminths are the common groups of microorganisms present in municipal wastewater (Akin *et al.* 1978). The enteric bacteria are commonly found in wastewater, with *Escherichia coli* and *Enterococcus faecalis* frequently at concentrations of ~10<sup>9</sup>/l and ~10<sup>8</sup>/l, respectively. However, concentrations of *E. coli* and total coliform are significantly 1 to 3 folds higher in the influent than the effluent (Ajonina *et al.* 2015). Salmonellae are the most prevalent pathogenic bacterial species in raw wastewater with a concentration ~ 5000 bacteria/l (Foster and Engelbrecht 1973). *Salmonella* concentrations averaged 130 bacteria/100 ml in the raw sewage water and 3 bacteria/100 ml in the treated sewage water (Langeland 1982). The presence and concentration of pathogenic microorganisms in sewage are determined by their prevalence among other populations and their ability to persist treatment processes. Biological wastewater treatment plants leave ~1-10% of *Salmonella*, *Mycobacterium* and viruses in the treated wastewater (Sorber & Sagik 1979). The aerosols containing pathogenic and non-pathogenic microorganisms are generated during wastewater treatment processes, as wastewater undergoes turbulent mixing or mechanical agitation (Sorber & Guter 1975, Sánchez-Monedero *et al.* 2008, Liu *et al.* 2020).

The transportation of bioaerosols is a function of time and distance (Pepper & Gerba 2015). Concentrations of total aerobic bacteria, total and fecal coliforms, fecal enterococci and coliphage were reported to significantly increase in the air within the perimeter of a WWTP in USA after operation (Fannin *et al.* 1985). Aeration tanks and trickling filters at WWTPs are the main sources emitting microorganisms into the ambient air (Han *et al.* 2020). High airborne microbial concentrations were found near aeration tanks and mechanical agitation, ranging within few to more than 8x10<sup>4</sup> CFU/m<sup>3</sup> (Korzeniewska *et al.* 2008). Concentrations of mesophilic bacteria, bacteria-associated certain waterborne virulence factors, mesophilic fungi and thermophilic fungi were 1.7 × 10<sup>4</sup> CFU/m<sup>3</sup>, 2 × 10<sup>3</sup> CFU/ m<sup>3</sup>, 1.7 × 10<sup>3</sup> CFU/m<sup>3</sup> and 4.5 × 10<sup>1</sup> CFU/m<sup>3</sup>, respectively in aerosols emitted by aeration tanks of an activated sludge plant (Bauer *et al.* 2002).

Airborne Gram negative bacteria, fecal indicators (*E. coli* & *Clostridia*), *Salmonella* and *P. aeruginosa* were detected at WWTP but in lower counts than Gram positive bacteria. *Escherichia coli* and *Salmonella* were found up to a distance of 300m and 10 m downwind of the aeration tanks, respectively, and a higher number of positive findings were observed during higher wind velocity and low sunshine (Müller 1980). Coliforms were found up in the air to a distance of 0.8 mile downwind of trickling-filter sewage treatment plant (Adams & Spendlove 1970). The dissemination of *Salmonella* by the air was low in relation to *Salmonella* content of the sewage itself (Müller 1980).

A microorganism can be released into the air from aerated sewage only when its concentration exceeds 10<sup>3</sup> cells/cm<sup>3</sup> in the sewage (Teltsch *et al.* 1980), a higher number of a given microorganism in sewage has a higher aerosol emission rate (Sawyer *et al.* 1993). Composition of airborne microflora is closely related with the type and number of microorganisms present in sewage waste (Ossowska-Cypryk 1991). The majority of the released aerosols do not travel very far distances. However, smaller particles tend to travel a considerable distance away from the source point (Mckinney 2004). The composition and size of microbial aerosols are influenced by type of treated wastewater, treatment technology, ambient conditions and shear stress force (Heinonen-Tanski *et al.* 2009). The highest emission of *P. fluorescens*, *E. coli*, *Enterococcus* sp. and *Salmonella* was detected in the air at the first stage of the purification in a municipal wastewater plant, Toruń, Poland (Paluszak *et al.* 2003). *Salmonella* and *Shigella* were not isolated from the air samples despite their presence in sewage water (Sekla *et al.* 1980).

Airborne microbial contamination greatly differed in the vicinity of aeration tank, maturing composting plant and 100 m downwind of municipal treatment plant in Poland, where the concentrations of *E. coli*, *Enterobacter* and *Salmonella* were ~10<sup>1</sup>CFU/m<sup>3</sup> in average (Breza-Boruta & Paluszak 2007). The highest microbial air contamination was found in the pretreatment of wastewater (screening, aerated grit removal and pumping) in a WWTP in Finland, where somatic coliphage and

enterococci were found in higher numbers and no *Salmonella* bacteria were detected (Heinonen-Tanski *et al.* 2009). The ratio between *Salmonella* to coliphage densities in sewage aerosols was 1:100,000 (Grunnet & Tramsen 1974) and *Salmonella* bacteria were not recovered in any of the air samples collected at a WWTP in Egypt (Abdel Hameed 1992). Airborne pathogenic enteric bacteria (*S. Enteritidis* and *S. Boydii*), reovirus and enterovirus were isolated in 2%, 46% and 9%, respectively of the total samples collected at different sites in sewage sludge treatment plants in Italy (Carducci *et al.* 2000).

A given quantity of pathogens present in sewage aerosols could represent a source of a threat to workers who are daily exposed to aerosols associated with a variety of infectious microorganisms (Grisoli *et al.* 2009). Wastewater treatment processes bring the workers in contact with multiple pathogens and infectious agents such as viruses (*Hepatitis-A*, *Polio*, *Coxsackie*, *Echo*, *Rota* and *Adeno*), bacteria (*Salmonella* spp., *Shigella* spp., *Campylobacter jejuni*, *Yersinia enterocolitica*, *Legionella pneumophila*, *Helicobacter pylori*, *Listeria monocytogenes* and *Mycobacterium xenopi*) and protozoa (*Giardia lamblia*, *Entamoeba histolytica* and *Helminthes*) (Mulloy 2001).

Municipal sewage sludge is utilized worldwide on agricultural lands to solve the problem of sewage disposal, water scarcity and environmental contamination. However, the increase of wastewater in land application has magnified problems such as production of aerosols containing pathogens and contamination of crop and ground and surface waters (Bitton 1980). In the United States, ~33% of the produced municipal sludge is applied onto agricultural lands (Mclamarra & Pruitt 1995), increasing accumulation of pathogens and toxic substances that may be released into the air environment.

Climatic and environmental factors differently affect the survival of airborne pathogenic bacteria. Temperature, relative humidity, oxygen content, UV radiation and reactive chemical radicles are the main factors affecting viability of airborne microorganisms (Ruiz-Gil *et al.* 2020). *Salmonellae* in sewage sludge spread on grass and may survive up to 72 weeks, and neither aerobic stabilization nor anaerobic digestion significantly reduces the contamination with *Salmonellae* (Hess & Breer 1975). Raw sludge from municipal sewage may release more airborne pathogens than aerobic/or anaerobic digestion, lime stabilization and thermal drying sludge (Straub *et al.* 1993). The application of raw sludge on agricultural lands has been prohibited in many countries due to its hazardous effects that may be presented by direct contact/or inhalation of infectious aerosols (Cole *et al.* 1999).

Low concentrations of *Salmonella*, coliforms and enteroviruses were detected in air samples collected downwind wastewater spray -irrigated fields. *Salmonella* was detected in 78% and 18% of wastewater and air samples ~40m downwind, respectively, and enteroviruses in 71% and 44% in wastewater and aerosols, respectively,

as an indication of the prevalence of enteroviruses than *Salmonella* in aerosols (Teltsch *et al.* 1980). This is attributed to viral contamination may be more resistance to inactivation processes than enteric bacteria and may be concentrated in aerosols than suspending fluid (Baylor *et al.* 1977). Table 1 shows the concentrations of *Salmonella* bacteria in wastewater and aerosols at WWTPs.

#### Biosolids and composts

Composting is used to stabilize biosolids, as organic substrates are subjected to microbial degradation. Composting produces substrates suitable for cultivation or aids in the disposal of wastes (Fig. 2). Application of composted sludge improves soil quality, but the microbiological safety should be considered (Brooks *et al.* 2005). The risk of infection posed to biosolid handlers reached 34% and 2% annually from exposure to Coxsackievirus A21 and *Salmonella*, respectively (Tanner 2004).

Growth and death rate of pathogens in biosolids, including *Salmonella*, depend on several factors such as moisture content, temperature, available nutrient, associated flora and indigenous microorganisms (Sidhu *et al.* 2001). Most of enteric pathogenic bacteria are non-spore formers and relatively sensitive to environmental factors (Vilanova & Blanch 2005). *Salmonella*, *E. coli* and fecal coliforms can regrow in moist conditions after treatment (Lang *et al.* 2007).

**Table 1.** Concentrations of *Salmonella* in wastewater and aerosols at WWTPs.

Environment	Concentration	Reference
Wastewater	2-60 MPN/100 ml	Katzenelson & Teltsch (1976)
Wastewater	<i>Salmonella</i> : coliforms 2:60 MPN/100 ml	
Aerosols	<i>Salmonella</i> : coliforms $3.2 \times 10^{-2}$ : $5.410^{-2}$ MPN/m <sup>3</sup> 43:1076 CFU/m <sup>3</sup>	Teltsch <i>et al.</i> (1980)
Dry sewage sludge	140-14000 CFU/100gm	Langeland (1982)
Raw wastewater	130 bacteria/100 ml	
Treated wastewater	3 bacteria/100 ml	
Bulk sludge	0.3-17000 CFU/gm	Hussong <i>et al.</i> (1985)
Raw sewage	5000 CFU/ml	Prazmo (1980)
Aerosols/aeration tank	$\leq 10^1$ CFU/m <sup>3</sup>	Breza-Boruta & Paluszak (2007)
Aerosols	<i>Salmonella</i> : Coliphage 1:100,000	Grunnet & Tramsen (1974)
Aerosols	$\leq 1$ CFU/m <sup>3</sup>	Heinonen-Tanski <i>et al.</i> (2009), Abdel Hameed (1992), Pillai <i>et al.</i> (1996)

The bacterial concentrations were reported to range between  $10^4$ - $10^6$  CFU/g in a well-managed compost, decreased over time to 150 CFU/g and increased over 6 weeks in poorly managed composts (Ogden *et al.* 2001). Wastewater biosolids generally contain *Salmonella* at a range of  $10^2$ - $10^3$  CFU/g dry weight (Epstein 1997) and  $\sim 10^5$  CFU/g in dewatered anaerobically digested sludge (Russ & Yanko 1981). *Salmonella* bacteria are known to survive composting process in low concentration (Gibbs *et al.* 1997) and can form filaments under moderately low-water conditions and upon rehydration can achieve high bacterial loads within a short period of time (Stackhouse *et al.* 2012). The active indigenous flora of compost establishes a homeostatic barrier against *Salmonella* which is considered an invader. However, in the absence of indigenous compost flora, the inoculated *Salmonella* may grow to potentially hazardous levels (Sidhu *et al.* 2001).

Microorganisms are released into the air when compost piles are formed or dismantled. The potential of aerosolization of pathogenic microorganisms from biosolids has become an important debated issue worldwide. The nature of the airborne microflora depends on the existing contamination of the starting materials and microbial development between disposal and composting (Lacey *et al.* 1996). The elevated temperature in composting kills-off coliforms and pathogens, however inadequate compost turning leads to temperature stratification and survival of pathogens (*Salmonella*) in cooler layers which may be emitted into the air during mechanical agitation/ or by wind action (Millner *et al.* 1980).

The biosolid land application generates bioaerosols through soil agitation and weathering of biosolid. Biosolids left on the soil surface are subjected to drying; rendering it friable, becoming airborne with the associated pathogens (Pillai 2007). At a municipal solid waste recycling and composting plant stations in Quebec, Canada, the concentrations of airborne total culturable bacteria and Gram-negative bacteria were above  $10^4$  CFU/m<sup>3</sup> and  $10^3$  CFU/m<sup>3</sup>, respectively, at six of the nine work stations (Marchand *et al.* 1995). *Salmonella* and *Enterobacter* bacteria were found in the air samples only in the vicinity of the compost piles in Poland (Brezab-Boruta & Paluszak 2007). The generation and disposal of bio-wastes potentially increase aerosolization of a wide variety of microbial pathogens.



**Fig. 2.** Photograph of a drying sewage sludge used as fertilizer.

### Livestock houses

Livestock houses have significant hazards to biocontamination of food (Hutchison *et al.* 2004), water (Devane *et al.* 2018) and soil (Nolan *et al.* 2020). Pathogenic microorganisms are shed in animals' excretions, secretions or exhaled in breath, litter (e.g. straw, sawdust or wood chippings) and feed (Chien *et al.* 2011). Poultry litter and manure can pose a serious threat to environmental and human health and need to be managed properly (Gržinić *et al.* 2023). *Salmonella* bacteria are ubiquitous in farm environment, and bioaerosols may be released into the air environment as free/ or associated dust particles (Zhao *et al.* 2014). In agricultural livestock farming, bioaerosols account for well over 90% of airborne dust (Aengst 1984), reaching  $\sim 10^7$  CFU/m<sup>3</sup> (Dungan 2010). The concentration of airborne total bacteria was 6.43 log CFU/m<sup>3</sup> in broiler houses, 5.1 log CFU/m<sup>3</sup> in pig buildings and 4.3 log CFU/m<sup>3</sup> in cattle buildings, and the overall concentrations of *Enterobacteriaceae* ranged between 3 and 4 log CFU/m<sup>3</sup> (Seedorf *et al.* 1998). In animal houses, the majority of airborne microbial composition is non-pathogenic and Gram-negative bacteria constituted 0.02 and 5.2% of the total amount of aerobic bacteria (Zucker *et al.* 2000).

There is evidence that enteric pathogens are important in airborne transmission of diseases among animals (Pepper & Gebra 2015). *Salmonella* Typhimurium aerosols are transmitted among calve houses (Hinton *et al.* 1983). *Salmonella* Typhimurium could survive for long periods in the air, and calves and mice exposed to *Salmonella* developed gastrointestinal symptoms, proving that pathogens could be spread by aerosolization (Wathes *et al.* 1988). *Bordetella bronchiseptica*, *Brucella suis*, *Haemophilus* spp., *Corynebacterium equi*, *Listeria monocytogenes*, *Mycobacterium* spp., *Mycoplasma* spp., *Pasteurella* spp., *Escherichia coli*, *Salmonella* spp., *Staphylococcus aureus*, *Streptococcus suis* and *Leptospira Pomona* are some pathogenic bacteria in pigs and poultry houses that can be airborne/or aerosol transmitted (Wathes 1995).

The aerosolized *Salmonella* Enteritidis could infect laying hens (Baskerville *et al.* 1992). *Salmonella* colonized and persisted in different tissues in broilers following exposure to aerosolized *Salmonella* (Pal *et al.* 2021). Chickens and animals are direct or indirect sources of *Salmonella* through feces and dust (Venter *et al.* 2004, Gale & Velazquez 2020). *Salmonella* infection has been experimentally proven to occur by oral, intraloacal, intratracheal, intraocular, navel and aerosol administration (Cox *et al.* 1990). The hatchery is the most significant contributor of *Salmonella* with a prevalence of 48.5%. Litter, feces, and indoor environment of poultry house are the other 3 major contributing factors with prevalences of 25.4, 16.3, and 7.9%, respectively (Wang *et al.* 2023b).

Cross-contamination of *Salmonella* from contaminated to uncontaminated eggs could be spread by fan-driven air (Berrang *et al.* 1995). *Salmonella* Typhimurium DT104 strain could be efficiently

transmitted to eggs up to 15 times more when laying hens were inoculated via aerosol route than crop route (Leach *et al.* 1999). Airborne transmission of *Salmonella enterica* serovar Typhimurium was demonstrated in chicks hatching in a cabinet containing infected hatchmates (Cason *et al.* 1994). *Salmonella* Enteritidis from infected chicks in an isolation cabinet rapidly transmitted to chicks physically separated from the infected seeder chicks but sharing the same air (Gast *et al.* 1998). However, the transmission mechanism of airborne *S. Enteritidis* has not been fully defined in chick's cabinets. *Salmonella* move through the air by itself or via water droplets, dust, dander or feathers need further studies (Cox *et al.* 1990, Davies & Wray 1996, Holt *et al.* 1999). These types of media may have important role in the transmission process of Salmonellae. Higher *Salmonella* and *E. coli* concentrations were detected in litter samples with water activity  $\geq 0.90$  and moisture  $\geq 35\%$  in a broiler litter (de Rezende *et al.* 2001). Airborne bacterial levels were linked to their densities in litter/or dust, *Salmonella* and *E. coli* averaged  $10^4$  MPN/g and  $\sim 10^8$  CFU/g of litter, respectively (Davies & Wray 1994). *Salmonella* concentrations ranged between  $10^3$ - $10^5$  MPN/g in litter and  $2.2 \times 10^{-1}$ -  $44 \times 10^{-1}$  MPN/m<sup>3</sup> inside the air environment of a poultry house (Chinivasagam *et al.* 2009). The prevalence of Salmonellae isolated from both poultry farm and processing plant environments were 5.4% and 4.7%, respectively with no *Salmonella* bacteria detected in the air samples (Alzenki *et al.* 2007). *Salmonella* Typhimurium was detected in the air at dehairing and evisceration locations in an Irish pig slaughtering plant (Pearce *et al.* 2006).

The number of total airborne aerobic bacteria and Gram negative bacteria varied between 780 and 20100 CFU/m<sup>3</sup> and 39 and 1030 CFU/m<sup>3</sup>, respectively, in Chinese rabbit houses (Duan *et al.* 2006). The median of airborne mesophilic bacteria at the processing area of the moving rail was  $1.7 \times 10^6$  CFU/m<sup>3</sup> with no *Salmonella* species detected from the air samples at a poultry house in Styria, Australia (Haas *et al.* 2005). On the other hand *Salmonella* bacteria only represented  $\sim 0.56\%$  of the total airborne bacterial colonies in a small poultry house in Egypt (Abdel Hameed *et al.* 2010). *Salmonella* were identified in 10% of total airborne bacterial colonies, with *S. Choleraesuis*, *S. Typhi* and *S. Typhimurium*

constituting 5.5% of the total bacterial counts in three pig and three beef plants in USA (Cosenza-Sutton 2004). The concentration of airborne *Salmonella* in a poultry production unit constituted up 3.3% of total bacterial cell counts measured by 4, 6-diamidino 2-phenylindole, ranging from  $2.2 \times 10^1$  to  $3 \times 10^6$  *Salmonella* targets/m<sup>3</sup> using *Salmonella*-specific *invA* genes of DNA (Fallschissel *et al.* 2009).

*Salmonella* bacteria have been isolated from the settled dust within unoccupied poultry shed as a result of the residual effect (Chinivasagam *et al.* 2009). *Salmonella* can survive for  $\sim 53$  weeks in dust (Davies & Wray 1994) and 26 months in thin layers of litter of dried feces and feed (Davies & Breslin 2003). Pathogenic microorganisms were found in low counts, suggesting that air environment is not a significant source of enteric biocontamination. However, the transmission of airborne *Salmonella* within the livestock environment may impact the bird's and worker's health. Table 2 shows levels of *Salmonella* in raw and aerosols at livestock facilities.

#### Factors influencing the survival of *Salmonella* aerosols

The air environment is not an optimal medium for the survival of microorganisms. Aerosolized pathogenic bacteria are subject to considerable stressors leading to cell injury and/or death in both Gram positive and negative bacteria (Heidelberg *et al.* 1997). The persistence of airborne microorganisms depends on their tenacity. The tenacity (the ability to survive the airborne) of different microbial species depends on meteorological factors (temperature and humidity, UV radiation, solar radiation), air pollution, free radicals and ozone-olefin reaction products (Open Air Factor, OAF) (Stärk 1999, Clauss *et al.* 2016). These factors lethally affect microbial viability and infectivity through chemical, physical and biological modifications to phospholipid, protein and nucleic acid moieties (Karra & Katsivella, 2007).

Gram negative bacteria, including Salmonellae, are rapidly die-off in the airborne state (Cox 1995). Some bacteria (anaerobic species) are highly sensitive and cannot grow in the presence of oxygen (Tang 2009). Desiccation is experienced by Gram-negative bacteria,  $\sim 90\%$  immediately loss their viability after aerosolization, due to denaturation of outer phospholipid bilayer membranes (Cox 1989).

**Table 2.** Levels of *Salmonella* in raw and aerosols at livestock facilities

Environment	Level	Reference
Swine house units	No <i>Salmonella</i>	Elliott <i>et al.</i> (1976)
Chick dust	$10^4$ CFU/g	Davies & Wray (1994)
Chick dust	$10^3$ - $10^5$ MPN/g	Chinivasagam <i>et al.</i> (2009)
Poultry house- air	0.22 - 4.4 MPN/m <sup>3</sup>	
Poultry house-air (DAPI)	$2.8 \times 10^5 \pm 1.9 \times 10^5$ cell/ m <sup>3</sup>	Fallschissel <i>et al.</i> (2009)
Poultry house-air (culture method)	$3.3 \times 10^2 \pm 1.2 \times 10^2$ CFU/m <sup>3</sup>	
Duck stalls (molecular method)	$2.5 \times 10^1$ - $3 \times 10^6$ genes/m <sup>3</sup>	
Small poultry house	$\sim 0.56\%$ of total bacterial isolates	Abdel Hameed <i>et al.</i> (2010)
Poultry house- picking area	2 - 598 CFU/m <sup>3</sup>	Heber <i>et al.</i> (2006)

Microorganisms generated from liquid suspension undergo desiccation (loss of water) and those generated as dust particles partially rehydrated (Cox 1995, Cox & Wathes 1995).

Long distance transport of microorganisms in the air depends on atmospheric dispersion, dilution, deposition, particle size and meteorological conditions (Gregory 1973). The immission concentrations of bioaerosols decreased exponentially with increasing distance from the source of emission. In the air environment, bioaerosols are exposed to wind and weather and their extent depends on the tenacity, size and composition of bioaerosol particles (Clauß 2020). The behavior of *Salmonella* in the air environment remains unpredictable (Carrique-Mas & Davis 2008). Temperature affects the molecular structure of the microorganism and consequently its inherent thermodynamic instability (Maillard reaction), involving the elimination of water molecules (Stärk 1999). At warmer temperatures, phospholipid membranes of Gram negative bacteria undergo many complex transition, separation and aggregation phases, leading to changes in biological functions. However, at cooler temperature, exothermic crystallization of lipid moieties together with protein subunit formation leads to loss of viability (Cox 1989). The effect of relative humidity on airborne microorganisms is difficult to determine, however surface damage (inactivation at high RH) and rehydration (inactivation at low RH) are the most influential factors (de Jong *et al.* 1973).

Airborne *Salmonella* are affected by sunlight and other environmental factors, because *Salmonella* bacteria are enteric microbes, adapting to live in a protected environment (Müller 1980). Aerosol particles play a crucial role in the transmission of airborne bacteria, as particles may protect microorganisms from harsh environmental conditions. A significant positive relationship was found between concentrations of aerosol sizes of 0.5-1.0µm and *Salmonella* species in a dairy house (Aminul Islam *et al.* 2020).

Climate change and global warming have contributed to the spread of pathogens. Several studies have recognized the importance of increased ambient temperature and precipitation in the spread and persistence of *Salmonella* in soil and food. The impact of extreme weather events on *Salmonella* infection rates among the most prevalent serovars has not been evaluated worldwide (Jiang *et al.* 2015; Morgado *et al.* 2021). Dust storms have positive (e.g. fertilization of aquatic and terrestrial ecosystems) and negative (e.g. transport of toxins and pathogenic microorganisms) effects. *Salmonella* proliferate rapidly at higher temperature, increasing their spread through different environmental media (Akil *et al.* 2014). Emergence or resurgence of numerous infectious diseases is influenced by environmental factors such as climate or land use change (Mills *et al.* 2010). However, the impact of extreme weather events on *Salmonella* growth and persistence in the air environment should be fully evaluated.

### Airborne Salmonella: Sampling and analysis techniques

The detection of airborne pathogenic bacteria is of great concern. The efficiency of collection depends on the sampling strategy, analysis technique and media used. Different air sampler types exist and not all are suitable for collecting a specific microorganism. The ideal air sampler is efficiently able to recover all microorganisms from the air and allow all the required analysis to be performed. Currently, there is a lack of standardized techniques to quantify airborne microorganisms. The advantages and drawbacks of different sampling methods (filtration, impingement, impaction, and sedimentation) have been previously reviewed (Buttner *et al.* 1997, Griffin *et al.* 2011, Adell *et al.* 2014).

The collection and analysis methods may represent a stress factor on microbial viability. Sampling technique, type of medium, cut-off diameter of sampling device and its detection limits play important stress factors on the survivability of microbial aerosols. Non-detection of *Salmonella* bacteria from the air environment could be attributed to their low concentrations at point sources (Kocwa-Haluch 1996). Moreover, the presence of many competing bacteria limits isolation of *Salmonella* in air samples (Carrique-Mas & Davies 2008). Several official organizations for standardization have developed reference methods for the isolation of *Salmonella*. Conventional detection methods for *Salmonella* bacteria are based on culturing techniques, using pre-enrichment broths, and selective enrichment media, followed by biochemical and serological reactions. Liquid impinge sampler using pre-enrichment broths (buffered peptone, selenite, tetrathionate brilliant green, Muller-Kauffmann tetrathionate and Rappaport-Vassiliadis soya) have been preferred to collect airborne *Salmonella* (ISO 2002). The efficiency of the enrichment broths depends on type of sample, addition of antibiotics, portion of the inoculum used and incubation temperature (35-37°C). Isolation of Salmonellae is enhanced by incubation of pre-enrichment broth into selective enrichment media (Carrique-Mas & Davis 2008) to detect low levels of pathogens; enabling reproduction of the injured cells and subsequently overestimate pathogens density (Sidhu & Toze, 2009).

Salmonellae can be isolated using numerous low-selective media (MacConkey agar, deoxycholate agar), intermediate-selective media (*Salmonella-Shigella* [SS] agar, Hektoen [HE] agar) and highly selective media (selenite agar with brilliant green), (Cooke *et al.* 1999). Most of the conventional plating media (e.g. brilliant green agar) are non-specific, developing a large number of false positive Salmonellae (*Citrobacter* and *Proteus*). XLD and HE agar are the most popular media for isolating *Salmonella* and their differentiation abilities rely on the characteristics of *Salmonella* (Rambach 1990).

*Salmonella* colonies are isolated and screened using different biochemical reactions. The main biochemical reactions are Triple sugar iron (TSI) agar (alkaline slant,

with acid, gas and H<sub>2</sub>S in the butt), lysine iron agar (Alkaline slant with alkaline, rare gas and H<sub>2</sub>S in the butt), oxidase reaction (-ve), predominantly lactose-negative and urease reaction (-ve) and confirm with polyvalent anti-sera (Table 3).

Airborne microbial concentrations cannot be accurately determined using only culture-dependent method; because microorganisms could be viable but non cultivable (Alvarez *et al.* 1995). The selective enrichment media may not restrict the growth of undesirable microorganisms (Albrecht & Kämpfer 2006). The majority of naturally occurring pathogenic microorganisms cannot be cultivated using the traditional cultivation techniques (Amann *et al.* 1995). A range of chromogenic media has been developed for the detection of *Salmonella*, based on combination of chromogenic substrate and conventional biochemical reactions. These media produce distinctive colonies; making *Salmonella* identification easier and faster. Rambach agar and *Salmonella* detection media (O'Neill *et al.* 2003) and BBL<sup>TH</sup> CHROM agar are the common chromogenic media used (Eigner *et al.* 2001). Chromogenic media offer a much higher degree of specificity than conventional media which are based on absence of lactose fermentation within *Salmonella* and/or their ability to generate hydrogen sulphide.

The culture independent technique, based-on DNA amplification by polymerase chain reaction (PCR) is used to complement /or replace culture based technique (Gugliandolo *et al.* 2011). The qualitative ISO 6579:2002 technique is the most sensitive and specific method among presence /absence PCR/ or ELISA for detecting *Salmonella* in the environmental samples (Eriksson & Aspan 2007). Molecular base methods offer advantages of a more rapid, sensitive and specific detection of pathogenic microorganisms (Kolb *et al.* 2005).

qPCR is a potential method for specific/ or genus specific quantification of aerosol samples (Dutil *et al.* 2007, Oppliger *et al.* 2008, Fallschissel *et al.* 2009). The qPCR analysis of airborne microorganisms gives higher counts than conventional cultivation methods as molecular method determines cultivable and non-cultivable cells. The accuracy and detection limit of qPCR are influenced by DNA extraction and analytical phases (Hospodsky *et al.* 2010). The drawback of the PCR is

related to its inability to provide information on pathogen viability which is necessary to investigate microbial infectivity (Zeng *et al.* 2016). The most frequently target species-specific and virulence associated genes used in the PCR of *Salmonella* are shown in Table 4.

#### *Salmonella as air bioindicator*

The criteria considered in selecting a microbial indicator include 1) the ability of a microorganism to survive in the environment of concern, 2) the correlation between the presence of the indicator and pathogens, 3) ease and speed of detection, and 4) non-pathogenicity of the indicator. The presence of fecal coliform is a good indicator of the possible presence of associated pathogenic bacteria, particularly *Salmonella*. However, pathogens are difficult to assay and seldom occur at readily detectable concentrations but high levels of coliforms and total bacterial counts may indicate the existence of enteric pathogens (Sorber & Sagik 1979). In contrast to Gram positive bacteria, Gram negative bacteria have a thinner cell wall; therefore they are more sensitive to dehydration and not viable in the air state for a long time. Gram negative bacteria represent ~1 - 10% of the airborne total bacteria (Matković *et al.* 2007), however Gram-negative bacteria may include pathogens such as Salmonellae. As a result of their thicker cell wall and the accompanying greater "robustness" towards the airborne state, most of the bacteria from the air detected via cultivation methods are Gram-positive bacteria (Zhao 2011).

**Table 3.** Appearance of *Salmonella* bacteria on different selective media.

Selective medium	Appearance
Bismuth sulfite agar	Fully developed colonies, convex, 1-3 mm in diameter, black with lustrous surface, form a shallow, soft, black pit with light edge
Brilliant green agar	Transparent pink colonies surrounded by a brilliant color
MacConkey and SS agar	Colonies usually colorless, transparent with light tan, light pinkish or yellow appearance tan centers, 1-5 mm
XLD medium	Pink to red with black center colonies
Hektoen enteric agar	Green or blue green colonies

**Table 4.** The target species specific and virulence associated genes used in the PCR of *Salmonella*

Target gene	Primer	Sequence (5' -3')	Amplicon size (bP)	Reference
Invasion plasmid Antigen-B ( <i>ipaB</i> )	<i>ipaB</i> -F <i>ipaB</i> -R	GGACTTTTTAAAAGCGGCGG GCCTCTCCCAGAGCCGTCTGG	314 429	Kaniga <i>et al.</i> (1995)
-	ST11 ST15	AGCCAACCATTGCTAAATTGGCGCA GGTAGAAATTCCAGCGGGTACTG	-	Adell <i>et al.</i> (2014)
-	Sef.B127L	5'-AGATTGGGCACTACACGTGT-3'	535	Wang <i>et al.</i> (2009)
-	SefB661R	(5'-TGTACTCCACCAGGTAATTG-3'	535	Santos <i>et al.</i> (2021)



*Enterobacteriaceae* are sensitive towards the airborne state, as they already die before/or during sampling and thus are barely detectable. The survival of coliforms in the air environment is still controversial. Coliforms have lower survivability in the air environment than *Salmonella* (Teltsch *et al.* 1980) and do not fulfill the main requirement of microbial indicator "its ability to survive in the environment is equal to/or more than the tested pathogenic microorganism". The stability of coliform in the air environment appears to be lower than certain viruses (Scarpino 1975). *Salmonella*, *Citrobacter*, *Clostridium*, *Proteus*, *Edwardsiella* and *Klebsiella* species have been associated with the presence of fecal contamination (Kromoredjo and Fujioka 1991) and Clostridia are better indicator of airborne pathogens (Hill *et al.* 1993).

#### Control measures

The practical control measures are crucial in livestock and waste applications to prevent release and spread of pathogenic microorganisms into the air environment (Hendriksen *et al.* 2004). Biosecurity management includes a set of practical measures to prevent and limit the spread of infections to humans and animals (Amass 2005). Biosecurity includes replacement of animal and husbandry (Andres & Davies 2015), dust reduction, air filtration and proper air disinfectants (Stärk 1999), electrostatic filtration, fogging and oil based spray, negative air ionization, vacuum cleaning, ventilation and wet scrubbers (Holt *et al.* 1999, Ritz *et al.* 2006). Rodent and insect control and disinfection between flocks are recommended to reduce *Salmonella* in farms (Gosling *et al.* 2014). Pressurized steam followed by forced hot air reduces levels of *Salmonella* and *Campylobacter* in transport cage flooring and reduce cross-contamination of broilers (Reina *et al.* 2024). Assessing biosecurity includes measuring the potential routes for disease transmission. Air is yet another vector by which pathogens can contaminate the final products. The adjacent nearby residential areas require higher standards of amenity. The width of a buffer zone > 400m between waste and livestock applications and residential areas should be taken in consideration during city planning.

The International Life Sciences Institute (ILSI) outlined a number of measures that should be considered with regard to air entering production floors (Beuchat *et al.* 2011), including a positive pressure air system to prevent the contaminated air infiltrating controlled production areas and eliminating residual moisture (Podolak *et al.* 2010). Filtering air entering production zones may also be effective as well as continuous monitoring of *Salmonella* in the air is important to maintain the appropriate state of the environment.

#### References

1. Abdel Hameed, A. 1992. Comparison of salmonellae and coliform bacteria in aerosols emitted from wastewater

#### Conclusion

*Salmonella* bacteria are potentially generated into the air from livestock farms and waste application facilities. These facilities are hotspots associated with high infection risks of aerosols- containing *Salmonella*. Salmonellae are found in aerosols in detectable counts. Transmission of *Salmonella* via the air pathway is less pathogenic and rarely occurs, however airborne *Salmonella* may represent a threat to public health, but no greater than that of the same count of pathogens ingested. The low count of *Salmonella* in the air is attributed to their enteric adapted to living in the protected environment, short time survives and occurrence is sporadic related to the incidence of disease infection. The efficiency of sampler, analytical technique and nutrient medium in use are important factors in detecting airborne *Salmonella*. The qPCR is fast, rapid and accurate for quantification of *Salmonella* in air samples. More sensitive laboratory methodological techniques should be created. The absence of correlation between the presence of Salmonellae and fecal coliforms make them fail to fulfill one of the main requirements of microbial indicator for air biocontamination. There is an urgent to identify more reliable alternative indicators which could be used for potential public health risk assessment. The development of new diagnostic tools (less labour and more rapid and sensitive) and vaccines targeting specific pathogenesis factors could be used in comparative investigations and control *Salmonella* transmission and infection. Finally, the presence of *Salmonella* in the air may have a hypothetical potential to cause infection.

#### Acknowledgement

The author would like to thank Prof. Dr. Helmy El-Zanfaly, Professor of Water and Sewage Microbiology, NRC, Egypt, for his support and valuable comments.

**Ethics Committee Approval:** Since the article does not contain any studies with human or animal subject, its approval to the ethics committee was not required.

**Data Sharing Statement:** All data are available within the study.

**Conflict of Interest:** The author has no conflicts of interest to declare.

**Funding:** The authors declared that this study has received no financial support.

**Editor-in-Chief note:** Abdel Hameed A. Awad is a member of Editorial Board of Trakya University Journal of Natural Sciences. However, he was't involved in the decision process during manuscript evaluation.

treatment plants and related irrigation systems. MSc. Thesis, Botany Dept., Faculty of Science, Mansoura Univ., Egypt.

2. Abdel Hameed, A., Elmorsy, T., Tarwarer, P., Green, C. & Gibbs, S. 2010. Air biocontamination in a variety of agricultural industry environments in Egypt: a pilot study. *Aerobiologia*, 26(3): 223-232. <https://doi.org/10.1007/s10453-010-9158-y>
3. Adams, A. & Spendlove, J. 1970. Coliform aerosols emitted by sewage treatment plants. *Science*, 169(3951): 1218-1220. <https://doi.org/10.1126/science.169.3951.1218>
4. Adell, E., Moset V., Zhao, Y., Jiménez Belenguier, A., Cerisuelo, A. & Cambra-López, M. (2014). Comparative performance of three sampling techniques to detect airborne Salmonella species in poultry farms. *Annals of Agricultural and Environmental Medicine*, 21(1): 15-24.
5. Aengst, C. 1984. The composition of dust in a pig fattening house [Zur Zusammensetzung des Staubes in einem Schweinemaststall]. Dissertation, Tierärztliche Hochschule Hannover [In German].
6. Akil, L., Ahmad, H. & Reddy, R. 2014. Effects of climate change on Salmonella infections. *Foodborne Pathogens and Disease*, 11(12): 974-80. <https://doi.org/10.1089/fpd.2014.1802>
7. Akin, E.; Jakubowski, J.; Lucas, B. & Pahren, H. 1978. Health hazards associated with wastewater effluents and sludge: Microbiological considerations, pp. 9-25. In: Sagik, B. & Sorber, C. (eds), *Proc. Conf. Risk Assessment and Health Effects of Land Application of Municipal Wastewater and Sludge*. Center for Applied Research and Technology, University of Texas at San Antonio, San Antonio.
8. Ajonina, C., Buzie, C., Rubiandini, R. & Otterpohl, R. 2015. Microbial pathogens in wastewater treatment plants (WWTP) in Hamburg. *Journal of Toxicology and Environmental Health, Part A*, 78(6): 381-387. <https://doi.org/10.1080/15287394.2014.989626>
9. Albrecht, A. & Kämpfer, P. 2006. Potential overestimation of the actinomycetes concentrations by growth of thermophilic species of Bacillus and Geobacillus on selective media for thermophilic actinomycetes. *Gefahrstoffe Reinhaltung der Luft*, 66(9): 365-367
10. Alvarez, A., Buttner, M. & Stetzenbach, L. 1995. PCR for bioaerosol monitoring: sensitivity and environmental interference. *Applied and Environmental Microbiology*, 61(10): 3639-3644. <https://doi.org/10.1128/aem.61.10.3639-3644.1995>
11. Alzenki, S., Al-Nasser, A., Al-Safar, A. Alomirah, H., Al-Haddad, A., Hendriksen, R. & Aarestrup, F. 2007. Prevalence and antibiotic resistance of *Salmonella* isolated from a poultry farm and processing plant environment in the state of Kuwait. *Foodborne Pathogens and Disease*, 4(3): 363-373. <https://doi.org/10.1089/fpd.2007.0017>
12. Amann, R., Ludwig, W. & Schleifer, K. 1995. Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiological Reviews*, 59 (1): 143-169. <https://doi.org/10.1128/mr.59.1.143-169.1995>
13. Amass, S. 2005. Biosecurity: stopping the bugs from getting in. *The Pig Journal*, 55: 104-114.
14. Aminul Islam, M., Ikeguchi, A. & Naide, T. 2020. Influence of temperature and humidity on the dynamics of aerosol numbers and airborne bacteria in a dairy calf house. *Biosystems Engineering*, 194(7): 213-226. <https://doi.org/10.1016/j.biosystemseng.2020.04.003>
15. Andres, V. & Davies, R. 2015. Biosecurity measures to control Salmonella and other infectious agents in Pig farms: a Review. *Comprehensive Reviews in Food Science and Food Safety*, 14(4): 317-335. <https://doi.org/10.1111/1541-4337.12137>
16. Andino, A. & Hanning, I. 2015. *Salmonella enterica*: survival, colonization, and virulence differences among serovars. *Scientific World Journal*, 2015: 520179. <https://doi.org/10.1155/2015/520179>
17. Ashurst, J., Truong, J. & Woodbury, B. 2022. *Salmonella Typhi*. In: *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, 2022.
18. Baskerville, A., Humphrey, T., Fitzgeorge, R., Cook, R., Chart, H., Rowe B. & Whitehead, A. 1992. Airborne infection of laying hens with *Salmonella enteritidis* phage type 4. *Veterinary Record*, 130(18): 395-398. <https://doi.org/10.1136/vr.130.18.395>
19. Bauer, H., Fuerhacker, M., Zibuschka, F., Schmid, H. & Puxbaum, H. 2002. Bacteria and fungi in aerosols generated by two different types of wastewater treatment plants. *Water Research*, 36(16): 3965-70. [https://doi.org/10.1016/S0043-1354\(02\)00121-5](https://doi.org/10.1016/S0043-1354(02)00121-5)
20. Baylor, E., Peters, V. & Baylor, M. 1977. Water to air transfer of virus. *Science*, 197(4305): 763-764. <https://doi.org/10.1126/science.329413>
21. Berrang, M., Cox, N. & Bailey, J. 1995. Measuring airborne microbial contamination of broiler hatching cabinets. *Journal of Applied Poultry Research*, 4(1): 83-87. <https://doi.org/10.1093/japr/4.1.83>
22. Beuchat, L., Komitopoulou, E., Beckers, H., Betts, R., Bourdichon, F. & Joosten, H. 2011. Persistence and survival of pathogens in dry food processing environments. Available at: <http://ils.i.org/Europe/documents/Persistence%20and%20Survival%20report.pdf> (Date accessed: January 10, 2024).
23. Blaser, M. and Lee, S. 1982. A review of human salmonellosis: I. Infective dose. *Reviews of Infectious Diseases* 4(6): 1096-1106.
24. Bitton, G. 1980. Introduction to environmental virology, pp. 200-242. John Wiley & Sons, Inc., New York.
25. Breza-Boruta, B., Paluszak, Z. 2007. Influence of water treatment on microbiological composition of air bioaerosols. *Polish Journal of Environmental Studies*, 16(5): 663-670.
26. Brooks, J., Tanner, B., Josephson, K., Haas, C., Gebra, C. & Pepper, I. 2005. A national study on the residential impacts of biological aerosols from the land application of biosolids. *Journal of Applied Microbiology*, 99(2): 310-322. <https://doi.org/10.1111/j.1365-2672.2005.02604.x>
27. Brooks, J., Tanner, B., Josephson, K., Gerba, C. & Pepper, I. 2004. Bioaerosols from land application of biosolids in the desert south west USA. *Water Science and Technology*, 50(1): 7-12. PMID: 15318479
28. Buttner, M., Willeke, K. & Grinshpun, S. 1997. Sampling and analysis of airborne microorganisms, pp. 629-640. In: Hurst C., Knudsen G., McInerney M., Stetzenbach L. &

Walter M. (eds), *Manual of environmental microbiology*, Washington D.C., American Society for Microbiology Press.

29. Cambra-López, M., Aarnink, A., Zhao, Y., Calvet, S. & Torres, A. 2010. Airborne particulate matter from livestock production systems: A review of an air pollution problem. *Environmental Pollution*, 158(1): 1-17. <https://doi.org/10.1016/j.envpol.2009.07.011>
30. Carducci, A., Tozzi, E., Rubulotta, E., Casini, B., Cantiani, L., Rovini, E., Muscillo, M. & Pacinim R. 2000. Assessing airborne biological hazard from urban wastewater treatment. *Water Research*, 34(4), 1173-1178. [https://doi.org/10.1016/S0043-1354\(99\)00264-X](https://doi.org/10.1016/S0043-1354(99)00264-X)
31. Carrique-Mas, J. & Davies, R. 2008. Sampling and bacteriological detection of Salmonella in poultry and poultry premises: a review. *Revue scientifique et technique*, 27(3): 665-677. <https://doi.org/10.20506/rst.27.3.1829>
32. Cason, J., Cox, N. & Bailey, J., 1994. Transmission of salmonella typhimurium during hatching of broiler chicks. *Avian Diseases*, 38 (3): 583-58. PMID: 7832712
33. Chien, Y., Chen, C., Lin, T., Chen, S. & Chien, Y. 2011. Characteristics of microbial aerosols released from chicken and swine feces. *Journal of the Air & Waste Management Association*, 61(8): 882-889. <https://doi.org/10.3155/1047-3289.61.8.882>
34. Chinivasagam, H., Tran, T., Maddock, L., Gale, A. & Blackall, P. 2009. Mechanically ventilated broiler sheds: a possible source of aerosolized *Salmonella*, *Campylobacter* and *Escherichia coli*. *Applied and Environmental Microbiology*, 75(23): 7417-7415. <https://doi.org/10.1128/AEM.01380-09>
35. Clauß, M. 2020. *Emission of bioaerosols from livestock facilities: methods and results from available bioaerosol investigations in and around agricultural livestock farming*. Thünen Institute of Agricultural Technology Bundesallee 47, Braunschweig/Germany, January.
36. Clauss, M., Huf, A. & Clauß, A. 2016. An unconventional bioaerosol chamber for tenacity studies of airborne microorganisms under almost ambient air conditions, Conference: Bioaerosol Chamber Expert Meeting, 14-15 April, Vienna, Austria.
37. Colbeck, I. & Lazaridis, M. 2010. Aerosols and environmental pollution. *Naturwissenschaften*, 97(2): 117-131. <https://doi.org/10.1007/s00114-009-0594-x>
38. Cole, D., Hill, V., Humenick, F. & Sobsey, M. 1999. Health, safety and environmental concerns of farm animal waste. *Occupational Medicine*, 14(2): 423-448. PMID:10329913
39. Cooke, V., Miles, R., Price, R. & Richardson, A. 1999. A novel chromogenic ester agar medium for detection of salmonellae. *Applied and Environmental Microbiology*, 65(2): 807-812. <https://doi.org/10.1128/AEM.65.2.807-812.1999>
40. Cosby, D., Cox, Ne., Harrison, M., Wilson, J., Jeff Buhr, R. & Fedorka-Cray, P. 2015. *Salmonella* and antimicrobial resistance in broilers: A review. *Journal of Applied Poultry Research*, 24(13): 408-426. <https://doi.org/10.3382/japr/pfv038>
41. Cosenza-Sutton, G. 2004. Enumeration of total airborne bacteria, yeast and mold contaminants and identification of *Escherichia coli* 0157:H7, *Listeria spp*, *Salmonella spp* and *Staphylococcus spp* in beef and pork slaughter facility. A dissertation presented to the graduate school of the University of Florida in partial fulfillment of the requirement for degree of doctor of philosophy, University of Florida.
42. Cox, C. 1989. Airborne bacteria and viruses. *Science Progress*, 73(4): 469-499. <https://www.jstor.org/stable/43421049>
43. Cox, C. 1995. Stability of airborne microbes and allergens, pp. 77-99. In: Cox C. & Wathes C. (eds), *Bioaerosols Handbook*. Lewis Publisher, NY.
44. Cox, C. & Wathes, C. 1995. *Bioaerosols Handbook*. Lewis Publishers, Boca Raton, FL., USA.
45. Cox, N., Bailey J., Mauldin, J. & Blankenship L. 1990. Presence and impact of *Salmonella* contamination in commercial broiler hatcheries. *Poultry Science*, 69(9): 1606-1609. <https://doi.org/10.3382/ps.0691606>
46. Dai, P., Shen, D., Tang, Q., Huang, K. & Li, C. 2020. PM2.5 from a broiler breeding production system: The characteristics and microbial community analysis. *Environmental Pollution*, 256: 113368. <https://doi.org/10.1016/j.envpol.2019.113368>
47. Darlow, H., Bale, W. & Carter, G. 1961. Infection of mice by the respiratory route with *Salmonella typhimurium*. *Journal of Hygiene (Camb.)*, 59(3): 303-308. <https://doi.org/10.1017/s0022172400038961>
48. Davies, R. & Breslin, M. 2003. Investigation of *Salmonella* contamination and disinfection in farm egg-packing plants. *Journal of Applied Microbiology*, 94(2): 191-196. <https://doi.org/10.1046/j.1365-2672.2003.01817.x>
49. Davies, R. & Wray, C. 1994. An approach to reduction of Salmonella infection in broiler chicken flocks through intensive sampling and identification of cross-contamination hazards in commercial hatcheries. *International Journal of Food Microbiology* 24(1-2): 147-160. [https://doi.org/10.1016/0168-1605\(94\)90114-7](https://doi.org/10.1016/0168-1605(94)90114-7)
50. Davies, R.H. & Wray, C. 1996. Persistence of *Salmonella* in poultry units and poultry food. *British Poultry Science*, 37(3): 589-596. <https://doi.org/10.1080/00071669608417889>
51. de Jong, J., Trouwborst, T. & Winkler, K. 1973. *The mechanism of virus decay in aerosols*, pp. 124-130. In: Hers, J. & Winkler, K. (eds), *Airborne transmission and airborne infection*. John Wiley & Sons, New York, NY.
52. de Rezende, C., Mallinson, E., Tablante, N., Morales, R. & Park, A. 2001. Effect of dry litter and airflow in reducing *Salmonella* and *Escherichia coli* populations in the broiler production environment. *Journal of Applied Poultry Research*, 10(3): 245-251. <https://doi.org/10.1093/japr/10.3.245>
53. Devane, M., Weaver, L., Singh, S. & Gilpin, B. 2018. Fecal source tracking methods to elucidate critical sources of pathogens and contaminant microbial transport through New Zealand agricultural watersheds – a review. *Journal of Environmental Management*, 222: 293-303. <https://doi.org/10.1016/j.jenvman.2018.05.033>
54. Duan, H., Chai, T., Müller, W. & Zucker, B. 2006. Concentration of airborne endotoxins and airborne bacteria in Chinese rabbit houses. *Berliner und Münchener*

- Tierärztliche Wochenschrift*, 119(1-2): 40-44, PMID:16450707
55. Dungan, R. 2010. Board-invited review: fate and transport of bioaerosols associated with livestock operations and manures. *Journal of Animal Science*, 88(11): 3693-3706. <https://doi.org/10.2527/jas.2010-3094>
  56. Dutil, S., Veillette, M., Meriaux, A., Lazure, L., Barbeau, J. & Duchaine, C. 2007. Aerosolization of *Mycobacteria* and *Legionella* during dental treatment: low exposure despite dental unit contamination. *Environmental Microbiology*, 9(11): 2836-2843. <https://doi.org/10.1111/j.1462-2920.2007.01395.x>
  57. Eigner, U., Reissbrodt, R., Hammann, R. & Fahr, A. 2001. Evaluation of a new chromogenic medium for the isolation and presumptive identification of *Salmonella* species from stool specimens. *European Journal of Clinical Microbiology & Infectious Diseases*, 20(8): 558-565. <https://doi.org/10.1007/s100960100546>
  58. Elliott, L., McCalla, T. & Deshazer, J. 1976. Bacteria in the air of housed swine units. *Applied and Environmental Microbiology*, 32(2): 270-273. <https://doi.org/10.1128/aem.32.2.270-273.1976>
  59. Epstein, E. 1997. The science of composting. CRC Press LLC, Florida, 504 p.
  60. Eriksson, E. & Aspan, A. 2007. Comparison of culture, ELISA and PCR techniques for *Salmonella* detection in faecal samples for cattle, pig and poultry. *BMC Veterinary Research*, 3: 21. <https://doi.org/10.1186/1746-6148-3-21>
  61. Fallschissel, K., Kämpker, P. & Jäckel, M. 2009. Direct detection of *Salmonella* cells in the air of livestock stables by real time PCR. *The Annals of Occupational Hygiene*, 53(8): 859-868. <https://doi.org/10.1093/annhyg/mep060>
  62. Fannin, K, Vana, S & Jakubowski, W. 1985. Effect of an activated sludge wastewater treatment plant on ambient air densities of aerosols containing bacteria and viruses. *Applied and Environmental Microbiology*, 49(5): 1191-1196. <https://doi.org/10.1128/aem.49.5.1191-1196.1985>
  63. Foster, D & Engelbrecht, R. 1973. Microbial hazard of disposing of wastewater on soil, pp. 247-270. In: Sopper, W. & Kardos, L. (eds), *Recycling treated municipal wastewater and sludge through forest and cropland*, Pennsylvania State University Press.
  64. Forgie, D., Sasser, L. & Neger, M. 2004. Compost facility requirements guideline: How to comply with part 5 of the organic matter recycling regulation, <http://www.env.gov.bc.ca/epd/codes/omr/pdf/compost.pdf>
  65. Gast, R., Mitchell B. & Holt, P. 1998. Airborne transmission of *Salmonella enteritidis* infection between groups of chicks in controlled environment isolation cabinets. *Avian Diseases*, 42(2): 315-320. PMID: 9645323
  66. Gibbs, R., Hu, C., Ho, G. & Unkovich, I. 1997. Regrowth of faecal coliforms and salmonellae in stored biosolids and soil amended with biosolids. *Water Science and Technology*, 35(11-12): 269-275. <https://doi.org/10.2166/wst.1997.0745>
  67. Gordon, M. 2011. Invasive nontyphoidal *Salmonella* disease: epidemiology, pathogenesis and diagnosis. *Current Opinion in Infectious Diseases*, 24(5): 484-489. <https://doi.org/10.1097/QCO.0b013e32834a9980>
  68. Gray, J. & Fedorka-Cray, P. 2002. *Salmonella*, pp. 55-68, In: Cliver, D. & Riemann H. (eds.), *Foodborne diseases* (2<sup>nd</sup> ed), Academic Press, ISBN 0121765598
  69. Gregory, P. 1973. *The microbiology of the atmosphere*, 2<sup>nd</sup> edition. Leonard Hill Books, Aylesbury, Bucks, England.
  70. Griffin, D., Gonzalez, C., Teigell, N., Petrosky, T., Northup, D. & Lyles, M. 2011. Observations on the use of membrane filtration and liquid impingement to collect airborne microorganisms in various atmospheric environments. *Aerobiologia*, 27(1): 25-35. <https://doi.org/10.1007/s10453-010-9173-z>
  71. Grisoli, P., Rodolfi, M., Villani, S., Grignan, E., Cottica, D., Beiri, A., Picco, A. & Dacarro, C. 2009. Assessment of airborne microorganism contamination in an industrial area characterized by an open composting facility and wastewater treatment plant. *Environmental Research*, 109(2): 135-142. <https://doi.org/10.1016/j.envres.2008.11.001>
  72. Grunnet, K. & Tramsen, C. 1974. Emission of airborne bacteria from a sewage treatment plant. *Revue Internationale d'Océanographie Médicale*, 34: 117-126.
  73. Gugliandolo, C., Lentini, V., Spanò, A. & Maugeri, T. 2011. Conventional and molecular methods to detect bacterial pathogens in mussels. *Letters in Applied Microbiology*, 52(1): 15-21. <https://doi.org/10.1111/j.1472-765X.2010.02959.x>
  74. Gale C. & Velazquez E., 2020. *Salmonella* spp in pigs: an update on diagnostics and control. *Livestock*, 25(1): 38-43. <https://doi.org/10.12968/live.2020.25.1.38>
  75. Gržinić, G, Piotrowicz-Cieślak, A., Klimkowicz-Pawlas, A., Górny, R., Ławniczek-Wałczyk, A, Piechowicz L, Olkowska E, Potrykus M, Tankiewicz M, Krupka M, Siebielec G & Wolska, L. 2023. Intensive poultry farming: A review of the impact on the environment and human health. *Science of The Total Environment*, 858, Part 3: 160014. <https://doi.org/10.1016/j.scitotenv.2022.160014>
  76. Gosling, R., Martelli, F., Wintrip, A., Sayers, A., Wheeler, K. & Davies, R. 2014. Assessment of producers response to *Salmonella* biosecurity issues and uptake of advice on laying hen farms in England and Wales. *British Poultry Science*, 55(5): 559-568. <https://doi.org/10.1080/00071668.2014.949620>
  77. Haas, D., Posch, J., Schmidt, S., Wüst, G., Sixl, W., Feierl, G., Marth, E. & Reinthaler, F. 2005. A case study of airborne culturable microorganisms in a poultry slaughter house in Styria, Austria. *Aerobiologia*, 21: 193 -201. <https://doi.org/10.1007/s10453-005-9003-x>
  78. Han, Y., Li, L., Wang, Y., Ma, J., Li, P., Han, C. & Liu, J. 2020. Composition, dispersion, and health risks of bioaerosols in wastewater treatment plants: A review. *Frontiers of Environmental Science and Engineering*, 15(3): 38. <https://doi.org/10.1007/s11783-020-1330-1>
  79. Heaton, J. C. & Jones, K. 2008. Microbial contamination of fruit and vegetables and the behaviour of enteropathogens in the phyllosphere: a review. *Journal of Applied Microbiology*, 104 (3): 613-626. <https://doi.org/10.1111/j.1365-2672.2007.03587.x>
  80. Heber, A., Peugh M., Lutgring K., Zimmerman N. & Linton R. 2006. Poultry slaughtering plants: concentrations of

microbial aerosols in poultry slaughtering and processing plants. *ASHRAE Transactions*, 112: 644-655.

81. Heidelberg, J., Shahamat, M., Levin, M., Rahman, I., Stelma, G., Grim, C. & Colwell R.R. 1997. Effect of aerosolization on culturability and viability of gram-negative bacteria. *Applied and Environmental Microbiology*, 63(9): 3585-3588. <https://doi.org/10.1128/aem.63.9.3585-3588.1997>
82. Heinonen-Tanski, H., Reponen, T. & Koivunen, J. 2009. Airborne enteric coliphages and bacteria in sewage treatment plants. *Water Research*, 43(9): 2558-2566. <https://doi.org/10.1016/j.watres.2009.03.006>
83. Hendriksen, S., Orsel, K., Wagenaar, J., Miko, A. & Van Duijkeren, E. 2004. Animal-to-human transmission of *Salmonella Typhimurium* DT104A variant. *Emerging Infectious Diseases*, 10(12): 2225-27. <https://doi.org/10.3201/eid1012.040286>
84. Hess, E. & Breer, C. 1975. Epidemiology of salmonellae and fertilizing of grassland with sewage sludge. *Zentralbl Bakteriol Orig B*, 161(1): 54-60. PMID:1189797
85. Hickey, J. & Reist, P. 1975. Health significance of airborne microorganisms from wastewater treatment processes. Part II: Health significance and alternatives for action. *Journal of the Water Pollution Control Federation*, 47(12): 2758-2773. PMID: 1107606
86. Hill, R., Knight, I., Anikis, M. & Colwell, R. 1993. Benthic distribution of sewage sludge indicated by *Clostridium perfringens* at a deep ocean dump site. *Applied and Environmental Microbiology*, 59(1): 47-51. <https://doi.org/10.1128/aem.59.1.47-51.1993>
87. Hinton, M., Ali, E., Allen, V. & Linton, A. 1983. The excretion of *Salmonella typhimurium* in the feces of cows fed milk substitute. *Journal of Hygiene (Camb)*, 91(1): 33-45. <https://doi.org/10.1017/s0022172400060009>
88. Holt, P., Mitchell, B., Seo, K-H. & Gast, R. 1999. Use of negative air ionization for reducing airborne levels of *Salmonella enterica* serovar Enteritidis in a room containing infected caged layer. *Journal of Applied Poultry Research*, 8: 440-446.
89. Hospodsky, D., Yamamoto, N. & Peccia, J. 2010. Accuracy, precision, and method detection limits of quantitative PCR for airborne bacteria and fungi. *Applied and Environmental Microbiology*, 76(21): 7004-7012. <https://doi.org/10.1128/AEM.01240-10>
90. Hussong, D., Burge, W. & Enkiri, N. 1985. Occurrence, growth and suppression of Salmonellae in composted sewage sludge. *Applied and Environmental Microbiology*, 50(4): 887-893. <https://doi.org/10.1128/aem.50.4.887-893.1985>
91. Hutchison, M.L., Gittins, J., Walker, A., Sparks, N., Humphrey, T.J., Burton, C. & Moore, A. 2004. An assessment of the microbiological risks involved with egg washing under commercial conditions. *Journal of food protection*, 67(1): 4-11.
92. ISO: 6579 2002. International Organization for Standardization, Microbiology of food and animal feeding stuffs — horizontal method for the detection of *Salmonella* spp., ISO 6579:2002.
93. Jiang, C., Shaw, K., Upperman, C., Blythe, D., Mitchell, C., Murtugudde, R., Sapkota A. & Sapkota, A. 2015. Climate change, extreme events and increased risk of salmonellosis in Maryland, USA: Evidence for coastal vulnerability. *Environment International*, 83: 58-62. <https://doi.org/10.1016/j.envint.2015.06.006>
94. Jones, A. & Harrison, R. 2004. The effects of meteorological factors on atmospheric bioaerosol concentrations- a review. *Science of The Total Environment*, 326(1-3): 151-180. <https://doi.org/10.1016/j.scitotenv.2003.11.021>
95. Kallapura, G., Morgan, M., Pumford, N., Bielke, L., Wolfenden, A., Faulkner, O., Latorre, J., Menconi, A., Hernandez-Velasco, X., Kuttappan, V., Hargis, B. & Tellez, G. 2014. Evaluation of the respiratory route as a viable portal of entry for *Salmonella* in poultry via intratracheal challenge of *Salmonella Enteritidis* and *Salmonella Typhimurium*. *Poultry Science*, 93(2): 340-346. <https://doi.org/10.3382/ps.2013-03602>
96. Kaniga, K., Tucker, S., Trollinger, D. & Galán, J. 1995. Homologs of the Shigella IpaB and IpaC invasins are required for *Salmonella typhimurium* entry into cultured epithelial cells. *Journal of Bacteriology*, 177(14): 3965-3971. <https://doi.org/10.1128/jb.177.14.3965-3971.1995>
97. Karra, S. & Katsivella, E. 2007. Microorganisms in bioaerosol emission from wastewater treatment plants during summer at Mediterranean site. *Water Research*, 41(6): 1355-1365. <https://doi.org/10.1016/j.watres.2006.12.014>
98. Katzenelson, E. & Teltsch, B. 1976. Dispersion of enteric bacteria by spray irrigation. *Journal Water Pollution Control Federation*, 48(4): 710-716. <https://www.jstor.org/stable/25038569>
99. Kim, K., Kabir, E. & Jahan, S. 2018. Airborne bioaerosols and their impact on human health. *Journal of Environmental Sciences*, 67: 23-35. <https://doi.org/10.1016/j.jes.2017.08.027>
100. Kocwa-Haluch, R. 1996. Comparison of the airborne spread of coliform and hemolytic bacteria around a sewage treatment plant. *Annals of Agricultural and Environmental Medicine*, 3(1): 13-17.
101. Kolb, S., Carbrera, A., Kammann, C., Kämpfer, P. & Conrad, R. 2005. Quantitative impact of CO<sub>2</sub> enriched atmosphere on abundance of methanotrophic bacteria in a meadow soil. *Biology and Fertility of Soils*, 41(5): 337-342. <https://doi.org/10.1007/s00374-005-0842-z>
102. Korzeniewska, E., Filipkowska, Z., Gotkowska-Ptachtka, A., Janczukowicz, W. & Rutkowski, B. 2008. Bacteriological pollution of the atmospheric air at the municipal and dairy wastewater treatment plant area and its surroundings. *Archives of Environmental Protection*, 34(4): 13-23.
103. Krishnamoorthy, S., Muthalagu, A., Priyamvada, H., Akkal, S., Valsan, A., Raghunathan, R., Kanawade, P. & Gunthe, S. 2020. On distinguishing the natural and human-induced sources of airborne pathogenic viable bioaerosols: characteristic assessment using advanced molecular analysis. *SN Applied Sciences*, 2: 1162. <https://doi.org/10.1007/s42452-020-2965-z>
104. Kromoredjo, P. & Fujioka, R. 1991. Evaluating three simple methods to assess the microbial quality of drinking water in Indonesia. *Environmental Toxicology & Water Quality*, 6(2): 259-270. <https://doi.org/10.1002/tox.2530060214>
105. Lacey, J., Williamson, P. & Crook, B. 1996. Microbial emission from composts and associated risks- trials and

- tribulations of occupational aerobiologist, Chapter- 1, In: Muilenberg, M. & Burge, H. (eds), *Aerobiology*, Lewis Publishers, NY.
106. Langeland, G. 1982. *Salmonella* spp. in the working environment of sewage treatment plants in Oslo, Norway. *Applied and Environmental Microbiology*, 43(5): 1111-1115. <https://doi.org/10.1128/aem.43.5.1111-1115.1982>
  107. Lang, N., Bellett-Travers M. & Smith S. 2007. Field investigations on the survival of *Escherichia coli* and presence of other enteric microorganisms in biosolids amended agricultural soil. *Journal of Applied Microbiology*, 103(5): 1868-1882. <https://doi.org/10.1111/j.1365-2672.2007.03489.x>
  108. Leach, S., Williams, A., Davies, A., Wilson, J., Marsh, P. & Humphery, T. 1999. Aerosol route enhances the contamination of intact eggs and muscle of experimentally infected laying hens by *Salmonella typhimurium* DT104. *FEMS Microbiology Letters*, 171(2): 203-207. <https://doi.org/10.1111/j.1574-6968.1999.tb13433.x>
  109. Liu, M., Nobu, M., Ren, J., Jin, X., Hong, G. & Yao, H. 2020. Bacterial compositions in inhalable particulate matters from indoor and outdoor wastewater treatment processes. *Journal of Hazardous Materials*, 385: 121515. <https://doi.org/10.1016/j.jhazmat.2019.121515>
  110. López, F., de las Mercedes Pescaretti, M., Morero, R. & Delgado M. 2012. *Salmonella Typhimurium* general virulence factors: A battle of David against Goliath? *Food Research International*, 45(2): 842-851. <https://doi.org/10.1016/j.foodres.2011.08.009>
  111. Marchand, G., Lavoie, J. & Lazure, L. 1995. Evaluation of bioaerosols in a municipal solid waste recycling and composting plant. *Journal of the Air & Waste Management Association*, 45(10): 778-781. <https://doi.org/10.1080/10473289.1995.10467406>
  112. Matković K., Vucemilo, M., Vinkovic, B., Seol, B., Pavicic, Z., Matkovic, S. 2007. Qualitative structure of airborne bacteria and fungi in dairy barn and nearby environment. *Czech Journal of Animal Science*, 52(8): 249-254. <https://doi.org/10.17221/2280-CJAS>
  113. McKinney, R. 2004. *Environmental Pollution Control Microbiology* (Ed), Chapter- 12, Air microbiology, Marcel Dekker Inc, New York, USA.
  114. McLamarra, J. & Pruitt, J. 1995. Beneficial reuse in the southeast. *Industrial Wastewater*, 3(2): 22-24.
  115. Michalkiewicz, M. 2019. Wastewater treatment plants as a source of bioaerosols. *Polish Journal of Environmental Studies*, 28(4): 2261-2271. <https://doi.org/10.15244/pjoes/90183>
  116. Millner, P., Bassett, D. & Marsh, P. 1980. Dispersal of *Aspergillus fumigatus* from Sewage Sludge Compost Piles Subjected to Mechanical Agitation in Open Air. *Applied and Environmental Microbiology*, 39(5): 1000-1009. <https://doi.org/10.1128/aem.39.5.1000-1009.1980>
  117. Mills, J., Gage, K. & Khan A. 2010. Potential influence of climate change on vector-borne and zoonotic diseases: A review and proposed research plan. *Environ Health Perspect*, 118 (11): 1507-1514. <https://doi.org/10.1289/ehp.0901389>
  118. Morgado, M., Jiang, C., Zambrana, J., Upperman, C., Mitchell, C. Boyle, M., Sapkota, A. & Sapkota, A. 2021. Climate change, extreme events, and increased risk of salmonellosis: foodborne diseases active surveillance network (FoodNet), 2004-2014. *Environmental Health*, 20: 105. <https://doi.org/10.1186/s12940-021-00787-y>
  119. Müller, G. 1980. Airborne dissemination of bacteria from sewage treatment plants. *Environmental International*, 3(4): 283-291. [https://doi.org/10.1016/0160-4120\(80\)90139-7](https://doi.org/10.1016/0160-4120(80)90139-7)
  120. Mulloy, K. 2001. Sewage workers: toxic hazards and health effects. *Occupational Medicine*, 16(1): 23-38. PMID: 11107222
  121. Mummy, L. K. 2014. *Salmonella*, Editor (s), Philip Wexler, Encyclopedia of Toxicology (Third Edition), Academic Press, pp. 211-212. <https://doi.org/10.1016/B978-0-12-386454-3.00537-6>
  122. Nolan, S., Thorn, C., Ashekuzzaman, S., Kavanagh, I., Nag, R., Bolton, D., Cummins, E., O'Flaherty, V., Abram, F., Richards, K. & Fenton, O. 2020. Land spreading with co-digested cattle slurry, with or without pasteurization, as a mitigation strategy against pathogen, nutrient and metal contamination associated with untreated slurry. *Science of The Total Environment*, 744(140841), 1-15. <https://doi.org/10.1016/j.scitotenv.2020.140841>
  123. Ogden, L., Fenlon, D., Vinten, A. & Lewis, D. 2001. The fate of *Escherichia coli* O157 in soil and its potential to contaminate drinking water. *International Journal of Food Microbiology*, 66(1-2): 111-7. [https://doi.org/10.1016/S0168-1605\(00\)00508-0](https://doi.org/10.1016/S0168-1605(00)00508-0)
  124. Oliveira, C., Carvalho, L. & Garcia, T. 2006. Experimental airborne transmission of *Salmonella* Agona and *Salmonella* Typhimurium in weaned pigs. *Epidemiology and Infection*, 134(1): 199-209. <https://doi.org/10.1017/S0950268805004668>
  125. O'Neill, W., Cooke, R.P., Plumb, H. & Kennedy, P. 2003. ABC chromogenic agar: a cost-effective alternative to standard enteric media for *Salmonella* spp. isolation from routine stool samples. *British Journal of Biomedical Science*, 60(4): 187-190. <https://doi.org/10.1080/09674845.2003.11783697>
  126. Oppliger, A., Charriere, N., Droz, P. & Rinsoz, T. 2008. Exposure to bioaerosols in poultry houses at different stages of fattening use of real time PCR for airborne bacterial quantification. *The Annals of Occupational Hygiene*, 52 (5): 405-412. <https://doi.org/10.1093/annhyg/men021>
  127. Ossowska-Cypryk, K. 1991. Application of indicator microorganisms for the assessment of air pollution level in the vicinity of the industrial waste water treatment plant. *Gaz, Woda i Technika Sanitarna*, 5:105 [in Polish].
  128. Pal, A., Riggs, M., Urrutia, A., Osborne, R., Jackson, A., Bailey, M., Macklin, K., Price, S., Buhret, R. & Bourassa, D. 2021. Investigation of the potential of aerosolized *Salmonella* Enteritidis on colonization and persistence in broilers from day 3 to 21. *Poultry Science*, 100(12): 101504. <https://doi.org/10.1016/j.psj.2021.101504>
  129. Paluszak, Z., Ligocka, A. & Breza-Boruta B. 2003. Effectiveness of sewage treatment based on selected fecal bacteria elimination in municipal wastewater treatment plant in Toruń. *Polish Journal of Environmental Studies*, 12(3): 345-349.

130. Pearce, R., Sheridan, J. & Bolton, D. 2006. Distribution of airborne microorganisms in commercial pork slaughter process. *International Journal of Food Microbiology*, 107(2): 186-19. <https://doi.org/10.1016/j.ijfoodmicro.2005.08.029>
131. Pepper, I. & Gebra, C. 2015. Aeromicrobiology, Chapter 5, pp. 89-110. In: Pepper I., Gerba C. & Gentry T. *Environmental Microbiology*, 3<sup>rd</sup> edition, Academic Press Publishers, Waltham, MA. <https://doi.org/10.1016/B978-0-12-394626-3.00005-3>
132. Pillai, S. 2007. Bioaerosols from land-applied biosolids: issues and needs. *Water Environment Research*, 79(3): 270-278. <https://doi.org/10.2175/106143007x156763>
133. Pillai, S., Widmer, K., Dowd, S. & Ricke, S. 1996. Occurrence of airborne bacteria and pathogen indicators during land application of sewage sludge. *Applied and Environmental Microbiology*, 62(1): 296-299. <https://doi.org/10.1128/aem.62.1.296-299.1996>
134. Podolak, R., Enache, E., Stone, W., Black, D. & Elliott, P. 2010. Sources and risk factors for contamination, survival, persistence, and heat resistance of Salmonella in low-moisture foods. *Journal of Food Protection*, 73(10): 1919-1936. <https://doi.org/10.4315/0362-028X-73.10.1919>
135. Prazmo, Z. 1980. *Salmonella* in municipal sewage for agricultural use. *Przegląd Epidemiologiczny*, 34: 147-153 [in Polish].
136. Putaud, J., Van Dingenen, R., Alastuey, A., Bauer, H., Birmili, W., Cyrys, J., Flentje, H., Fuzzi, S., Gehrig, R.H., Hansson, C., Harrison, R.M., Herrmann, H., Hitznerberger, R., Hüglin, C., Jones, A.M., Kasper-Giebl, A., Kiss, G., Kousa, A., Kuhlbusch, T.A.J., Löschau, G., Maenhaut, W., Molnar, A., Moreno, T., Pekkanen, J., Perrino, C., Pitz, M., Puxbaum, H., Querol, X., Rodriguez, S., Salma, I., Schwarz, J., Smolik, J., Schneider, J., Spindler, G., ten Brink, H., Tursic, J., Viana, M., Wiedensohler, A. & Raes, F. 2010. A European aerosol phenomenology – 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe. *Atmospheric Environment*, 44(10): 1308-1320. <https://doi.org/10.1016/j.atmosenv.2009.12.011>
137. Rambach, A. 1990. New plate medium for facilitated differentiation of *Salmonella* spp. from *Proteus* spp. and other enteric bacteria. *Applied and Environmental Microbiology*, 56(1): 301-303. <https://doi.org/10.1128/aem.56.1.301-303.1990>
138. Reina, M., Urrutia, A., Figueroa J., Riggs M, Macklin K., Buhr R., Price S. & Bourassa D. 2024. Application of pressurized steam and forced hot air for cleaning broiler transport container flooring. *Poultry Science*, 103(2): 103276. <https://doi.org/10.1016/j.psj.2023.103276>
139. Ritz, C., Mitchell, B., Fairchild, B., Czarick, M. & Worley, J. 2006. Improving in-house air quality in broiler production facilities using an electrostatic space charge system. *Journal of Applied Poultry Research*, 15(2): 333-340. <https://doi.org/10.1093/japr/15.2.333>
140. Ruiz-Gil, T., Acuña, J., Fujiyoshi, S., Tanaka, D., Noda, J., Maruyama, F. & Jorquera, M. 2020. Airborne bacterial communities of outdoor environments and their associated influencing factors. *Environment International*, 145: 106156. <https://doi.org/10.1016/j.envint.2020.106156>
141. Russ, C. & Yanko, W. 1981. Factors affecting salmonellae repopulation in composted sludges. *Applied and Environmental Microbiology*, 41(3): 597-602. <https://doi.org/10.1128/aem.41.3.597-602.1981>
142. Sánchez-Monedero, M., Aguilar, M., Fenoll, R. & Roig, A. 2008. Effect of the aeration system on the levels of airborne microorganisms generated at wastewater treatment plants. *Water Research*, 42(14): 3739-3744. <https://doi.org/10.1016/j.watres.2008.06.028>
143. Santos, P., Figueiredo, H., da Silva, L., da Silva, R, Cardoso, G., Moraes, C. & Rodrigues, A. 2021. Evaluation of a rapid detection method of Salmonella in comparison with the culture method and microbiological quality in fish from the Brazilian Amazon. *Food Science & Technology (Campinas)*, 41(1): 151-157. <https://doi.org/10.1590/fst.38719>
144. Sawyer, B., Elenbogen, G., Rao, K., O'Brien, P., Zenz, D. & Lue-Hing, C. 1993. Bacterial aerosol emission rates from municipal wastewater aeration tanks. *Applied and Environmental Microbiology*, 59(10): 3183-3186. <https://doi.org/10.1128/aem.59.10.3183-3186.1993>
145. Scarpino, P. 1975. Human enteric viruses and bacteriophages as indicators of sewage pollutions, P. 49-61, In: Gamegon ALH (Ed.), *Discharge of sewage from sea outfalls*, Pergamon Press, Oxford.
146. Seedorf, J., Hartung, J., Schroder, M., Linkert, K., Phillips, V., Holden, M., Sneath, R., Short, J., White, R., Pedersen, S., Takai, H. Johnsen, J.O., Metz, J.H.M., Groot Koerkamp, P.W.G., Uenk, G.H., & Wathes, C.M. 1998. Concentrations and emissions of airborne endotoxins and microorganisms in livestock buildings in Northern Europe. *Journal of Agricultural Engineering Research*, 70(1): 97-109. <https://doi.org/10.1006/jaer.1997.0281>
147. Sekla, L., Gemmill, D., Monfreda, J., Ilysyk, M., Stackiw, W., Kay, C., Hopper C., van Buckenhoutm, L. & Eibisch, G. 1980. Sewage treatment plant workers and their environment; a health study, pp. 281-294, In: Pahren, H. & Jakubowski, W. (eds), *Wastewater Aerosols and Diseases*, US-EPA, Ohio, Cincinnati, Sept. 19-21.
148. Shaji, S., Selvaraj, R. & Shanmugasundaram, R. 2023. *Salmonella* infection in poultry: A review on the pathogen and control strategies. *Microorganisms*, 11(11): 2814. <https://doi.org/10.3390/microorganisms11112814>
149. Shuval, H., Fattal, B. & Yekutieli, P. 1986. State of the art review: An epidemiological approach to the health effects of wastewater reuse. *Water Science and Technology*, 18(9): 147-162. <https://doi.org/10.2166/wst.1986.0087>
150. Sidhu, J., Gibbs, R., Ho, G. & Unkovich, I. 2001. The role of indigenous microorganisms in suppression of *Salmonella* regrowth in composted biosolids. *Water Research*, 35(4): 913-920. [https://doi.org/10.1016/S0043-1354\(00\)00352-3](https://doi.org/10.1016/S0043-1354(00)00352-3)
151. Sidhu, J. & Toze S. 2009. Human pathogens and their indicators in biosolids: a literature review. *Environment International*, 35(1): 187-201. <https://doi.org/10.1016/j.envint.2008.07.006>
152. Skórska, C., Sitkowska, J., Krysińska-Traczyk, E., Cholewa, G. & Dutkiewicz, J. 2005. Exposure to airborne microorganisms, dust and endotoxin during processing of peppermint and chamomile herbs on farms. *Annals of Agricultural and Environmental Medicine*, 12(2): 281-288. PMID: 16457486

153. Smets, W., Moretti, S., Denys, S. & Lebeer, S. 2016. Airborne bacteria in the atmosphere: Presence, purpose, and potential. *Atmospheric Environment*, 139: 214-221. <https://doi.org/10.1016/j.atmosenv.2016.05.038>
154. Sorber, C. & Guter, K. 1975. Health and hygiene aspects of spray irrigation. *American Journal of Public Health*, 65(1): 47- 52. <https://doi.org/10.2105/ajph.65.1.47>
155. Sorber, C. & Sagik, B. 1979. Indicators and pathogens in wastewater aerosols and factors affecting survivability, pp 23-35, In: Pahren, H. & Jakubowski, W. (eds), *Wastewater Aerosols and Diseases*, US-EPA, Ohio, Cincinnati, Sept. 19-21.
156. Stackhouse, R., Faith, N., Kaspar, C., Czuprynski, C. & Wong, A. 2012. Survival and virulence of *Salmonella enterica* serovar Enteritidis filaments induced by reduced water activity. *Applied and Environmental Microbiology*, 78(7): 2213-2220. <https://doi.org/10.1128/AEM.06774-11>
157. Stärk, K. 1999. The role of infectious aerosols in disease transmission in pigs. *The Veterinary Journal*, 158(3): 164-181. <https://doi.org/10.1053/tvjl.1998.0346>
158. Stetzenbach, L. 2009. Airborne infectious microorganisms. *Encyclopedia of Microbiology*, 175-182. <https://doi.org/10.1016/B978-012373944-5.00177-2>
159. Straub, T., Pepper, I. & Gerba, G. 1993. Hazards from pathogenic microorganisms in land deposited sewage sludge. *Reviews of Environmental Contamination and Toxicology*, 132: 55-91. [https://doi.org/10.1007/978-1-4684-7065-9\\_3](https://doi.org/10.1007/978-1-4684-7065-9_3)
160. Tang, J. 2009. The effect of environmental parameters on the survival of airborne infectious agents. *Journal of the Royal Society Interface*, 6(Suppl 6): S737-46. <https://doi.org/10.1098/rsif.2009.0227.focus>
161. Tanner, B. 2004. Aerosolization of microorganisms and risk of infection from reuse wastewater residuals. Ph.D Dissertation, graduate school of medicine, University of Arizona, Tucson, AZ.
162. Teltsch, B., Kedmi, S., Bonnet, L., Borenzstajn-Rotem Y. & Katzenelson, E. 1980. Isolation and identification of pathogenic microorganisms at wastewater irrigated fields: ratio in air and wastewater. *Applied and Environmental Microbiology*, 39(6): 1183-1190. <https://doi.org/10.1128/aem.39.6.1183-1190.1980>
163. Thomas, G., Paradell Gil, T., Müller, C., Rogers, H., Berger, C. 2024. From field to plate: How do bacterial enteric pathogens interact with ready-to-eat fruit and vegetables, causing disease outbreaks? *Food Microbiology*, 117: 104389. <https://doi.org/10.1016/j.fm.2023.104389>
164. Venter, P., Lues J. F. R., Theron H. 2004. Quantification of bioaerosols in automated chicken egg production plants. *Poultry Science*, 83(7): 1226-1231. <https://doi.org/10.1093/ps/83.7.1226>
165. Vilanova, X. & Blanch, A. 2005. Distribution and persistence of fecal bacterial populations in liquid and dewatered sludge from a biological treatment plant. *The Journal of General and Applied Microbiology*, 51(6): 361-368. <https://doi.org/10.2323/jgam.51.361>
166. Vought, K. & Tatini, S. 1998. *Salmonella enteritidis* contamination of ice cream associated with a 1994 multistate outbreak. *Journal of Food Protection*, 61(1): 5-10. <https://doi.org/10.4315/0362-028x-61.1.5>
167. Wang, S., Yeh, D. & Wei, C. 2009. Specific PCR primers for the identification of *Salmonella enterica* serovar enteritidis in chicken-related samples. *Journal of Food and Drug Analysis*, 17(3): 183-189. <https://doi.org/10.38212/2224-6614.2612>
168. Wang, B., Butler D., Hamblin M. & Monack D. 2023a. One species, different diseases: the unique molecular mechanisms that underlie the pathogenesis of typhoidal *Salmonella* infections. *Current Opinion in Microbiology* 72: 102262. <https://doi.org/10.1016/j.mib.2022.102262>
169. Wang, J., Vaddu, S., Bhumanapalli, S., Mishra, A., Applegate, T., Singh, M. & Thippareddi, H. 2023b. A systematic review and meta-analysis of the sources of *Salmonella* in poultry production (pre-harvest) and their relative contributions to the microbial risk of poultry meat. *Poultry Science*, 102(5): 102566. <https://doi.org/10.1016/j.psj.2023.102566>
170. Wathes, C. 1995. Bioaerosols in animal houses, pp. 547-577. In: Cox, C. & Wathes, C. (eds). *Bioaerosols Handbook*, Boca Roton, FL: CRC press.
171. Wathes, C., Zidan, W., Pearson, G., Hinton, M & Todd, N. 1988. Aerosol infections of calves and mice with *Salmonella typhimurium*. *Veterinary Research*, 123(23): 590-594. PMID: 3062881
172. Werber, D., Dreesman, J., Feil, F., Van Treeck, U., Fell, G., Ethelberg, S., Hauri, A.M., Roggentin, P., Prager, R., Fisher, I.S.T., Behnke, S.C., Bartelt, E., Weise, E., Ellis, A., Siitonen, A., Andersson, Y., Tschäpe, H., Kramer, M.H. & Ammon, A. 2005. International outbreak of *Salmonella* Oranienburg due to German chocolate. *BMC Infectious Diseases*, 5: 7. <https://doi.org/10.1186/1471-2334-5-7>
173. Wéry, N. 2014. Bioaerosols from composting facilities – a review. *Frontiers in Cellular and Infection Microbiology*, 4: 42. <https://doi.org/10.3389/fcimb.2014.00042>
174. Xie, W., Li, Y., Bai, W., Hou, J., Ma, T., Zeng, X., Zhang, L. & An, T. 2021. The source and transport of bioaerosols in the air: A review. *Frontiers of Environmental Science & Engineering*, 15(3): 44. <https://doi.org/10.1007/s11783-020-1336-8>
175. Zeng, D., Chen, Z., Jiang, Y., Xue, F. & Li, B. 2016. Advances and challenges in viability detection of foodborne Pathogens. *Frontiers in Microbiology*, 7: Article 1833. <https://doi.org/10.3389/fmicb.2016.01833>
176. Zhang, B. 2020. The effect of aerosols to climate change and society. *Journal of Geoscience and Environment Protection*, 8(8): 55-78. <https://doi.org/10.4236/gep.2020.88006>
177. Zhang, J, Li, Y., Xu, E., Jiang, L., Tang, J., Li, M., Zhao, X., Chen, G., Zhu, H., Yu, X, & Zhang, X. 2019. Bacterial communities in PM2.5 and PM10 in broiler houses at different broiler growth stages in spring. *Polish Journal of Veterinary Sciences*, 22(3): 495-504. <https://doi.org/10.24425/pjvs.2019.129957>
178. Zhao, Y. 2011. Effectiveness of multi-stage scrubbers in reducing emissions of air pollutants from pig houses. *Transactions of the ASABE*, 54(1): 285-293. <https://doi.org/10.13031/2013.36256>
179. Zhao, Y., Aarnink, A., De Jong, M. & Groot Koerkamp, P. 2014. Airborne microorganisms from livestock production systems and their relation to dust. *Critical Reviews in*



*Environmental Science and Technology*, 44(10): 1071-1128.  
<https://doi.org/10.1080/10643389.2012.746064>

*Veterinary Medicine Series B-Infectious Diseases and  
Veterinary Public Health*, 47(1): 37-46.  
<https://doi.org/10.1046/j.1439-0450.2000.00308.x>

180. Zucker, B., Trojan, S. & Muller, W. 2000. Airborne gram negative bacterial flora in animal houses. *Journal of*