



EFFECTS OF SALICYLIC ACID SEED PRIMING ON GERMINATION OF LENTIL (*Lens culinaris* Medik.) EXPOSED TO SALT STRESS

Gizem KAMÇI TEKİN^{1*}, Behiye Tuba BİÇER², Fatma BAŞDEMİR³

¹Şırnak University, Faculty of Agriculture, Department of Field Crops, 73000, Şırnak, Türkiye

²Dicle University, Faculty of Agriculture, Department of Field Crops, 21000, Diyarbakır, Türkiye


³Harran University, Ceylanpınar Agriculture Vocational High School, 63000, Şanlıurfa, Türkiye


Abstract: Lentil (*Lens culinaris* Medik.) is an essential crop globally, particularly in Türkiye, West Asia, Southern Europe, India, and Africa. Lentil is a valuable food source, rich in proteins, carbohydrates, minerals, and vitamins. However, lentil production faces challenges due to salinity stress, which hampers water uptake and causes toxic effects on plants. The study aimed to investigate the effects of salicylic acid (SA) seed priming and varying salt (NaCl) concentrations on the germination and development of lentil seeds. The research was conducted in the Field Crops Biotechnology Laboratory, University of Dicle, Faculty of Agriculture, Türkiye. In the research, control, two salt (NaCl), four salicylic acid (SA) doses and their combinations were used on *Lens culinaris* variety called İlke. Germination percentage, energy, rate index, mean germination time, peak value, and vigor index were calculated. The results demonstrated that SA had a significant impact on improving lentil germination under salt stress conditions. Specifically, lower concentrations of NaCl combined with SA showed positive effects on germination and growth parameters compared to higher NaCl concentrations. The highest germination percentage, energy, and rate index values were in the control and 0.25 SA dose treatment. Salicylic acid doses under 50 NaCl concentration exhibited higher germination percentage, energy, and rate index values than under 100 NaCl. Increasing SA and salt doses negatively affected lentil seed germination. This finding is significant for lentil cultivation in salt-affected areas, offering a potential strategy to improve crop resilience and yield in challenging environmental conditions.


Keywords: Lentil, *Lens culinaris* Medik., Salt stress, Salicylic acid, Germination

*Corresponding author: Şırnak University, Faculty of Agriculture, Department of Field Crops, 73000, Şırnak, Türkiye

E mail: gzmkm2@gmail.com (G. KAMÇI TEKİN)

Gizem KAMÇI TEKİN  <https://orcid.org/0000-0001-6551-8669>

Behiye Tuba BİÇER  <https://orcid.org/0000-0001-8357-8470>

Fatma BAŞDEMİR  <https://orcid.org/0000-0002-1086-5628>

Received: March 25, 2024

Accepted: April 19, 2024

Published: May 15, 2024

Cite as: Kamçi Tekin G, Biçer BT, Başdemir F. 2024. Effects of salicylic acid seed priming on germination of lentil (*Lens culinaris* Medik.) exposed to salt stress. *BSJ Agri*, 7(3): 280-286.

1. Introduction

Lentil is a legume crop with $2n=14$ chromosomes and an annual growth cycle. It is an important crop in Türkiye, following peas and chickpeas (Ladizinsky, 1979; Jha and Halder, 2016; Sehgal et al., 2021). In Türkiye, lentils are produced on 81,741 hectares of land, yielding 353,631 tons of product (FAO, 2019). Lentils are also cultivated in West Asia, Southern Europe, India, and Africa.

Lentils are a nutritious food source high in carbohydrates, protein, minerals, and vitamins (Migliozzi et al., 2015). The protein content varies depending on growing conditions, techniques, and varietal characteristics, but it averages 22.7-31.9% (Toklu et al., 2009). Lentils are essential for human and animal nutrition, especially for those who consume vegan and vegetarian products.

Plants are exposed to biotic and abiotic stress factors (Georgieva and Vassileva, 2023). Biotic stress conditions refer to the impact of bacteria, fungi, plants, and animals, while abiotic stress conditions include radiation, drought, salinity, and variations in temperature and light intensity. Reactive oxygen species (ROS) are produced and accumulated by plants under different stress conditions

(Guo et al., 2023). Disruptions in metabolic events lead to oxidative stress, which causes damage to DNA, enzyme inactivation, and lipid peroxidation (Tounekti et al., 2013).

Legumes are highly sensitive to salinity stress, which can reduce yield (Farooq et al., 2017). Salinity stress can reduce plant quality and yield and negatively affect soil structure. Salinity can increase due to evaporation at high temperatures and uncontrolled irrigation. Physiological drought is caused by salinity stress, which prevents water uptake from plant roots and can have a toxic effect on glycophytic plants due to the accumulation of salt ions (Tester and Davenport, 2003). Salinity stress can reduce the germination rate and cause yield losses by preventing germination (Ondrasek et al., 2022). According to AL-Tawaha et al. (2013), salinity hurts the growth and development of lentils. Munns (2002) found that plant cells lose water and shrink in volume shortly after salt is applied.

Several growth regulators have been shown in numerous studies to effectively prevent salt stress. Studies, including the use of salicylic acid, a phenolic compound that belongs to the hydroxyl group. Salicylic acid helps



prevent oxidative damage caused by water stress from salt accumulation in plant roots and protects protein structure (Kaydan et al., 2007; Vicente and Plasencia, 2011). Choudhary et al. (2021), Idrees et al. (2011) and Lee et al. (2010) found that applying salicylic acid increased germination rates by preventing salt stress. Jain and Srivastava (1981) and Ramanujam et al. (1998) reported that salicylic acid, when applied at low concentrations, accelerates vegetative growth, and increases nodule formation, flowering, and pod number, particularly in legumes, resulting in a positive effect on grain yield. Additionally, Senaratna et al. (2000) and Kaydan and Yagmur (2006) found that salicylic acid applications in beans and tomatoes increase plant tolerance to drought and frost stress.

There is study aimed was to investigate the effects of different salicylic acid seed priming and salt concentrations on the germination and development of lentil.

2. Materials and Methods

The research was conducted in the Field Crops Biotechnology Laboratory, Faculty of Agriculture, Dicle University.

In the research, control, 2 salt (NaCl), 4 salicylic acid (SA) doses and their combinations were used on *Lens culinaris* variety called İlke (Table 1).

Table 1. Salt (NaCl), Salicylic acid (SA) and NaCl + SA combinations

Treatment	Combinations	
Control	50 NaCl +	0.25
NaCl (mM /L)		0.50
50		0.75
100		1.0
SA (mM/L)	100 NaCl +	0.25
0.25		0.50
0.50		0.75
0,75		1.0
1.0		

SA= salicylic acid, NaCl= salt, NaCl + SA= salt + salicylic acid.

The experiment followed the ISTA (1996) rules. To ensure surface sterilization of seeds, they were shaken in a 5% sodium hypochlorite solution for three minutes and then washed with distilled water. Excess water from the surface-sterilized seeds was removed with sterile filter papers. The experiment was arranged in randomized plots with three replications, using 40 lentil seeds for each treatment.

For salicylic acid pretreatment, seeds were soaked for 24 h in solutions of different concentrations (0.25, 0.50, 0.75, 1.0 mM/L). For the salt treatment, irrigation water with different salt concentrations (50 and 100 mM NaCl/L) was used. The seeds were soaked in distilled water for the control treatment. The Petri dishes were then incubated in a germination cabinet at 25±2 °C

(Doruk Kahraman and Topal, 2024). The germination paper was changed every two days for the duration of the experiment. Germinating seeds were defined as those with a length of the radicle of more than 2 mm (Doruk Kahraman and Topal, 2024). Measurements of the radicle and shoot traits were made after the seeds had been transferred to containers of 20 x 5 x 10 cm on the fourth day. Radicle and shoot lengths were measured with a ruler in cm, and radicle and shoot fresh weights were measured with a precision balance with an accuracy of 0.0001 g.

Germination percentage (GP) (%) (Equation 1), germination energy (GE) (%) (Equation 2), germination rate index (GRI) (Equation 3), mean germination time (MGT) (Equation 4), peak value (PV) (Equation 5), and vigour index (VI) (Equation 6) values were calculated using the following formulas:

$$GP = \frac{\text{Number of seeds germinated}}{\text{Total number of seeds tested}} \times 100 \quad (1)$$

$$GE = \frac{\text{No. of seeds germinated on day (4. day)}}{\text{Total number of seeds tested}} \times 100 \quad (2)$$

$$GRI = \left(\frac{G1}{1}\right) + \left(\frac{G2}{2}\right) + \left(\frac{G3}{3}\right) + \dots + \left(\frac{Gi}{i}\right) \quad (3)$$

$$MGT = \frac{\sum Fx}{\sum (F * x +, \dots)} \quad (4)$$

$$PV = \frac{\text{Highest seed germinated}}{\text{Number of days}} \quad (5)$$

$$VI = (\text{Average root length} + \text{Average hypocotyle length}) * GP \quad (6)$$

2.1. Statistical Analysis

The data obtained from the study were subjected to ANOVA analysis in the JMP 14 PRO statistical program. Differences between means were determined according to 0.05 in the Duncan test.

3. Results and Discussion

The effects of NaCl, salicylic acid and their combinations on *Lens culinaris* Medik (İlke cv.) germinated seeds were observed during for seven days (Figure 1). In the control group, 17 out of 40 seeds were recorded as germinated on the first day, and the germination of 36 seeds was completed on the 3rd day of the experiment, and 4 seeds weren't germinated during the experiment. The highest number of germinated seeds on the first day was observed at 0.25 SA dose and 0.50 SA dose. Under 50 NaCl concentration, 5 germinated seeds were recorded on the first day, and the highest number of germinated seeds was reached on the 5th day with 13.7 seeds. Under 100 NaCl concentration, seed germination occurred on the first day, but no germinated seeds were observed on the following days. SA treatment doses at 100 NaCl concentration regressed germination compared to the 50 NaCl concentration and significantly reduced the number

of germinating seeds.

The effects of NaCl, salicylic acid and their combinations on the germination percentage, germination energy and germination rate index of *Lens culinaris* Medik (İlke cv.) seeds were significant (Table 2, Figure 2).

The highest germination percentage, energy and rate index values were in the control (90.0%, 90.00% and 76.53, respectively). 100 NaCl concentration was exhibited the lowest germination percentage, energy and rate index values (3.33%, 2.50% and 1.32%, respectively). The values for germination percentage, energy and rate index under 50 NaCl concentration were 34.2%, 30.0% and 22.8%, respectively. 50 NaCl concentration showed lower germination values than the

control, SA, and salt + SA combinations, although not as much as 100 NaCl.

The values for germination percentage, energy and rate index for the 0.25 SA dose treatment (89.2%, 89.2% and 76.5%, respectively) were higher than those for the 1.0 SA dose treatment (80.8%, 78.3% and 54.3%, respectively). Salicylic acid doses, under 50 NaCl concentration exhibited higher germination percentage, energy and rate index values than under 100 NaCl. 0.25 SA dose, under 50 NaCl and 100 NaCl concentrations exhibited higher germination percentage, energy and rate index values than other doses. It was determined that increasing SA and salt doses negatively affected lentil germination.




Figure 1. Effect of salt and SA treatments for germinated seeds and days.




Figure 2. Effect of salt and SA treatments on germination percentage (GP), germination energy (GE), and germination rate index (GRI).

Table 2. Anova analysis for germination traits

Source	DF	GP	GE	GRI	MGT	PV
Treatment	14	3169.31**	3215.32**	2258.71**	0.293428**	14.3092**
Error	30	57.36	45.97	18.11	0.034870	0.3222
C. Total	44					

GP= Germination percentage %, GE= Germination energy %, GRI= germination rate index, MGT= Mean germination time, PV= Peak Value, VI= Vigour index %, *= P<0.05, **= P<0.001.

Ekmekci et al. (2005) reported that the decrease in germination rate with increasing salt dose was due to the toxicity of Na and Cl ions and the increase in osmotic pressure, which prevented the entry of water into the seed needed for germination. Uyanik et al. (2014) and Foolad and Lin (1997) reported that some metabolic disorders and inhibition of germination regulatory protein synthesis in plants exposed to salt stress caused a decrease in germination rate. It was also found by Mahdavi and Sanavy (2007), Fallahi et al. (2015), Gheidary et al. (2017), Haileselasie and Gselasie (2012), Tsegay and Gebreslassie (2014) and others that germination rate decreased with increasing salinity.

Mean Germination Time (MGT) and Peak Value (PV) were given in Table 2, Figure 3. MGT ranged from 4.0 to 5.5 days across all treatments. Mean germination time was more than 4.5 days for 100 NaCl concentration and all SA doses under 100 NaCl concentration. The highest mean germination time (5.26 days) was observed in 1.0 SA doses under 100 NaCl concentration. The control

group and salicylic acid doses had the highest peak values. Specifically, 0.25 SA (5.94) and 1.0 SA (6.27) had the highest peak values. Altuner et al. (2022) found that salt applications decreased the germination index in all varieties and decreased to the lowest level in 150 mM salt applications. Additionally, the average germination time increased with increasing salt doses. Similarly, Petrović et al. (2016) and Anaya et al. (2018) reported negative effects of increasing salt stress on germination in legumes. It is worth noting that SA can act as a signaling molecule, activating defense mechanisms in seeds that help them tolerate salt stress. High concentrations of SA can inhibit germination or have no significant effect, although it can indirectly lead to improved water uptake and faster germination in salt-stressed environments. The effectiveness of SA can vary depending on the type of seed, the level of salt stress, and other environmental factors.

Radicle length, hypocotyl length and vigor index were given in Table 3 and Figure 4.

Table 3. Analysis of variance for radicle and hypocotyl length, radicle and hypocotyl fresh weight and vigor index

Variation Sources	DF	Radicle fresh weight	Hypocotyl fresh weight	DF	Radicle length	Hypocotyl length	Vigor Index
Treatment	4	0.000090	0.000073	9	11.5764**	5.29765**	2655.53**
Error	10	0.000061	0.000025	18	0.4355	0.06657	45.31
C. Total	14			27			

Treatment	Radicle fresh weight (g)	Hypocotyl fresh weight (g)
Control	0.058	0.041
0.25 SA	0.058	0.031
0.50 SA	0.054	0.028
0.75 SA	0.066	0.031
1SA	0.066	0.035

SA= Salicylic acid, *= P<0.05, **= P<0.001.




Figure 3. Effect of salt and SA treatments on radicle length, hypocotyl length, and vigor index on 8th days.




Figure 4. Effect of salt and SA treatments on radicle length, hypocotyl length, and vigor index on 8th days.

Radicle length, hypocotyl length, and vigor index were measured for the control, 50 NaCl concentration, SA doses, and salicylic acid doses under 50 NaCl concentration. The results showed that high salt stress had a negative impact on radicle and hypocotyl lengths, even with different SA doses. The control group had the highest radicle length, hypocotyl length, and vigor index (5.25 cm, 3.45 cm, and 7.8, respectively). Salicylic acid doses appeared to increase radicle and hypocotyl length under 50 NaCl salt stress. Radicle and hypocotyl fresh weight were only measured at different doses of salicylic acid. According to Haileselassie and Gselassie (2012), radicle and hypocotyl lengths are crucial parameters for assessing salt stress. This is because radicles absorb salt from the soil and water and contribute to hypocotyl development. Therefore, measuring radicle and hypocotyl lengths provides initial information on the level of salt exposure experienced by plants. El-Tayeb (2005) and Khodary (2004) found that pre-treating plants with SA increased their weight. Fallahi et al. (2015) reported that an increase in salt content led to a decrease in the fresh weights of rootlets and stems.

5. Conclusion

The study indicated that different combinations of salicylic acid (SA) and salt (NaCl) concentrations on lentil germination had a crucial impact. The results showed that the germination percentage, energy, and rate index values were differently affected by the combination of SA and salt concentrations. The group exposed to the combination of salt and SA had different germination values compared to the control group. However, the germination values were not as low as those observed in the group exposed to 100 NaCl concentration alone. High salt doses hinder nutrient uptake and water absorption, further compromising the overall growth and vigor of the seeds.

and rate index values were higher for the 0.25 SA dose treatment compared to the 1.0 SA dose treatment. This suggests that a low dose of SA had a more positive effect on lentil germination under 50 NaCl concentration. The study also revealed that the effects of SA doses varied depending on the salt concentration. Under 50 NaCl concentration, the germination percentage, energy, and rate index values were higher for all SA doses compared to under 100 NaCl concentration. This indicates that under the lower salt concentration, SA has a more beneficial impact on lentil germination. Overall, increasing doses of SA and salt had a negative influence on lentil germination. However, the combination of SA and salt concentrations, as well as the specific doses used, played a significant role in determining the extent of this negative impact. Higher SA doses lead to reduced radicle growth and shoot development in lentil seedlings. In addition to the negative effect on the germination of lentil seeds, it was worth noting that the increase in SA (salicylic acid) and salt doses can also impact other aspects of seed development.

These findings showed that significance the treatments of SA under salt stress conditions in lentil cultivation to ensure optimal seed germination and subsequent plant growth.

It was found that the germination percentage, energy,

Author Contributions

The percentage of the author(s) contributions is presented below. All authors reviewed and approved the final version of the manuscript.

	G.K.T.	B.T.B.	F.B.
C	35	35	30
D	35	35	30
S		100	
DCP	50		50
DAI	50	25	25
L	50	30	20
W	50	30	20
CR	35	35	30
SR	50	30	20
PM	35	35	30
FA	45	10	45

C=Concept, D= design, S= supervision, DCP= data collection and/or processing, DAI= data analysis and/or interpretation, L= literature search, W= writing, CR= critical review, SR= submission and revision, PM= project management, FA= funding acquisition.

Conflict of Interest

The authors declared that there is no conflict of interest.

Ethical Consideration

Ethics committee approval was not required for this study because of there was no study on animals or humans.

References

Al-Tawaha ARM, Al-Tawaha M, Latief A, Al-Ghzawi ALA. 2013. Effect of chitosan coating on seed germination and salt tolerance of Lentil (*Lens culinaris* L.). ROC, 14(2): 489-491.

Altuner F, Oral E, Baran İ. 2022. Determination of the effects of salt (NaCl) stress on germination in some barley (*Hordeum vulgare* L.) varieties. JOTAF, 19(1): 39-50.

Anaya F, Pelagio-Flores R, López-Gómez M. 2018. Salt stress affects germination and early seedling growth of faba bean (*Vicia faba* L.). Agronomy, 8(11): 272.

Choudhary S, Zehra A, Mukarram M, Wani KI, Naeem M, Khan MMA, Aftab T. 2021. Salicylic acid-mediated alleviation of soil boron toxicity in *Mentha arvensis* and *Cymbopogon flexuosus*: Growth, antioxidant responses, essential oil contents and components. Chemosphere, 276: 130153. <https://doi.org/10.1016/j.chemosphere.2021.130153>

Doruk Kahraman N, Topal A. 2024. Tuz stresine maruz kalan makarnalık buğday çeşitlerinde tohum çimlenmesinin fizyolojik göstergelerindeki farklılıklar. Mustafa Kemal Üniv Tar Bil Derg, 29(1): 148-157. <https://doi.org/10.37908/mkutbd.1385772>

Ekmekçi E, Apan M, Kara T. 2005. The effect of salinity on plant growth. OMU J Fac of Agric, 20(3): 118-125.

El-Tayeb MA. 2005. Response of barley grains to the interactive effect of salinity and salicylic acid. Plant Growth Regul, 45: 215-224.

Fallahi HR, Fadaeian G, Gholami M, Daneshkhah O, Hosseini FS, Aghhavani-Shajari M, Samadzadeh A. 2015. Germination response of grass pea (*Lathyrus sativus* L.) and arugula (*Eruca sativa* L.) to osmotic and salinity stress. Plant Breed Seed Sci, 71: 97-108.

FAO. 2019. <http://www.fao.org/faostat/en/#data/QCL>. (accessed date: March 1, 2024).

Farooq M, Gogoi N, Hussain M, Barthakur S, Paul S, Bharadwaj N, Migdadi HM, Alghamdi SS, Siddique KHM. 2017. Effects, tolerance mechanisms and management of salt stress in grain legumes. Plant Physiol Biochem, 118: 199-217.

Foolad MR, Lin GY. 1997. Genetic potential for salt tolerance during germination in *Lycopersicon* species. Hortscience, 32: 296-300.

Georgieva M, Vassileva V. 2023. Stress management in plants: examining provisional and unique dose-dependent responses. Int J Mol Sci, 24(6): 5105. <https://doi.org/10.3390/ijms24065105>

Gheidary S, Akhzari D, Pessarakli M. 2017. Effects of salinity, drought, and priming treatments on seed germination and growth parameters of *Lathyrus sativus* L. J Plant Nutr, 40 (10): 1507-1514.

Guo W, Xing Y, Luo X, Li F, Ren M, Liang Y. 2023. Reactive oxygen species: a crosslink between plant and human eukaryotic cell systems. Int J Mol Sci, 24(17): 13052. <https://doi.org/10.3390/ijms241713052>

Haleselasia TH, Gselasia B. 2012. The effect of salinity (NaCl) on germination of selected grass pea (*Lathyrus sativus* L.) landraces of tigray. Asian J Agric Sci, 4(2): 96-101.

Idrees M, Naeem M, Aftab T, Khan M. 2011. Salicylic acid mitigates salinity stress by improving antioxidant defence system and enhances vincristine and vinblastine alkaloids production in periwinkle (*Catharanthus roseus* (L.) G. Don). Acta Physiol Plant, 33(3): 987-999.

Jain A, Srivastava HS. 1981. Effect of salicylic acid on nitrate reductase activity in maize seedlings. Physiol Plant, 51: 339-342

Jha TB, Halder M. 2016. Searching chromosomal landmarks in Indian lentils through EMA-based Giemsa staining method. Protoplasma, 253(5): 1223-1231. <https://doi.org/10.1007/s00709-015-0873-7>

Kaydan D, Yagmur M, Okut N. 2007. Effects of salicylic acid on the growth and some physiological characters in salt stressed wheat (*Triticum aestivum* L.). JAS, 13:114-119.

Kaydan D, Yağmur M. 2006. Effects of different salicylic acid doses and treatments on wheat (*Triticum aestivum* L.) and lentil (*Lens culinaris* Medik.) yield and yield components. JAS, 12(03): 285-293. https://doi.org/10.1501/Tarimbil_0000000463

Khodary SEA. 2004. Effect of salicylic acid on the growth, photosynthesis and carbohydrate metabolism in salt stressed maize plants. Int J Agric Biol, 6(1): 5-8.

Ladizinsky G. 1979. The origin of lentil and its wild genepool. Euphytica, 28(1): 179-187.

Lee S, Kim SG, Park CM. 2010. Salicylic acid promotes seed germination under high salinity by modulating antioxidant activity in arabidopsis. New Phytol, 188: 626-637.

Mahdavi B, Sanavy SAMM. 2007. Germination and seedling growth in grass pea (*Lathyrus sativus*) Cultivars under salinity conditions. Pak J Biol Sci, 10 (2): 273-279.

Migliozzi M, Thavarajah D, Thavarajah P, Smith P. 2015. Lentil and kale: complementary nutrient-rich whole food sources to combat micronutrient and calorie malnutrition. Nutrients, 7(11): 9285-9298. <https://doi.org/10.3390/nu7115471>

Munns R. 2002. Comparative physiology of salt and water stress. Plant Cell Environ, 25: 239-250. <http://dx.doi.org/10.1046/j.0016-8025.2001.00808.x>

Ondrasek G, Rathod S, Manohara KK, Gireesh C, Anantha MS, Sakhare AS, Parmar B, Yadav BK, Bandumula N, Raihan F, Zielińska-Chmielewska A, Meriño-Gergichevich C, Reyes-Díaz

- M, Khan A, Panfilova O, Seguel Fuentealba A, Romero SM, Nabil B, Wan CC, Shepherd J, Horvatinec J. 2022. Salt stress in plants and mitigation approaches. *Plants*, 11(6): 717. <https://doi.org/10.3390/plants11060717>
- Petrović S, Đorđević S, Mikić A, Cvijanović P. 2016. Salinity stress affects seed germination and early seedling growth of pea (*Pisum sativum* L.). *Emir J Food Agric*, 28(12): 902-908.
- Ramanujam MP, Jaleel VA, Kumaravelu G. 1998. Effect of salicylic acid on nodulation, nitrogenous compounds, and related enzymes of *Vigna mungo*. *Biol Plant*, 41: 307-311.
- Sehgal A, Sita K, Rehman A, Farooq M, Kumar S, Yadav R, Siddique KH. 2021. Lentil. In *Crop Physiology Case Histories for Major Crops*. Academic Press, New York, US, pp: 408-428.
- Senaratna T, Touchell D, Bunn E, Dixon K. 2000. Acetyl salicylic acid induces multiple stress tolerance in bean and tomato plants. *Plant Growth Regul*, 30: 157-161.
- Tester M, Davenport R. 2003. Na⁺ tolerance and Na⁺ transport in higher plants. *Ann Bot*, 91: 503-507.
- Toklu F, Biçer BT, Karaköy T. 2009. Agro-morphological characterization of the Turkish lentil landraces. *Afr J Biotechnol*, 8(17): 4121-4127.
- Tounekti T, Hernández I, Munné-Bosch S. 2013. Salicylic acid biosynthesis and role in modulating terpenoid and flavonoid metabolism in plant responses to abiotic stress. Hayat S, et al. (eds.), *Salicylic acid*. Springer, New York, US, pp: 141-162.
- Tsegay BA, Gebreslassie B. 2014. The effect of salinity (NaCl) on germination and early seedling growth of *Lathyrus sativus* and *Pisum sativum* var. *abyssinicum*. *Afr J Plant Sci*, 8(5): 225-231.
- Uyanık M, Kara ŞM, Korkmaz K. 2014. Determination of responses of some winter canola (*Brassica napus* L.) cultivars to salt stress at germination period. *J Agric Sci*, 20: 368-375.
- Vicente MRS, Plasencia J. 2011. Salicylic acid beyond defense: its role in plant growth and development. *J Exp Bot*, 62: 3321-3338.