

## Investigation of Microplastic Contamination in *Diadema setosum* Obtained from a Fishing Barn

### Balıkçı Barınağından Elde Edilen *Diadema setosum*'da Mikroplastik Kontaminasyonunun Araştırılması

Türk Denizcilik ve Deniz Bilimleri Dergisi

Cilt: XX Sayı: XX (20XX) XX-XX

Ece KILIÇ<sup>1,\*</sup> , Erkan UĞURLU<sup>1</sup> <sup>1</sup>Marine Science and Technology Faculty, Iskenderun Technical University, Iskenderun, Hatay, Türkiye

#### ABSTRACT

This study is undertaken to evaluate microplastic contamination levels in *Diadema setosum* specimens obtained from a fishing barn. Microplastic (MP) pollution levels and their potential impacts on marine biota are still unknown compared to coastal and offshore environments. For this purpose, 19 individuals of *D. setosum* were collected and microplastic abundance in their gastrointestinal tract (GIT) and gonad were investigated. Mean microplastic abundance in GITs was  $3.0 \text{ MPs} \pm 3.1 \text{ MPs}$  per individual and  $0.9 \pm 1.0 \text{ MPs}$  per g wet weight. Mean microplastic abundance in the gonads was  $0.3 \pm 0.6 \text{ MPs}$  per individual and  $0.08 \pm 0.2 \text{ MPs}$  per g wet weight. Among all MPs, 45% of extracted MPs were fibers, followed by fragments (44%) and pellets (11%). Regarding size, the majority of the MPs extracted from GITs and all of the MPs extracted from gonads were small size MPs (less than 1 mm in size). FTIR analysis validated the plastic nature of suspected particles. Polyethylene (PE) (50%) and polypropylene (PP) (50%) were the most common type of polymers. These are the main polymers used in production of fishing nets; therefore, this result seems to validate the anthropogenic influence in the study area. This study contributes to the knowledge of the transfer of microplastics to the marine food web and highlights the need for protective measures.

**Keywords:** Iskenderun Bay, Microplastics, Microplastic ingestion, Pollution, Türkiye

#### Article Info

Received: 07 May 2024

Revised: 22 July 2024

Accepted: 22 July 2024

\* (corresponding author)

E-mail: ece.kilic@iste.edu.tr

**To cite this article:** Kılıç, E., Uğurlu E. (20XX). Investigation of Microplastic Contamination in *Diadema setosum* Obtained from a Fishing Barn, *Turkish Journal of Maritime and Marine Science* XX (XX): XX-XX.  
doi: 10.52998/trjmms.1479819.

## ÖZET

Bu çalışma, bir balıkçı barınağından toplanan *Diadema setosum*'u örneklerindeki mikroplastik kontaminasyonunun varlığını araştırmak amacıyla yapılmıştır. Mikroplastik (MP) kirlilik seviyeleri ve bunların deniz biyotası üzerindeki potansiyel etkileri, kıyı ve açık deniz ortamlarıyla karşılaştırıldığında hala bilinmemektedir. Bu amaçla 19 *D. setosum* bireyi toplanmış ve gastrointestinal sistem (GİS) ve gonaddaki mikroplastik bolluğu araştırılmıştır. GİS'teki ortalama mikroplastik bolluğu kişi başına  $3,0 \text{ MP} \pm 3,1 \text{ MP}$  ve ıslak ağırlığın gramı başına  $0,9 \pm 1,0 \text{ MP}$  olarak bulundu. Gonaddaki ortalama mikroplastik miktarı birey başına  $0,3 \pm 0,6 \text{ MP}$  ve yaş ağırlık başına  $0,08 \pm 0,2 \text{ MP}$  olarak bulundu. Tüm MP'ler arasında, ekstrakte edilen MP'lerin %45'inin fiber, %44'ünün fragment ve %11'inin pellet olduğu bulunmuştur. Boyutla ilgili olarak, GIT'ten çıkarılan MP'lerin çoğunluğu ve gonaddan çıkarılan MP'lerin tümü küçük boyutlu MP'lerdi (boyutu 1 mm'den küçük). FTIR analizi, şüpheli parçacıkların plastik yapısını doğruladı ve yaygın polimer türü olarak polietilen (PE) (%50) ve polipropilen (PP) (%50) bulundu. Bu polimerler balık ağlarının üretiminde kullanılan başlıca polimerlerdir, dolayısıyla bu sonuç, çalışma alanındaki antropojenik etkiyi doğrulamaktadır. Bu çalışma, mikroplastiklerin deniz ürünleri ağına aktarımı konusundaki bilgi birikimine katkıda bulunmakta ve koruyucu ölçümlerin gerekliliğini vurgulamaktadır.

**Anahtar sözcükler:** İskenderun Körfezi, Mikroplastik, Mikroplastik alımı, Kirlilik, Türkiye

## 1. INTRODUCTION

The plastic industry continues its significant growth due to the common usage of plastics in almost every area of our daily lives. Yet, low recycling rates and the absence of clear efficient management systems (Jambeck *et al.*, 2015) cause the entrance of plastic waste into marine environments. Once these persistent pollutants enter marine environments, they break down into smaller particles (Manzoor *et al.*, 2022).

Microplastics are defined as plastic particles which are less than 5 mm in size. Even though their first appearance in the aquatic environment was back in the 1970s (Carpenter *et al.*, 1972), their significant threat to marine biota was recently understood. Recent studies showed that microplastic pollution is a paramount concern even in the deepest part of the ocean (e.g., Mariana Trench (Peng *et al.*, 2018)).

The Mediterranean Sea is known as one of the most important hotspots of microplastic contamination since it has a semi-enclosed water circulation system with limited water flow (Everaert *et al.*, 2020). The contamination level in the coastal regions is more alarming due to river discharges, anthropogenic activities such as industrial and domestic wastewater discharges, shipping, tourism, fishing industry, etc. (Suaria *et al.*, 2016). Thus, marine organisms living in

coastal regions are at more risk than those living in offshore waters (Compa *et al.*, 2019; Lebreton and Andrady, 2019).

Sea urchins live in the rock and reef habitats of coastal environments (Nyawira and McClanahan, 2020). They are important benthic grazers and they have a crucial function in "material circulation and energy flow" in benthic environments (Dethiera *et al.*, 2019; Feng *et al.*, 2020). These species are commonly omnivorous grazers and detritus feeders, feeding on the algae or seagrass by scraping from hard substratum (Nyawira and McClanahan, 2020). Since seaweeds are potentially sink of microplastic particles (Gutow *et al.*, 2016), these small home range species are more susceptible to microplastic contamination in their natural habitat (Murano *et al.*, 2020). Laboratory studies showed that microplastic ingestion causes oxidative stress (Richardson *et al.*, 2021), decreases fertilization rate (Lyons *et al.*, 2021), deforms morphology (Bertucci and Bellas, 2021), and impacts the immune system (Murano *et al.*, 2023). Besides, microplastic particles act like a gate for toxic compounds (Rendell-Bhatti *et al.*, 2021; Di Natale *et al.*, 2022) and increase the toxicity risk of hazardous chemicals like heavy metals (Rial *et al.*, 2023). It needs to be emphasized that these species are potential candidates for protection according to the

International Union for Conservation of Nature (IUCN) (Manzo and Schiavo, 2022). Therefore, microplastic contamination have become more alarming and threatening issue for sea urchins.

For more than three decades, sea urchins have been used as model organisms for ecotoxicological studies (Manzo and Schiavo, 2022). These species are used as an effective bioindicators for the monitoring of several pollutants including microplastics (Lawrence, 2001, 2013; Bayed *et al.*, 2005; Soualili *et al.*, 2008; Hennicke *et al.*, 2021; Huseini *et al.*, 2021), since they have low mobility and good response to benthic biota (Huseini *et al.*, 2021). Previous studies have demonstrated that microplastic concentration in the sea urchins' bodies is strongly correlated with the sediment (Hennicke *et al.*, 2021; Huseini *et al.*, 2021; Rahmawati *et al.*, 2023), indicating that these organisms are more likely to reflect the pollution status of the surrounding environment into their body (Pinheiro *et al.*, 2020).

In addition to their ecological role, sea urchins are used for human consumption due to their delicious gonad. Stefánsson *et al.* (2017) reported that 50000 tons of sea urchins were harvested and consumed in Japan, Chile, New Zealand, and the Philippines, globally. They also reported that their consumption become popular in European countries such as Italy, France, and Spain, and 3000-3500 tons of sea urchins are harvested from the Mediterranean Sea annually (Stefánsson *et al.*, 2017).

Current literature mainly focuses on microplastic occurrence in vertebrate species i.e. fish. Yet, information related to microplastic occurrence in invertebrate species is scant (de Sa *et al.*, 2018). Among invertebrate species, most of the studies have focused on mollusks especially the bivalvae class (Kögel *et al.*, 2020), and the presence of MPs in other taxons seems to be ignored. On the other hand, information regarding microplastic presence in sea urchins is highly valuable considering their ecological role, market value, nutrition value, and endangered potential.

This study was conducted to examine the microplastic abundance in the gastrointestinal tract and gonad of *Diadema setosum* specimens collected from a fishing barn in İskenderun Bay. Sampling location has particular importance

because microplastic pollution levels in fishing barns are particularly unknown compared to other coastal and offshore environments. Today, it is a well-known fact that the degradation of fishing nets/ropes in shallow marine environments like fishing barns causes the release of microplastics (Welden and Covie, 2017). Similarly, a recent study has demonstrated that a significant portion of microplastics found in the sediment of fishing barns originated from the abrasion of fishing gears (Xue *et al.*, 2020). This study is the first report that evaluates the MPs presence in a sea urchin species from Türkiye.

## 2. MATERIALS AND METHODS

### 2.1 Study area and sampling

Morphologically undamaged 19 *D. setosum* specimens were collected from the İskenderun fisheries barn (36°35'34"N 36°10'33"E) in the Hatay province of Türkiye, northeastern Mediterranean Sea (Figure 1).

Field observation revealed that the study area is severely contaminated with plastics. For that reason, it is hypothesized that the study area is severely contaminated with microplastics; since, the fisheries industry is one of the major sources of microplastics (An *et al.*, 2020; Xue *et al.*, 2020).



**Figure 1.** The sampled area and samples of *D. setosum* specimens.

### 2.2 Sampling dissection and digestion

The methodology was adopted from Feng *et al.* (2020). Each specimen was weighed, thawed, and washed with distilled water to remove any sediment or outdoor contamination. The diameter length was measured by a vernier

caliper. Next, each specimen was dissected from the mouth of the sea urchin carefully to avoid harm to internal tissue. Then, the gastrointestinal tract and gonad of each individual were extracted, separately weighted, placed into glass beakers, and covered with aluminum foil immediately.

After the dissection process, 50 mL of H<sub>2</sub>O<sub>2</sub> was added to the beakers. H<sub>2</sub>O<sub>2</sub> was selected as a digestion agent due to environmental concerns. The beakers were placed on a hot plate and kept at 50 °C for 12 h until the solution was homogenized, and tissues were completely digested. Later, the remaining solution was filtered through 0.47 µm glass filters through a glass filter system at low vacuum. Then, filters were placed into sterile Petri dishes until microscopic examination.

### 2.3 Microscopic examination

Filters were examined for the existence of microplastic particles by Olympus SZX7 microscope. Plastic nature was pre-validated by the hot needle method (Hanke *et al.*, 2013; De Witte *et al.*, 2014). Identified particles were classified according to type (fiber, fragment), color (black, red, blue, white, transparent, green, brown, and yellow), and their size. Filters that have MPs larger than 1 mm were placed in glass Petri dishes and set aside for Fourier transform infrared (FTIR) analysis.

### 2.4 Polymer identification

Fourier transform infrared spectroscopy (FTIR) was employed to validate the plastic nature of identified MPs. At this stage, out of 62 MPs, 6 MPs suitable in size (>1 mm) were taken as subsamples and used in spectroscopic analyses. FTIR analysis was carried out on a SHIMADZU QATR10 FTIR spectrophotometer equipped with single reflection attenuated total reflectance (ATR) accessory. The spectrum range was arranged as 4000–400 cm<sup>-1</sup> and a resolution was set to 4.0 cm<sup>-1</sup> with 32 scans for each measurement. The polymer type was identified by comparing absorbance spectra to reference libraries of SHIMADZU library.

### 2.5 Quality assurance and control

Collected specimens were quickly placed into pre-washed (i.e. three times with prefiltered distilled water) glass jars to eliminate airborne contamination.

Extensive precautions were taken in the closed laboratories with restricted access. Windows and air conditioners were always kept closed and turned off to minimize the airflow (Bessa *et al.*, 2019). Before each analysis, laboratory surfaces, dissection equipment, and glass beakers were cleaned three times with pre-filtered distilled water (Torre *et al.*, 2016). Used chemicals and distilled water were filtered before use at all times. Only authorized personnel were allowed to enter the laboratories and they wore nitrile gloves and cotton aprons at all times. Filters were checked under the microscope for the presence of MPs before use. Three wet blank filters were placed in the laboratory during the dissection and microscopic examination steps. Only one fiber sample was detected in the blank samples which suggests that the results are scientifically acceptable. The dataset was corrected by extracting the contamination data.

### 2.6 Statistical analysis

The normality of the data was checked by Shapiro-Wilk test. Since normality was not validated, Spearman correlation analysis was used to evaluate the correlations between morphological parameters (test diameter, test height, total weight, weight of GIT and gonad) and MPs abundance in the GIT and gonad.

The Kurskal-Wallis test was employed to examine the variations in microplastic abundance between organs. Calculations were carried out using PAST program version 4.03 (Hammer *et al.*, 2001).

## 3. RESULTS

In this study, 19 individuals of *D. setosum* were employed to investigate the impact of fishing barns on marine biota by examining the microplastic abundance in the gastrointestinal tract and gonad of sea urchin. Mean diameter, mean height and mean weight of sampled individuals were 118.3±12.2 mm, 88.2±4.4 mm,

and  $50.3 \pm 19.1$  g, respectively (Table A1).

Statistical analysis showed that there is no correlation between morphological parameters and microplastic abundance (i.e. MP amount in the GIT & total weight, MP amount in the gonad & total weight, MP amount in the GIT & test diameter, MP amount in the gonad & test diameter, MP amount in the GIT & test height, MP amount in the gonad & test height, MP amount in the GIT & wet weight of GIT, MP amount in the gonad and wet weight of gonad)(Table A9).

In total 62 MPs were extracted from the specimens. Among all, 57 of them were extracted from GIT and 5 of them were extracted from gonad. Mean microplastic abundance was estimated as  $3.0 \text{ MPs} \pm 3.1 \text{ MPs}$  per individual (ind) and  $0.9 \pm 1.0 \text{ MPs}$  per g wet weight in GIT. When it comes to gonads, mean microplastic abundance was found as  $0.3 \pm 0.6 \text{ MPs}$  per individual and  $0.08 \pm 0.2 \text{ MPs}$  per g wet weight. Among 19 samples, the GIT of 15 specimens and the gonad of 5 specimens were contaminated with MPs.

The results showed that both microplastic presence rate and the extracted microplastic amount are higher in GIT than in gonad tissue. Kruskal Wallis test confirmed a significant variance in sample medians of MPs extracted from GIT and gonad ( $H(\chi^2):13,11$ ;  $H_c(\text{tie corrected}):15,1$ ;  $p:0.0001021$ )

Fiber-shaped MPs were found to be dominant in both organs. Among all MPs, 45% of extracted MPs were fiber and that were followed by fragments (44%) and pellets (11%). Extracted MPs were divided into 5 categories in terms of color black, blue, green, red color, and colorless (white and transparent MPs). Blue-colored MPs were dominant (Figure 2). Regarding size, the majority of the MPs extracted from GIT and all of the MPs extracted from gonad were small-size MPs (less than 1 mm in size) (Figure 2).




Figure 2. Characteristics of extracted

microplastic particles from gastrointestinal tract (GIT) and gonad of *D. setosum*

Six of the MPs extracted from the GIT were validated by FTIR analysis. Analysis showed that all of the suspected particles were plastic in nature and polymer types were determined as 3 polyethylene MPs (50%) and 3 polypropylene MPs (50%) (Appendix Figure A1-A6).

#### 4. DISCUSSIONS

İskenderun Bay is located inside one of the biggest plastic hotspots of the Mediterranean Sea (Papadimitriou and Allinson, 2022) and it is severely contaminated with microplastics (Gündoğdu and Çevik, 2017). Besides, contamination levels become more alarming in the coastal regions due to anthropogenic activities and intense fishing activities. Even though many studies investigate the microplastic pollution levels in the surface water and sediment of the northeastern Mediterranean Sea, there is no information regarding the microplastic pollution levels in fishing barns. Yet, fishing activities have particular importance in the formation and distribution of microplastics. In fact, they are defined as a major source of secondary microplastics (An *et al.*, 2020; Xue *et al.*, 2020). In this study, microplastic presence in the sea urchin *Diadema setosum* sampled from the fishing barn was investigated to highlight the danger potential of MPs pollution levels in the fishing barns on the marine biota. Besides, this study is the first report regarding microplastic accumulation in *D. setosum* from Türkiye.

In this study, among 19 examined specimens, 15 of them contained microplastic in their GIT (79%) and 5 of them contained MPs in their gonad (22%) (Table A1). The high frequency of occurrence in the GIT points out that these species are highly confronted with microplastics. High frequency of microplastic occurrence in GIT found in this study is in line with previous reports (Table A2). Feng *et al.* (2020) investigated the microplastic presence ( $n=210$ ) in the 4 different sea urchin species from the Yellow Sea and reported that 89.5% of examined specimens contained microplastic in their GIT. Other studies reported a 100% microplastic occurrence rate in the *Paracentrotus lividus*

sampled from the Eastern Aegean Sea, Greece. (Hennicke *et al.*, 2021), *D. setosum* sampled from Barranglombo Island, Indonesia (Savannah *et al.*, 2021), *Tetrapygnus niger* sampled from Peru (De-la-Torre *et al.*, 2020).

Microplastic particles deposited on the sediment cover the top layer of sea grass or rocky bottom and they are eventually consumed by grazer organisms. There seems to be quite a difference in the average amount of microplastics extracted from GIT in the literature. The highest MPs abundance in the GIT of *D. setosum* was reported from Untung Jawa, Jakarta as  $2175.55 \pm 584.26$  MPs per ind (Huseini *et al.*, 2021). Followed by a study conducted in Indonesia, which reported the mean MPs abundance as  $23.70 \pm 2.99$  MPs per ind (Sawalan *et al.*, 2021). Results obtained in this study ( $3.0 \pm 3.1$  MPs per ind) were significantly lower than previous reports which employed *D. setosum* as biomonitoring organisms; however, similar to those employed *P. lividus* (Raguso *et al.*, 2022; Murano *et al.*, 2022), *Strongylocentrotus intermedius*, *Temnopleurus hardwickii*, *Temnopleurus reevesii*, *Hemicentrotus pulcherrimus* (Feng *et al.*, 2020), *Diadema africanum* (Sevillano-González *et al.*, 2022), *Tetrapygnus niger* (De-la-Torre *et al.*, 2020) from different parts of the ocean (Table A2). Previous reports showed that microplastic concentration in the sediment severely impacts the microplastic abundance in the GIT (Hennicke *et al.*, 2021; Huseini *et al.*, 2021; Rahmawati *et al.*, 2023). Therefore, variations in the reported mean abundance seem to primarily result from microplastic concentration in the ambient environment rather than species type.

Only five specimens contained MPs in gonads with a mean value of  $0.9 \pm 1.0$  MPs per ind, which is coherent with the previous reports (Feng *et al.*, 2020; Murano *et al.*, 2020). Statistical analysis showed that MPs abundance in the organs was significantly different ( $p < 0.05$ ). Previous research speculated that MPs accumulated in the gonads were not transfer from the intestine (Leddy and Johnson, 2000; Feng *et al.*, 2020). Gonads receive MPs found in the coelomic fluid while they fill the whole-body cavity which accumulates MPs originate from peristomal gill and tube feed (Leddy and Johnson, 2000; Feng *et*

*al.*, 2020). Our results support this idea because of their significantly smaller size and lower amount compared to GIT.

Ingested MPs size is restricted to the animal size (Jäms *et al.*, 2020). Coherent to this, almost all ingested MPs extracted from the GIT were smaller than 1 mm. In addition, the size of MPs extracted from gonads were varied between 218  $\mu\text{m}$  to 658  $\mu\text{m}$ . Pyl *et al.* (2022) reported that MPs which are greater than 10  $\mu\text{m}$  could not originate from intestinal wall of *P. lividus*. Therefore, extracted MPs could originate from ceramic fluid (Leddy and Johnson, 2000; Feng *et al.*, 2020) or an alternative route that needs to be tested in the future.

Microplastic type (fiber, fragment or pellet) is an important parameter in determining exposure risk and bioavailability of MPs. Fiber-shaped MPs were considered to be the most harmful type for marine invertebrates (Wright *et al.*, 2013). Fiber microplastics are commonly attributed to the fragmentation of fishing nets (Koongolla *et al.*, 2020) and it is the most commonly ingested type of MPs in the marine biota of the Mediterranean Sea (Koraltan *et al.*, 2022; Kılıç and Yücel, 2022; Yücel and Kılıç, 2023a, b). Similarly, both this study and previous studies conducted in sea urchins reported the dominance of fiber microplastic (Hennicke *et al.*, 2021; Sawalman *et al.*, 2021; Sevillano-González *et al.*, 2022; De-la-Torre *et al.*, 2020; Raguso *et al.*, 2022; Murano *et al.*, 2022; Huseini *et al.*, 2021; Feng *et al.*, 2020). Therefore, sea urchins seem to have a higher ingestion preference for fiber microplastics compared to other types. Apart from this, sea urchins generate secondary microplastics while grazing on plastic surfaces (Porter *et al.*, 2019). So, it is possible that urchins fed on newly generated fiber or fragment-type microplastics while grazing on the contaminated seaweed of fishing nets on the sea bottom. This condition is significantly important for fishing barns. Because, in a high microplastic-containing region like fishing barns, they accelerate the biodegradation of macroplastics (Boudouresque and Verlaque, 2007; De-la-Torre *et al.*, 2020). While rasping on the surface of the plastic materials by calcionic teeth, they leave scratchers and generate secondary microplastics (Boudouresque and Verlaque, 2007; De-la-Torre

*et al.*, 2020). That in turn increases the ecotoxicological risk of microplastics for themselves and other marine animals.

Blue was the most dominant color in the extracted MPs, which overlaps with the previous reports in sea urchins (Hennicke *et al.*, 2021; Sawalman *et al.*, 2021; Sevillano-González *et al.*, 2022; De-la-Torre *et al.*, 2020; Murano *et al.*, 2022; Huseini *et al.*, 2021; Feng *et al.*, 2020), fish (Güven *et al.*, 2017; Giani *et al.*, 2019). Previous reports demonstrate that *P. lividus* might select blue-colored particles due to their similarity to the main food source. Besides, the study area is filled with the remaining, blue-colored fishing nets, which seems to increase the ingestion rate of MPs.

Microplastic abundance in the fishing area is linked to the fishing effort (Wright *et al.*, 2021), and higher MPs abundance is observed in the regions where fisheries activities are dominant (Xue *et al.*, 2017). Today, the majority of fishing gears are produced from synthetic polymers such as polyethylene and polypropylene (Nelms *et al.*, 2021). In this study, half of the examined MPs were PE, and the remaining were PP. Since the polymer is an important factor that is used to link between contamination source and ecotoxicity of animals, the results presumably imply that the majority of the extracted MPs were derive from fishing activities.

It is a well-known fact that microplastics entering the food chain might cause adverse effects on the upper trophic levels including humans (Hennicke *et al.*, 2021). Since sea urchins are a traditional food source for highly populated countries such as Japan, China, and Mediterranean European countries (Stefánsson *et al.*, 2017). Therefore, microplastics found in sea urchins might create a pathway for microplastics to the human body. Species that are eaten with gastrointestinal tracts and gills such as small-size fish and bivalves may pose a greater risk to human health (Smith *et al.*, 2018; Baechle *et al.*, 2020; Hennicke *et al.*, 2021). Previous studies in the İskenderun Bay reported  $2.9\pm 2.7$  MPs per ind in fish *Mullus barbatus*,  $3.4\pm 2.7$  MPs per ind in fish *Saurida undosquamis* (Kılıç and Yücel, 2022),  $0.26\pm 0.5$  MPs per ind in patella *Patella caerulea* (Yücel and Kılıç, 2023a),  $0.2\pm 0.5$  MPs/ind in mussel *Brachidontes pharaonis* (Yücel and Kılıç,

2023b). Compared to other species like fish, the edible parts of this species contained fewer MPs. Thus, it is reasonable to assume that consumption of the gonads of *D. setosum* is safe in terms of microplastic accumulation. In other words, this species poses a considerably low to moderate threat to human health. More detailed studies are required to evaluate the potential hazards of consumption of microplastic-containing aquatic products.

## 5. CONCLUSIONS

This study is conducted to investigate the microplastic contamination levels in sea urchin *D. setosum* since sea urchin species are a good reflector of microplastic pollution in the ambient environment. Results validate the omnipresence of microplastics since the majority of the specimens contained microplastics in the bodies. Polypropylene and polyethylene were the main types of polymers extracted from the GIT which might reveal the impact of the fishing industry since these are the main types of polymers used in fishing nets. As far as we know, this is the first focusing on the marine animals habituated in a fishing barn in Türkiye; therefore, the results demonstrate the importance of the fisheries industry in the formation and distribution of microplastics. There is still little known in the transfer of microplastics throughout food web, yet studies like this will provide insight to the microplastic contamination in different trophic levels and different environments.

## ACKNOWLEDGEMENTS

Authors would like to thank Dr. Nebil Yücel for his wisdom in article editing.

## AUTHORSHIP STATEMENT

## CONTRIBUTION

**Ece KILIÇ:** Conceptualization, Methodology, Validation, Formal Analysis, Resources, Writing - Original Draft.

**Erkan UĞURLU:** Validation, Formal Analysis, Resources, Writing-Review and Editing, Data Curation, Software, Visualization, Writing-Review and Editing.

## CONFLICT OF INTERESTS

The author(s) declare that for this article they have no actual, potential or perceived conflict of interests.

## ETHICS COMMITTEE PERMISSION

No ethics committee permissions is required for this study.

## FUNDING

No funding was received from institutions or agencies for the execution of this research.

## ORCID IDs

Ece KILIÇ:

 <https://orcid.org/0000-0003-1953-5008>

Erkan UĞURLU:

 <https://orcid.org/0000-0001-8940-8421>

## 6. REFERENCES

- An, L., Liu, Q., Deng, Y., Wu, W., Gao, Y., Ling, W. (2020). Sources of microplastic in the environment. In: Microplastics in terrestrial environments: *Emerging Contaminants and Major Challenges* 143-159. doi:10.1007/978\_2020\_449.
- Bayed, A., Quiniou, F., Benrha, A., Guillou, M. (2005). The *Paracentrotus lividus* populations from the northern Moroccan Atlantic coast: growth, reproduction and health condition. *Journal of the Marine Biological Association of the United Kingdom* 85(4): 999-1007. doi:10.1017/S0025315405012026.
- Bertucci, J.I., Bellas, J. (2021). Combined effect of microplastics and global warming factors on early growth and development of the sea urchin (*Paracentrotus lividus*). *Science of the Total Environment* 782: 146888. doi:10.1016/j.scitotenv.2021.146888.
- Bessa, F., Frias, J., Kögel, T., Lusher, A., Andrade, J.M., Antunes, J., Gerdts, G. (2019). Harmonized protocol for monitoring microplastics in biota. doi:10.13140/RG.2.2.28588.72321/1.
- Boudouresque, F.C., Verlaque, M. (2007). Ecology of *Paracentrotus lividus*. In: *Developments in Aquaculture and Fisheries Science*. Elsevier. 2007. pp. 243-85.
- Carpenter, E.J., Anderson, S.J., Harvey, G.R., Miklas, H.P., Peck, B.B. (1972). Polystyrene spherules in coastal waters. *Science* 178(4062): 749-750. doi:10.1126/science.178.4062.749.
- Compa, M., Alomar, C., Wilcox, C., Van Sebille, E., Lebreton, L., Hardesty, B.D., Deudero, S. (2019). Risk assessment of plastic pollution on marine diversity in the Mediterranean Sea. *Science of The Total Environment* 678: 188-196. doi:10.1016/j.scitotenv.2019.04.355.
- de Sa, L.C., Oliveira, M., Ribeiro, F., Rocha, T.L., Futter, M.N. (2018). Studies of the effects of microplastics on aquatic organisms: what do we know and where should we focus our efforts in the future? *Science of The Total Environment* 645: 1029-1039. doi:10.1016/j.scitotenv.2018.07.207.
- De Witte, B., Devriese, L., Bekaert, K., Hoffman, S., Vandermeersch, G., Cooreman, K., Robbens, J. (2014). Quality assessment of the blue mussel (*Mytilus edulis*): Comparison between commercial and wild types. *Marine Pollution Bulletin* 85(1): 146-155. doi:10.1016/j.marpolbul.2014.06.006.
- De-la-Torre, G.E., Dioses-Salinas, D.C., Huamantupa-Aybar, S., Davila-Carrasco, J. (2020). Preliminary observations of plastic debris in the gastrointestinal tract of sea urchin *Tetrapygus niger*. *Brazilian Journal of Natural Sciences* 3(2): 316-320. doi:10.31415/bjns.v3i2.94.
- Dethlefsen, M.N., Hoinsb, G., Kobelt, J., Lowed, A.T., Galloway, A.W.E., Schrame, J.B., Raymore, M., Duggins, D.O. (2019). Feces as food: the nutritional value of urchin feces and implications for benthic food webs. *Journal of Experimental Marine Biology and Ecology* 514-515: 95-102. doi:10.1016/j.jembe.2019.03.016.
- Di Natale, M.V., Carroccio, S.C., Dattilo, S., Cocca, M., Nicosia, A., Torri, M., Bennici, C.D., Musco, M., Masullo, T., Russo, S., Mazzola, A., Cuttitta, A. (2022). Polymer aging affects the bioavailability of microplastics-associated contaminants in sea urchin embryos. *Chemosphere* 309(P1): 136720. doi:10.1016/j.chemosphere.2022.136720.
- Everaert, G., De Rijcke, M., Lonneville, B., Janssen, C.R., Backhaus, T., Mees, J., van Sebille, E., Koelmans, A.A., Catarino, A.I., Vandegheuchte, M.B. (2020). Risks of floating microplastic in the global ocean. *Environmental Pollution* 267: 115499. doi:10.1016/j.envpol.2020.115499.
- Feng, Z., Wang, R., Zhang, T., Wang, J., Huang, W., Li, J., Xu, J., Gao, G. (2020). Microplastics in specific tissues of wild sea urchins along the coastal areas of northern China. *Science of The Total Environment* 728. doi:10.1016/j.scitotenv.2020.138660.



- Giani, D., Bains, M., Galli, M., Casini, S., Fossi, M.C. (2019).** Microplastics occurrence in edible fish species (*Mullus barbatus* and *Merluccius merluccius*) collected in three different geographical sub-areas of the Mediterranean Sea. *Marine Pollution Bulletin* 140: 129-137. doi:10.1016/j.marpolbul.2019.01.005.
- Gündoğdu, S., Çevik, C. (2017).** Micro- and mesoplastics in Northeast Levantine coast of Turkey: The preliminary results from surface samples. *Marine Pollution Bulletin* 118(1-2): 341-347. doi:10.1016/j.marpolbul.2017.03.002.
- Gutow, L., Eckerlebe, A., Gimenez, L., Saborowski, R. (2016).** Experimental evaluation of seaweeds as a vector for microplastics into marine food webs. *Environmental Science & Technology* 50: 915-923. doi:10.1021/acs.est.5b02431.
- Güven, O., Gokdag, K., Jovanovic, B., Kideys, A.E. (2017).** Microplastic litter composition of the Turkish territorial waters of the Mediterranean Sea, and its occurrence in the gastrointestinal tract of fish. *Environmental Pollution* 223: 286-294. doi:10.1016/j.envpol.2017.01.025.
- Hammer, Ø., Harper, D.A.T., Ryan, P.D. (2001).** PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4(1): 1-9.
- Hanke, G., Galgani, F., Werner, S., Oosterbaan, L., Nilsson, P., Fleet, D., Kinsey, S., Thompson, R., Van Franeker, J.A., Vlachogianni, T., Palatinus, A., Scoullou, M., Veiga, J.M., Matiddi, M., Alcaro, L., Maes, T., Korpinen, S., Budziak, A., Leslie, H., Gago, J., Liebezeit, G. (2013).** Guidance on Monitoring of Marine Litter in European Seas. European Commission. Accessed Date: 18.06.2023, <http://hdl.handle.net/10508/1649> is retrieved.
- Hennicke, A., Macrina, L., Malcolm-Mckay, A., Miliou, A. (2021).** Assessment of microplastic accumulation in wild *Paracentrotus lividus*, a commercially important sea urchin species, in the Eastern Aegean Sea, Greece. *Regional Studies in Marine Science* 45: 101855. doi:10.1016/j.rsma.2021.101855.
- Huseini, D.R., Suryanda, A., Patria, M.P. (2021).** Comparative analysis of microplastic content in water, sediments, and digestive traces of sea urchin *Diadema setosum* (Leske, 1778) on Untung Jawa Island and Tidung Island, Seribu Islands, Jakarta. IOP Conference Series: *Materials Science and Engineering* 1098(5): 052051. doi:10.1088/1757-899x/1098/5/052051.
- Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan, R., Law, K.L. (2015).** Plastic waste inputs from land into the ocean. *Science* 347(6223): 768-771. doi:10.1126/science.1260352.
- Jâms, I.B., Windsor, F.M., Poudevigne-Durance, T., Ormerod, S.J., Durance, I. (2020).** Estimating the size distribution of plastics ingested by animals. *Nature Communications* 11: 1-7. doi:10.1038/s41467-020-15406-6.
- Kılıç, E., Yücel, N. (2022).** Microplastic occurrence in the gastrointestinal tract and gill of bioindicator fish species in the northeastern Mediterranean. *Marine Pollution Bulletin* 177: 113556. doi:10.1016/j.marpolbul.2022.113556.
- Kögel, T., Bjørøy, Ø., Toto, B., Bienfait, A.M., Sanden, M. (2020).** Micro- and nanoplastic toxicity on aquatic life: determining factors. *Science of The Total Environment* 709: 136050. doi:10.1016/j.scitotenv.2019.136050.
- Koongolla, J.B., Lin, L., Pan, Y.F., Yang, C.P., Sun, D.R., Liu, S., Xu, X.R., Maharana, D., Huang, J.S., Li, H.X. (2020).** Occurrence of microplastics in gastrointestinal tracts and gills of fish from Beibu Gulf, South China Sea. *Environmental Pollution* 258: 113734. doi:10.1016/j.envpol.2019.113734.
- Koralan, İ., Mavruk, S., Güven, O. (2022).** Effect of biological and environmental factors on microplastic ingestion of commercial fish species. *Chemosphere* 303: 135101. doi:10.1016/j.chemosphere.2022.135101.
- Lawrence, J.M. (2001).** Edible Sea Urchins: Biology and Ecology. Elsevier. Lawrence, J.M., 2013. Sea Urchins: Biology and Ecology. Academic Press.
- Lebreton, L., Andrady, A. (2019).** Future scenarios of global plastic waste generation and disposal. *Palgrave Communications* 5: 6. doi:10.1057/s41599-018-0212-7.
- Leddy, H.A., Johnson, A.S. (2000).** Walking versus breathing: mechanical differentiation of sea urchin podia corresponds to functional specialization. *Biology Bulletin* 198(1): 88-93. doi:10.2307/1542806.
- Lyons, D.M., Thomas, P.J., Oral, R., Pagano, G., Tez, S., Toscanesi, M., Ranier, P., Trifuoggi, M. (2021).** Impact of polystyrene and polymethylmethacrylate microplastics on sea urchin *Paracentrotus lividus* embryogenesis. *Nanotoxicology and microplastics Arh Hig Rada Toksikol* 72: 75.
- Manzo, S., Schiavo, S. (2022).** Physical and chemical threats posed by micro (nano) plastic to sea urchins. *Science of The Total Environment* 808: 152105. doi:10.1016/j.scitotenv.2021.152105.
- Manzoor, S., Naqash, N., Rashid, G., Singh, R. (2022).** Plastic material degradation and formation of microplastic in the environment: a review. *Materials Today: Proceeding* 56: 3254-3260.

- Murano, C., Agnisola, C., Caramiello, D., Castellano, I., Casotti, R., Corsi, I., Palumbo, A. (2020).** How sea urchins face microplastics: Uptake, tissue distribution and immune system response. *Environmental Pollution* 264: 114685. doi:10.1016/j.envpol.2020.114685.
- Murano, C., Nonnis, S., Grassi, F., Maffioli, E., Corsi, I., Tedeschi, G., Palumbo, A. (2023).** Response to microplastic exposure : An exploration into the sea urchin immune cell proteome. *Environmental Pollution*, 320: 121062. doi:10.1016/j.envpol.2023.121062.
- Nelms, S.E., Duncan, E.M., Patel, S., Badola, R., Bhola, S., Chakma, S., Chowdhury, G.W., Godley, B.J., Haque, A.B., Johnson, J.A., Khatoon, H., Kumar, S., Napper, I.E., Niloy, M.N.H., Akter, T., Badola, S., Dev, A., Rawat, S., Santillo, D., Sarker, S., Sharma, E., Koldewey, H. (2021).** Riverine plastic pollution from fisheries: insights from the Ganges River system. *Science of The Total Environment* 756: 143305. doi:10.1016/j.scitotenv.2020.143305.
- Nyawira, A.M., McClanahan, T.R. (2020).** Chapter 3: Diedama. Elsevier B.V. doi:10.1016/B978-0-12-819570-3.00023-8.
- Papadimitriou, M., Allinson, G. (2022).** Microplastics in the Mediterranean marine environment: a combined bibliometric and systematic analysis to identify current trends and challenges. *Microplastics Nanoplastics* 2(1): 1-25. doi:10.1186/s43591-022-00026-2.
- Peng, X., Chen, M., Chen, S., Dasgupta, S., Xu, H., Ta, K., Du, M., Li, J., Guo, Z., Bai, S. (2018).** Microplastics contaminate the deepest part of the world's ocean. *Geochemical Perspectives Letters* 9(1): 1-5. doi:10.7185/geochemlet.1829.
- Pinheiro, L.M., do Sul, J.A.I., Costa, M.F. (2020).** Uptake and ingestion are the main pathways for microplastics to enter marine benthos: A review. *Food Webs* 24: e00150. doi:10.1016/j.fooweb.2020.e00150.
- Porter, A., Smith, K.E., Lewis, C. (2019).** The sea urchin *Paracentrotus lividus* as a bioindicator of plastic. *Science of The Total Environment* 693: 133621. doi:10.1016/j.scitotenv.2019.133621.
- Pyl, M., Taylor, A., Oberhänsli, F., Swarzenski, P., Hussamy, L., Besson, M., Danis, B., Metian, M. (2022).** Size-dependent transfer of microplastics across the intestinal wall of the echinoid *Paracentrotus lividus*. *Aquatic Toxicology* 250: 106235. doi:10.1016/j.aquatox.2022.106235.
- Rahmawati Krisanti, M., Riani, E., Cordova, M.R. (2023).** Microplastic contamination in the digestive tract of sea urchins (Echinodermata: Echinoidea) in Kepulauan Seribu, Indonesia. *Environmental Monitoring and Assessment* 195(9): 1103. doi:10.1007/s10661-023-11655-2.
- Rendell-Bhatti, F., Paganos, P., Pouch, A., Mitchell, C., D'Aniello, S., Godley, B.J., Pazdro, K., Arnone, M.I., Jimenez-Guri, E. (2021).** Developmental toxicity of plastic leachates on the sea urchin *Paracentrotus lividus*. *Environmental Pollution* 269: 115744. doi:10.1016/j.envpol.2020.115744.
- Rial, D., Bellas, J., Vidal-Liñán, L., Santos-Echeandía, J., Campillo, J.A., León, V.M., Albentosa, M. (2023).** Microplastics increase the toxicity of mercury, chlorpyrifos and fluoranthene to mussel and sea urchin embryos. *Environmental Pollution* 336 (August): 122410. doi:10.1016/j.envpol.2023.122410.
- Richardson, C.R., Burritt, D.J., Allan, B.J.M., Lamare, M.D. (2021).** Microplastic ingestion induces asymmetry and oxidative stress in larvae of the sea urchin *Pseudechinus huttoni*. *Marine Pollution Bulletin* 168: 112369. doi:10.1016/j.marpolbul.2021.112369.
- Smith, M., Love, D.C., Rochman, C.M., Neff, R.A. (2018).** Microplastics in seafood and the implications for human health. *Current Environmental Health Reports* 5(3): 375-386. doi:10.1007/s40572-018-0206-z.
- Soualili, D., Dubois, P., Gosselin, P., Pernet, P., Guillou, M. (2008).** Assessment of seawater pollution by heavy metals in the neighbourhood of Algiers: use of the sea urchin, *Paracentrotus lividus*, as a bioindicator. *ICES Journal of Marine Science* 65(2): 132-139. doi:10.1093/icesjms/fsm183.
- Stefánsson, G., Kristinsson, H., Ziemer, N., Hannon, C., James, P. (2017).** Markets for sea urchins: a review of global supply and markets. *Skýrsla Matis* 45: 10-17. doi:10.13140/RG.2.2.12657.99683.
- Suaría, G., Avio, C.G., Mineo, A., Lattin, G.L., Magaldi, M.G., Belmonte, G., Moore, C.J., Regoli, F., Aliani, S. (2016).** The Mediterranean plastic soup: synthetic polymers in Mediterranean surface waters. *Scientific Reports* 6: 37551. doi:10.1038/srep37551.
- Torre, M., Digka, N., Anastasopoulou, A., Tsangaris, C., Mytilineou, C. (2016).** Anthropogenic microfibres pollution in marine biota. A new and simple methodology to minimize airborne contamination. *Marine Pollution Bulletin* 113(2016): 55-61. doi:10.1016/j.marpolbul.2016.07.050.

- Welden, N.A., Cowie, P.R. (2017).** Degradation of common polymer ropes in a sublittoral marine environment. *Marine Pollution Bulletin* 118(2017): 248-253. doi:10.1016/j.marpolbul.2017.02.072.
- Wright, S.L., Thompson, R.C., Galloway, T.S. (2013).** The physical impacts of microplastics on marine organisms: a review. *Environmental Pollution* 178: 483-492. doi:10.1016/j.envpol.2013.02.031.
- Wright, L.S., Napper, I.E., Thompson, R.C. (2021).** Potential microplastic release from beached fishing gear in Great Britain's region of highest fishing litter density. *Marine Pollution Bulletin* 173: 113115. doi:10.1016/j.marpolbul.2021.113115.
- Xue, B., Zhang, L., Li, R., Wang, Y., Guo, J., Yu, K., Wang, S. (2020).** Underestimated microplastic pollution derived from fishery activities and "hidden" in deep sediment. *Environmental Science & Technology* 54(4): 2210-2217. doi:10.1021/acs.est.9b04850.
- Yücel, N., Kılıç, E. (2023a).** Presence of microplastic in the *Patella caerulea* from the northeastern Mediterranean Sea. *Marine Pollution Bulletin* 188: 114684. doi:10.1016/j.marpolbul.2023.114684.
- Yücel, N., Kılıç, E. (2023b).** Spatial Distribution of Microplastic Contamination in the Invasive Red Sea Mussel *Brachidontes pharaonis* (Fischer P., 1870) Around the İskenderun Bay. *Journal of Agricultural Production* 4(1): 7-15. doi:10.56430/japro.1232650.

ON PRESS

## Appendix I

**Table A1.** Morphological statistics i.e. test diameter, test height, GIT weight, gonad weight (mean  $\pm$  standard deviation) and microplastic abundance and frequency of occurrence (%) in the examined organs

Test diameter (mm)	Test height (mm)	GIT weight (g ww)	Gonad weight (g ww)	# of MP's	# of fiber MP's	# of fragment MP's	# of pelet MP's	MPs per ind	MPs per g ww	FO (%)
118.3 $\pm$ 12.2	88.2 $\pm$ 4.4	0.37 $\pm$ 0.16	5.25 $\pm$ 3.5	GIT 57	24	26	7	3 $\pm$ 3.1	0.8 $\pm$ 1.0	79
				Gonad 5	4	1	0	0.3 $\pm$ 0.6	0.08 $\pm$ 0.2	22

## Appendix II

**Table A2.** Recent literature indicating mean microplastic abundance in organs as well as predominant type of microplastic in similar species

Species	Location	GIT MP's per ind	MPs per g ww	Gonad MP's per ind	MPs per g ww	Common type	Common color	References
<i>Diadema setosum</i>	İskenderun Bay, Türkiye	3.0 MP's $\pm$ 3.1	0.9 $\pm$ 1.0	0.3 $\pm$ 0.6	0.08 $\pm$ 0.2	Fiber	Blue	This study
<i>Paracentrotus lividus</i>	Greece	26	1.9			Fiber	Blue	Hennicke <i>et al.</i> , 2021
<i>Diadema setosum</i>	Barranglombo Island, Indonosia	23.70 $\pm$ 2.99	-			Fiber	Blue	Sawalman <i>et al.</i> , 2021
<i>Diadema africanum</i>	Canary Island, Spain	9.2 $\pm$ 3.0				Fiber	Blue	Sevillano-González <i>et al.</i> , 2022
<i>Diadema africanum</i>	El Poris	10.0 $\pm$ 4.5				Fiber	Blue	Sevillano-González <i>et al.</i> , 2022
<i>Tetrapygyus niger</i>	Peru	3.22 $\pm$ 0.49				Fiber	Blue	De-la-Torre <i>et al.</i> , 2020
<i>Paracentrotus lividus</i>	Sardinia, Italy	1.0 $\pm$ 0.3				Fiber	Grey	Raguso <i>et al.</i> , 2022
<i>Diadema setosum</i>	Untung Jawa, Jakarta	2,175.55 $\pm$ 584.26				Fiber	-	Huseini <i>et al.</i> , 2021
	Tidung, Jakarta	1,786.66 $\pm$ 451.17				Fiber	-	
<i>Paracentrotus lividus</i>	Southern Italy,	1.2		0.83		Fiber	Black and blue	Murano <i>et al.</i> , 2022
<i>4 different species</i>	Yellow Sea, China	3.04		0.82		Fiber	Blue, gray	Feng <i>et al.</i> , 2020

### Appendix III




Figure A1. FTIR spectra of microplastic particle 1 (*Polypropylene*)

### Appendix IV

Microplastic Particle 2:




Figure A2. FTIR spectra of microplastic particle 2 (*Polypropylene*)

### Appendix V




Figure A3. FTIR spectra of microplastic particle 3 (*Polyethylene*)

## Appendix VI




Figure A4. FTIR spectra of microplastic particle 4 (*Polyethylene*)

## Appendix VII




Figure A5. FTIR spectra of microplastic particle 5 (*Polyethylene*)

## Appendix VIII




Figure A6. FTIR spectra of microplastic particle 6 (*Polypropylene*)

**Appendix IX**

**Table A9.** Results of correlation analysis (Below column shows statistical value, while upper column show p value)

	Test diameter	Test height	Weight	GIT weight	Gonad weight	MP in GIT	MP in Gonad
Test diameter		1.30E-07	6.63E-05	0.054845	0.00018	0.14943	0.9042
Test height	0.90215		0.000104	0.13054	0.000317	0.61684	0.66162
weight	0.78596	0.77315		0.1659	0.000744	0.1793	0.75556
GIT weight	0.44728	0.35956	0.33128		0.097013	0.57951	0.36812
Gonad weight	0.75614	0.73717	0.70526	0.39192		0.15055	0.96001
MP in GIT	-0.34388	-0.12268	-0.32166	-0.13574	-0.34299		0.72934
MP in Gonad	-0.02962	0.10741	0.076512	-0.21881	0.012341	-0.085	

**IN PRESS**