

Sağlık Bilimlerinde Yapay Zeka Dergisi

Journal of Artificial Intelligence in Health Sciences

A New Approach to Objective Monitoring of Sympathetic Blocks for Chronic Pain Treatment through Infrared Thermal Imaging

Kronik Ağrı Tedavisinde Kullanılan Sempatik Blokların Objektif Monitörizasyonuna Kızılötesi Termal Görüntüleme ile Yeni Yaklaşım

Burcu Candan*

To the Editor;

In a recent article published at this journal, Ervural and Ceylan have discussed the use of artificial intelligence methods and thermal imaging in medical applications [1]. In their concluding statement, the authors of the article conjecture that infrared thermal imaging can potentially be used in the diagnosis and treatment of Complex Regional Pain Syndrome (CRPS).

Sympathetic blocks (SB) are used for both diagnosis and treatment of sympathetically mediated pain (SMP), including CRPS [2,3]. While successful SB can improve tissue perfusion, resulting in a reduction of edema and pain, these blocks may also have a high failure rate. In particular, while they are effective for sympathetically mediated pain, they are not effective in the treatment of sympathetically independent pain (SIP) [6].

Despite the frequent use of SB, monitoring and assessing the success of SB using conventional techniques is known to be difficult [3]. Commonly used methods for assessing the success of SB include observation of clinical signs, skin temperature monitoring, and pulse amplitude monitoring [4,5]. However, these methods often demonstrate unpredictable or delayed response. A monitor which can objectively, simply, rapidly, and accurately assess the success of SB would not only improve the diagnosis, but would also contribute to better patient care by increasing procedural accuracy and efficiency. In particular, an objective confirmation of sympathectomy created by an attempted SB could potentially help differentiate SMP from SIP and help better plan the prospective treatment options.

In a recent work, however, we have shown that comparison of pre- and post-operation thermal profiles of a limb can help objectively assess the success of the SB in clinical settings [7].

Received / Geliş	20.08.2023
Accepted / Kabul	27.12.2023
Publication Date	31.12.2023

Burcu Candan*,

* Hospital for Special Surgery,
Musculoskeletal and Interventional Pain
Management Department of Anesthesiology, Critical
Care and Pain Management

***Corresponding Author:**

Hospital for Special Surgery,
Musculoskeletal and Interventional Pain
Management Department of Anesthesiology,
Critical Care and Pain
Management

E-posta: burcu.candan@gmail.com

ORCID: <https://orcid.org/0000-0003-2241-1225>

Conflict of Interest

No conflict of interest was declared by the author.

Financial Disclosure

The authors declared that this study received no financial support.

Acknowledgements

The author thanks Hospital for Special Surgery, NY, USA and Dr. Semih Gungor for enabling the study referred to in this letter

Author Contributions

Motivation / Concept: Burcu Candan
Design: Burcu Candan
Control / Supervision: Burcu Candan
Writing: Burcu Candan
Critical Review: Burcu Candan

Unfortunately, commercial infrared cameras are limited in their use in this context. Firstly, they provide point readings, while we need the temperature profile of the whole extremity. Moreover, the thermal images provided by the cameras include not only the limb, but also the background; therefore, a background elimination process has to be applied before a thermal profile can be created. Especially given the difficulty of collecting highly normalized thermal images in clinical settings, creating masks that separate the foreground (i.e. the feet) from the background is a nontrivial task that can benefit from artificial intelligence techniques.

Once the pre-and post-operation thermal profiles of the affected limb has been obtained, these can be compared using various methods. The simplest (and potentially easiest to interpret) approach would be to compare the average temperatures directly. Alternatively, temperature histograms can be created for both pre-and post-operation thermal profiles and these can be compared using histogram difference measures. A third alternative would be to create pre- and post-operation heatmaps and compare the spatio-thermal distributions of these heatmaps to pinpoint where the temperature difference is higher. These spatio-thermal patterns and their temporal evaluation after the operation can help better inform the clinician.

In conclusion, [7] has shown that infrared thermal imaging, coupled with image processing and analysis techniques, can be highly effective in assessing SB, addressing an urgent need in the treatment of CRPS patients.

REFERENCES

1. Ervural S, Ceylan M. “Kızılötesi Termal Görüntülemenin Yapay Zekâ Tabanlı Biyomedikal Uygulamalarına Bakış”. JAIHS 2021; 1(1):24-34. doi: 10.52309/jai.2021.5
2. Day M. “Sympathetic blocks: the evidence”, Pain Practice 2008; 8(2):98–109.
3. Krumova EK, Gussone C, Regeniter S, Westermann A, Zenz M, Maier C. "Are sympathetic blocks useful for diagnostic purposes?", Regional Anesthesia and Pain Medicine 2011; Nov-Dec;36(6):560-7.
4. Park SY, Nahm FS, Kim YC, Lee SC, Sim SE, Lee SJ. The cut-off rate of skin temperature change to confirm successful lumbar sympathetic block. J Int Med Res. 2010 Jan-Feb;38(1):266-75. doi: 10.1177/147323001003800131. PMID: 20233538.
5. Gungor S, Rana B, Fields K, et al. 2017. "Changes in the Skin Conductance Monitor as an End Point for Sympathetic Nerve Blocks", Pain Medicine,18(11):2187-2197.
6. Goebel A, Barker CH, Turner-Stokes L et al. 2018. "Complex Regional Pain Syndrome in Adults: UK guidelines for diagnosis, referral and management in primary and secondary care." London: Royal College of Physicians, Available from: <https://www.rcplondon.ac.uk/guidelines-policy/complex-regional-pain-syndrome-adults>
7. Gungor S, Candan B. “Infrared (FLIR) Imaging as a Monitor for Sympathetic Blocks in Complex Regional Pain Syndrome (CRPS)” accepted to be presented (Best Free Paper Session-Chronic Pain) at “6th World Congress on Regional Anesthesia & Pain Medicine”, 6-9 September 2023.