

# Body Weight Parameters at the Initial Visit for Eating Disorders in Adolescents: Are These the Markers of Serious Complications?

Ata Yiğit ÇÖLLÜ<sup>1</sup>, Demet TAŞ<sup>2</sup>, Alkım ÖDEN AKMAN<sup>3</sup>, Esra ÇÖP<sup>4</sup>

<sup>1</sup>Department of Child Health and Diseases, Ankara Bilkent City Hospital, Ankara, Türkiye

<sup>2</sup>Division of Adolescent Health, Ankara Yıldırım Beyazıt University, Ankara Bilkent City Hospital, Ankara, Türkiye

<sup>3</sup>Division of Adolescent Health, University of Health Sciences, Ankara Bilkent City Hospital, Ankara, Türkiye

<sup>4</sup>Department of Child and Adolescent Mental Health and Diseases, Ankara Bilkent City Hospital, University of Health Sciences, Ankara, Türkiye

## ABSTRACT

**Objective:** Eating disorders (ED) are psychiatric disorders often accompanied by medical complications. This study aimed to identify the complications in ED patients during hospital admission and their relationship with body weight (BW).

**Material and Methods:** Patients diagnosed with ED per DSM-5 criteria seen at least once in the adolescent department were included. Digital medical records were used to obtain patient information. Body mass index (BMI), BMI percentile, and percentage of BW by height (IBW%) were calculated to analyze their association with medical complications.

**Results:** Our study included 144 patients, 140 (97.2%) females and 4 (2.8%) males. Of the patients; 94 were analysed as Anorexia Nervosa (AN), 28 as Atypical AN and 17 as Bulimia Nervosa (BN). At least one cardiac complication was detected in 29.3% (39), gastrointestinal complications in 54.2% (78), hypoglycaemia in 18.2% (26), secondary amenorrhoea in 34.5% (48) and 45.8% (66) received inpatient treatment. Patients with IBW% < 75 exhibited significantly higher rates of hypoglycemia, bradycardia, low sT3, amenorrhea, and hospitalization compared to those with IBW% ≥ 75 (p < 0.001). Bone mineral density (BMD) was significantly inversely associated with Z-score, time to ED diagnosis and duration of amenorrhoea (p = 0.006; p = 0.044; p = 0.032, respectively).

**Conclusion:** Our findings show that at least one medical complication frequently develops in ED patients at the initial evaluation. Patients with AN and BN, which usually start in adolescence, may frequently present to paediatric outpatient clinics with medical complaints. The role of internal physicians in the early diagnosis of ED is important to prevent serious complications in these patients.

**Key Words:** Adolescent, Anorexia nervosa, Bulimia nervosa, Eating disorder, Medical complication

## INTRODUCTION

Anorexia nervosa (AN) and bulimia nervosa (BN) are types of ED that often onset in adolescence and progress with significant complications involving all body systems (1). According to DSM-5 diagnostic criteria, lifetime AN was reported by 0.8-6.3%, BN by 0.8-2.6%, and binge eating disorder (BED) by 0.6-6.1% of young women. Avoidant/Restrictive Food Intake Disorder (ARFID) is the new definition in DSM-5 of the type

formerly called "Feeding Disorder in Infancy or Early Childhood". Its incidence in children and adults varies between 0.5%-5% (2). AN is the most common cause of death among psychiatric disorders. Although some of the deaths are suicides, half of them are caused by medical complications (3).

Although the body weight (BW) of AN patients is very low compared to their peers, they are constantly preoccupied with the thought and behavior of losing weight. AN patients are dissatisfied with their weight and/or their body, sometimes

**Conflict of Interest:** On behalf of all authors, the corresponding author states that there is no conflict of interest.

**Ethics Committee Approval:** This study was conducted in accordance with the Helsinki Declaration Principles. The study was approved by Ankara Bilkent City Hospital, Ethics Committee (23.11.2022-2860).

**Contribution of the Authors:** **ÇÖLLÜ YA:** Taking responsibility in patient follow-up, collection of relevant biological materials, data management and reporting, execution of the experiments, Taking responsibility in necessary literature review for the study, Taking responsibility in the writing of the whole or important parts of the study. **TAŞ D:** Constructing the hypothesis or idea of research and/or article, Planning methodology to reach the conclusions, Organizing, supervising the course of progress and taking the responsibility of the research/study, Taking responsibility in logical interpretation and conclusion of the results, Taking responsibility in necessary literature review for the study, Taking responsibility in the writing of the whole or important parts of the study, Reviewing the article before submission scientifically besides spelling and grammar. **ÖDEM AKMAN A:** Constructing the hypothesis or idea of research and/or article, Planning methodology to reach the conclusions. **ÇÖP E:** Planning methodology to reach the conclusions, Reviewing the article before submission scientifically besides spelling and grammar.

**How to cite:** Çöllü YA, Taş D, Öden Akman A and Çöp E. Body Weight Parameters at the Initial Visit for Eating Disorders in Adolescents: Are These the Markers of Serious Complications?. Turkish J Pediatr Dis. 2025;19(1): 12-18.



0009-0007-2314-4188 : ÇÖLLÜ YA  
0000-0003-2586-5740 : TAŞ D



0000-0001-8080-7127 : ÖDEN AKMAN A  
0000-0001-8451-0099 : ÇÖP E

Correspondence Address:

**Demet TAŞ**

Department of Child Health and Diseases,  
Adolescent Health, Ankara Yıldırım Beyazıt University,  
Ankara Bilkent City Hospital, Ankara, Türkiye  
E-posta: demettas19691@hotmail.com

Received : 30.06.2024

Accepted : 07.08.2024

DOI:10.12956/tchd.1499664

part of their body. They usually have a restricted diet, some of them also have a one-way diet. BN and AN binge-purge subtype engage in compensatory behaviors such as vomiting and excessive exercise after consuming inappropriate food (4). Inappropriate restricted feeding, compensatory behaviors and rapid weight loss can lead to serious morbidity. Electrolyte disturbances, metabolic disorders, bradycardia and other cardiovascular pathologies are serious acute complications that should be closely monitored in patients with eating disorders (5). Common endocrine complications in patients with AN include amenorrhoea, oligomenorrhoea, cessation of puberty, hypothyroidism, and hypercortisolism. In patients with amenorrhoea, osteopenia and osteoporosis are long-term complications leading to clinically significant fractures and increased risk of fractures throughout life (6).

In a review, it was stated that the degree of low BW or rapid loss of BW predicts medical complications (7). Although ED is a psychiatric disorder, patients often present to health centers due to medical complications. Adolescents with EDs may present to the pediatrician with symptoms related to weight loss, malnutrition or vomiting behaviors. Symptoms such as constipation, early feeling of satiety, oesophagitis due to vomiting may be the first reason for presentation (8). They may also present with common symptoms such as fatigue, hair loss, chills and menstrual irregularities. Life-threatening complications such as bradycardia, hypotension and hypoglycemia may be detected on examination (9).

Patients with AN and BN with onset in adolescence are more likely to present to the pediatric outpatient clinic with one of the above symptoms. Recognising and early management of medical complications accompanying eating disorders is critical for the treatment and recovery of EDs (10). This study aimed to determine the complications in eating disorder patients at first presentation and how they were related to BW.

## MATERIAL and METHODS

Between April 2019 and January 2023, 144 adolescents aged 12-18 years who were diagnosed with eating disorders (AN, BN and ARFID) according to DSM-5, and admitted to Adolescent Health Department of Ankara Bilkent City Children's Hospital were included. The study was approved by Ankara Bilkent City Hospital, Ethics Committee (23.11.2022-2860). In this retrospective cross-sectional study, the symptoms and findings of the patients at the time of admission were taken from the medical records. A digital patient file containing detailed information of all systemic examination findings, water consumption and systemic symptoms such as constipation, vomiting and bowel movements characteristics was examined. Constipation was determined by using Rome-III criteria (11).

Some medical complications were evaluated in relation to BW at the time of diagnosis; BMI (weight (kg)/height<sup>2</sup> (m<sup>2</sup>)); BMI-percentile,

percentage of ideal body weight (%IBW) (weight (kg)/weight-for-height in the 50% percentile (kg)x100) (12).

Orthostatic blood pressure, pulse rate and body temperature (°C) were evaluated. A peak heart rate <60 beats/sec was considered bradycardia (13).

Patients' daily water intake was assessed according to the Turkey 2015 Dietary Guidelines (14). Urine density between 1007 and 1030 was considered normal. Leukocyturia was considered as 5 leukocytes per Hpf (high power field) (X40) and more in complete urine examination.

Patients who did not menstruate between 21-45 days despite 2 years after menarche, whose menstruation period lasted less than 3 days and more than 10 days, and whose daily number of pads was less than 1-2 and more than 7 were considered as irregular menstruation. Those who did not menstruate for at least 3 consecutive cycles were considered as secondary amenorrhoea (15).

Electrocardiography (ECG) was performed and pulse rate and QTc values were calculated. Echocardiographic findings (mitral valve prolapse (MVP), mitral regurgitation (MR), pericardial effusion) were obtained from echocardiography and cardiological evaluation. The results of the lumbar spine BMD measurements were assessed. Normal values of complete blood count were considered according to the guidelines of the Turkish Society of Haematology for the diagnosis and treatment of erythrocyte diseases and hemoglobin disorders (16).

According to the 2022 guidelines of the International Pediatric and Adolescent Diabetes Association, blood sugar levels below 70 mg/dL are considered hypoglycemia (17).

Other disease-specific biochemistry and hormone results were evaluated on the basis of the hospital laboratory references according to the age of the patients.

### Statistical Analysis

The IBM Statistical Package for Social Sciences (SPSS), Windows version 23.0 (SPSS Inc. Chicago, USA) was used for the statistical analysis of the research data. In the descriptive statistics section, categorical variables were presented as numbers and percentages; continuous variables were presented as mean  $\pm$  standard deviation and median (minimum-maximum value). The conformity of continuous variables to normal distribution was assessed using visual (histogram and probability plots) and analytical methods (Kolmogorov-Smirnov/Shapiro-Wilk tests).

As a result of the normality analyses, the paired sample t-test was used for comparison analyses between two dependent groups for data of continuous variables that were found to be normally distributed between groups.

For comparison analyses of categorical variables between independent groups, the Pearson chi-square test, continuity correction test and Fisher's exact test were used.

The relationship between BMD values and independent predictors was assessed using Pearson correlation analysis for normally

distributed data and Spearman correlation analysis for non-normally distributed data. When the absolute value of the correlation coefficient ( $\rho$ ) is  $\leq 0.30$ , there is a weak relationship,  $0.30-0.50$  is a moderate relationship, and  $r \geq 0.50$  is a strong relationship. In this study, the level of statistical significance was accepted as  $p < 0.050$ .

## RESULTS

A total of 144 patients were included in our study, of whom 4 (2.8%) were male and 140 (97.2%) were female. Patient diagnoses were as follows: 65.3% (94) AN (restrictive and binge types), 19.4% (28) atypical AN, 11.8% (17) BN and 3.5% (5) ARFID. Hospitalisation was an indication in 45.8% (66) of the patients.

Of the hospitalised patients, 80.3% (53) were diagnosed with AN, 9.1% (6) with atypical AN, 9.1% (6) with BN and one patient with ARFID.

At admission, 81.9% (118) of the patients had lost weight, 13.9% (20) had vomited, 3.5% (5) had lost their appetite, and 0.7% (1) had binge eating. The mean age at presentation was  $14.96 \pm 1.52$  (11.0-18.0) years. The mean difference between age at presentation and age at diagnosis was  $11.75 \pm 9.18$  (6.0-48.0) months. The mean BW and IBW at presentation were  $47.2 \pm 9.4$  (28.0-82.0) and  $88.4 \pm 17.3\%$  (56.0-150.0), respectively. Details of the patients' height, BMI, BMI(p), blood pressure (BP), pulse parameters are shown in Table I.

Orthostatic pulse difference was found in 6.1% (8) of patients, orthostatic systolic blood pressure difference in 3.8% (5) and orthostatic diastolic blood pressure difference in 9.2% (12).

Secondary amenorrhoea was found in 34.5% (48) and primary amenorrhoea in 2.9% (4) of the 139 patients whose menstruation was evaluated. Irregular menstruation was found in 38.8% (33) of the patients. The mean duration of amenorrhoea was  $7.9 \pm 3.4$  (4.0-18.0) months.

Of the four patients with primary amenorrhoea, three were AN restrictive and one was ARFID. Of the 48 patients with secondary amenorrhoea, 40 were AN (restrictive), two were AN (restrictive and binge type), two were BN and four were AN (atypical). The mean value of BMD Z score was  $-0.88 \pm 1.31$ .

**Table I: Height and Body Weight Characteristics of the Patients at Admission**

| Parameters                      | (n=144)            |
|---------------------------------|--------------------|
| Body Weight, kg*                | 47.2 (28.0-82.0)   |
| Height, cm†                     | 161.2±6.9          |
| BMI, kg/m <sup>2</sup> *        | 18.1 (12.78-31.63) |
| BMI-Percentile*                 | 20.1 (0.02-99.8)   |
| IBW %*                          | 88.4 (56.0-150.0)  |
| Body Temperature °C†            | 36.46±0.21         |
| Pulse (n=133)*                  | 80.7 (45.0-120.0)  |
| Systolic Blood Pressure, mmHg*  | 101.6 (80.0-130.0) |
| Diastolic Blood Pressure, mmHg* | 65.7 (40.0-90.0)   |

\*: median (min-max), †: mean±SD

It was found that 47.2% (67) of the patients had dry skin, 43% (61) had hair loss and 33.8% (48) had lanugo hair growth. A total of 45 patients (31.3%) consumed less than one litre of water per day.

In total, 29.3% (39) of patients presented with at least one cardiac complication. Bradycardia was found in 23.1% (30), mitral regurgitation in 21.1% (15), pericardial effusion in 8.5% (6) and MVP in 7% (5) of all patients. The mean heart rate was  $69.2 \pm 14.5$  (41.0-124.0)/min and the mean corrected QT interval (QTc) was  $0.39 \pm 0.05$  (0-0.44)ms. None of the patients had a pathologically long QTc interval.

Of the participants, 54.2% (78) had at least one gastrointestinal complication. Vomiting was observed in 29.9% (43) and constipation

**Table II: Laboratory Parameters of Patients**

| Parametreler                                  | n (%)     |
|---|-----------|
| Creatinin, mg/dL (n=144)                      |           |
| Low   | 4 (2.8)   |
| High  | 16 (11.1) |
| Ure, mg/dL (n=140)                            |           |
| Low   | 1 (0.7)   |
| High  | 10 (7.1)  |
| Uric Acid, mg/dL (n=140)                      |           |
| Low   | 1 (0.7)   |
| High  | 29 (20.7) |
| Total Protein, g/L (n=139)                    |           |
| Low   | 15 (10.8) |
| High  | 1 (0.7)   |
| Albumin, g/dL (n=141)                         |           |
| Low   | 2 (1.4)   |
| High  | 56 (39.7) |
| Aspartate aminotransferase (AST), U/L (n=143) |           |
| High  | 16 (11.2) |
| Alanine aminotransferase (ALT), U/L (n=139)   |           |
| High  | 18 (12.9) |
| TSH, mU/L (n=139)                             |           |
| Low   | 4 (2.9)   |
| High  | 2 (1.4)   |
| rT4, ng/dl (n=139)                            |           |
| Low   | 5 (3.6)   |
| High  | 4 (2.9)   |
| rT3, ng/L (n=102)                             |           |
| Low   | 65 (62.5) |
| High  | 1 (1.0)   |
| Leucocyte, x 10 <sup>9</sup> /L (n=144)       |           |
| Low   | 23 (16.0) |
| High  | 2 (1.4)   |
| Hemoglobin, g/dL (n=144)                      |           |
| Low   | 13 (9.0)  |
| High  | 5 (3.5)   |
| Neutrophile, x 10 <sup>9</sup> /L (n=144)     |           |
| Low   | 16 (11.1) |
| High  | 1 (0.7)   |
| Trombocyte, x 10 <sup>9</sup> /L (n=141)      |           |
| Low   | 4 (2.8)   |
| High  | 2 (1.4)   |

**Table III: Body Weight parameters according to the presence of hypoglycaemia and bradycardia in patients**

|                        | Hypoglycemia |             |        | Bradycardia  |              |        |
|------------------------|--------------|-------------|--------|--------------|--------------|--------|
|                        | Yes n=26 (%) | No n=117(%) | p      | Yes n=30 (%) | No n=100 (%) | p      |
| BMI, kg/m <sup>2</sup> |              |             |        |              |              |        |
| ≥18                    | 6 (9.1)      | 60 (90.9)   | 0.017* | 9 (15.3)     | 50 (84.7)    | 0.085* |
| <18                    | 20 (26.0)    | 57 (74.0)   |        | 21 (29.6)    | 50 (70.4)    |        |
| BMI-p                  |              |             |        |              |              |        |
| ≥5                     | 10 (13.2)    | 66 (86.8)   | 0.149* | 10 (15.2)    | 56 (84.8)    | 0.049* |
| <5                     | 16 (23.9)    | 51 (76.1)   |        | 20 (31.2)    | 44 (68.8)    |        |
| IBW                    |              |             |        |              |              |        |
| ≥ %75                  | 14 (13.1)    | 93 (86.9)   | 0.008* | 15 (16.0)    | 79 (84.0)    | 0.006* |
| < %75                  | 12 (35.3)    | 22 (64.7)   |        | 14 (41.2)    | 20 (58.8)    |        |

\*: Continuity Correction test

**Table IV: BMI, BMI-p and IBW % according to the presence of amenorrhea and sT3 levels in patients**

| Parameters             | Amenorrhea (Primer and Seconder) |             |         | sT3             |              |         |
|------------------------|----------------------------------|-------------|---------|-----------------|--------------|---------|
|                        | Yes n=52 (%)                     | No n=87 (%) | p       | Normal n=38 (%) | Low n=65 (%) | p       |
| BMI, kg/m <sup>2</sup> |                                  |             |         |                 |              |         |
| ≥18                    | 11 (21.2)                        | 55 (63.2)   | <0.001* | 29 (65.9)       | 15 (34.1)    | <0.001* |
| <18                    | 41 (78.8)                        | 32 (36.8)   |         | 9 (15.3)        | 50 (84.7)    |         |
| BMI-p                  |                                  |             |         |                 |              |         |
| ≥5                     | 17 (32.7)                        | 59 (67.8)   | <0.001† | 28 (54.9)       | 23 (45.1)    | <0.001* |
| <5                     | 35 (67.3)                        | 28 (32.2)   |         | 10 (19.2)       | 42 (80.8)    |         |
| IBW                    |                                  |             |         |                 |              |         |
| ≥%75                   | 29 (55.8)                        | 77 (88.5)   | <0.001† | 36 (49.3)       | 37 (50.7)    | <0.001* |
| < %75                  | 23 (44.2)                        | 10 (11.5)   |         | 2 (7.1)         | 26 (92.9)    |         |

\*: Pearson Chi-kare test, †: Continuity Correction test

in 51.4% (74) of the patients. Three patients had oesophagitis, two had rectal bleeding and one had hemorrhoids. At least one endocrinological complication was found with 68.1% (98) of the patients. Hypoglycemia occurred in 18.2% (26) of the patients and the mean glucose level was 79.06±13.95 (47.0-152.0) mg/dl.

The creatinine level was found to be high in 11.1% (16), the urea level in 7.1% (10) and the uric acid level in 20.7% (29) of the patients. Albumin level was found to be low in 1.4% (2) and high in 39.7% (56) of patients (Table II).

In our study, AST levels were found to be high in 11.2% (16) and ALT levels were found to be high in 12.9% (18) of patients. White blood cell count was low in 16% (23), hemoglobin was low in 9% (13), high in 3.5% (5) and neutrophil count was low in 11.1% (16) of the patients. Among the patients' thyroid function tests, TSH levels were low in 2.9% (4) and high in 1.4% (2); sT4 levels were low in 3.6% (5) and high in 2.9% (4); sT3 levels were low in 62.5% (62) and high in 1% (1) (Table II).

When the electrolytes of the patients were evaluated: Serum chlorine level was low in 2.9% (4) and high in 8.8% (12), potassium level was low in 4.3% (6), calcium level was low in 5.7% (8) and high in 12.9% (18), phosphorus level was low in 3.6% (5) and magnesium level was normal in all patients.

It was found that 31.3% (44) of all participants drank less water and the mean urine density was 1017.3±9.54 (1000-1042). Urine density was low in 15.6% (22), high in 7% (10) and 77.3% (109) normal

range of these patients. Sterile pyuria was found in 31.4% (44) of the patients.

Thirty three (41.3%) of patients with restrictive type AN drank less water, and urine density was low in 18.7% (15) and high in 10% (10) of these patients.

The rate of sterile pyuria was not significantly higher in patients with BMI <18, BMI-p<5 and IBW <75% (p=0.227; p=0.999; p=0.221, respectively).

A statistically significant increase in the incidence of hypoglycaemia was observed in patients with a BMI of less than 18 and an IBW of less than 75% (p=0.049; p=0.006, respectively). Bradycardia was statistically significantly more common in patients with BMI-p<5 and IBW <75% and (p=0.049; p=0.006, respectively) (Table III).

Patients with BMI<18, BMI-p<5 and IBW≥75% had statistically significantly more amenorrhoea (p<0.001). The statistical analysis yielded significant results, indicating a higher prevalence of low sT3 levels among patients with a BMI less than 18, a BMI-p less than 5, and an IBW less than 75% (p<0.001) (Table IV).

There was a statistically significant higher number of hospitalisations in patients with BMI <18, BMI-p <5 and IBW <75% (p<0.001; p<0.001; p=0.015, respectively). (Table V).

The relationship between urea, creatinine and uric acid levels and body weight parameters was evaluated using Fisher's exact test. In patients with BMI <18, BMI-p<5 and IBW<75%, no significant

**Table V: Body Weight parameters according to patients' hospitalisation status**

| Parameters             | Hospitalization |             | p                   |
|------------------------|-----------------|-------------|---------------------|
|                        | Yes n=66 (%)    | No n=78 (%) |                     |
| BMI, kg/m <sup>2</sup> |                 |             |                     |
| ≥18                    | 17 (25.4)       | 50 (74.6)   | <0.001 <sup>†</sup> |
| <18                    | 49 (63.6)       | 28 (36.4)   |                     |
| BMI-p                  |                 |             |                     |
| ≥5                     | 24 (31.2)       | 53 (68.8)   | <0.001 <sup>†</sup> |
| <5                     | 42 (62.7)       | 25 (37.3)   |                     |
| IBW                    |                 |             |                     |
| ≥ %75                  | 42 (38.9)       | 66 (61.1)   | 0.015 <sup>†</sup>  |
| < %75                  | 22 (64.7)       | 12 (35.3)   |                     |

\*: Pearson Chi-kare test, †: Continuity Correction test

**Table VI: Relationship between mean BMD Z-score and BW**

|                              | r                   | p     |
|------------------------------|---------------------|-------|
| BMD Z score*Age of diagnosis | -0.401 <sup>†</sup> | 0.006 |
| BMD Z score*IBW %            | 0.302 <sup>‡</sup>  | 0.044 |
| BMD Z score* BMI             | 0.054 <sup>‡</sup>  | 0.726 |
| BMD Z score*BMI-Percentil    | 0.183 <sup>‡</sup>  | 0.228 |
| BMD Z score*Amenorrhoea Time | -0.363 <sup>‡</sup> | 0.032 |

**BMD:** Bone Mineral Density, **r:** Correlation coefficient, **†:** Pearson correlation  
**‡:** Spearman's correlation

increases or decreases in creatinine, urea and uric acid levels were observed (for urea,  $p=0.751$ ;  $p=0.999$ ;  $p=0.451$ , for creatinine,  $p=0.301$ ;  $p=0.274$ ;  $p=0.535$ , for uric acid,  $p=137$ ;  $p=498$ ;  $p=0.511$ , respectively).

A significant negative correlation was found between BMD Z-score values and time to disease diagnosis ( $p<0.006$ ). A moderately significant relationship was found between BMD Z score values and IBW% ( $p<0.044$ ). A significant negative correlation was found between BMD values and duration of amenorrhoea ( $p<0.032$ ) (Table VI).

## DISCUSSION

This study investigated the relationship between the presenting signs, symptoms and complications of ED patients and their BW at the time of presentation. A total of 144 patients, of whom 4 (2.8%) male and 140 (97.2%) female were included in our study. Among the male patients, three patients were identified as AN (restrictive) type and one patient was identified as ARFID.

Anorexia nervosa represented the majority of patients followed in adolescence. This, predictably, was due to a high rate of weight loss and associated medical complications in patients with AN. The mean age at diagnosis was 14.6 years and was similar in boys and girls; however, we could not compare the ages at diagnosis because the number of male patients was very small. The peak age for ED diagnosis was 15 years for girls and 16 years for boys in a study conducted in the UK (18).

In our study, the mean difference between the ages of diagnosis and first symptoms was 11.75 months. In another study, the difference between the onset of symptoms and admission to the clinic was 11 months (19). In a long-term study, 48% of adolescents with AN were found to have growth curve abnormalities on average 9.7 months before the onset of AN-related symptoms on retrospective observation (20). These studies show that there is a delay in diagnosis in patients with AN, resulting in serious long-term morbidity. These data suggest that there may be some reasons why patients with ED are diagnosed with an average delay of 1 year. Due to the distorted body image of ED patients, they do not see themselves as sick. Diagnosis may be delayed because it takes time for parents to notice the changed eating behaviour and for medical signs to appear. As with other chronic diseases, one of these reasons may be that their access to health centers was disrupted during the Covid-19 pandemic (21).

It was observed that about half of our patients were hospitalized and the majority of the hospitalized patients were AN patients. This is because, as expected, more medical complications developed with increased BW loss. One study reported that 55% of patients with AN had at least one hospital admission (19).

In our study, significantly more hypoglycaemia was observed in patients with low BW. This finding suggests that we should be more cautious about hypoglycemia in patients with low BW because of the decrease in glycogen stores due to weight loss. A published case study reported that clinically severe hypoglycaemia may be a fatal complication of AN, although it is not very common (22). We found hypoglycaemia in 18.2% of our patients.

Elevated blood creatinine levels were found in 11.1%, urea in 7.1% and uric acid in 20.7% of our patients. In patients with anorexia nervosa, urea and creatinine levels are expected to be low due to malnutrition and low muscle tissue. However, elevated levels are a sign of severe kidney damage or dehydration. Impaired renal function was found to be more predictive of disease severity than low BMI in eating disorder patients (23).

Although the mechanism is not well understood, studies have shown that albumin levels are normal in patients with eating disorders. It has been suggested that this may be due to increased albumin transfer from the extravascular to the intravascular space by adaptive mechanisms (24). Mehler et al. (25) explain that one reason for the preservation of serum albumin levels in patients with anorexia nervosa is the absence of inflammation in these patients, and that it is not a good marker in the follow-up of patients. In our study, 39.7% of patients had a mildly elevated albumin level.

Liver transaminase levels have been found to be elevated in approximately 10% of patients, and all of these patients had a diagnosis of restrictive AN. One study showed that BMI and transaminase elevation were inversely proportional in AN patients (26).

Iron deficiency anemia is not expected in patients with AN. The main cause of anemia and other blood cell deficiencies in patients with AN is bone marrow suppression. However, abnormalities in the complete blood count are common. In our study, 16% of patients had a low white blood cell count and 11.1% had

neutropenia. Hemoglobin was found to be low in 9% of patients. As expected, iron deficiency anemia was not found. A review reported that the rate of anemia was 21-39%, leucopenia 29-39% and thrombocytopenia 5-11% (27). In a study of 60 adolescent girls diagnosed with AN, the rate of neutropenia and anemia was 22% (28). Compared to the literature, blood cell disorders were detected less frequently in our patients.

Although 41% of adolescents with AN drank less water, only 7% were found to have increased urine density. It is well known that osmoregulation is impaired in patients with AN (29). Sterile pyuria was found in 31.4% of patients.

Three of the 6 patients with hypokalemia on admission were diagnosed with AN, two with atypical AN and one with BN. Four of the five patients with hypophosphatemia were diagnosed with AN and one with atypical AN. Electrolyte imbalances are a serious complication of eating disorders. One study drew attention to this issue and recommended caution regarding undiagnosed eating disorders in patients presenting to a health center with electrolyte imbalances (30).

Secondary amenorrhoea and irregular menstruation were observed in approximately one third and one quarter of our patients, respectively. It is well known that low BW is associated with amenorrhoea, which is one of the most common complications in ED patients (31). In a study of 251 ED patients, the incidence of amenorrhoea was found to be 61% in AN patients and 11% in BN patients, and the same study found that the frequency and duration of amenorrhoea were associated with BMI. The same study reported that amenorrhoea developed in 82% of those with a BMI below 18 who exercised (32).

Cardiac complications are the most common cause of sudden death in patients with AN. Numerous cardiac abnormalities have been described in the literature, including pericardial and valvular pathologies, changes in left ventricular muscle and function, conduction abnormalities, bradycardia, hypotension, and peripheral vascular contractility abnormalities. At least one cardiac complication was observed in 29.3% of our patients. The most common cardiac complication was bradycardia and mitral regurgitation was observed with the second frequency. Sinus bradycardia is the most common cardiovascular complication in patients with AN. Bradycardia improves with nutrition and weight gain. We found that BW was inversely associated with the occurrence of bradycardia in our patients. Pericardial effusion and MVP were less common. QTc prolongation was not observed in any of our patients. Although some studies have shown QTc prolongation, most studies have reported no QTc prolongation (33).

At least one GIS complication was observed in 54.2% of patients. The most common objective GIS finding was constipation. Endoscopic oesophagitis was also found in three patients, rectal bleeding in two patients and hemorrhoids in one patient. One study reported that constipation was found in 35% of patients diagnosed with AN and in 15% of patients diagnosed with BN (34). Constipation may be expected in patients with AN in association with food and fluid restriction. However, impaired gastrointestinal

motility in patients with AN and altered gut microbiota in relation to their food preferences increase the susceptibility to constipation (35).

At least one endocrine complication was observed in 68.1% of patients, the most common being hypothyroidism and menstrual disorders. In our study, sT3 levels were found to be low in 62.5% of patients. In our study, low sT3 levels were found to be significantly higher in patients with low BMI and IBW%. One study reported that the amount and rate of weight loss affected low sT3 concentrations. In addition, it was reported that low sT3 was an important determinant of weight loss in the premenopausal period, whereas low sT3 could not predict weight loss in the postmenopausal period. In the same study, serum TSH concentrations were found to be within normal limits in ED patients and were not affected by changes in BW. It was found that blood sT3 levels also increased with increasing BW and reached normal levels (36).

As seen in our study, amenorrhoea has a negative effect on bone mineral density. This increases the risk of osteopenia and osteoporosis in the long term. The severity of long-term complications also increases with the duration of the disease and its early onset. A study comparing spinal bone densities in adolescent females found that prolonged amenorrhoea was associated with decreased BMD Z-scores (37). The data from our study support this finding. A significant, negative and moderate correlation was found between BMD Z score values and age at diagnosis.

In conclusion, according to our data, medical complications affecting all body systems are common in patients with AN and BN. Although Eating Disorders are psychiatric disorders, somatic symptoms or medical complications are likely to be the reason for patients to consult a health center. Early identification of patients with weight loss or compensatory behaviors secondary to ED may prevent morbidity and mortality. Recognising possible symptoms and signs of AN and BN, which often begin in adolescence, in pediatric outpatient clinics will protect against acute and chronic complications and prevent delayed multidisciplinary treatment.

## REFERENCES

1. Volpe U, Tortorella A, Manchia M, Monteleone AM, Albert U, Monteleone P. Eating disorders: What age at onset? *Psychiatry Res* 2016;238:225-7.
2. Silén Y, Keski-Rahkonen A. Worldwide prevalence of DSM-5 eating disorders among young people. *Curr Opin Psychiatr* 2022;35:362-71.
3. Welch E, Ghaderi A. Mortality in anorexia nervosa – a look back at and beyond one of the most cited papers in the field. *Advances in Eating Disorders* 2015;3:221-9.
4. Harrington BC, Jimerson M, Haxton C, Jimerson DC. Initial evaluation, diagnosis, and treatment of anorexia nervosa and bulimia nervosa. *Am Fam Physician* 2015;91:46-52.
5. Sachs KV, Harnke B, Mehler PS, Krantz MJ. Cardiovascular complications of anorexia nervosa: A systematic review. *Int J Eat Disord* 2016;49:238-48.
6. Hung C, Muñoz M, Shibli-Rahhal A. Anorexia Nervosa and Osteoporosis. *Calcif Tissue Int* 2022;110:562-75.

7. Gibson D, Stein A, Khatri V, Wesselink D, Sitko S, Mehler PS. Associations between low body weight, weight loss, and medical instability in adults with eating disorders. *Int J Eat Disord* 2024;57:869-78.
8. Sato Y, Fukudo S. Gastrointestinal symptoms and disorders in patients with eating disorders. *Clin J Gastroenterol* 2015;8:255-63.
9. Campbell K, Peebles R. Eating disorders in children and adolescents: state of the art review. *Pediatrics* 2014;134:582-92.
10. Klein DA, Sylvester JE, Schvey NA. Eating Disorders in Primary Care: Diagnosis and Management. *Am Fam Physician* 2021;103:22-32.
11. Baaleman DF, Velasco-Benítez CA, Méndez-Guzmán LM, Benninga MA, Saps M. Functional gastrointestinal disorders in children: agreement between Rome III and Rome IV diagnoses. *Eur J Pediatr* 2021;180:2297-03.
12. Ogden CL, Kuczmarski RJ, Flegal KM, Mei Z, Guo S, Wei R, et al. Centers for Disease Control and Prevention 2000 growth charts for the United States: improvements to the 1977 National Center for Health Statistics version. *Pediatrics* 2002;109:45-60.
13. Shen WK, Sheldon RS, Benditt DG, Cohen MI, Forman DE, Goldberger ZD, et al. ACC/AHA/HRS Guideline for the Evaluation and Management of Patients With Syncope A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Heart Rhythm* 2017;14:e155-e217.
14. T. C. Sağlık Bakanlığı Türkiye Beslenme Rehberi (TÜBER). 2015; 46-9.
15. Diaz A, Laufer MR, Breech LL. Menstruation in girls and adolescents: using the menstrual cycle as a vital sign. *Pediatrics* 2006;118:2245-50.
16. Eritrosit Hastalıkları ve Hemoglobin Bozuklukları Tanı ve Tedavi Kılavuzu. 2019;22-6. Available from: [www.thd.org.tr](http://www.thd.org.tr)
17. Abraham MB, Karges B, Dovc K, Naranjo D, Ana Arbelaez M, Mbogo J, et al. ISPAD Clinical Practice Consensus Guidelines 2022: Assessment and management of hypoglycemia in children and adolescents with diabetes. *Pediatr Diabetes* 2022;23:1322-40.
18. Petkova H, Simic M, Nicholls D, Ford T, Prina AM, Stuart R, et al. Incidence of anorexia nervosa in young people in the UK and Ireland: a national surveillance study. *BMJ Open* 2019 1;9: e027339.
19. Steinhausen HC, Grigoriu-Serbanescu M, Boyadjieva S, Neumärker KJ, Metzke CW. Course and predictors of rehospitalization in adolescent anorexia nervosa in a multisite study. *Int J Eat Disord* 2008;41:29-36.
20. Swanson SA, Crow SJ, Le Grange D, Swendsen J, Merikangas KR. Prevalence and correlates of eating disorders in adolescents. Results from the national comorbidity survey replication adolescent supplement. *Arch Gen Psychiatry* 2011;68:714-23.
21. Singh K, Kaushik A, Johnson L, Jaganathan S, Jarhyan P, Deepa M, et al. Patient experiences and perceptions of chronic disease care during the COVID-19 pandemic in India: a qualitative study. *BMJ Open* 2021;11: e048926.
22. Yanai H, Yoshida H, Tomono Y, Tada N. Severe hypoglycemia in a patient with anorexia nervosa. *Eat Weight Disord* 2008;13:e1-3.
23. Gurevich E, Steiling S, Landau D. Incidence of Impaired Kidney Function Among Adolescent Patients Hospitalized With Anorexia Nervosa. *JAMA Netw Open* 2021;4:e2134908.
24. Winston AP. The clinical biochemistry of anorexia nervosa. *Ann Clin Biochem* 2012;49:132-43.
25. Mehler PS, Koutsavlis A. Anorexia Nervosa, Albumin, and Inflammation. *Am J Med* 2021 ;134:e401.
26. Speranza E, Marra M, De Filippo E, De Caprio C, Sammarco R, Morlino D, Contaldo F, Pasanisi F. Nutritional indicators and metabolic alterations in outpatients with anorexia nervosa: a retrospective study. *Eat Weight Disord* 2021;26:2693-99.
27. Hütter G, Ganepola S, Hofmann WK. The hematology of anorexia nervosa. *Int J Eat Disord* 2009;42:293-300.
28. Misra M, Aggarwal A, Miller KK, Almazan C, Worley M, Soyka LA, et al. Effects of anorexia nervosa on clinical, hematologic, biochemical, and bone density parameters in community-dwelling adolescent girls. *Pediatric* 2004;114:1574-83.
29. Stheneur C, Bergeron S, Lapeyraque A. Renal complications in anorexia nervosa. *Eating and Weight Disord* 2014;19:455-60.
30. Hundemer GL, Clarke A, Akbari A, Bugeja A, Massicotte-Azarniouch D, Knoll G, et al. Analysis of Electrolyte Abnormalities in Adolescents and Adults and Subsequent Diagnosis of an Eating Disorder. *JAMA Netw Open* 2022;5:e2240809.
31. Haines MS. Endocrine complications of anorexia nervosa. *J Eat Disord* 2023;11:24.
32. Abraham SF, Pettigrew B, Boyd C, Russell J, Taylor A. Usefulness of amenorrhoea in the diagnoses of eating disorder patients. *J Psychosom Obstet Gynaecol* 2005;26:211-5.
33. Sachs KV, Harnke B, Mehler PS, Krantz MJ. Cardiovascular complications of anorexia nervosa: A systematic review. *Int J Eat Disord* 2016;49:238-48.
34. Riedlinger C, Schmidt G, Weiland A, Stengel A, Giel KE, Zipfel S, et al. Which Symptoms, Complaints and Complications of the Gastrointestinal Tract Occur in Patients With Eating Disorders? A Systematic Review and Quantitative Analysis. *Front Psychiatry* 2020;11:195.
35. Herman A, Bajaka A. The role of the intestinal microbiota in eating disorders - bulimia nervosa and binge eating disorder. *Psychiatry Res* 2021;300:113923.
36. Swenne I, Stridsberg M, Thurfjell B, Rosling A. Triiodothyronine is an indicator of nutritional status in adolescent girls with eating disorders. *Horm Res* 2009;71:268-75.
37. Fazeli PK. Low bone mineral density in anorexia nervosa: Treatments and challenges. *Clin Rev Bone Miner Metab* 2019;17:65-76.